

## Homework 7

- (1) Suppose the manifold  $M$  is the disjoint union of manifolds  $M_1$  and  $M_2$ . Then prove that  $H_{dR}^k(M) = H_{dR}^k(M_1) \oplus H_{dR}^k(M_2)$ .
- (2) Let  $\phi : M \rightarrow N$  be a smooth map between manifolds. If  $\omega \in \Omega^k(N)$ , then prove that  $d(\phi^*\omega) = \phi^*(d\omega)$ .
- (3) Compute the de Rham cohomology of  $S^2$ . Then inductively compute the de Rham cohomology of  $S^n$  for all  $n \geq 2$ .
- (4) Compute the de Rham cohomology of a compact 2-dimensional manifold (i.e., surface) of genus  $g \geq 1$  without boundary. [You can do this informally, by drawing pictures of the surface of genus  $g$  and its decomposition into pieces.]
- (5) Compute the de Rham cohomology of  $\mathbb{R}P^n$  for all  $n \geq 2$ .
- (6) (30 points) Complete the proof that the short exact sequence of cochain maps

$$0 \rightarrow \mathcal{A} \xrightarrow{\phi} \mathcal{B} \xrightarrow{\psi} \mathcal{C} \rightarrow 0$$

gives rise to a long exact sequence on cohomology. (This problem should be done carefully and completely — it's worth 30 points.)

- (7) (20 points) Prove the existence of a partition of unity subordinate to an open cover  $\{U_\alpha\}$  of a noncompact manifold  $M$ .
- (8) Let  $M$  be an oriented manifold of dimension  $n$  and  $\omega \in \Omega^n(M)$ . Let  $\{\phi_\alpha : U_\alpha \rightarrow \mathbb{R}^n\}$  be an oriented atlas and  $\{f_\alpha\}$  be a partition of unity subordinate to  $\{U_\alpha\}$ . We defined

$$\int_M \omega = \sum_\alpha \int_{\phi_\alpha(U_\alpha)} (\phi_\alpha^{-1})^*(f_\alpha \omega).$$

Prove that this definition does not depend on the choice of oriented atlas as well as on the choice partition of unity.