

1. Let $f : \mathbb{C} \rightarrow \mathbb{C} \cup \{\infty\}$ be a non-constant doubly periodic function with periods 1 and τ as in HW3. (Recall $\tau \in \mathbb{C}$ has $\text{Im } \tau > 0$.)

(a) Show that the sum of the residues in any fundamental parallelogram (i.e., with vertices $\{z, z+1, z+1+\tau, z+\tau\}$) whose sides miss the poles of f is equal to zero.

(b) Show that $f(z) = w$ and $f(z) = \infty$ have an equal number of solutions in each fundamental parallelogram (for any $w \in \mathbb{C}$ and counting with multiplicity). From HW3 we know that this number is non-zero.

(c) Show that

$$\wp(z) := \frac{1}{z^2} + \sum_{n \in \mathbb{Z}^2 \setminus \{0\}} \frac{1}{(z - n_1 - n_2\tau)^2} - \frac{1}{(n_1 - n_2\tau)^2}$$

defines a meromorphic function and that

$$\wp'(z) := -2 \sum_{n \in \mathbb{Z}^2} \frac{1}{(z - n_1 - n_2\tau)^3}.$$

Once you prove convergence, it is clear that $\wp'(z)$ is doubly periodic.

(d) Use the fact that \wp is even (i.e. $\wp(z) = \wp(-z)$) to deduce that \wp is doubly periodic.

2. (a) Let A be a $n \times n$ matrix and let $f(z) := \det(z\text{Id} - A)$. Show

$$\frac{f'(z)}{f(z)} = \text{Tr}\{(z\text{Id} - A)^{-1}\}.$$

(b) Apply Rouché's Theorem to f to show that the eigenvalues of A (repeated according to algebraic multiplicity) depend continuously on the entries of A . Here, we say that the distance between two multi-sets is the sum of the distances under the shortest matching.

(c) Now suppose $A(t)$ depends holomorphically on a parameter $t \in \mathbb{D}$ and that λ_0 is a simple (i.e. multiplicity one) eigenvalue of $A(0)$. Show that there is a holomorphic function $t \mapsto \lambda(t)$ defined in an open neighbourhood of 0 so that $\lambda(0) = \lambda_0$ and $\lambda(t)$ is an eigenvalue of $A(t)$.

Remark: The matrix function $t \mapsto \begin{bmatrix} 0 & 1 \\ t & 0 \end{bmatrix}$ shows that the eigenvalues may only be Hölder continuous as functions of the entries in A . In the $n \times n$ case, they may be merely Hölder $\frac{1}{n}$ continuous.

3. Suppose $f : \mathbb{C} \rightarrow \mathbb{C}$ is bijective and holomorphic.

(a) Show that f^{-1} is holomorphic.

(b) Use the continuity of f^{-1} to show that $|f(z)| \rightarrow \infty$ as $|z| \rightarrow \infty$.

(c) Show that $f(z) = O(|z|)$ as $z \rightarrow \infty$. [*Hint:* Look at $g(z) = 1/f(1/z)$.]

(d) Deduce that $f(z) = az + b$ for some $a \in \mathbb{C} \setminus \{0\}$ and $b \in \mathbb{C}$.

4. Show that every meromorphic bijection f from $\mathbb{C} \cup \{\infty\}$ to itself is of the form

$$f(z) = \frac{az + b}{cz + d}$$

for some quadruplet $a, b, c, d \in \mathbb{C}$ obeying $ad - bc = 1$.

5. (a) Use Möbius transformations to determine the analogue of the Schwarz Reflection Principle for holomorphic functions f defined in an open neighbourhood of an arc of the circle $\{|z| = 1\}$ that obey $|f(e^{i\theta})| = 1$.

(b) Use Möbius transformations to determine the analogue of Schwarz Lemma for mappings f of the half-plane $\{\operatorname{Re} z > 0\}$ to itself that obey $f(1) = 1$.

(c) Use Möbius transformations and Schwarz Lemma to prove the *Borel–Carathéodory Theorem*: Let f be holomorphic in an open neighbourhood of the closed unit disk, $\bar{\mathbb{D}}$. For each $z \in \mathbb{D}$,

$$|f(z)| \leq \frac{1 + |z|}{1 - |z|} |f(0)| + \frac{2|z|}{1 - |z|} \sup_{w \in \mathbb{D}} \operatorname{Re} f(w).$$