

HOMEWORK 6

1. Let G be a finite group and C the center of G . Let $\mu : C \rightarrow F^\times$ be a character of C . Prove that there is an irreducible representation $\rho : G \rightarrow \text{GL}(V)$ such that $\rho(c)(v) = \mu(c)v$ for all $c \in C$ and $v \in V$.
2. Let F be a field of characteristic $p > 0$ and G a finite p -group. Prove that $\text{rad}(F[G]) = \{\sum a_g g \in F[G] \mid \sum a_g = 0\}$. Determine all simple (left) $F[G]$ -modules.
3. Let V be the kernel of $F^n \rightarrow F$ taking (a_1, \dots, a_n) to $\sum a_i$. The symmetric group S_n acts on V by permutations of the coordinates. Prove that if the characteristic of F does not divide n , then the corresponding representation is irreducible.
4. Let ρ be a representation of a finite group G and V the corresponding G -space. Show that the dual space V^* has the structure of a G -space via $(g\varphi)(v) = \varphi(g^{-1}v)$ for $g \in G$, $\varphi \in V^*$ and $v \in V$. Prove that $\chi_{\rho^*}(g) = \chi(g^{-1})$, where ρ^* is the representation corresponding to the G -space V^* .
5. Show that ρ^* is irreducible if and only if so is ρ .

For all problems below the base field F is algebraically closed of characteristic zero.

6. Let G be a finite group. Define the abelian group $\text{Rep}(G)$ by generators and relations as follows. The generators are the isomorphism classes $[\rho]$ of representations ρ of G . The relations are $[\rho \oplus \rho'] = [\rho] + [\rho']$ for all representations ρ and ρ' . Prove that $\text{Rep}(G)$ is a free abelian group with basis the set of isomorphism classes of irreducible representations of G . Prove that the tensor product yields the structure of a commutative ring on $\text{Rep}(G)$.
7. Let G be a finite group. Prove that the map $\text{Rep}(G) \rightarrow \text{Ch}(G)$ taking the class $[\rho]$ to the character χ_ρ is a well defined injective ring homomorphism. Write the multiplication table for $\text{Rep}(S_3)$.
8. Find all groups that have exactly 2 nonisomorphic representations.
9. Find an irreducible 2-dimensional representation of the symmetric group S_4 .
10. Let $C(g)$ be the conjugacy class of an element g in a finite group G and χ be the character of an irreducible representation ρ . Prove that if $|C(g)|$ is

relatively prime to $\dim(\rho)$ and $\chi(g) \neq 0$, then $\rho(g)$ is the multiplication by a scalar.