

# Homework 2 for Math 214B Algebraic Geometry

Burt Totaro

Spring 2016, UCLA

Due on Friday, May 6.

(1) For a hypersurface  $X$  of degree  $d$  in  $\mathbf{P}^{n+1}$  over a field  $k$ ,  $n \geq 1$ , compute the Hilbert series of the graded ring  $\bigoplus_j H^0(X, \mathcal{O}(j))$ .

(2) Let  $X$  be a hypersurface of degree  $d$  in projective space  $\mathbf{P}^{n+1}$  over a field  $k$ . Compute the cohomology groups  $H^i(X, \mathcal{O}_X)$  for all  $i$ . Deduce that  $X$  is not isomorphic to  $\mathbf{P}^n$  for  $d$  large enough; what range of  $d$  do you get?

(3) Let  $X$  be the affine line over a field  $k$ . We know that  $H^i(X, E) = 0$  for every quasi-coherent sheaf  $E$  on  $X$  (or any affine scheme) and every  $i > 0$ . Does this vanishing hold for every sheaf of  $\mathcal{O}_X$ -modules, not necessarily quasi-coherent?

(4) Let  $U = A^2 - 0$  over a field  $k$ . Using a suitable cover of  $U$  by affine open subsets, show that  $H^1(U, \mathcal{O})$  is isomorphic to the  $k$ -vector space with basis  $\{x^i y^j : i, j < 0\}$ . In particular, it is a  $k$ -vector space of infinite dimension. Use this calculation to show that the scheme  $U$  is not affine.

(5) Let  $X$  be a noetherian separated scheme. Define the *cohomological dimension* of  $X$ , denoted  $\text{cd}(X)$ , to be the least integer  $n$  such that  $H^i(X, F) = 0$  for all quasi-coherent sheaves  $F$  and all  $i > n$ . For example, Serre's Theorem III.3.7 in Hartshorne says that  $\text{cd}(X) = 0$  if and only if  $X$  is affine. Grothendieck's Theorem III.2.7 implies that  $\text{cd}(X) \leq \dim(X)$ .

(a) In the definition of  $\text{cd}(X)$ , show that it is sufficient to consider only coherent sheaves on  $X$ . Use exercise II.5.15 and Prop. III.2.9.

(b) If  $X$  is quasi-projective over a field  $k$ , then it is even sufficient to consider vector bundles on  $X$ . Use Cor. II.5.18.

(c) Suppose that  $X$  has a covering by  $r + 1$  open affine subsets. Use Čech cohomology to show that  $\text{cd}(X) \leq r$ .

(d) If  $X$  is quasi-projective scheme of dimension  $r$  over a field  $k$ , show that  $X$  can be covered by  $r + 1$  open affine subsets. Conclude (independent of Grothendieck's theorem) that  $\text{cd}(X) \leq \dim(X)$ .

(e) Let  $Y$  be a set-theoretic complete intersection (exercise I.2.17) of codimension  $r$  in  $X = A_k^n$ . Show that  $\text{cd}(X - Y) \leq r - 1$ .

(6) Let  $X = \text{Spec } k[x_1, x_2, x_3, x_4]$  be affine 4-space over a field  $k$ . Let  $Y_1$  be the plane  $x_1 = x_2 = 0$  and let  $Y_2$  be the plane  $x_3 = x_4 = 0$ . Show that  $Y = Y_1 \cup Y_2$

is not a set-theoretic complete intersection in  $X$ . Therefore the projective closure  $\overline{Y} \subset \mathbf{P}_k^4$  is also not a set-theoretic complete intersection. [Hint: Use problem 5(e) above. Then show that  $H^2(X - Y, \mathcal{O}_X) \neq 0$ , by using exercises III.2.3 (cohomology with support) and III.2.4 (Mayer-Vietoris).]