

**Problem 1** True or false?

- (a) The integral  $\int_0^{2\pi} \int_0^1 \int_r^1 dz dr d\theta$  represents the volume of a right cone.
- (b) The jacobian of the transformation given by  $x = u^2 - 2v, y = 3v - 2uv$  is given by  $-4u^2 + 6u + 4v$ .
- (c) Is the vector field  $\vec{F} = \langle x^2y, xy^2 \rangle$  conservative?
- (d) The divergence of a vector field is a vector field.
- (e) If  $\nabla \times \vec{F} = 0$  then  $\vec{F}$  is conservative.

**Problem 2** Evaluate the integral

$$\int_0^1 \int_{-\sqrt{1-x^2}}^{\sqrt{1-x^2}} \int_{-\sqrt{1-x^2-y^2}}^{\sqrt{1-x^2-y^2}} dz dy dx$$

**Problem 3** Calculate  $\iiint_B \sqrt{x^2 + y^2} dV$  where  $B$  is the region bounded above by the half sphere  $x^2 + y^2 + z^2 = 9$  with  $z \geq 0$  and below by the cone  $z^2 = 3(x^2 + y^2)$ .

**Problem 4** Use a change of variables to evaluate the integral  $\iint_R (x-y)e^{x^2-y^2} dA$  where  $R$  is the region bound by the lines  $x+y=1, x+y=3$  and curves  $x^2-y^2=-1, x^2-y^2=1$ .

**Problem 5** Let  $E$  be the region bounded below by the cone  $z = \sqrt{x^2 + y^2}$  and above by the sphere  $x^2 + y^2 + z^2 = z$ . Set up a triple integral in spherical coordinates and find the volume of the region using the following orders of integration:

- (a)  $d\rho d\phi d\theta$
- (b)  $d\phi d\rho d\theta$ .

**Problem 6** Let  $\vec{F} = \langle 2x \ln(y), \frac{x^2}{y} + z^2, 2yz \rangle$  and let  $C$  be the curve parameterised by

$$\vec{r}(t) = \langle t^2, t, t \rangle \text{ for } 1 \leq t \leq e. \text{ Calculate } \int_C \vec{F} \cdot d\vec{r}$$

- (a) without using the Fundamental Theorem of Line Integrals and
- (b) using the Fundamental Theorem of Line Integrals.

**Problem 7** Calculate the line integral  $\int_C (x^2 + y^2 + z^2) ds$  where  $C$  is the part of the helix parameterised by  $\vec{r}(t) = \langle \cos t, \sin t, t \rangle$  for  $0 \leq t \leq 2\pi$ .

**Problem 8** A solid  $Q$  has the form  $D \times I$  where  $D$  is some finite region in the  $xy$ -plane and  $I = [a, b]$  is a finite interval. The density  $\rho(x, y, z)$  of the solid  $Q$  doesn't depend on the variable  $z$ . Show that the center of mass of  $Q$  lies on the plane  $z = \frac{a+b}{2}$ .

**Problem 9** True or False?

(a) If  $C$  is parameterised by  $\vec{r}(t) = (t, t)$  for  $0 \leq t \leq 1$ , then

$$\int_C xy ds = \int_0^1 t^2 dt.$$

(b) If vector field  $\vec{F}$  has zero curl on the open and connected region  $D$ , then line integrals of  $\vec{F}$  are path independent on  $D$ .

(c) If a vector field  $\vec{F}$  is path independent on an open connected region  $D$ , then the vector field  $\vec{F}$  is conservative on  $D$ .

**Problem 10** Use a change of variables to calculate  $\iint_R (x^2 + 25y^2)^2 dA$  where  $R$  is the ellipse with boundary  $x^2 + 25y^2 = 1$ .

**Problem 11** Find a potential function for  $\vec{F} = \langle 12x^2, \cos y \cos z, 1 - \sin y \sin z \rangle$ .

**Problem 12** Compute  $\int_C \cos x \cos y dx - \sin x \sin y dy$  where  $C$  is parameterised by  $\vec{r}(t) = (t, t^2)$  for  $0 \leq t \leq 1$ .

**Problem 13** Prove that if  $\vec{F}$  is a conservative vector field in  $\mathbb{R}^3$  with continuously differentiable component functions, then  $\nabla \times \vec{F} = 0$ .

**Problem 14** Let  $\vec{F} = \left\langle \frac{y}{x^2 + y^2}, \frac{-x}{x^2 + y^2} \right\rangle$ , the vortex field.

(a) Consider the function  $g(x, y) = \arctan(x/y)$ . Show that  $\vec{F} = \nabla g$ .

(b) Is  $\vec{F}$  conservative?

(c) Consider the path  $P$  that is the two line segments from  $(1, 1)$  to  $(1, 7)$  and then to  $(2, 2)$ . Evaluate  $\int_P \vec{F} \cdot d\vec{r}$ .