

1 Mathematical Models and Methods in Applied Sciences  
 2 Vol. 22, Suppl. (2012) 1140005 (39 pages)  
 3 © World Scientific Publishing Company  
 4 DOI: 10.1142/S0218202511400057



5 **AGGREGATION AND SPREADING VIA**  
 6 **THE NEWTONIAN POTENTIAL:**  
 7 **THE DYNAMICS OF PATCH SOLUTIONS**

8 ANDREA L. BERTOZZI  
 9 *Department of Mathematics,*  
 10 *University of California Los Angeles,*  
 11 *Los Angeles, CA 90095, USA*  
 12 *bertozzi@math.ucla.edu*

13 THOMAS LAURENT  
 14 *Department of Mathematics,*  
 15 *University of California Riverside,*  
 16 *Riverside, CA 92521, USA*  
 17 *laurent@math.ucr.edu*

18 FLAVIEN LÉGER  
 19 *Department of Mathematics,*  
 20 *University of California Los Angeles,*  
 21 *Los Angeles, CA 90095, USA*  
 22 *flavien@math.ucla.edu*

23 Received 27 July 2011  
 24 Communicated by N. Bellomo and F. Brezzi

25 This paper considers the multidimensional active scalar problem of motion of a function  
 26  $\rho(x, t)$  by a velocity field obtained by  $v = -\nabla N * \rho$ , where  $N$  is the Newtonian potential.  
 27 We prove well-posedness of compactly supported  $L^\infty \cap L^1$  solutions of possibly mixed  
 28 sign. These solutions include an important class of solutions that are proportional to  
 29 characteristic functions on a time-evolving domain. We call these *aggregation patches*.  
 30 Whereas positive solutions collapse on themselves in finite time, negative solutions spread  
 31 and converge toward a self-similar spreading circular patch solution as  $t \rightarrow \infty$ . We give  
 32 a convergence rate that we prove is sharp in 2D. In the case of positive collapsing  
 33 solutions, we investigate numerically the geometry of patch solutions in 2D and in 3D  
 34 (axisymmetric). We show that the time evolving domain on which the patch is supported  
 35 typically collapses on a complex skeleton of codimension one.

36 *Keywords:* Active scalar; nonlocal transport; aggregation; Newtonian potential.

37 AMS Subject Classification: 35Q35, 35Q70, 35Q92, 31B10, 34G25, 76B03

\*Current address: CMLA, ENS Cachan, 61 avenue du Président Wilson 94235 Cachan Cedex.

*A. L. Bertozzi, T. Laurent & F. Léger*

## 1 Introduction

Consider the aggregation equation

$$\frac{\partial \rho}{\partial t} + \operatorname{div}(\rho v) = 0, \quad v = -\nabla N * \rho, \quad (1.1)$$

where  $N$  is the Newtonian potential in  $\mathbb{R}^d$  for  $d \geq 2$ . For more general interaction kernels  $K$ , this problem has been a very active area of research in the literature.<sup>7,5,7,9,10,12,14–21,25,28,30–32,38,39,45–49,56,57,65–67</sup> These models arise in a number of applications including aggregation in materials science,<sup>39,59,60</sup> cooperative control,<sup>38</sup> granular flow,<sup>26,27,67</sup> biological swarming models,<sup>55,56,65,66</sup> evolution of vortex densities in superconductors<sup>1,2,35,36,54</sup> and bacterial chemotaxis.<sup>7,15,16,19</sup> A body of recent work has focused on the problem of finite time singularities and local versus global well-posedness in multiple space dimensions for both the inviscid case (1.1)<sup>7,9–13,17,25,32,40,45</sup> and the cases with various kinds of diffusion.<sup>4,13,15,46,47</sup> The highly studied Keller–Segel problem typically has a Newtonian potential and linear diffusion. For the pure transport problem (1.1), of particular interest is the transition from smooth solutions to weak and measure solutions with mass concentration.

In two space dimensions, Eq. (1.1) with the Newtonian potential arises as a model for the evolution of vortex densities in superconductors,<sup>1,2,35,36,50–52,54,62,63</sup> and also in models for adhesion dynamics.<sup>59,60</sup> These problems are known to develop finite time singularities and several papers have considered the existence of solutions with measured initial data.<sup>1,35,60</sup> Among others, our paper develops a complete theory of weak solution in arbitrary dimension for the case where the density function  $\rho$  is bounded, compactly supported and has mixed sign. Our theory includes sharp well-posedness for  $L^\infty$  data including the maximal time interval of existence.

### 22 1.1. Description of the problem and well-posedness theory

Here we use the convention that  $\Delta N = \delta$ , with this convention the interaction kernel is attractive for positive  $\rho$ . Using the fact that  $\operatorname{div} v = -\Delta N * \rho = -\rho$ , Eq. (1.1) can be rewritten

$$\frac{\partial \rho}{\partial t} + \nabla \rho \cdot v = \rho^2, \quad v = -\nabla N * \rho \quad (1.2)$$

and therefore the values of  $\rho$  satisfy the ODE  $\dot{y} = y^2$  along the “characteristics” defined by

$$\frac{d}{dt} X^t(\alpha) = v(X^t(\alpha), t), \quad X^0(\alpha) = 0. \quad (1.3)$$

Solving  $\dot{y} = y^2$  gives the formulas

$$\rho(X^t(\alpha), t) = \left( \frac{1}{\rho_0(\alpha)} - t \right)^{-1} \quad \text{and} \quad \rho(x, t) = \left( \frac{1}{\rho_0(X^{-t}(x))} - t \right)^{-1}, \quad (1.4)$$

*Aggregation via the Newtonian Potential and Aggregation Patches*

where  $X^{-t} : \mathbb{R}^d \rightarrow \mathbb{R}^d$  is the inverse of the mapping  $X^t$ . It is clear from (1.4) that if  $\rho_0$  is strictly positive somewhere, then the first blow-up occurs at time

$$t = \frac{1}{\sup_{x \in \mathbb{R}^d} \rho_0(x)}.$$

On the other hand, if  $\rho_0(x) \leq 0$  for all  $x$ , the values on every characteristic converge to zero as  $t \rightarrow +\infty$  and no blowup occurs. These argument will be made rigorous in Sec. 2, where we prove the existence and uniqueness of compactly supported bounded solutions of mixed sign on the time interval  $[0, T]$  with

$$T < (\sup \rho_0)^{-1}.$$

1 If the initial data is negative, then  $(\sup \rho_0)^{-1} = +\infty$  and therefore solutions exist  
 2 for all  $t > 0$ . The theory is first carried out in Lagrangian variables for functions  
 3  $\rho$  that are Hölder continuous, using the classical Picard theory on a Banach space,  
 4 thus proving both existence and uniqueness. Then we prove the existence for  $L^\infty$   
 5 functions by passing to the limit in smooth approximations, also using Lagrangian  
 6 variables for the passage to the limit. However, the resulting solution is a classical  
 7 solution in the sense of distributions, in Eulerian variables. Uniqueness of  $L^\infty$  solu-  
 8 tions is proved using well-known  $H^{-1}$  energy estimates for the Eulerian form of the  
 9 problem, we do not include the proof but rather refer the reader to the extensive  
 10 literature using these techniques, dating back to the seminal paper of Yudovich<sup>68</sup>  
 11 for  $L^\infty$  vorticity for the 2D Euler problem.

12 We use the term “aggregation equation” in a generalized sense. While non-  
 13 negative solutions collapse on themselves in finite time,<sup>10,25</sup> nonpositive  
 14 solutions spread. The second case is equivalent to the first case with the  
 15 evolution backward in time. The reader should keep this in mind when  
 16 reading Sec. 3, which is devoted to nonpositive “spreading” solutions.

## 17 1.2. Aggregation patches

We now discuss aggregation patches, solutions of (1.1) with initial data

$$\rho_0(x) = c\chi_{\Omega_0}(x),$$

where  $c$  is a (possibly negative) constant and  $\chi_{\Omega_0}(x)$  is the characteristic func-  
 tion of the bounded domain  $\Omega_0 \subset \mathbb{R}^d$ . Note that such initial data are compactly  
 supported and belong to  $L^1 \cap L^\infty$  therefore the well-posedness theory from Sec. 2  
 applies. These solutions are analogues of the famous *vortex patch* solutions of the 2D  
 incompressible Euler equations.<sup>6,23,29,33,34,69</sup> However, for the aggregation problem,  
 the patch solutions exist in any dimension. From (1.4) we see that

$$\rho(x, t) = \left(\frac{1}{c} - t\right)^{-1} \chi_{\Omega_t}(x), \quad (1.5)$$

*A. L. Bertozzi, T. Laurent & F. Léger*

where  $\Omega_t = X^t(\Omega_0)$  is a time evolving domain and the mapping  $X^t : \mathbb{R}^d \rightarrow \mathbb{R}^d$  is defined by (1.3). As mass is conserved, we necessarily have

$$|\Omega_t| = (1 - ct)|\Omega_0|.$$

where  $|\Omega_t|$  stands for the Lebesgue measure of  $\Omega_t$ . Therefore positive patches (i.e.  $c > 0$ ) collapse to a domain of Lebesgue measure 0 at time  $t = 1/c$ . In contrast, the Lebesgue measure of negative patches (i.e.  $c < 0$ ) is increasing linearly in time. The simplest patch solutions are the circular patches

$$\rho(x, t) = \left(\frac{1}{c} - t\right)^{-1} \chi_{\Omega_t}(x), \quad \Omega_t = B(0, R(t)), \quad R(t) = R_0(1 - ct)^{1/d}. \quad (1.6)$$

They are explicit self-similar solutions to (1.1). In Sec. 3 we show that in the case of negative  $L^\infty$  spreading solutions, these self-similar circular patches are global attractor of the dynamics. Moreover we prove that the convergence rate toward these global attractors is given by

$$\|\rho(\cdot, t) - \Phi(\cdot, t)\|_{L^1} \leq Ct^{-\lambda}, \quad \lambda = \frac{1}{2d-1}$$

1 where  $\rho(x, t)$  is a negative solution and  $\Phi(x, t)$  is a spreading negative circular patch.  
2 Note in particular that this give a convergence rate in  $1/\sqrt{t}$  in two dimensions.

In Sec. 4 we give an explicit formula for an elliptical patch in two dimensions. In the collapsing case, the finite time singularity results in convergence to a weighted measure along an interval of length  $2(a_0 - b_0)$  where  $a_0$  is the length of the semi-major axis of the initial data and  $b_0$  is the length of the semi-minor axis (Theorem 4.1). We then show that the  $L^1$  difference between an elliptic spreading patch and a circular one decays like  $1/\sqrt{t}$  (Theorem 4.2), therefore proving the sharpness of our convergence rate in 2D. The remainder of Sec. 4 is devoted to a numerical study of the different patch evolutions in 2D and 3D for both the spreading and collapsing problem. The 3D solutions shown are axisymmetric which allows for a reduction of the computational complexity to that of the 2D problem. In both cases we reduce the numerical simulation to a self-deforming curve in the plane by using a contour dynamics formalism, similar to what has been done for vortex patches in 2D. Indeed, for patch solutions, by integrating by parts we can express  $v$  as a boundary integral:

$$v(x, t) = - \left(\frac{1}{c} - t\right)^{-1} \nabla N * \chi_{\Omega_t}(x) = \left(\frac{1}{c} - t\right)^{-1} \int_{\partial\Omega_t} N(x - y)n(y)d\sigma(y).$$

3 Therefore we just need to compute the motion of the curve  $\partial\Omega_t$  deforming according  
4 to the above velocity field.

5 Whereas an elliptical patch collapses toward a singular measure supported on  
6 a line segment, more general 2D aggregation patches are observed numerically to  
7 converge toward singular measures supported on more complex domains, also of  
8 codimension one, often consisting of the union of several curves. We refer to this

1 union of intersecting curves as the skeleton of the patch at the collapse time. How-  
 2 ever this shape is not the topological skeleton of the initial data. In 3D, aggregation  
 3 patches typically converge toward singular measure uniformly distributed on the  
 4 union of intersecting surfaces, see Fig. 6.

5 In the spreading case we numerically observe a “pinching” phenomena in which  
 6 the solution clearly converges in  $L^1$  to the circular patch solution as predicted by  
 7 the analysis, however the boundary does not necessarily converge to a circle. For  
 8 some initial data the boundary “pinches”, creating defects which consist of slits  
 9 cutting into the circular shape, see Fig. 3. These defects may not disappear in  
 10 the long run. For 3D toroidal initial data the spreading problem deforms to fill  
 11 a sphere in 3D, while preserving the toroidal topology of the initial data. In this  
 12 case there is a natural slit that forms related to the preserved topology of the  
 13 torus.

### 14 **1.3. Relationship with the existing literature**

15 In an earlier paper, two of the authors<sup>41</sup> have shown that the problem considered  
 16 here reduces to the inviscid Burgers equation for the case of radially symmetric data  
 17 in any space dimension. This observation has been made for the one-dimensional  
 18 problem in earlier papers, but not for multiple dimensions. The connection to Burg-  
 19 ers equation allows us to understand many features of this problem via exact solu-  
 20 tions with radial symmetry. The general theory of the equations, considered here,  
 21 has roots in many previous papers in the literature. Section 2 is largely inspired by  
 22 Chaps. 4 and 8 of the book on *Vorticity and Incompressible Flow*<sup>53</sup> however some  
 23 of these ideas and other ideas have been used to study this aggregation problem  
 24 in prior papers. The  $C^{0,\gamma}$  Hölder results are also proved in a recent manuscript.<sup>37</sup>  
 25 Lin and Zhang<sup>50</sup> have studied our problem for spreading case (corresponding to  
 26 negative sign initial data in our model) in two dimensions and develop general exis-  
 27 tence theory for data a measure of a fixed sign. Their techniques are similar to  
 28 the ones we use for the existence of  $L^\infty$  solutions of mixed sign and come from  
 29 well-known methods in incompressible fluid dynamics. Our work generalizes their  
 30 existence results to the mixed sign problem in general dimension and also con-  
 31 sider asymptotic behavior of the spreading problem in general dimension. Nieto,  
 32 Poupaud, and Soler<sup>59</sup> consider our problem in general dimension for data that is in  
 33  $W^{1,\infty}(\mathbb{R}^d)$  — these results being most relevant to our existence theory for Hölder  
 34 continuous solutions. They also consider  $L^\infty$  data for the spreading problem in  
 35 general dimension and for the collapsing problem in one space dimension — corre-  
 36 sponding to the case of Lipschitz solutions of the inviscid Burgers equation. For the  
 37 spreading problem, they use methods involving entropies and energy estimates in  
 38 Eulerian variables — these methods are best applied to the spreading problem. Our  
 39 results for  $L^\infty$  data involve solutions with mixed sign, and the existence theory is  
 40 done mainly in Lagrangian variables using uniform estimates for the particle paths.  
 41 Masmoudi and Zhang<sup>54</sup> consider a closely related problem, where the sign of the

*A. L. Bertozzi, T. Laurent & F. Léger*

1 density changes the sign of the velocity field. They establish well-posedness theory  
 2 in  $W^{1,p}$ ,  $p > 2$ , and in  $C^\gamma$  for  $0 < \gamma < 1$ . Mainini<sup>52</sup> considers our problem in  $L^\infty$   
 3 but without collapse and uses optimal transport theory. Another very related prob-  
 4 lem is the Chapman–Rubinstein–Schatzman model<sup>1,2</sup> which involves an interaction  
 5 kernel that is the fundamental solution of  $(I - \Delta)$  on a bounded domain with Dirich-  
 6 let boundary condition. That literature makes extensive use of optimal transport  
 7 theory.

8 Regarding the global attractor for the spreading problem, Caffarelli and  
 9 Vazquez<sup>24</sup> recently consider the dynamics of fractional diffusion, which can be  
 10 written in our formulation, only with more singular potentials corresponding to  
 11  $(-\Delta)^{-s}$  where  $0 < s < 1$  (our problem corresponds to the case  $s = 1$ ). Note  
 12 that the corresponding kernel operators are of the form  $|x|^{-p}$  where  $p = d - 2s$ .  
 13 For these more singular kernels there is a gain of regularity in the solution as the  
 14 problem is diffusive going forward in time. It has some properties similar to the  
 15 porous media equation. They prove convergence to the global attractor for these  
 16 more singular kernels using entropy methods. Compared to this work, we prove  
 17 convergence to the global attractor via a direct estimate on the size of the support  
 18 of the solution and we obtain a convergence rate in  $L^1$ . In the case of radial symme-  
 19 try, the problem reduces to Burgers equation<sup>11</sup> and thus can be understood using  
 20 classical scalar law theory. For an attractive–repulsive problem<sup>37</sup> that is equiva-  
 21 lent to ours after a change of variables, one can prove convergence to the patch  
 22 solution for initial data with radial symmetry using an explicit computation along  
 23 characteristics.

24 Models of vortex patches date back to the classical example of the Kirchoff  
 25 ellipse<sup>43</sup> which has been extensively studied in straining flows.<sup>8,42,58</sup> Numerical  
 26 simulations of the general vortex patch problem date back to work of Zabusky,  
 27 Hughs, and Roberts<sup>69</sup> and resulted in speculation about the global regularity of  
 28 the vortex patch boundary<sup>23,34</sup> which was finally settled analytically by Chemin<sup>29</sup>  
 29 and reproved using potential theory estimates by Bertozzi and Constantin.<sup>6</sup> To the  
 30 best of our knowledge, the numerical simulation of the aggregation problem and the  
 31 observation that aggregation patches collapse to complex skeletons of codimension  
 32 one are new.

33 Finally we remark that a flow with related properties, but different bound-  
 34 ary conditions, is considered in the context of a level set method used to  
 35 solve an elliptic obstacle problem.<sup>22</sup>

## 36 2. General Well-Posedness Theory for Compactly Supported 37 $L^\infty$ Solutions of Mixed Sign

38 To avoid complications at infinity, for which we know of no physically inter-  
 39 esting cases, we simplify the problem by considering compactly supported data,  
 40 so that the data is also automatically in  $L^1$  due to the essential supremum  
 41 bound.

1 **2.1. Well-posedness theory for Hölder continuous**  
 2 **densities  $C_0^\gamma(\mathbb{R}^d)$**

In this section we derive the well-posedness theory for densities  $\rho \in C_0^\gamma(\mathbb{R}^d)$ , functions that are Hölder continuous, and that have compact support. We note that the problem is studied in Ref. 59 for the case of  $W^{1,\infty}$  densities, which would suffice for our purposes here. However, for completeness we present an argument in Hölder spaces and will use this, including the compact support, to prove well-posedness of general  $L^\infty$  solutions and also to prove a result on long time behavior, which require estimates on the support of the solution. The proof uses ODEs on Banach spaces in Lagrangian coordinates. The ODE (1.3) defining the particle path  $t \mapsto X^t(\alpha)$  can be written as

$$\frac{d}{dt}X^t(\alpha) = - \int_{\mathbb{R}^d} \nabla N(X^t(\alpha) - y) \rho(y, t) dy.$$

The change of variables  $y = X^t(\alpha')$  gives

$$\frac{d}{dt}X^t(\alpha) = - \int_{\mathbb{R}^d} \nabla N(X^t(\alpha) - X^t(\alpha')) \rho(X^t(\alpha'), t) \det(\nabla_\alpha X^t(\alpha')) d\alpha'.$$

However, note that if we define  $Q(\alpha, t) = \det(\nabla_\alpha X^t(\alpha))$ , then  $Q$  satisfies

$$\frac{d}{dt}Q(\alpha, t) = \operatorname{div} v(X^t(\alpha), t) Q(\alpha, t) = -\rho(X^t(\alpha), t) Q(\alpha, t).$$

But since  $\frac{d}{dt}\rho(X^t(\alpha), t) = \rho(X^t(\alpha), t)^2$  we see that  $\rho(X^t(\alpha), t)Q(\alpha, t)$  is constant in time. Using the fact that  $Q(\alpha, 0) = 1$  we obtain the following Lagrangian formulation of (1.1):

$$\begin{aligned} \frac{d}{dt}X^t(\alpha) &= - \int_{\mathbb{R}^d} \nabla N(X^t(\alpha) - X^t(\alpha')) \rho(\alpha', 0) Q(\alpha', 0) d\alpha' \\ &= - \int_{\mathbb{R}^d} \nabla N(X^t(\alpha) - X^t(\alpha')) \rho_0(\alpha') d\alpha'. \end{aligned} \quad (2.1)$$

We then define the Banach space on which to work:

$$B = \{X : \mathbb{R}^d \rightarrow \mathbb{R}^d \text{ such that } \|X\|_{1,\gamma} < \infty\}, \quad (2.2)$$

$$\|X\|_{1,\gamma} = |X(0)| + \|\nabla_\alpha X\|_{L^\infty} + |\nabla_\alpha X|_\gamma, \quad (2.3)$$

where  $|\cdot|_\gamma$  stands for the standard Hölder semi-norm. We write our problem as

$$\frac{d}{dt}X^t = F(X^t), \quad X^0 = \operatorname{Id}, \quad (2.4)$$

where  $F : B \rightarrow B$  is defined by

$$F(X)(\alpha) = - \int_{\mathbb{R}^d} \nabla N(X(\alpha) - X(\alpha')) \rho_0(\alpha') d\alpha'. \quad (2.5)$$

3 **Theorem 2.1.** (Local existence and continuation of  $C^{1,\gamma}$  particle paths) *Consider*  
 4 *an initial density  $\rho_0$  that is Hölder continuous and with compact support. Then the*

*A. L. Bertozzi, T. Laurent & F. Léger*

1 *initial value problem (2.4)–(2.5) has a unique solution on a maximal time interval*  
 2 *of existence  $[0, T^*)$ . Either  $T^*$  is infinite or the Banach norm (2.3) blows up as*  
 3  *$t \rightarrow T^*$ .*

4 **Remark 2.1.** Although we prove well-posedness of the problem in the Lagrangian  
 5 formulation, this result implies the existence of a solution to the problem in original  
 6 Eulerian variables (1.1). This is discussed in detail for the Euler problem in Chap. 4  
 7 of Ref. 53 and is proved rigorously for the weaker case of  $L^\infty$  data in the next  
 8 subsection.

**Proof.** The proof is similar to the local existence proof of the vorticity formulation  
 of Euler equation in Chap. 4 of Ref. 53 and therefore here we just discuss a few key  
 points. In order to apply the Picard theorem on a Banach space, the hardest part  
 is to show that  $F : B \rightarrow B$  is Lipschitz. We focus on the calculation of the first  
 variation with respect to  $X$ . That results in a linear operator which we write as

$$\begin{aligned} \frac{\delta F}{\delta X} Y &:= \frac{d}{d\epsilon} F(X + \epsilon Y)|_{\epsilon=0} \\ &= -\frac{d}{d\epsilon} \int \nabla N(X(\alpha) - X(\alpha') + \epsilon[Y(\alpha) - Y(\alpha')]) \rho_0(\alpha') d\alpha' |_{\epsilon=0} \\ &= -\int D^2 N(X(\alpha) - X(\alpha')) [Y(\alpha) - Y(\alpha')] \rho_0(\alpha') d\alpha'. \end{aligned}$$

Following the arguments in Chap. 4 of Ref. 53 (see, for example, Proposition  
 4.2) we obtain

$$\left\| \frac{\delta F}{\delta X} Y \right\|_{1,\gamma} \leq C(\|X\|_{1,\gamma}) \|\rho_0\|_\gamma \|Y\|_{1,\gamma}, \quad (2.6)$$

9 where  $\|X\|_\gamma = \|X\|_{L^\infty} + |X|_\gamma$ . This shows the Lipschitz continuity of  $F$ . We refer the  
 10 reader to Chap. 4 of Ref. 53 for the details of this calculation.  $\square$

Define  $|\cdot|_{\max}$  and  $|\cdot|_{\min}$  as follows:

$$|f|_{\max} = \sup_{x \in \mathbb{R}^d} f(x) \quad \text{and} \quad |f|_{\min} = -\inf_{x \in \mathbb{R}^d} f(x). \quad (2.7)$$

Let us emphasize that  $|f|_{\min} \geq 0$ . Also we obviously have

$$\frac{1}{2}(|f|_{\min} + |f|_{\max}) \leq \|f\|_{L^\infty} \leq |f|_{\min} + |f|_{\max}. \quad (2.8)$$

11 **Theorem 2.2.** *Let  $[0, T^*)$  be the maximal time interval of existence of the ini-*  
 12 *tial value problem (2.4)–(2.5) with Hölder continuous, compactly supported, initial*  
 13 *density  $\rho_0$ . Then  $T^* = 1/|\rho_0|_{\max}$ .*

14 Note that if  $\rho_0 \leq 0$  and compactly supported then  $|\rho_0|_{\max} = 0$  and  $T^* = +\infty$ . The  
 15 rest of this subsection is devoted to the proof of Theorem 2.2. We follow arguments

*Aggregation via the Newtonian Potential and Aggregation Patches*

1 similar to the classical Beale–Kato–Majda theorem for the 3D Euler equation,<sup>3</sup> in  
2 particular the alternate Hölder space proof from Chap. 4 of Ref. 53.

Assume that the classical solution  $\rho(x, t)$  given by Theorem 2.1 exists on a time interval  $[0, T)$ . Since it is a classical solution it satisfies (1.4) along the particle path. We therefore see that  $T$  cannot be strictly greater than  $1/|\rho_0|_{\max}$ , otherwise the solution would blow up and leave the space of continuous functions. This implies that  $T^* \leq 1/|\rho_0|_{\max}$ . Then we show that if the time interval  $[0, T)$  on which the solution is defined is such that  $T < 1/|\rho_0|_{\max}$ , then there exists a constant  $C$  such that

$$\|X^t\|_{1,\gamma} < C \quad \text{for all } t \in [0, T), \quad (2.9)$$

3 therefore giving  $T^* \geq 1/|\rho_0|_{\max}$  and concluding the proof.

We now prove (2.9) under the condition  $T < 1/|\rho_0|_{\max}$ . Define the constant

$$k = 1 - T|\rho_0|_{\max} \quad \text{and} \quad K = 1 + T|\rho_0|_{\min}. \quad (2.10)$$

Clearly  $0 < k \leq 1$  and  $K \geq 1$ . Note that because of (1.4) we have the uniform bounds

$$|\rho_0|_{\max} \leq |\rho(\cdot, t)|_{\max} = \left( \frac{1}{|\rho_0|_{\max}} - t \right)^{-1} \leq \frac{|\rho_0|_{\max}}{k}, \quad (2.11)$$

$$\frac{|\rho_0|_{\min}}{K} \leq |\rho(\cdot, t)|_{\min} = \left( \frac{1}{|\rho_0|_{\min}} + t \right)^{-1} \leq |\rho_0|_{\min} \quad (2.12)$$

for all  $t \in [0, T]$ . Therefore, from (2.8),

$$\begin{aligned} \frac{\|\rho_0\|_{L^\infty}}{2K} &\leq \frac{|\rho_0|_{\min} + |\rho_0|_{\max}}{2K} \leq \|\rho(\cdot, t)\|_{L^\infty} \\ &\leq \frac{|\rho_0|_{\min} + |\rho_0|_{\max}}{k} \leq \frac{2\|\rho_0\|_{L^\infty}}{k} \end{aligned} \quad (2.13)$$

4 for all  $t \in [0, T]$ . We then prove the following elementary estimate:

**Lemma 2.1.** *For all  $t \in [0, T)$*

$$|\rho(\cdot, t)|_\gamma \leq \frac{|\rho_0|_\gamma}{k^2} \|\nabla_x X^{-t}\|_{L^\infty}^\gamma. \quad (2.14)$$

**Proof.** From (1.4), we have:

$$\rho(x, t) = \frac{\rho_0(X^{-t}(x))}{1 - \rho_0(X^{-t}(x))t}. \quad (2.15)$$

A. L. Bertozzi, T. Laurent & F. Léger

Then we write:

$$\begin{aligned} |\rho(x, t) - \rho(y, t)| &= \left| \frac{\rho_0(X^{-t}(x)) - \rho_0(X^{-t}(y))}{(1 - \rho_0(X^{-t}(x))t)(1 - \rho_0(X^{-t}(y))t)} \right| \\ &\leq \frac{1}{k^2} |\rho_0(X^{-t}(x)) - \rho_0(X^{-t}(y))| \\ &\leq \frac{1}{k^2} |\rho_0|_\gamma \|\nabla_x X^{-t}\|_{L^\infty}^\gamma |x - y|^\gamma. \quad \square \end{aligned}$$

1 We then need the following potential theory lemma from Ref. 53 (Lemmas 4.5  
2 and 4.6 on p. 144):

**Lemma 2.2.** *Consider  $f \in C^\gamma(\mathbb{R}^d; \mathbb{R}^d)$ ,  $0 < \gamma < 1$ , be a compactly supported function with supported in the ball of radius  $R$ . Then there exists a constant  $C$  independent of  $R$  and  $f$  such that*

$$\|PVN_{ij} * f\|_{L^\infty} \leq C \left\{ |f|_\gamma \epsilon^\gamma + \max\left(1, \ln \frac{R}{\epsilon}\right) \|f\|_{L^\infty} \right\}, \quad \forall \epsilon > 0, \quad (2.16)$$

$$|PVN_{ij} * f|_\gamma \leq C |f|_\gamma. \quad (2.17)$$

3 Here  $PVN_{ij} * f$  denotes the principal value singular integral of the Hessian matrix  
4 of the Newtonian potential convolved with  $f$ .

5 We note that all entries of the convolution kernel  $N_{ij}$  have mean zero on spheres  
6 (see e.g. Proposition 2.18 p. 74 of Ref. 53) because they are each exact derivatives  
7 of  $\nabla N$  which is homogeneous of degree  $1 - d$ . Hence the PV integral is well-defined  
8 on the Hölder spaces. We also note that the full velocity gradient matrix can be  
9 expressed in terms of principal value integrals of the form  $PVN_{ij} * \rho$  plus a constant  
10 matrix times  $\rho$  (see e.g. Proposition 2.17, p. 74 of Ref. 53). This fact will be used  
11 below in estimating  $\nabla v$  in both sup and Hölder norms.

12 Next we derive a crude estimate on the size of the support of  $\rho(x, t)$ :

**Lemma 2.3.** *Suppose that  $\rho_0$  is supported in a ball of radius  $R_0$ . Then for all  $t \in [0, T)$ ,  $\rho(\cdot, t)$  is supported in a ball of radius*

$$R(t) = R_0 + \frac{C}{k} (\|\rho_0\|_{L^1} + \|\rho_0\|_{L^\infty}) t, \quad (2.18)$$

13 where  $C > 0$  is a constant depending only on the dimension

**Proof.** This lemma is a direct consequence of a simple potential theory estimate that gives a bound on the velocity of the particle path:

$$\|v(\cdot, t)\|_{L^\infty} = \|\nabla N * \rho(\cdot, t)\|_{L^\infty} \leq \frac{C}{k} (\|\rho_0\|_{L^1} + \|\rho_0\|_{L^\infty}) \quad (2.19)$$

14 for some constant  $C$  depending only on the dimension. This potential theory esti-  
15 mate is proven in the next subsection, see (2.36).  $\square$

*Aggregation via the Newtonian Potential and Aggregation Patches*

Differentiating the particle equation gives

$$\frac{d}{dt} \nabla_{\alpha} X(\alpha, t) = \nabla v(X(\alpha, t), t) \nabla_{\alpha} X(\alpha, t). \quad (2.20)$$

This equation directly implies

$$\frac{d}{dt} \|\nabla_{\alpha} X(\cdot, t)\|_{L^{\infty}} \leq \|\nabla v(X(\cdot, t), t)\|_{L^{\infty}} \|\nabla_{\alpha} X(\cdot, t)\|_{L^{\infty}},$$

and hence

$$\|\nabla_{\alpha} X(\cdot, t)\|_{L^{\infty}} \leq \exp\left(\int_0^t \|\nabla v(\cdot, s)\|_{L^{\infty}} ds\right). \quad (2.21)$$

To bound  $|\nabla_{\alpha} X(\cdot, t)|_{\gamma}$ , we note that Eq. (2.20) implies that

$$\begin{aligned} \frac{d}{dt} |\nabla_{\alpha} X(\cdot, t)|_{\gamma} &\leq |\nabla v(X(\cdot, t), t)|_{\gamma} \|\nabla_{\alpha} X(\cdot, t)\|_{L^{\infty}} + \|\nabla v(X(\cdot, t), t)\|_{L^{\infty}} |\nabla_{\alpha} X(\cdot, t)|_{\gamma} \\ &\leq |\nabla v(\cdot, t)|_{\gamma} \|\nabla_{\alpha} X(\cdot, t)\|_{L^{\infty}}^{1+\gamma} + \|\nabla v(X(\cdot, t), t)\|_{L^{\infty}} |\nabla_{\alpha} X(\cdot, t)|_{\gamma} \\ &\leq C |\rho(\cdot, t)|_{\gamma} e^{(1+\gamma) \int_0^t \|\nabla v(\cdot, s)\|_{L^{\infty}} ds} + \|\nabla v(\cdot, t)\|_{L^{\infty}} |\nabla_{\alpha} X(\cdot, t)|_{\gamma}. \end{aligned} \quad (2.22)$$

The last estimate is a direct application of (2.17) and (2.21). Using Lemma 2.1 combined with the fact that the inverse characteristic map satisfies  $\frac{d}{dt} X^{-t} = -v(X^{-t}, t)$ , and thus satisfies an inequality like (2.21), we obtain:

$$|\rho(\cdot, t)|_{\gamma} \leq \frac{|\rho_0|_{\gamma}}{k^2} \exp\left(\gamma \int_0^t \|\nabla v(\cdot, s)\|_{L^{\infty}} ds\right). \quad (2.23)$$

Finally applying (2.16) with  $\epsilon = (\|\rho(\cdot, t)\|_{L^{\infty}} / |\rho(\cdot, t)|_{\gamma})^{1/\gamma}$  we have

$$\|\nabla v(\cdot, t)\|_{L^{\infty}} \leq C \left\{ \|\rho(\cdot, t)\|_{L^{\infty}} + \max\left(1, \ln\left(\frac{R(t) |\rho(\cdot, t)|_{\gamma}^{1/\gamma}}{\|\rho(\cdot, t)\|_{L^{\infty}}^{1/\gamma}}\right)\right) \|\rho(\cdot, t)\|_{L^{\infty}} \right\}.$$

Note that due to Lemma 2.3 and (2.13) the logarithm term can be bounded as follows:

$$\ln\left(\frac{R(t) |\rho(\cdot, t)|_{\gamma}^{1/\gamma}}{\|\rho(\cdot, t)\|_{L^{\infty}}^{1/\gamma}}\right) \leq C + \frac{1}{\gamma} \ln |\rho(\cdot, t)|_{\gamma}$$

for some constants  $C$  which depend only on the initial data  $\rho_0$  and the time  $T$ . And therefore, using (2.13) again we find

$$\|\nabla v(\cdot, t)\|_{L^{\infty}} \leq c_1 + c_2 \ln |\rho(\cdot, t)|_{\gamma} \quad (2.24)$$

for some constants  $c_1$  and  $c_2$  which depend only on the initial data  $\rho_0$  and the time  $T$ . Substituting (2.23) in (2.24) we obtain:

$$\|\nabla v(\cdot, t)\|_{L^{\infty}} \leq \tilde{c}_1 + \tilde{c}_2 \int_0^t \|\nabla v(\cdot, s)\|_{L^{\infty}} ds, \quad (2.25)$$

1 where  $\tilde{c}_1$  and  $\tilde{c}_2$  are again two constants depending only on the initial data  $\rho_0$  and  
2 the time  $T$ . By Gronwall lemma this gives us a bound for  $\|\nabla v(\cdot, t)\|_{L^{\infty}}$  on the time

*A. L. Bertozzi, T. Laurent & F. Léger*

1 interval  $[0, T)$ , which in turn, from (2.23), gives a bound for  $|\rho(\cdot, t)|_\gamma$  on the same  
 2 time interval  $[0, T)$ . We can then use these two bounds on  $\|\nabla v(\cdot, t)\|_{L^\infty}$  and  $|\rho(\cdot, t)|_\gamma$   
 3 together with Gronwall lemma to conclude from (2.22) that  $|\nabla_\alpha X^t|_\gamma$  is bounded  
 4 on the time interval  $[0, T)$ .

5 Similar but easier estimates can be obtained for the other terms in the Banach  
 6 norm  $\|X^t\|_{1,\gamma}$  defined in (2.3). This concludes the proof of Theorem 2.2.

## 7 **2.2. Existence of $L^\infty$ solutions**

8 This subsection is devoted to the proof of the following theorem:

**Theorem 2.3.** *Let  $\rho_0 \in L^1 \cap L^\infty(\mathbb{R}^d)$  with compact support and let  $T$  be such that:*

$$0 < T < \frac{1}{|\rho_0|_{\max}}. \quad (2.26)$$

*Then there exists a function*

$$\rho \in C([0, T], L^1(\mathbb{R}^d)) \cap L^\infty(\mathbb{R}^d \times (0, T))$$

*satisfying Eq. (1.1) in the sense of distribution and satisfying  $\rho(\cdot, 0) = \rho_0(\cdot)$ .  
 Moreover, for all  $t \in [0, T]$  we have the following equalities:*

$$\int_{\mathbb{R}^d} \rho(x, t) dx = \int_{\mathbb{R}^d} \rho_0(x) dx, \quad (2.27)$$

$$|\rho(\cdot, t)|_{\max} = \left( \frac{1}{|\rho_0|_{\max}} - t \right)^{-1} \quad \text{and} \quad |\rho(\cdot, t)|_{\min} = \left( \frac{1}{|\rho_0|_{\min}} + t \right)^{-1}. \quad (2.28)$$

Note that if the initial data is negative, then  $|\rho_0|_{\max} = 0$  and  $T$  can be chosen as large as we want. Therefore when the equation is “spreading”, we have global existence. We recall that a function  $\rho(x, t)$  satisfies Eq. (1.1) in the sense of distribution if for all  $\phi \in C_0^\infty(\mathbb{R}^d \times (0, T))$ ,

$$\int_0^T \int_{\mathbb{R}^d} \left( \frac{\partial \phi}{\partial t}(x, t) + v(x, t) \cdot \nabla \phi(x, t) \right) \rho(x, t) dx dt = 0, \quad (2.29)$$

where  $v(x, t) = -(\nabla N * \rho(\cdot, t))(x)$ .

Let us introduce the norm

$$\|\cdot\| = \|\cdot\|_{L^1} + \|\cdot\|_{L^\infty}.$$

Since  $\rho \in C([0, T], L^1(\mathbb{R}^d)) \cap L^\infty(\mathbb{R}^d \times [0, T])$  we clearly have that

$$\|\rho(\cdot, t)\| \leq C \quad \text{for all } t \in [0, T]$$

9 and as we will see, from classical potential theory estimates, this implies that the  
 10 velocity field  $v = -\nabla N * \rho$  is bounded on  $\mathbb{R}^d \times (0, T)$ . Therefore Eq. (2.29) makes  
 11 sense.

*Aggregation via the Newtonian Potential and Aggregation Patches*

1 To prove Theorem 2.3 we follow the ideas in Refs. 9, 53 and 68 by approximating  
 2 the initial data by convolving with a mollifier. As usual let  $\eta \in C_0^\infty(\mathbb{R}^d)$  be a positive  
 3 function of mass 1 and define  $\eta^\epsilon(x) = \epsilon^{-d}\eta(x/\epsilon)$ .

4 **Proposition 2.1.** *Consider a compactly supported initial density  $\rho_0 \in L^1 \cap L^\infty(\mathbb{R}^d)$   
 5 and let  $\rho^\epsilon, v^\epsilon$  be the corresponding smooth solution of the same evolution equation  
 6 with regularized initial data  $\rho_0^\epsilon := \eta^\epsilon * \rho_0$  on the time interval  $[0, T]$  where  $T <$   
 7  $1/|\rho_0|_{\max}$ . Then we have the following:*

- 8 (i)  $\rho^\epsilon \in C([0, T], L^1(\mathbb{R}^d))$ .  
 (ii) There exist constants  $c_1 > 0$  and  $c_2 > 0$  such that

$$\|v^\epsilon(\cdot, t)\|_{L^\infty} \leq c_1 \|\rho^\epsilon(\cdot, t)\| \leq c_2 \|\rho_0\| \quad \text{for all } \epsilon > 0 \text{ and } t \in [0, T].$$

- (iii) There exist functions  $\rho$  and  $v = -\nabla N * \rho$  such that

$$\sup_{t \in [0, T]} \|\rho^\epsilon(\cdot, t) - \rho(\cdot, t)\|_{L^1(\mathbb{R}^d)} \rightarrow 0 \quad \text{as } \epsilon \rightarrow 0, \quad (2.30)$$

$$\|v^\epsilon - v\|_{L^\infty(\mathbb{R}^d \times (0, T))} \rightarrow 0 \quad \text{as } \epsilon \rightarrow 0. \quad (2.31)$$

**Proof.** By mollifying the initial data at time zero, we consider continuous solutions of the above problem

$$\rho^\epsilon(x, t) = \frac{\rho_0^\epsilon(X_\epsilon^{-t}(x))}{1 - \rho_0^\epsilon(X_\epsilon^{-t}(x))t}, \quad (2.32)$$

where the particle trajectories satisfy

$$\frac{d}{dt} X_\epsilon^t = v^\epsilon(X_\epsilon^t(\alpha, t), t), \quad X_\epsilon^0(\alpha) = \alpha,$$

and the velocity  $v^\epsilon = \nabla N * \rho^\epsilon$ . Recall from the previous section that the function  $t \mapsto \rho^\epsilon(X_\epsilon^t(\alpha), t) \det \nabla_\alpha X_\epsilon^t(\alpha)$  is constant and therefore, using (2.32), we have that

$$\det(\nabla_\alpha X_\epsilon^t(\alpha)) = 1 - \rho_0^\epsilon(\alpha)t. \quad (2.33)$$

As in the previous subsection define the constants  $0 < k < 1$  and  $K \geq 1$  by

$$k = 1 - T|\rho_0|_{\max} \quad \text{and} \quad K = 1 + T|\rho_0|_{\min}. \quad (2.34)$$

We then have

$$k \leq \det(\nabla_\alpha X_\epsilon^t(\alpha)) \leq K \quad (2.35)$$

9 for all  $\epsilon > 0$ ,  $t \in [0, T]$  and  $\alpha \in \mathbb{R}^d$ .

We first prove (ii). Note that using the fact that  $\det(\nabla_\alpha X_\epsilon^t(\alpha)) = 1 - \rho_0^\epsilon(\alpha)t > 0$  and doing the change of variable  $\alpha = X_\epsilon^{-t}(x)$  we have

$$\|\rho^\epsilon(\cdot, t)\|_{L^1} = \int_{\mathbb{R}^d} \frac{|\rho_0^\epsilon(X_\epsilon^{-t}(x))|}{1 - \rho_0^\epsilon(X_\epsilon^{-t}(x))t} dx = \int_{\mathbb{R}^d} |\rho_0^\epsilon(\alpha)| d\alpha = \|\rho_0^\epsilon\|_{L^1} \leq \|\rho_0\|_{L^1}.$$

A. L. Bertozzi, T. Laurent & F. Léger

From (2.32) we easily obtained  $\|\rho^\epsilon(\cdot, t)\|_{L^\infty} \leq \frac{2}{k}\|\rho_0^\epsilon\|_{L^\infty} \leq \frac{2}{k}\|\rho_0\|_{L^\infty}$  for all  $t \in [0, T]$  (such an estimate was derived in the previous section, see (2.11)–(2.13)). We therefore obtain

$$\|\rho^\epsilon(\cdot, t)\| \leq \frac{2}{k} \|\rho_0\| \quad \forall t \in [0, T].$$

Now let  $\omega(x)$  be the characteristic function of the unit ball,  $\omega(x) = \chi_{B(0,1)}(x)$ , then

$$\begin{aligned} \|v^\epsilon(\cdot, t)\|_{L^\infty} &\leq \|\omega \nabla N\|_{L^1} \|\rho^\epsilon\|_{L^\infty} + \|(1 - \omega) \nabla N\|_{L^\infty} \|\rho^\epsilon\|_{L^1} \\ &\leq C \|\rho^\epsilon(\cdot, t)\| \\ &\leq \frac{2C}{k} \|\rho_0\| \end{aligned} \quad (2.36)$$

1 which concludes the proof of (ii).

2 We now turn to the proof of (iii). We first need the following potential theory  
3 estimate that establishes a uniform log-Lipschitz estimate for the velocity field:

**Lemma 2.4.** (Potential theory estimate for the velocity) *Given an initial density  $\rho_0 \in L^1 \cap L^\infty(\mathbb{R}^d)$  and let  $\rho^\epsilon$  and  $v^\epsilon$  be the smooth solutions as defined above. Then  $v^\epsilon$  satisfies the following estimate independent of  $\epsilon$*

$$\sup_{t \in [0, T]} |v^\epsilon(x^1, t) - v^\epsilon(x^2, t)| \leq c \|\rho(\cdot, t)\| |x^1 - x^2| (1 - \ln^- |x^1 - x^2|), \quad (2.37)$$

4 where  $\ln^-$  is the negative part (near field) of the natural log.

5 This lemma is proved for the case of the 2D Biot–Savart kernel for the incom-  
6 pressible Euler equations in Ref. 53. The extension to our problem is straightforward  
7 because our velocity kernel satisfies the same conditions as the Biot–Savart kernel,  
8 in fact the Biot–Savart kernel is a special case of the orthogonal flow in 2D. We  
9 leave the proof to the reader, noting that the estimate is well known for potential  
10 theory in  $\mathbb{R}^d$  for the Poisson equation. This lemma directly yields the following  
11 estimates for the particle paths:

**Lemma 2.5.** (Potential theory estimates for characteristics) *Given the assumptions of Lemma 2.4 then there exist  $C > 0$  and an exponent  $\beta(t) = \exp(-C \|\rho_0\| t)$  such that for all  $\epsilon > 0$  and  $t \in [0, T]$ ,*

$$|X_\epsilon^{-t}(x^1) - X_\epsilon^{-t}(x^2)| \leq C |x^1 - x^2|^{\beta(t)}, \quad (2.38)$$

$$|X_\epsilon^t(\alpha^1) - X_\epsilon^t(\alpha^2)| \leq C |\alpha^1 - \alpha^2|^{\beta(t)} \quad (2.39)$$

and for all  $t_1, t_2 \in [0, T]$ ,

$$|X_\epsilon^{-t_1}(x) - X_\epsilon^{-t_2}(x)| \leq C |t_1 - t_2|^{\beta(t)}, \quad (2.40)$$

$$|X_\epsilon^{t_1}(\alpha) - X_\epsilon^{t_2}(\alpha)| \leq C |t_1 - t_2|^{\beta(t)}. \quad (2.41)$$

*Aggregation via the Newtonian Potential and Aggregation Patches*

This lemma is proved in Ref. 53 (see Lemma 8.2) assuming the conditions of Lemma 2.4 above. We note that the *a priori* bound for the  $L^\infty$  norm of the velocity field actually gives Lipschitz continuity in time, although this is not needed for the proof below. By the Arzelà–Ascoli theorem there exist two functions  $X^t(\alpha)$  and  $Y^t(x)$  and a sequence  $\epsilon_k \rightarrow 0$  such that

$$X_{\epsilon_k}^t(\alpha) \rightarrow X^t(\alpha) \quad \text{and} \quad X_{\epsilon_k}^{-t}(x) \rightarrow Y^t(x) \quad \text{as } \epsilon_k \rightarrow 0 \quad (2.42)$$

and these convergences are uniform both in time and space on compact set of  $\mathbb{R}^d \times [0, T]$ . By passing to the limit in

$$X_\epsilon^t(X_\epsilon^{-t}(x)) = x; \quad X_\epsilon^{-t}(X_\epsilon^t(\alpha)) = \alpha$$

1 we obtain that  $Y^t$  is the inverse of  $X^t$ . So we will write  $Y^t = X^{-t}$ .

Given the above limiting particle paths  $X^{-t}(x)$  we define the limiting density according to the formula

$$\rho(x, t) = \frac{\rho_0(X^{-t}(x))}{1 - \rho_0(X^{-t}(x))t}. \quad (2.43)$$

We also define the functions

$$\tilde{\rho}_0^\epsilon(\alpha, t) \equiv \frac{\rho_0^\epsilon(\alpha)}{1 - \rho_0^\epsilon(\alpha)t} \quad \text{and} \quad \tilde{\rho}_0(\alpha, t) \equiv \frac{\rho_0(\alpha)}{1 - \rho_0(\alpha)t}$$

so that  $\rho^\epsilon$  and  $\rho$  can be rewritten as:

$$\rho^\epsilon(x, t) = \tilde{\rho}_0^\epsilon(X_\epsilon^{-t}(x), t) \quad \text{and} \quad \rho(x, t) = \tilde{\rho}_0(X^{-t}(x), t). \quad (2.44)$$

2 The tilde notation is used to denote the time evolving densities in Lagrangian coordinates. Our goal next is to prove that  $\rho^\epsilon$  converge in  $L^1$  uniformly in time toward  
 3  $\rho$ . The proof here differs from the arguments in<sup>53</sup> for the 2D vorticity problem due  
 4 to the fact that the particle paths are not volume preserving, in contrast to the  
 5 incompressibility of the 2D vorticity problem, and the fact that the density evolves  
 6 in a prescribed way along particle paths rather than being conserved. We first derive  
 7 some straightforward estimates on  $\tilde{\rho}_0^\epsilon$  and  $\tilde{\rho}_0$ :  
 8

**Lemma 2.6.** *For all  $\epsilon > 0$  and  $t, s \in [0, T]$  we have:*

$$\sup_{t \in [0, T]} \|\tilde{\rho}_0^\epsilon(\cdot, t) - \tilde{\rho}_0(\cdot, t)\|_{L^1} \leq \frac{1}{k^2} \|\rho_0^\epsilon(\cdot) - \rho_0(\cdot)\|_{L^1}, \quad (2.45)$$

$$\|\tilde{\rho}_0^\epsilon(\cdot, t) - \tilde{\rho}_0^\epsilon(\cdot, s)\|_{L^1} \leq \frac{\|\rho_0\|_{L^\infty} \|\rho_0\|_{L^1}}{k^2} |t - s|, \quad (2.46)$$

$$\|\nabla_\alpha \tilde{\rho}_0^\epsilon(\cdot, t)\|_{L^\infty} = \frac{1}{k^2} \|\nabla_\alpha \rho_0^\epsilon(\cdot)\|_{L^\infty}. \quad (2.47)$$

A. L. Bertozzi, T. Laurent & F. Léger

**Proof.** Estimate (2.45) is a simple consequence of the fact that:

$$\frac{\rho_0^\epsilon(\alpha)}{1 - \rho_0^\epsilon(\alpha)t} - \frac{\rho_0(\alpha)}{1 - \rho_0(\alpha)t} = \frac{\rho_0^\epsilon(\alpha) - \rho_0(\alpha)}{(1 - \rho_0^\epsilon(\alpha)t)(1 - \rho_0(\alpha)t)}.$$

Estimate (2.46) come from the equality

$$\frac{\rho_0^\epsilon(\alpha)}{1 - \rho_0^\epsilon(\alpha)t} - \frac{\rho_0^\epsilon(\alpha)}{1 - \rho_0^\epsilon(\alpha)s} = \frac{\rho_0^\epsilon(\alpha)^2}{(1 - \rho_0^\epsilon(\alpha)t)(1 - \rho_0^\epsilon(\alpha)s)}(t - s).$$

Estimate (2.47) comes from the fact that:

$$\nabla_\alpha \tilde{\rho}_0^\epsilon(\alpha, t) = \frac{1}{(1 - \rho_0^\epsilon(\alpha)t)^2} \nabla_\alpha \rho_0^\epsilon(\alpha). \quad \square$$

Then we note that as a direct consequence of (2.35) and the change of variable formula, we have that for any nonnegative integrable function  $f$ ,

$$k \int_{\mathbb{R}^d} f(\alpha) d\alpha \leq \int_{\mathbb{R}^d} f(X_\epsilon^{-t}(x)) dx \leq K \int_{\mathbb{R}^d} f(\alpha) d\alpha. \quad (2.48)$$

1 To conclude the proof of (iii) we will need a similar estimate for the limiting particle  
2 path  $X^{-t}(x)$ :

**Lemma 2.7.** For all  $t \in [0, T]$ , and nonnegative  $f \in L^1(\mathbb{R}^d)$ ,

$$\int_{\mathbb{R}^d} f(X^{-t}(x)) dx \leq K \int_{\mathbb{R}^d} f(\alpha) d\alpha. \quad (2.49)$$

The proof, which involves some technical arguments from real analysis, can be found in the appendix. We are now ready to prove that  $\rho^\epsilon$  converge in  $L^1$  uniformly in time toward  $\rho$ . Using formula (2.44) we write:

$$\begin{aligned} \|\rho^\epsilon(\cdot, t) - \rho(\cdot, t)\|_{L^1} &\leq \|\tilde{\rho}_0^\epsilon(X_\epsilon^{-t}, t) - \tilde{\rho}_0(X_\epsilon^{-t}, t)\|_{L^1} \\ &\quad + \|\tilde{\rho}_0(X_\epsilon^{-t}, t) - \tilde{\rho}_0(X^{-t}, t)\|_{L^1}. \end{aligned}$$

Using (2.48) and (2.45) we see that the first term is bounded by  $\frac{K}{k^2} \|\rho_0^\epsilon - \rho_0\|_{L^1}$  and therefore can be made small uniformly in time. Next we rewrite the second term as

$$\begin{aligned} \|\tilde{\rho}_0(X_\epsilon^{-t}, t) - \tilde{\rho}_0(X^{-t}, t)\|_{L^1} &\leq \|\tilde{\rho}_0(X_\epsilon^{-t}, t) - \tilde{\rho}_0^{\epsilon_1}(X_\epsilon^{-t}, t)\|_{L^1} \\ &\quad + \|\tilde{\rho}_0^{\epsilon_1}(X_\epsilon^{-t}, t) - \tilde{\rho}_0^{\epsilon_1}(X^{-t}, t)\|_{L^1} \\ &\quad + \|\tilde{\rho}_0^{\epsilon_1}(X^{-t}, t) - \tilde{\rho}_0(X^{-t}, t)\|_{L^1}. \end{aligned}$$

Note that the function  $\tilde{\rho}_0^{\epsilon_1}$  is different from  $\tilde{\rho}_0^\epsilon$ . Let us fix a  $\delta > 0$ . Once again, because of the bound (2.48) and the bound (2.49) from Lemma 2.7 we can choose an  $\epsilon_1$  such that the first and the last terms are smaller than  $\delta/3$  for all  $\epsilon > 0$  and for all  $t \in [0, T]$ . We then claim that as  $\epsilon \rightarrow 0$ , the middle term converges to 0

*Aggregation via the Newtonian Potential and Aggregation Patches*

uniformly in time. Indeed, because of (2.47), there exists a constant  $C(\epsilon_1) > 0$  such that

$$|\tilde{\rho}_0^{\epsilon_1}(X_\epsilon^{-t}(x), t) - \tilde{\rho}_0^{\epsilon_1}(X^{-t}(x), t)| \leq C(\epsilon_1)|X_\epsilon^{-t}(x) - X^{-t}(x)|.$$

Using the uniform convergence of the backward particle path and the fact that there exists a compact set  $K_0$  such that

$$X_\epsilon^t(\text{supp } \rho_0^{\epsilon_1}) \subset K_0 \quad \text{for all } t \in [0, T] \text{ and all } \epsilon > 0,$$

1 we can then choose  $\epsilon$  small enough to make the  $\|\tilde{\rho}_0^{\epsilon_1}(X_\epsilon^{-t}, t) - \tilde{\rho}_0^{\epsilon_1}(X^{-t}, t)\|_{L^1}$  less  
2 than  $\delta/3$  for all  $t \in [0, T]$ . This concludes the proof of (2.30).

We now show (2.31). The proof follows Ref. 53. Let  $\omega_\delta(x)$  be the characteristic function of the ball of radius  $\delta$ ,  $\omega_\delta(x) = \chi_{B(0, \delta)}(x)$ , then

$$\begin{aligned} \|v^\epsilon(\cdot, t) - v(\cdot, t)\|_{L^\infty} &\leq \|\omega_\delta \nabla N\|_{L^1} (\|\rho^\epsilon - \rho\|_{L^\infty}) + \|(1 - \omega_\delta) \nabla N\|_{L^\infty} \|\rho^\epsilon - \rho\|_{L^1} \\ &\leq \|\omega_\delta \nabla N\|_{L^1} \frac{2\|\rho_0\|_{L^\infty}}{k} + \|(1 - \omega_\delta) \nabla N\|_{L^\infty} \|\rho^\epsilon - \rho\|_{L^1}, \end{aligned}$$

3 where we have used the bounds  $\|\rho^\epsilon(\cdot, t)\|_{L^\infty} \leq \frac{1}{k}\|\rho_0\|_{L^\infty}$  and  $\|\rho(\cdot, t)\|_{L^\infty} \leq \frac{1}{k}\|\rho_0\|_{L^\infty}$   
4 which can be directly read from the explicit formulas (2.32) and (2.43). Since  $\nabla N$   
5 is locally integrable the first term can be made small by choosing  $\delta$  small. Then  
6 we let  $\epsilon$  go to zero and we use the fact that  $\rho^\epsilon$  converges to  $\rho$  strongly in  $L^1$  and  
7 uniformly in time. This concludes the proof of (iii).

Finally we show (i). Using formula (2.44) we write:

$$\begin{aligned} \|\rho^\epsilon(\cdot, t) - \rho^\epsilon(\cdot, s)\|_{L^1} &\leq \|\tilde{\rho}_0^\epsilon(X_\epsilon^{-t}, t) - \tilde{\rho}_0^\epsilon(X_\epsilon^{-t}, s)\|_{L^1} \\ &\quad + \|\tilde{\rho}_0^\epsilon(X_\epsilon^{-t}, s) - \tilde{\rho}_0^\epsilon(X_\epsilon^{-s}, s)\|_{L^1} \end{aligned}$$

8 and we conclude using (2.48) and (2.46) for the first term, and (2.47) and (2.40)  
9 for the second term. The proof of Proposition 2.1 is complete.

10 We now prove Theorem 2.3:

**Proof of the Theorem 2.3.** Convergences (iii) and bounds (ii) allow one to pass to the limit in (2.29)–(2.29) and therefore prove the existence of bounded compactly supported solutions. The uniform convergence (2.30) together with (i) give the continuity of  $\rho(t)$  as a function taking values in  $L^1(\mathbb{R}^d)$ . To prove conservation of mass (2.27), choose the test function  $\phi$  in (2.29) to be

$$\phi(x, t) = \chi_{[t_1, t_2]}^\epsilon(t) \chi_{B(0, R)}^\epsilon(x),$$

where  $0 < t_1 < t_2 < T$  and  $R$  is large enough so that  $\text{supp } \rho(\cdot, t) \subset B(0, R)$  for all  $t \in [0, T]$ . The smooth function  $\chi_{[t_1, t_2]}^\epsilon$  is equal to one inside  $[t_1 + \epsilon, t_2 - \epsilon]$ , equal to zero outside of  $[t_1 - \epsilon, t_2 + \epsilon]$ , increasing on  $[t_1 - \epsilon, t_1 + \epsilon]$  and decreasing on

*A. L. Bertozzi, T. Laurent & F. Léger*

$[t_2 - \epsilon, t_2 + \epsilon]$ . The function  $\chi_{B(0,R)}^\epsilon$  is similarly defined. Then letting  $\epsilon$  go to zero in (2.29) and using the fact that

$$t \mapsto \int_{\mathbb{R}^d} \rho(x, t) dt \quad (2.50)$$

is continuous on  $[0, T]$  we obtain that

$$\int_{\mathbb{R}^d} \rho(x, t_1) dt = \int_{\mathbb{R}^d} \rho(x, t_2) dt \quad \text{for all } 0 < t_1 < t_2 < T.$$

1 Using again (2.50) we can let  $t_1 \rightarrow 0$  to obtain the desired result. Equalities (2.28)  
 2 can directly be read from (2.43) together with the fact that for  $t$  fixed, the function  
 3  $f(z) = (1/z - t)^{-1}$  is increasing on  $(-\infty, +\infty)$ .  $\square$

### 4 **2.3. Uniqueness of $L^\infty$ solution**

5 Uniqueness of solutions follows by an energy estimate involving the primitive of  
 6  $\rho$  — or in other words an  $H^{-1}$  inner product for a comparison of two densities. The  
 7 argument requires several steps and involves some estimates with singular integral  
 8 operators. It is the same argument that was used in Ref. 68 for the classical vorticity  
 9 problem and more recently in a number of papers for the aggregation problem with  
 10 and without diffusion, see Refs. 59, 9, 4 and 61.

11 **Theorem 2.4.** *The solution from Theorem 2.3 is unique.*

12 We omit the proof because it has already been done in a number of references  
 13 above. We note that several papers already prove partial existence results combined  
 14 with uniqueness. The main contribution of our paper is to fully develop the sharp  
 15 existence theory for signed data.

### 16 **3. Convergence to Self-Similarity and Estimate** 17 **of the Size of the Support**

In this section we consider solutions of Eq. (1.1) with negative initial data, or equivalently, solutions of

$$\frac{\partial \rho}{\partial t} + \operatorname{div}(\rho v) = 0, \quad v = \nabla N * \rho \quad (3.1)$$

18 with positive initial data (note the change of sign in the formula defining the veloc-  
 19 ity). Recall that for this problem solutions are spreading and exist globally in time,  
 20 see Theorem 2.3. We prove in the theorem below that any positive solution of (3.1)  
 21 with initial data in  $L^\infty$  and with compact support converges in  $L^1$  to a spread-  
 22 ing circular patch. The proof relies on an estimate for all time  $t$  of the size of

*Aggregation via the Newtonian Potential and Aggregation Patches*

1 the support of  $\rho(\cdot, t)$ . For simplicity we normalize everything to mass one and we  
2 consider solutions belonging to  $\mathcal{P}(\mathbb{R}^d)$ , the space of probability measures.

**Theorem 3.1.** *Suppose  $\rho_0 \in \mathcal{P}(\mathbb{R}^d)$  is compactly supported and belongs to  $L^\infty(\mathbb{R}^d)$ . Choose  $r_0$  and  $h_0$  such that*

$$\text{supp } \rho_0 \subset \overline{B(0, r_0)} \quad \text{and} \quad \|\rho_0\|_{L^\infty} \leq h_0. \quad (3.2)$$

Let  $\Phi(x, t)$  be the circular patch of mass 1 with initial height  $h_0$ , that is

$$\Phi(x, t) = \frac{h_0}{1 + h_0 t} \chi_{B(0, R(t))}(x), \quad \text{where } R(t) = R_0(1 + h_0 t)^{1/d} \quad \text{and}$$

$$R_0 = (\omega_d h_0)^{-1/d}.$$

Note that since  $\rho \in \mathcal{P}(\mathbb{R}^d)$  we necessarily have  $R_0 \leq r_0$ . Define

$$E(t) = \frac{E_0}{(1 + h_0 t)^{1/2^{d-1}}} \quad \text{where } E_0 = \frac{r_0^d}{R_0^d} - 1 \geq 0.$$

Let  $\rho(x, t)$  be a solution of the PDE (3.1) with initial data  $\rho_0(x)$ . Then for all  $t \geq 0$  we have:

$$\text{supp } \rho(\cdot, t) \subset \overline{B(0, r(t))} \quad \text{where } r(t) = R(t)(1 + E(t))^{1/d} \quad (3.3)$$

and

$$\|\rho(\cdot, t) - \Phi(\cdot, t)\|_{L^1} \leq 2E(t). \quad (3.4)$$

3 **Remark 3.1.** We prove in Theorem 4.2 in the next section that in the two-  
4 dimensional case, the above  $1/\sqrt{t}$  convergence rate is sharp (this is done by comput-  
5 ing a family of exact elliptical solutions). Moreover, for the special case of radially  
6 symmetric data, one has the sharper rate of convergence  $1/t$  in all dimensions,  
7 because equations reduce to a version of the inviscid Burgers equation (see the last  
8 section of Ref. 11).

**Remark 3.2.** It is enough to prove Theorem 3.1 for the Hölder continuous solution given by Theorem 2.1. Indeed assume that the result of the theorem holds for all compactly supported initial data which are Hölder continuous and assume that we are given non-smooth initial data  $\rho_0 \in L^\infty$  satisfying (3.2). Let  $\eta(x)$  be a smooth probability measure supported in the ball of radius 1, define  $\eta^\epsilon(x) = \epsilon^{-d} \eta(x/\epsilon)$  and  $\rho_0^\epsilon = \rho_0 * \eta^\epsilon$ . Note that  $\rho_0^\epsilon$  is still a probability measure. Now fix a  $\delta > 0$ . For all  $0 < \epsilon < \delta$  the function  $\rho_0^\epsilon$  satisfy

$$\text{supp } \rho_0^\epsilon \subset \overline{B(0, r_0 + \delta)} \quad \text{and} \quad \|\rho_0^\epsilon\|_{L^\infty} \leq h_0. \quad (3.5)$$

Let  $\rho^\epsilon(x, t)$  be the classical solution given by Theorem 2.1 starting with initial data  $\rho_0^\epsilon$ ,  $0 < \epsilon < \delta$ . Since we have assumed the result of the theorem holds for these

*A. L. Bertozzi, T. Laurent & F. Léger*

solutions, from (3.5) we can conclude that

$$\text{supp } \rho^\epsilon(\cdot, t) \subset \overline{B(0, r_\delta(t))} \quad \text{where } r_\delta(t) = R(t)(1 + E_\delta(t))^{1/d}$$

for all  $0 < \epsilon < \delta$ , where  $R(t)$  is defined as in the theorem and  $E_\delta(t)$  is defined by replacing  $r_0$  by  $r_0 + \delta$ . Fix a time  $t$ , since the particle path converges uniformly, by letting  $\epsilon \rightarrow 0$  we obtain

$$\text{supp } \rho(\cdot, t) \subset \overline{B(0, r_\delta(t))} \quad \text{where } r_\delta(t) = R(t)(1 + E_\delta(t))^{1/d}.$$

1 Then by letting  $\delta \rightarrow 0$  we obtain the desired estimate (3.3). As will be pointed out  
 2 at the end of the proof of Theorem 3.1, estimate (3.4) is a direct consequence of  
 3 estimate (3.3).

4 The remainder of this section is devoted to the proof of Theorem 3.1. As noted in  
 5 Remark 3.2 we can assume without loss of generality that  $\rho_0$  is a Hölder continuous,  
 6 compactly supported, probability measure. From now on let us fix such a  $\rho_0$ , let us  
 7 also fix  $r_0$ ,  $h_0$ ,  $\Phi(x, t)$ ,  $R(t)$  and  $E(t)$  to be as defined in the theorem. Finally let  
 8  $\rho(x, t)$  be the Hölder continuous solution starting with initial data  $\rho_0$ .

### 9 **3.1. Change of variables**

We now move into the reference frame of  $\Phi(x, t)$ . We do the change of variable:

$$\tilde{x} = \frac{x}{R(t)} = \frac{x}{R_0(1 + h_0 t)^{1/d}}, \quad \tilde{t} = \ln(1 + h_0 t) \quad (3.6)$$

and we define

$$\tilde{\rho}(\tilde{x}, \tilde{t}) = R_0^d e^{\tilde{t}} \rho \left( R_0 e^{\tilde{t}/d} \tilde{x}, \frac{e^{\tilde{t}} - 1}{h_0} \right). \quad (3.7)$$

One can easily check that  $\tilde{\rho}(\tilde{x}, \tilde{t})$  satisfies

$$\frac{\partial \rho}{\partial t} + \text{div}(\rho v) = 0, \quad v = \omega_d \nabla N * \rho - \frac{x}{d}, \quad (3.8)$$

where we have dropped all the tilde in the above equation for better readability. One can also check that since  $\rho(x, t) \leq \frac{h_0}{1+h_0 t}$  for all  $x \in \mathbb{R}^d$  and  $t \geq 0$ , then

$$\tilde{\rho}(\tilde{x}, \tilde{t}) \leq \frac{1}{\omega_d} \quad \text{for all } \tilde{x} \in \mathbb{R}^d \text{ and } \tilde{t} \geq 0. \quad (3.9)$$

10 Note that the stationary state of (3.8) is the radially symmetric patch of radius  
 11 one, height  $1/\omega_d$  and mass one. Going back to the original variable, this stationary  
 12 state obviously corresponds to  $\Phi(x, t)$ .

1 **3.2. Estimate of the size of the support**

2 In this subsection we drop the tilde for convenience. The following general lemma  
3 is fundamental to our proof:

**Lemma 3.1.** (Frozen in time estimate of the velocity at the boundary) *Suppose  $\mu \in \mathcal{P}(\mathbb{R}^d)$  satisfies*

$$\text{supp } \mu \subset \overline{B(0, r)} \quad \text{and} \quad \|\mu\|_{L^\infty} \leq \frac{1}{\omega_d} \quad (3.10)$$

for some  $r > 0$  (note that since  $\mu \in \mathcal{P}(\mathbb{R}^d)$  we necessarily have  $r \geq 1$ ). Then the velocity field

$$v(x) = \omega_d (\nabla N * \mu)(x) - \frac{x}{d}$$

satisfies

$$v(x) \cdot x \leq -\frac{1}{2^{d-1}d^{d-2}}(r^d - 1) \quad \text{for all } x \in \partial B(0, r). \quad (3.11)$$

**Proof.** Choose  $x$  such that  $|x| = r$ . Using the fact that  $\nabla N * \chi_{B(0, r)}(x) = x/d$  we obtain

$$v(x) = -\omega_d \nabla N * \left[ \frac{1}{\omega_d} \chi_{B(0, r)} - \mu \right] (x).$$

Note that the function between square bracket is positive and has mass  $r^d - 1$  and is supported in  $B(0, r)$ . So

$$\begin{aligned} v(x) \cdot x &= -\omega_d \int_{B(0, r)} \nabla N(x - y) \cdot x \left[ \frac{1}{\omega_d} - \mu \right] (y) dy \\ &\leq -\omega_d \left( \inf_{y \in B(0, r)} \nabla N(x - y) \cdot x \right) (r^d - 1). \end{aligned}$$

Note that since  $|x| \geq |y|$  we have that

$$|x - y|^2 = |x|^2 - 2x \cdot y + |y|^2 \leq 2|x|^2 - 2x \cdot y = 2(x - y) \cdot x.$$

Therefore

$$\nabla N(x - y) \cdot x = \frac{(x - y) \cdot x}{d\omega_d \|x - y\|^d} \geq \frac{1}{2d\omega_d \|x - y\|^{d-2}} \geq \frac{1}{2d\omega_d (2r)^{d-2}} \quad (3.12)$$

4 which conclude the proof.  $\square$

Define

$$\Omega_t = \text{supp } \rho(\cdot, t) \quad \text{and} \quad L(t) = \sup_{x \in \Omega_t} |x|. \quad (3.13)$$

*A. L. Bertozzi, T. Laurent & F. Léger*

Also let  $X(\alpha, t)$  be the (rescaled) particle paths, i.e.

$$\frac{d}{dt}X(\alpha, t) = v(X(\alpha, t), t), \quad X(\alpha, 0) = \alpha, \quad (3.14)$$

where

$$v(x, t) = \omega_d(\nabla N * \rho(t))(x) - \frac{x}{d}. \quad (3.15)$$

Fix a time  $t \geq 0$  and choose  $\alpha$  so that  $X(\alpha, t) \in \Omega_t$  and  $|X(\alpha, t)| = L(t)$ . Then, heuristically,

$$\begin{aligned} \frac{d}{dt} \frac{1}{2} L(t)^2 &= \frac{d}{dt} \frac{1}{2} |X(\alpha, t)|^2 = \dot{X}(\alpha, t) \cdot X(\alpha, t) \\ &= v(X(\alpha, t), t) \cdot X(\alpha, t) \leq -\frac{1}{2^{d-1}d L(t)^{d-2}} (L(t)^d - 1), \end{aligned} \quad (3.16)$$

where we have used Lemma 3.1 to obtain the last inequality. Multiplying both sides by  $dL(t)^{d-2}$  leads to

$$\frac{d}{dt} L(t)^d \leq -\frac{1}{2^{d-1}} (L(t)^d - 1)$$

1 which provide us with an estimate on the size of the support. In order to make the  
2 above heuristic argument into a rigorous one, we will need the following amount of  
3 regularity on the particle paths:

- 4 (i) The particle paths  $X(\alpha, t)$  are differentiable in time and satisfy (3.14) pointwise.  
5 (ii)  $|v(x, t)|$  is bounded on any compact set of  $\mathbb{R}^d \times [0, +\infty)$ .

6 An elementary argument shows that condition (ii) guarantees that  $L(t)$  is contin-  
7 uous. Since we have chosen  $\rho_0$  to be Hölder continuous, Sec. 2 guarantees that  
8 the particle paths satisfy the regularity conditions (i) and (ii), see for instance  
9 Eq. (2.19). The above argument is in the same spirit as the proof of finite time  
10 blowup for  $L^\infty$  solutions of aggregation equations with less singular kernels by the  
11 first two authors and Carrillo,<sup>10</sup> in which they estimate the size of the support  
12 of the solution and show that it must collapse inside a ball with radius that goes  
13 to zero in finite time. We are now ready to prove rigorously the above heuristic  
14 argument.

**Lemma 3.2.** (Estimate of the size of the support) *Let  $\rho(x, t)$  be the solution of the rescaled problem (3.8) with initial data  $\rho_0 \in \mathcal{P}(\mathbb{R}^d) \cap C^\gamma(\mathbb{R}^d)$ . Assume that*

$$\text{supp } \rho_0 \subset \overline{B(0, r_0)} \quad \text{and} \quad \|\rho(\cdot, t)\|_{L^\infty} \leq \frac{1}{\omega_d} \quad \forall t \geq 0. \quad (3.17)$$

Note that since  $\rho_0 \in \mathcal{P}(\mathbb{R}^d)$  we necessarily have  $r_0 \geq 1$ . Let  $r(t)$  be such that:

$$\frac{d}{dt} r(t)^d = -\frac{1}{2^{d-1}} (r(t)^d - 1), \quad r(0) = r_0. \quad (3.18)$$

15 Then  $\rho(\cdot, t)$  is supported in  $\overline{B(0, r(t))}$  for all  $t \geq 0$ .

*Aggregation via the Newtonian Potential and Aggregation Patches*

**Proof.** Let  $\Omega_t$ ,  $L(t)$  and  $X(\alpha, t)$  be defined by (3.13) and (3.14). As  $\rho_0 \in C^\gamma$ , (i) and (ii) holds and the function  $L(t)$  is continuous. For  $0 < \epsilon < 1$  define the function  $r_\epsilon(t)$  by:

$$\frac{d}{dt}r_\epsilon(t)^d = -\frac{(1-\epsilon)}{2^{d-1}}(r_\epsilon(t)^d - 1), \quad r_\epsilon(0) = r_0. \quad (3.19)$$

Note that  $r_\epsilon(t)$  decays toward 1 slower than the function  $r(t)$  defined by (3.18). Note also that for  $t$  fixed,  $\lim_{\epsilon \rightarrow 0} r_\epsilon(t) = r(t)$ . We will prove that  $\Omega_t \subset \overline{B(0, r_\epsilon(t))}$  for all  $t \geq 0$  and all  $0 < \epsilon < 1$ . Taking the limit as  $\epsilon$  goes to 0 will then give the desired result. Fix  $0 < \epsilon < 1$ . We do the proof by contradiction. Assume that there exists a time  $t_1 > 0$  such that  $L(t_1) > r_\epsilon(t_1)$  and define

$$\mathcal{T} = \{t \geq 0 \mid L(t) \geq r_\epsilon(t)\}.$$

Clearly  $t_1 \in \mathcal{T}$  so  $\mathcal{T}$  is not empty. Since both  $L(t)$  and  $r_\epsilon(t)$  are continuous, the set  $\mathcal{T}$  is closed. Therefore there exists a time  $t^* \in \mathcal{T}$  such that

$$t^* = \min \mathcal{T}. \quad (3.20)$$

Using again the continuity of  $L(t)$  and  $r_\epsilon(t)$  we have that  $L(t^*) = r_\epsilon(t^*)$ . Choose  $x \in \Omega_{t^*}$  such that  $|x| = L(t^*) = r_\epsilon(t^*)$  and let  $\alpha = X^{-t^*}(x)$ . For this particle  $\alpha$ , we have:

$$|X(\alpha, t^*)| = r_\epsilon(t^*) \quad \text{and} \quad |X(\alpha, t)| < r_\epsilon(t) \quad \text{for all } t < t^*. \quad (3.21)$$

Let us prove the second statement. Since  $X(\alpha, t^*) \in \Omega_{t^*}$ , then  $X(\alpha, t) \in \Omega_t$  for all  $t \geq 0$ . This simply comes from the fact that  $\Omega_t = X^t(\Omega_0)$ . Since  $X(\alpha, t) \in \Omega_t$  for all  $t \geq 0$  we have that  $L(t) \geq |X(\alpha, t)|$  for all  $t \geq 0$  and therefore it is not possible for  $|X(\alpha, t)|$  to become greater than or equal to  $r_\epsilon(t)$  before time  $t^*$  otherwise  $t^*$  would not be the minimum of  $\mathcal{T}$ . To summarize,  $t \mapsto X(\alpha, t)$  is the particle path which first reaches the boundary of the ball of radius  $r_\epsilon(t)$  and this occurs at time  $t^*$ . Using (3.21) we then get that for  $h > 0$ ,

$$\begin{aligned} \frac{X(\alpha, t^*) - X(\alpha, t^* - h)}{h} \cdot X(\alpha, t^*) &\geq \frac{|X(\alpha, t^*)|^2 - |X(\alpha, t^* - h)||X(\alpha, t^*)|}{h} \\ &\geq \frac{r_\epsilon(t^*) - r_\epsilon(t^* - h)}{h} r_\epsilon(t^*). \end{aligned}$$

Since the particle path are differentiable with respect to time and satisfies (3.14) pointwise (see Sec. 2), letting  $h \rightarrow 0$  and then using (3.19) we obtain:

$$\begin{aligned} v(X(\alpha, t^*), t^*) \cdot X(\alpha, t^*) &\geq r'_\epsilon(t^*) r_\epsilon(t^*) \\ &= -\frac{1-\epsilon}{2^{d-1} d r_\epsilon(t^*)^{d-2}} (r_\epsilon(t^*)^d - 1) \\ &> -\frac{1}{2^{d-1} d r_\epsilon(t^*)^{d-2}} (r_\epsilon(t^*)^d - 1). \end{aligned}$$

1 This contradicts Lemma 3.1. □

*A. L. Bertozzi, T. Laurent & F. Léger*

### 1 3.3. Proof of the theorem

Let us now put back the tilde. Solving the ODE (3.18) we find that

$$\tilde{r}(\tilde{t})^d - 1 = (\tilde{r}_0^d - 1)e^{-\frac{\tilde{t}}{2^{d-1}}}$$

and going back to the original variable this translates into:

$$\frac{r(t)^d}{R(t)^d} - 1 = \left( \frac{r_0^d}{R_0^d} - 1 \right) \frac{1}{(1 + h_0 t)^{\frac{1}{2^{d-1}}}}.$$

2 The right-hand side of the above equation is  $E(t)$ . We therefore have proven  
3 estimate (3.3).

We now show that the  $L^1$ -estimate (3.4) is a direct consequence of the estimate of the size of the support (3.3). Clearly  $\rho(\cdot, t)$  is smaller than  $\Phi(\cdot, t)$  inside the ball of radius  $R(t)$  and greater outside. Using moreover the fact that both  $\Phi(\cdot, t)$  and  $\rho(\cdot, t)$  have mass 1 we easily find that

$$\begin{aligned} \|\Phi(\cdot, t) - \rho(\cdot, t)\|_{L^1} &= \int_{|x| \leq R(t)} \Phi(x, t) - \rho(x, t) dx + \int_{|x| > R(t)} \rho(x, t) dx \\ &= 1 - \int_{|x| \leq R(t)} \rho(x, t) dx + \int_{|x| > R(t)} \rho(x, t) dx \\ &= 2 \int_{|x| > R(t)} \rho(x, t) dx. \end{aligned}$$

Using (3.3) and the fact that

$$\|\rho(\cdot, t)\|_{L^\infty} \leq \frac{h_0}{1 + h_0 t} = \frac{1}{\omega_d R(t)^d}$$

it is then easy to see that the mass of  $\rho(\cdot, t)$  outside the ball of radius  $R(t)$  is bounded by  $E(t)$ :

$$\begin{aligned} \int_{|x| > R(t)} \rho(x, t) dx &= \int_{r(t) \geq |x| > R(t)} \rho(x, t) dx \leq \int_{r(t) \geq |x| > R(t)} \frac{1}{\omega_d R(t)^d} dx \\ &= \frac{r(t)^d}{R(t)^d} - 1 = E(t). \end{aligned}$$

4 This concludes the proof of estimate (3.4) as well as the proof of Theorem 3.1.

## 5 4. Exact Solutions and Numerical Examples

6 In this section we construct a family of exact solutions in 2D and present numerical  
7 results in 2D and 3D. The family of exact solutions consists of elliptical aggre-  
8 gation patches and is derived directly from the well-known theory for elliptical  
9 vortex patches. For the collapsing problem, these exact solutions provide us with  
10 an example of aggregation patches collapsing to a nontrivial singular measure (The-  
11 orem 4.1). For the spreading problem, elliptical patches allow us to prove that the

1 convergence rate toward selfsimilarity derived in the previous section is sharp in  
2 2D (Theorem 4.2).

3 We then present numerical results of more general aggregation patches in 2D and  
4 3D, the latter case having axial symmetry for ease of computation. The numerics  
5 can be made very accurate by reformulating the problem as a dynamics equation  
6 for the boundary of the patch alone and by doing a time change of variable so that  
7 the blow up occurs as  $s \rightarrow \infty$ . The numerics illustrate the phenomena of collapse  
8 toward a complex skeleton in the collapsing case (Figs. 2, 5 and 6), as well as the  
9 phenomena of “pinching” of the boundary in the spreading case (Figs. 3 and 7).

#### 10 4.1. Elliptical patches in 2D

11 There is a class of well-known vortex patch solutions to the 2D Euler equations con-  
12 sisting of rotating ellipses. These are known as the Kirchoff ellipses (see Kirchoff,<sup>43</sup>  
13 Chap. 20, Lamb<sup>44</sup> p. 232, and Majda and Bertozzi<sup>53</sup> Chap. 8). For these exact solu-  
14 tions, the velocity field associated with constant vorticity ellipse can be computed  
15 analytically along with the dynamics of the patch. By computing the orthogonal  
16 velocity field we immediately obtain an analytical formula for  $\nabla N * \rho$  for  $\rho$  the  
17 characteristic function of an ellipse.

Given a density  $\rho = \rho_0 \chi_{E(a,b)}$  where

$$E(a, b) = \left\{ x \mid \frac{x_1^2}{a^2} + \frac{x_2^2}{b^2} < 1 \right\},$$

the velocity field generated is precisely

$$v(x) = \begin{cases} -\frac{\rho_0}{a+b}(bx_1, ax_2), & x \in E(a, b), \\ -\frac{\rho_0}{\sqrt{a^2 + \lambda^2} + \sqrt{b^2 + \lambda^2}}((\sqrt{b^2 + \lambda})x_1, (\sqrt{a^2 + \lambda})x_2), & x \notin E(a, b), \end{cases}$$

where  $\lambda(x)$  satisfies

$$\frac{x_1^2}{a^2 + \lambda} + \frac{x_2^2}{b^2 + \lambda} < 1.$$

We note that inside the ellipse (and on the boundary) the flow field is linear. Linear flow fields map ellipses to ellipses and thus the elliptical shape is preserved under the flow, although the major and minor axes will change. For the vorticity problem this results in exact solutions that rotate at a constant angular velocity. For the aggregation problem it results in ellipses that collapse or spread and for which we obtain explicit ODEs for the major and minor axes:

$$\dot{a} = \dot{b} = -\frac{\rho(t)ab}{a+b}, \quad \rho(t) = \frac{\rho_0}{1 - \rho_0 t}.$$

18 So the quantity  $a - b$  is conserved and we obtain the following theorem about  
19 collapsing elliptical patches:

**Theorem 4.1.** (Collapse of positive Elliptical Patches toward singular measures)  
Let  $\rho(x, t)$  be the solution of (1.1) starting with initial  $\rho_0 \in \mathcal{P}(\mathbb{R}^2)$  being the uniform

A. L. Bertozzi, T. Laurent & F. Léger

distribution on the ellipse  $E_0 = \{x \mid \frac{x_1}{a_0} + \frac{x_2}{b_0} < 1\}$ ,  $0 < b_0 \leq a_0$ . As  $t \rightarrow T^*$ , where  $T^* = (\pi a_0 b_0)^{-1}$  is the maximal time of existence, the solution converges weakly- $*$  as a measure toward the weighted distribution on the segment

$$\mu(x_1) = \frac{2\sqrt{x_0^2 - x_1^2}}{\pi x_0^2} \delta_{I(x_0)},$$

1 where  $\delta_{I(x_0)}$  is the uniform measure on the interval  $x_1 \in [-x_0, x_0]$  and  $x_0 \equiv a_0 - b_0$   
 2 is the semi-major axis of the limiting ellipse, i.e.  $a_0 \rightarrow x_0$  as  $t \rightarrow T^*$ .

The proof of this theorem is straightforward. From the exact solution we have  $b_0 \rightarrow 0$  and  $a_0 \rightarrow x_0$  as  $t \rightarrow T^*$ . Clearly the limiting measure is a weighted measure on the interval from  $[-x_0, x_0]$ . So what is left is to compute the weight. We note that prior to the collapse, the  $x_2$  coordinate of the ellipse boundary satisfies  $x_2(x_1) = \pm b(t) \sqrt{1 - \frac{x_1^2}{a(t)^2}}$ . The weight function is the limit of the portion of mass that collapses onto the point  $x$  on the interval and this is simply  $\lim_{t \rightarrow T^*} 2\rho(t)x_2(x_1) = \lim_{t \rightarrow T^*} 2\rho(t)b(t) \sqrt{1 - \frac{x_1^2}{a(t)^2}}$ . Using the fact that  $\rho(t) = 1/\pi(a(t)b(t))$  we can substitute this into the limit to obtain the weight

$$\lim_{t \rightarrow T^*} \frac{2}{\pi a(t)} \sqrt{1 - \frac{x_1^2}{a(t)^2}} = \frac{2}{\pi x_0} \sqrt{1 - \frac{x_1^2}{x_0^2}}.$$

Note also that one can directly integrate the ODEs for  $a$  and  $b$ , in particular using the dynamics of the product  $ab$ . It is easy to solve for both variables:

$$a(t) = \frac{(a_0 - b_0) + \sqrt{(a_0 - b_0)^2 + 4a_0b_0(1 - \rho_0 t)}}{2}, \quad (4.1)$$

$$b(t) = \frac{-(a_0 - b_0) + \sqrt{(a_0 - b_0)^2 + 4a_0b_0(1 - \rho_0 t)}}{2}. \quad (4.2)$$

3 Using the above exact solution we can now prove that the convergence rate in  
 4 Theorem 3.1 is sharp in two space dimensions.

**Theorem 4.2.** ( $L^1$  convergence rate of negative Elliptical Patches toward selfsimilarity) Let  $\rho(\cdot, t) = \rho(t)\chi_{E(t)}$  be an elliptical patch solution with negative initial data, with  $\rho(t) = \frac{-1}{1+t}$  and  $E(t) = \{x \mid \frac{x_1}{a(t)} + \frac{x_2}{b(t)} < 1\}$ ,  $a(t)$  and  $b(t)$  being the semi-major axis and semi-minor axis of the ellipse  $E(t)$ . Let  $\Phi(x, t) = -(1+t)^{-1}\chi_{B(0, R(t))}$ , where  $R(t) = \sqrt{(1+t)/\pi}$ . Then

$$\begin{aligned} \|\rho(\cdot, t) - \Phi(\cdot, t)\|_{L^1} &= 4a_0b_0 \left[ 2 \arctan \left( \sqrt{\frac{a(t)}{b(t)}} \right) - \frac{\pi}{2} \right] \\ &= \frac{2\sqrt{a_0b_0}(a_0 - b_0)}{\sqrt{1+t}} + O\left(\frac{1}{1+t}\right). \end{aligned}$$

1 This is proved by a direct computation involving the above solutions.

2 **Remark 4.1.** There are analogous ellipsoid solutions in 3D of the form  $\frac{x_1^2}{a^2} + \frac{x_2^2}{b^2} +$   
 3  $\frac{x_3^2}{c^2} = 1$  where  $a, b, c$  satisfy ODEs involving elliptic integrals. This is due to a known  
 4 formula for the gradient of the Newtonian potential convolved with an ellipsoid in  
 5 3D, a classical result that comes from the Newtonian physics of self-gravitating  
 6 bodies. Computing these solutions is beyond the scope of this paper however we  
 7 expect them to have similar qualitative behavior to the 2D elliptical solutions.

8 **4.2. Collapsing aggregation patches, time rescaling,**  
 9 **and contour dynamics**

We now consider the case of a general collapsing patch. Without loss of generality  
 we take the initial density to be one and for simplicity of notation we assume that  
 the patch is simply connected with smooth boundary. Formally we compute a time  
 and amplitude rescaling for the equations of motion by defining

$$s = \ln \frac{1}{1-t}, \quad \tilde{\rho} = (1-t)\rho, \quad \tilde{v} = (1-t)v. \quad (4.3)$$

Substituting these changes into the original equation  $\rho_t + \nabla \cdot (\rho v) = 0$ , after some  
 calculus we formally obtain

$$\tilde{\rho}_s + \tilde{v} \cdot \nabla \tilde{\rho} = 0; \quad v = -\nabla N * \tilde{\rho}, \quad (4.4)$$

where in the above calculation we use the fact that  $-\nabla \cdot v = \rho$  and that term  
 cancels the additional term obtained from differentiating  $\rho = \frac{\tilde{\rho}}{1-t}$  with respect to  
 time. We also use the fact that  $\rho(x, t)$  is  $\frac{1}{1-t}$  times the characteristic function of  
 the domain  $\Omega_s = \Omega(t)$  for  $s = -\ln(1-t)$ . The upshot is that we have a new time  
 variable  $s$  and the patch singularity occurs as  $s \rightarrow \infty$ . Moreover, the patch remains  
 a characteristic function (the rescaled density does not grow or shrink) and it is  
 simply transported along characteristics of the rescaled velocity. The next step is  
 to rewrite (4.4) as a dynamic equation for the boundary of the patch. This can be  
 done following the classical contour dynamics formulation for vortex patches.<sup>53,69</sup>  
 Note that, integrating by parts, we can express  $\tilde{v}$  as a boundary integral:

$$\tilde{v}(x, s) = -\nabla N * \chi_{\Omega_s}(x) = \int_{\partial\Omega_s} N(x-y)n(y)d\sigma(y). \quad (4.5)$$

For the two-dimensional problem, we drop the tilde in Eq. (4.5) and parametrize  
 the curve  $\partial\Omega_s$  by  $z(\alpha, s)$ ,  $\alpha \in [0, 2\pi]$ , a Lagrangian variable. We obtain

$$v(x, s) = \int_0^{2\pi} N(x-z(\alpha, s)) \left[ \frac{\partial z}{\partial \alpha}(\alpha, s) \right]^\perp d\alpha$$

and since  $\Omega_s$  moves according to  $v$  we get

$$\frac{\partial z}{\partial s}(\alpha, s) = \frac{1}{2\pi} \int_0^{2\pi} \ln |z(\alpha, s) - z(\alpha', s)| \left[ \frac{\partial z}{\partial \alpha}(\alpha', s) \right]^\perp d\alpha'. \quad (4.6)$$

A. L. Bertozzi, T. Laurent & F. Léger

We discretize  $\partial\Omega_s$  with  $N$  particles  $X_1(s), \dots, X_N(s)$ . We use a midpoint rule to approximate the velocity field:

$$\begin{aligned} v(x, s) &= \frac{1}{2\pi} \int_{\partial\Omega_s} \ln|x-y|n(y)d\sigma(y) \\ &\approx \frac{1}{2\pi} \sum_{i=1}^N \ln \left| x - \frac{X_{i+1} + X_i}{2} \right| \frac{(X_{i+1} - X_i)^\perp}{|X_{i+1} - X_i|} |X_{i+1} - X_i| \\ &= \frac{1}{2\pi} \sum_{i=1}^N \ln \left| x - \frac{X_{i+1} + X_i}{2} \right| (X_{i+1} - X_i)^\perp \end{aligned} \quad (4.7)$$

1 with the convention that  $X_{N+1} := X_1$ . See Fig. 1(a) for a graphic explanation of  
 2 this formula.




Fig. 1. 2D collapsing aggregation patches. (a) schematic of quadrature. (b) and (c) show an exact elliptical solution with respectively 50 and 150 gridpoints compared with the analytic solution at times  $t = 0, 0.62, 0.86, 0.95$ .

## Aggregation via the Newtonian Potential and Aggregation Patches

This leads to the system of ODEs

$$\dot{X}_k = \frac{1}{2\pi} \sum_{i=1}^N \ln \left| X_k - \frac{X_{i+1} + X_i}{2} \right| (X_{i+1} - X_i)^\perp, \quad 1 \leq k \leq N \quad (4.8)$$

1 that we solve using an adaptive Runge–Kutta–Fehlberg algorithm order 4/5  
 2 (“ODE45” in MATLAB). These schemes use an adaptive timestep chosen by setting  
 3 the local time truncation error. In our simulations we take that to be smaller than  
 4  $10^{-5}$ . Figures 1(b) and 1(c) show the results of the numerics for the exact elliptical  
 5 solution starting with the initial ellipse  $x_1^2/5^2 + x_2^2/2^2 = 1$ . We compare results with  
 6 50 versus 150 grid points at times  $t = 0, 0.62, 0.86, 0.95$ . The results compare well  
 7 with the exact solution shown as a solid line. More information about the accuracy  
 8 of the numerical scheme can be found in the Appendix. In Fig. 2 we show that  
 9 aggregation patches of more general shapes typically collapse toward more complex  
 10 skeletons. Each has initial data that is the characteristic function of some region.  
 11 The singularity time is  $t = 1$ . At the collapse time, the patch becomes a singular  
 12 measure supported on a domain of Lebesgue measure zero which consists of the




Fig. 2. 2D collapsing aggregation patches. All examples show an initial patch that is the characteristic function of some region. At  $t = 1$  the solution converges to a set of measure zero. (a)–(c) show snapshots of the boundary evolution of such a patch. In this example, as  $t \rightarrow 1$ , the patch converges toward a singular measure supported on a three-branched star. Other high resolution simulations with various shapes are shown in (d)–(f). In each image, the dashed line is the boundary of the patch at time  $t = 0$  and the solid line is the boundary at time  $t = 0.99995$ . The number of grid points used is displayed in each subfigure.

*A. L. Bertozzi, T. Laurent & F. Léger*

1 union of intersecting curves. The patch skeleton shown is actually the solution of  
 2 true patch at time  $t = 0.99995$ ; the boundary has almost collapsed and the interior  
 3 is not visible. The annulus example in Fig. 2(f) retains its empty core at the collapse  
 4 time.

### 5 **4.3. Spreading case in two dimensions**

We now consider the case of a spreading patch solution, corresponding to a negative  
 solution of (1.1) or a positive solution of (3.1). Without loss of generality we assume  
 the initial density is the characteristic function of some region. We make a similar  
 change of variables to those in the previous subsection with some modifications:

$$s = \ln(1+t), \quad \tilde{\rho} = (1+t)\rho, \quad \tilde{v} = (1+t)v - \tilde{x}/d, \quad \tilde{x} = x/(1+t)^{1/d}. \quad (4.9)$$

Note the additional rescaling of space along the lines of the similarity variables  
 described in Sec. 3. Also note that  $\tilde{v}$  is divergence-free inside the patch, due to  
 the additional  $x/d$  term subtracted. Substituting these changes into the original  
 equation  $\rho_t + \nabla \cdot (\rho v) = 0$ , after some calculus we formally obtain

$$\tilde{\rho}_s + \tilde{v} \cdot \tilde{\nabla} \tilde{\rho} = 0; \quad \tilde{v} = \tilde{\nabla} \tilde{N} * \tilde{\rho} - \tilde{x}/d. \quad (4.10)$$

6 The patch remains a characteristic function (the rescaled density does not grow or  
 7 shrink) and it is simply transported along characteristics of the modified velocity  
 8 field. We write the equation for the boundary of the patch following the contour  
 9 dynamics formalism described above and compute the solution for several exam-  
 10 ples in 2D. The results are shown in Fig. 3 in the rescaled variables. Note that  
 11 initial regions of high curvature produce defects in the long time limit in which the  
 12 boundary folds on itself, despite the  $L^1$  convergence to the exact circular solution.  
 13 Note that we cannot have convergence in  $L^\infty$  because the solution and its limit are  
 14 both piecewise constant. The numerics suggest that one may not have pointwise  
 15 convergence of the boundary due to the defects.

16 Figure 4 shows a computational example involving interacting particles (point  
 17 masses) for the spreading problem. Note that they also self-organize in the long  
 18 time limit to the exact circular patch solution.

### 19 **4.4. 3D numerics — axisymmetric patches**

Here we compute some axisymmetric examples in 3D. We construct the boundary  
 of the patch as a surface of revolution. Let

$$(x(\alpha), y(\alpha)) \in ]0, 1[$$

be a curve in the  $x - y$  half-plane for  $x > 0$ . We rotate this curve in three dimen-  
 sions, about the  $y$ -axis, to obtain a surface of revolution. The surface can thus be  
 parametrized by

$$\phi(\alpha, \theta) = (x(\alpha) \cos(\theta), y(\alpha), x(\alpha) \sin(\theta)), \quad \alpha \in ]0, 1[, \theta \in ]0, 2\pi[.$$

*Aggregation via the Newtonian Potential and Aggregation Patches*




Fig. 3. Four examples of contour dynamics for the rescaled spreading problem (4.10) in two dimensions. (a)–(c) show snapshots of the boundary evolution of a patch. At time  $s = 10$  the patch has reached steady state. (d)–(f) show various shapes: the initial boundary is shown as a dashed line and the long time limit ( $s = 10$ ) is shown as a solid line.




Fig. 4. 2D spreading dynamics for initial data a sum of 2000 particles. The long time limit is shown on the right.

A. L. Bertozzi, T. Laurent & F. Léger

For the collapsing problem, the velocity at a point  $p$  is

$$\mathbf{v}(p) = \int_{\alpha} \int_{\theta=0}^{2\pi} -\frac{1}{4\pi\|p - \phi(\alpha, \theta)\|} \frac{\partial \phi}{\partial \theta} \times \frac{\partial \phi}{\partial \alpha}(\alpha, \theta) d\theta d\alpha.$$

Hence for a point  $p = (X, Y, 0)$  in the  $(Oxy)$  plane we have

$$\begin{aligned} & \vec{v}(X, Y, 0) \\ &= \int_{\alpha} \int_{\theta=0}^{2\pi} \frac{-x(\alpha) \begin{pmatrix} y'(\alpha) \cos(\theta) \\ -x'(\alpha) \\ y'(\alpha) \sin(\theta) \end{pmatrix}}{4\pi\sqrt{(x(\alpha) \cos(\theta) - X)^2 + (x(\alpha) \sin(\theta))^2 + (y(\alpha) - Y)^2}} d\theta d\alpha. \end{aligned}$$

By symmetry  $\mathbf{v}(X, Y, 0) \cdot \mathbf{e}_z = 0$  so we define  $v_X$  and  $v_Y$  by

$$\mathbf{v}(X, Y, 0) = v_X(X, Y)\mathbf{e}_x + v_Y(X, Y)\mathbf{e}_y.$$

After some integration we find

$$\begin{aligned} v_X(X, Y) &= \int_{\alpha} \frac{1}{\pi} \frac{x(\alpha)y'(\alpha)}{A\sqrt{A+B}} \left[ (A+B)\mathcal{E}\left(\frac{2A}{A+B}\right) - B\mathcal{F}\left(\frac{2A}{A+B}\right) \right] d\alpha, \\ v_Y(X, Y) &= \int_{\alpha} \frac{1}{\pi} \frac{x(\alpha)x'(\alpha)}{\sqrt{A+B}} \mathcal{F}\left(\frac{2A}{A+B}\right) d\alpha \end{aligned}$$

with

$$A(\alpha) = 2x(\alpha)X, \quad B(\alpha) = x^2(\alpha) + X^2 + (y(\alpha) - Y)^2$$

and  $\mathcal{E}, \mathcal{F}$  are the complete elliptic integral of the first and second kinds

$$\mathcal{E}(m) = \int_0^{\frac{\pi}{2}} \sqrt{1 - m \sin^2(x)} dx, \quad \mathcal{F}(m) = \int_0^{\frac{\pi}{2}} \frac{dx}{\sqrt{1 - m \sin^2(x)}}, \quad m \in [0, 1].$$

If the surface has a toroidal topology, then the section in the  $(Oxy)$  half-plane is a closed curve  $\Gamma$ . As in the 2D case, we discretize  $\Gamma$  with  $N$  particles  $(x_1(s), y_1(s)), \dots, (x_N(s), y_N(s))$  and use a midpoint rule to approximate the velocity field:

$$\begin{aligned} x_k &= \frac{1}{\pi} \sum_{i=1}^N \frac{x_{i+1} + x_i}{2} (y_{i+1} - y_i) \frac{1}{A_i \sqrt{A_i + B_i}} \\ &\quad \times \left[ (A_i + B_i)\mathcal{E}\left(\frac{2A_i}{A_i + B_i}\right) - B_i\mathcal{F}\left(\frac{2A_i}{A_i + B_i}\right) \right], \\ y_k &= \frac{1}{\pi} \sum_{i=1}^N \frac{x_{i+1} + x_i}{2} (x_{i+1} - x_i) \frac{1}{\sqrt{A_i + B_i}} \mathcal{F}\left(\frac{2A_i}{A_i + B_i}\right), \end{aligned}$$

*Aggregation via the Newtonian Potential and Aggregation Patches*

where  $x_{N+1} := x_1, y_{N+1} := y_1$  and

$$A_i = 2x_k \frac{x_{i+1} + x_i}{2},$$

$$B_i = \left( \frac{x_{i+1} + x_i}{2} \right)^2 + x_k^2 + \left( \frac{y_{i+1} + y_i}{2} - y_k \right)^2.$$

1 As in 2D we present some numerical results. In Figs. 5 and 7 we show a section  
 2 of the surface rather than the surface itself for readability. We also show a complete  
 3 3D surface in Fig. 6. We observe similar behavior in 2D and in 3D: in the collapsing  
 4 problem the patches converge to a skeleton of codimension 1, and in the expanding  
 5 problem the patches converge to a ball (in rescaled coordinates). Note that in the  
 6 expanding case, since the figures display the section of the surface in the half plane,  
 7 the limit is a half circle. Note also that when there is a high curvature the long time  
 8 limit presents the same defaults on the boundary as in 2D.




Fig. 5. Three examples of 3D collapsing patches showing the section of the surface in the  $x - y$  half-plane. Case (b) is also shown below in Fig. 6 as a surface of revolution. The dashed line is the initial data and the solid line shows the boundary at  $t = 0.99995$ .




Fig. 6. 3D visualization of Fig. 5(b), a toroidal patch with a square section.

A. L. Bertozzi, T. Laurent & F. Léger




Fig. 7. 3D rescaled expanding patches. These figures show the section of the surface in the  $x$ - $y$  half-plane. The dashed line is the initial data and the long time limit is shown as a solid line.

## 1 Appendix. Proof of Lemma 2.7

**Lemma A.1.** For all  $t \in [0, T]$ , and nonnegative  $f \in L^1(\mathbb{R}^d)$ ,

$$\int_{\mathbb{R}^d} f(X^{-t}(x)) dx \leq K \int_{\mathbb{R}^d} f(\alpha) d\alpha. \quad (\text{A.1})$$

**Proof.** Recall that  $X_\epsilon^{-t}(x)$  converge uniformly toward  $X^{-t}(x)$  on compact subset of  $\mathbb{R}^d \times (0, T)$  and that

$$k \leq \det(\nabla_\alpha X_\epsilon^t(\alpha)) \leq K \quad (\text{A.2})$$

for all  $\epsilon > 0$ ,  $t \in [0, T]$  and  $\alpha \in \mathbb{R}^d$ . First we are going to show that

$$|X^t(E)| \leq K|E| \quad (\text{A.3})$$

2 for all bounded measurable set  $E \subset \mathbb{R}^d$ . Here  $|E|$  denote the Lebesgue measure of  
3 the set  $E$ .

Assume first that  $E = B_{x_0, r}$  is a ball of radius  $r < \infty$  centered at  $x_0 \in \mathbb{R}^d$ . Given any  $\lambda > 1$  there exists an  $\epsilon > 0$  such that

$$|X^t(B_{x_0, r})| \leq |X_\epsilon^t(B_{x_0, \lambda r})| \leq K|B_{x_0, \lambda r}| \leq \lambda^d K|B_{x_0, r}|.$$

4 To see this, note that since  $X_\epsilon^{-t}(x)$  converge uniformly toward  $X^{-t}(x)$  on compact  
5 sets, we have that given  $\lambda > 1$ ,  $X^t(B_{x_0, r}) \subset X_\epsilon^t(B_{x_0, \lambda r})$  if  $\epsilon$  is small enough. This  
6 gives the first inequality. The second inequality comes from (A.2) and the third  
7 inequality is just a rescaling of the ball in  $\mathbb{R}^d$ . Letting  $\lambda \rightarrow 1$  we see that (A.3)  
8 holds for any ball of finite radius.

Assume now that  $E = \mathcal{O}$  is a bounded open set and let  $\lambda > 1$ . By a classical covering lemma (see e.g. Lemma 2, p. 15, in Ref. 64) there exists a finite number of pairwise disjoint balls  $B_{x_1, r_1}, \dots, B_{x_k, r_k}$  so that

$$\cup_i B_{x_i, r_i} \subset \mathcal{O} \subset \cup_i B_{x_i, \lambda r_i}.$$

*Aggregation via the Newtonian Potential and Aggregation Patches*

Then, since (A.3) holds for balls, we have:

$$|X^t(\mathcal{O})| \leq \sum_i |X^t(B_{x_i, \lambda r_i})| \leq K \sum_i |B_{x_i, \lambda r_i}| = \lambda^d K \sum_i |B_{x_i, r_i}| \leq \lambda^d K |\mathcal{O}|.$$

1 Letting  $\lambda \rightarrow 1$  we see that (A.3) holds for any bounded open set.

We now show that (A.3) holds for any bounded measurable set  $E$ . Let  $\mathcal{O}_n$  be a sequence of bounded open set such that  $E \subset \mathcal{O}_n$  and  $|\mathcal{O}_n| \rightarrow |E|$ . Thus we have

$$|X^t(E)| \leq |X^t(\mathcal{O}_n)| \leq K |\mathcal{O}_n| \rightarrow K |E|.$$

Finally we approximate  $f$  by simple functions. For  $\lambda > 1$  let:

$$f \leq \sum_{i=-\infty}^{\infty} \lambda^i \chi_{E_i} < \lambda f, \quad \text{where } E_i = \{x : \lambda^{i-1} < f(x) \leq \lambda^i\}.$$

Here  $\chi_{E_i}(x)$  is the characteristic function of the set  $E_i$ . Using this construction we have

$$\begin{aligned} \int_{\mathbb{R}^d} f(X^{-t}(x)) dx &\leq \int_{\mathbb{R}^d} \sum_i \lambda^i \chi_{E_i}(X^{-t}(x)) dx = \sum_i \lambda^i |X^t(E_i)| \\ &\leq K \sum_i \lambda^i |E_i| = K \int_{\mathbb{R}^d} \sum_i \lambda^i \chi_{E_i}(\alpha) d\alpha \leq \lambda K \int_{\mathbb{R}^d} f(\alpha) d\alpha. \end{aligned}$$

□

2 *Numerical convergence in 2D*

3 We perform a mesh refinement study to illustrate that the scheme is  $O(\frac{1}{N})$  where  
 4  $N$  is the number of particles which discretize the boundary of the patch. The  
 5 quadrature would ordinarily be  $O(\frac{1}{N^2})$  however the singularity of the kernel keeps  
 6 it at  $O(\frac{1}{N})$  and this is illustrated below.




Fig. A.1. Logarithm of the error with respect to the number of points, with linear fit  $y = px + q$ , shown are the Newtonian potential and a smooth kernel  $K(x) = e^{-\frac{x^2}{2}}$ .

A. L. Bertozzi, T. Laurent & F. Léger

1 We compute the time evolution of an elliptical patch with  $N = 2^k N$  particles on  
 2 the boundary for  $k = 1, \dots, 12$  from  $s = 0$  to  $s = 3$ . Then we compare at final time  
 3  $s = 3$  each solution  $f_k$  with  $2^k N$  points,  $k = 1, \dots, 11$  to the solution  $f_{12}$  that has  
 4 the most points. More specifically we compute the average distance between the  
 5 points of  $f_k$  and the points of  $f_{12}$  (subsampling). Then we repeated the calculation  
 6 with the nonsingular kernel  $K(x) = e^{-\frac{x^2}{2}}$  instead of the Newtonian kernel  $N$ . Then  
 7 we do a linear fit  $y = px + q$ . Figure A.1 displays both results. The slope of the line  
 8 for the Newtonian kernel is  $-1.06$  whereas the slope of the line for the exponential  
 9 kernel is  $-2.02$ .

## 10 Acknowledgments

11 We thank Betsy Stovall for helpful comments regarding Lemma 2.7. This research  
 12 is supported by NSF grants DMS-0907931 and DMS-1109805.

## 13 References

- 14 1. L. Ambrosio, E. Mainini and S. Serfaty, Gradient flow of the chapman-rubinstein-  
 15 schatzman model for signed vortices, *Ann. Inst. H. Poincaré (C) Non Linear Anal.*  
 16 **28** (2011) 217–246.
- 17 2. L. Ambrosio and S. Serfaty, A gradient flow approach to an evolution problem arising  
 18 in superconductivity, *Comm. Pure Appl. Math.* **61** (2008) 1495–1539.
- 19 3. J. T. Beale, T. Kato and A. Majda, Remarks on the breakdown of smooth solutions  
 20 for the 3-D Euler equations, *Comm. Math. Phys.* **94** (1984) 61–66.
- 21 4. J. Bedrossian, N. Rodriguez and A. L. Bertozzi, Local and global well-posedness  
 22 for aggregation equations and Patlak–Keller–Segel models with degenerate diffusion,  
 23 *Nonlinearity* **24** (2011) 1683–1714.
- 24 5. D. Benedetto, E. Caglioti and M. Pulvirenti, A kinetic equation for granular media,  
 25 *RAIRO Modél. Math. Anal. Numér.* **31** (1997) 615–641.
- 26 6. A. L. Bertozzi and P. Constantin, Global regularity for vortex patches, *Comm. Math.*  
 27 *Phys.* **152** (1993) 19–28.
- 28 7. A. L. Bertozzi and T. Laurent, Finite-time blow-up of solutions of an aggregation  
 29 equation in  $R^n$ , *Comm. Math. Phys.* **274** (2007) 717–735.
- 30 8. A. L. Bertozzi, Heteroclinic orbits and chaotic dynamics in planar fluid flows, *SIAM*  
 31 *J. Math. Anal.* **19** (1988) 1271–1294.
- 32 9. A. L. Bertozzi and J. Brandman, Finite-time blow-up of  $L^\infty$ -weak solutions of an  
 33 aggregation equation, *Commun. Math. Sci.* **8** (2010) 45–65.
- 34 10. A. L. Bertozzi, J. A. Carrillo and T. Laurent, Blow-up in multidimensional aggregation  
 35 equations with mildly singular interaction kernels, *Nonlinearity* **22** (2009) 683–710.
- 36 11. A. L. Bertozzi, J. B. Garnett and T. Laurent, Characterization of radially symmetric  
 37 finite time blowup in multidimensional aggregation equations, 2011.
- 38 12. A. L. Bertozzi, T. Laurent and J. Rosado,  $L^p$  theory for the multidimensional aggre-  
 39 gation equation, *Comm. Pure Appl. Math.* **64** (2011) 45–83.
- 40 13. A. L. Bertozzi and D. Slepčev, Existence and uniqueness of solutions to an aggregation  
 41 equation with degenerate diffusion, *Commun. Pure Appl. Anal.* **9** (2010) 1617–1637.
- 42 14. P. Biler and W. A. Woyczyński, Global and exploding solutions for nonlocal quadratic  
 43 evolution problems, *SIAM J. Appl. Math.* **59**(3) (1999) 845–869.

*Aggregation via the Newtonian Potential and Aggregation Patches*

- 1 15. A. Blanchet, J. Carrillo and N. Masmoudi, Infinite time aggregation for the critical  
2 Patlak-Keller-Segel model in  $\mathbb{R}^2$ , *Comm. Pure Appl. Math.* **61** (2008) 1449–1481.
- 3 16. A. Blanchet, J. Dolbeault and B. Perthame, Two-dimensional Keller-Segel model:  
4 Optimal critical mass and qualitative properties of the solutions, *Electron. J. Differ-*  
5 *ential Equations*, 44 (2006) 32 pp. (electronic).
- 6 17. M. Bodnar and J. J. L. Velazquez, An integro-differential equation arising as a limit  
7 of individual cell-based models, *J. Differential Equations* **222** (2006) 341–380.
- 8 18. S. Boi, V. Capasso and D. Morale, Modeling the aggregative behavior of ants of the  
9 species *polyergus rufescens*, *Nonlinear Anal. Real World Appl.* **1** (2000) 163–176.
- 10 19. M. P. Brenner, P. Constantin, L. P. Kadanoff, A. Schenkel and S. C. Venkataramani,  
11 Diffusion, attraction and collapse, *Nonlinearity* **12** (1999) 1071–1098.
- 12 20. M. Burger, V. Capasso and D. Morale, On an aggregation model with long and short  
13 range interactions, *Nonlinear Anal. Real World Appl.* **8** (2007) 939–958.
- 14 21. M. Burger and M. D. Francesco, Large time behavior of nonlocal aggregation models  
15 with nonlinear diffusion, *Netw. Heterog. Media* **3** (2008) 749–785.
- 16 22. M. Burger, N. Matevosyan and M.-T. Wolfram, A level set based shape optimization  
17 method for an elliptic obstacle problem, *Math. Models Methods Appl. Sci.* **21** (2011)  
18 619–649.
- 19 23. T. F. Buttke, A fast adaptive vortex method for patches of constant vorticity in two  
20 dimensions, *J. Comput. Phys.* **89** (1990) 161–186.
- 21 24. L. Caffarelli and J. Vázquez, Nonlinear porous medium flow with fractional potential  
22 pressure, *Arch. Rational Mech. Anal.* (2011) 1–29, DOI:10.1007/s00205-011-0420-4.
- 23 25. J. A. Carrillo, M. DiFrancesco, A. Figalli, T. Laurent and D. Slepčev, Global-in-time  
24 weak measure solutions and finite-time aggregation for nonlocal interaction equations,  
25 *Duke Math. J.* **156** (2011) 229–271.
- 26 26. J. A. Carrillo, R. J. McCann and C. Villani, Kinetic equilibration rates for granular  
27 media and related equations: Entropy dissipation and mass transportation estimates,  
28 *Rev. Mat. Iber.* **19** (2003) 971–1018.
- 29 27. J. A. Carrillo, R. J. McCann and C. Villani, Contractions in the 2-Wasserstein length  
30 space and thermalization of granular media, *Arch. Rational Mech. Anal.* **179** (2006)  
31 217–263.
- 32 28. J. Carrillo and J. Rosado, Uniqueness of bounded solutions to aggregation equations  
33 by optimal transport methods, in *European Congress of Mathematics* (Eur. Math.  
34 Soc., 2010), pp. 3–16.
- 35 29. J.-Y. Chemin, Persistance de structures géométriques dans les fluides incompressibles  
36 bidimensionnels, *Ann. Sci. École Norm. Sup. (4)* **26** (1993) 517–542.
- 37 30. J. Dolbeault and B. Perthame, Optimal critical mass in the two-dimensional Keller-  
38 Segel model in  $\mathbb{R}^2$ , *C. R. Math. Acad. Sci. Paris* **339** (2004) 611–616.
- 39 31. H. Dong, The aggregation equation with power-law kernels: Ill-posedness, mass con-  
40 centration and similarity solutions, *Comm. Math. Phys.* **304** (2011) 649–664.
- 41 32. H. Dong, On similarity solutions to the multidimensional aggregation equation, *SIAM*  
42 *J. Math. Anal.* **43** (2011) 1995–2008.
- 43 33. D. G. Dritschel, A fast contour dynamics method for many-vortex calculations in  
44 two-dimensional flows, *Phys. Fluids A* **5** (1993) 173–186.
- 45 34. D. G. Dritschel and N. J. Zabusky, A new, but flawed, numerical method for vortex  
46 patch evolution in two dimensions, *J. Comput. Phys.* **93** (1991) 481–484.
- 47 35. Q. Du and P. Zhang, Existence of weak solutions to some vortex density models,  
48 *SIAM J. Math. Anal.* **34**(6) (2003) 1279–1299.
- 49 36. E. Weinan, Dynamics of vortex liquids in Ginzburg–Landau theories with applications  
50 to superconductivity, *Phys. Rev. B* **50** (1994) 1126–1135.

A. L. Bertozzi, T. Laurent & F. Léger

- 1 37. R. C. Fetecau, Y. Huang and T. Kolokolnikov, Swarm dynamics and equilibria for a  
2 nonlocal aggregation model, *Nonlinearity* **24** (2011) 2681–2716.
- 3 38. V. Gazi and K. M. Passino, Stability analysis of swarms, *IEEE Trans. Automat.*  
4 *Control* **48** (2003) 692–697.
- 5 39. D. D. Holm and V. Putkaradze, Formation of clumps and patches in self-aggregation  
6 of finite-size particles, *Phys. D* **220** (2006) 183–196.
- 7 40. Y. Huang and A. L. Bertozzi, Self-similar blowup solutions to an aggregation equation  
8 in  $\mathbf{R}^n$ , *SIAM J. Appl. Math.* **70** (2010) 2582–2603.
- 9 41. E. F. Keller and L. A. Segel, Initiation of slime mold aggregation viewed as an insta-  
10 bility, *J. Theor. Biol.* **26** (1970) 399–415.
- 11 42. S. Kida, Motion of an elliptic vortex in a uniform shear flow, *J. Phys. Soc. Jpn* **50**  
12 (1981) 3517–3520.
- 13 43. G. Kirchoff, *Vorlesungen über Mechanik* (B. G. Teubner, 1897).
- 14 44. S. H. Lamb, *Hydrodynamics*, 6th edn. (Cambridge Univ. Press, 1932).
- 15 45. T. Laurent, Local and global existence for an aggregation equation, *Comm. Partial*  
16 *Differential Equations* **32**(10–12) (2007) 1941–1964.
- 17 46. D. Li and J. Rodrigo, Finite-time singularities of an aggregation equation in  $\mathbb{R}^n$  with  
18 fractional dissipation, *Comm. Math. Phys.* **287** (2009) 687–703.
- 19 47. D. Li and J. L. Rodrigo, Refined blowup criteria and nonsymmetric blowup of an  
20 aggregation equation, *Adv. Math.* **220** (2009) 1717–1738.
- 21 48. D. Li and X. Zhang, On a nonlocal aggregation model with nonlinear diffusion, *Disc.*  
22 *Cont. Dyn. Syst.* **27** (2010) 301–323.
- 23 49. H. Li and G. Toscani, Long-time asymptotics of kinetic models of granular flows,  
24 *Arch. Ration. Mech. Anal.* **172** (2004) 407–428.
- 25 50. F. Lin and P. Zhang, On the hydrodynamic limit of Ginzburg–Landau vortices, *Disc.*  
26 *Cont. Dyn. Syst.* **6** (2000) 121–142.
- 27 51. E. Mainini, A global uniqueness result for an evolution problem arising in supercon-  
28 ductivity, *Boll. Unione Mat. Ital. (9)* (2009).
- 29 52. E. Mainini, Well-posedness for a mean field model of Ginzburg–Landau vortices with  
30 opposite degrees, 2010.
- 31 53. A. J. Majda and A. L. Bertozzi, *Vorticity and Incompressible Flow*, Cambridge Texts in  
32 Applied Mathematics, Vol. 27 (Cambridge Univ. Press, 2002).
- 33 54. N. Masmoudi and P. Zhang, Global solutions to vortex density equations arising from  
34 sup-conductivity, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **22** (2005) 441–458.
- 35 55. A. Mogilner, L. Edelstein-Keshet, L. Bent and A. Spiros, Mutual interactions, poten-  
36 tials, and individual distance in a social aggregation, *J. Math. Biol.* **47** (2003) 353–389.
- 37 56. A. Mogilner and L. Edelstein-Keshet, A nonlocal model for a swarm, *J. Math. Biol.*  
38 **38** (1999) 534–570.
- 39 57. D. Morale, V. Capasso and K. Oelschläger, An interacting particle system mod-  
40 elling aggregation behavior: From individuals to populations, *J. Math. Biol.* **50** (2005)  
41 49–66.
- 42 58. J. C. Neu, The dynamics of a columnar vortex in an imposed strain, *Phys. Fluids* **27**  
43 (1984) 2397–2402.
- 44 59. J. Nieto, F. Poupaud and J. Soler, High-field limit for the Vlasov–Poisson–Fokker-  
45 Planck system, *Arch. Rational Mech. Anal.* **158** (2001) 29–59.
- 46 60. F. Poupaud, Diagonal defect measures, adhesion dynamics and Euler equation, *Methods Appl. Anal.* **9** (2002) 533–561.
- 47 61. R. Robert, Unicité de la solution faible à support compact de l'équation de Vlasov  
48 Poisson, *C. R. Acad. Sci. Sér. I* **324** (1997) 873–877.
- 49

*Aggregation via the Newtonian Potential and Aggregation Patches*

- 1 62. É. Sandier and S. Serfaty, A rigorous derivation of a free-boundary problem arising  
2 in superconductivity, *Ann. Sci. École Norm. Sup. (4)* **33** (2000) 561–592.
- 3 63. E. Sandier and S. Serfaty, *Vortices in the magnetic Ginzburg-Landau model*, Progress  
4 in Nonlinear Differential Equations and their Applications, Vol. 70 (Birkhäuser, 2007).
- 5 64. E. M. Stein, *Harmonic Analysis: Real-Variable Methods, Orthogolality, and Oscillatory*  
6 *Integrals* (Princeton Univ. Press, 1993).
- 7 65. C. M. Topaz and A. L. Bertozzi, Swarming patterns in a two-dimensional kinematic  
8 model for biological groups, *SIAM J. Appl. Math.* **65**(1) (2004) 152–174.
- 9 66. C. M. Topaz, A. L. Bertozzi and M. A. Lewis, A nonlocal continuum model for  
10 biological aggregation, *Bull. Math. Biol.* **68** (2006) 1601–1623.
- 11 67. G. Toscani, One-dimensional kinetic models of granular flows, *M2AN Math. Model.*  
12 *Numer. Anal.* **34** (2000) 1277–1291.
- 13 68. V. I. Yudovich, Non-stationary flow of an ideal incompressible liquid, *USSR Comput.*  
14 *Math. Math. Phys.* **3** (1963) 1407–1456.
- 15 69. N. J. Zabusky, M. H. Hughes and K. V. Roberts, Contour dynamics for the Euler  
16 equations in two dimensions, *J. Comput. Phys.* **30** (1979) 96–106.