

FUNCTIONAL CLT FOR RANDOM WALK AMONG BOUNDED RANDOM CONDUCTANCES

MAREK BISKUP AND TIMOTHY M. PRESCOTT

Department of Mathematics, University of California at Los Angeles

ABSTRACT. We consider the nearest-neighbor simple random walk on \mathbb{Z}^d , $d \geq 2$, driven by a field of i.i.d. random nearest-neighbor conductances $\omega_{xy} \in [0, 1]$. Apart from the requirement that the bonds with positive conductances percolate, we pose no restriction on the law of the ω 's. We prove that, for a.e. realization of the environment, the path distribution of the walk converges weakly to that of non-degenerate, isotropic Brownian motion. The quenched functional CLT holds despite the fact that the local CLT may fail in $d \geq 5$ due to anomalously slow decay of the probability that the walk returns to the starting point at a given time.

1. INTRODUCTION

Let \mathbb{B}_d denote the set of unordered nearest-neighbor pairs (i.e., edges) of \mathbb{Z}^d and let $(\omega_b)_{b \in \mathbb{B}_d}$ be i.i.d. random variables with $\omega_b \in [0, 1]$. We will refer to ω_b as the *conductance* of the edge b . Let \mathbb{P} denote the law of the ω 's and suppose that

$$\mathbb{P}(\omega_b > 0) > p_c(d), \tag{1.1}$$

where $p_c(d)$ is the threshold for bond percolation on \mathbb{Z}^d ; in $d = 1$ we have $p_c(d) = 1$ so there we suppose $\omega_b > 0$ a.s. This condition ensures the existence of a unique infinite connected component \mathcal{C}_∞ of edges with strictly positive conductances; we will typically restrict attention to ω 's for which \mathcal{C}_∞ contains a given site (e.g., the origin).

Each realization of \mathcal{C}_∞ can be used to define a random walk $X = (X_n)$ which moves about \mathcal{C}_∞ by picking, at each unit time, one of its $2d$ neighbors at random and passing to it with probability equal to the conductance of the corresponding edge. Technically, X is a Markov chain with state space \mathcal{C}_∞ and the transition probabilities defined by

$$P_{\omega,z}(X_{n+1} = y | X_n = x) := \frac{\omega_{xy}}{2d} \tag{1.2}$$

if $x, y \in \mathcal{C}_\infty$ and $|x - y| = 1$, and

$$P_{\omega,z}(X_{n+1} = x | X_n = x) := 1 - \frac{1}{2d} \sum_{y: |y-x|=1} \omega_{xy}. \tag{1.3}$$

© 2007 by M. Biskup and T.M. Prescott. Reproduction, by any means, of the entire article for non-commercial purposes is permitted without charge.

The second index on $P_{\omega,z}$ marks the initial position of the walk, i.e.,

$$P_{\omega,z}(X_0 = z) := 1. \quad (1.4)$$

The counting measure on \mathcal{C}_∞ is invariant and reversible for this Markov chain.

The $d = 1$ walk is a simple, but instructive, exercise in harmonic analysis of reversible random walks in random environments. Let us quickly sketch the proof of the fact that, for a.e. ω sampled from a translation-invariant, ergodic law on $(0, 1]^{\mathbb{B}^d}$ satisfying

$$\mathbb{E}\left(\frac{1}{\omega_b}\right) < \infty, \quad (1.5)$$

the walk scales to Brownian motion under the usual diffusive scaling of space and time. (Here and henceforth \mathbb{E} denotes expectation with respect to the environment distribution.) The derivation works even for unbounded conductances provided (1.2–1.3) are modified accordingly.

Abbreviate $C := \mathbb{E}(1/\omega_b)$. The key step of the proof is to realize that

$$\varphi_\omega(x) := x + \frac{1}{C} \sum_{n=0}^{x-1} \left(\frac{1}{\omega_{n,n+1}} - C \right) \quad (1.6)$$

is harmonic for the Markov chain. Hence $\varphi_\omega(X_n)$ is a martingale whose increments are, by (1.5) and a simple calculation, square integrable in the sense

$$\mathbb{E}E_{\omega,0}[\varphi_\omega(X_1)^2] < \infty. \quad (1.7)$$

Invoking the stationarity and ergodicity of the Markov chain on the space of environments “from the point of view of the particle” — we will discuss the specifics of this argument later — the martingale $(\varphi_\omega(X_n))$ satisfies the conditions of the Lindeberg-Feller martingale functional CLT and so the law of $t \mapsto \varphi_\omega(X_{\lfloor nt \rfloor})/\sqrt{n}$ tends weakly to that of a Brownian motion with diffusion constant given by (1.7). By the Pointwise Ergodic Theorem and (1.5) we have $\varphi_\omega(x) - x = o(x)$ as $|x| \rightarrow \infty$. Thus the path $t \mapsto X_{\lfloor nt \rfloor}/\sqrt{n}$ scales, in the limit $n \rightarrow \infty$, to the same function as the deformed path $t \mapsto \varphi_\omega(X_{\lfloor nt \rfloor})/\sqrt{n}$. As this holds for a.e. ω , we have proved a *quenched* functional CLT.

While the main ideas of the above $d = 1$ solution work in all dimensions, the situation in $d \geq 2$ is, even for i.i.d. conductances, significantly more complicated. Progress has been made under additional conditions on the environment law. One such condition is *strong ellipticity*,

$$\exists \alpha > 0 : \quad \mathbb{P}(\alpha \leq \omega_b \leq 1/\alpha) = 1. \quad (1.8)$$

Here an annealed invariance principle was proved by Kipnis and Varadhan [20] and its quenched counterpart by Sidoravicius and Sznitman [29]. Another natural family of environments are those arising from supercritical *bond percolation* on \mathbb{Z}^d for which (ω_b) are i.i.d. zero-one valued with $\mathbb{P}(\omega_b = 1) > p_c(d)$. For these cases an annealed invariance principle was proved by De Masi, Ferrari, Goldstein and Wick [11, 12] and the quenched case was established in $d \geq 4$ by Sidoravicius and Sznitman [29], and in all $d \geq 2$ by Berger and Biskup [6] and Mathieu and Piatnitski [25].

A common feature of the latter proofs is that, in $d \geq 3$, they require the use of heat-kernel upper bounds of the form

$$P_{\omega,x}(X_n = y) \leq \frac{c_1}{n^{d/2}} \exp\left\{-c_2 \frac{|x-y|^2}{n}\right\}, \quad x, y \in \mathcal{C}_\infty, \quad (1.9)$$

where c_1, c_2 are absolute constants and n is assumed to exceed a random quantity depending on the environment in the vicinity of x and y . These were deduced by Barlow [2] using sophisticated arguments that involve isoperimetry, regular volume growth and comparison of graph-theoretical and Euclidean distances for the percolation cluster. While the use of (1.9) is conceptually rather unsatisfactory — one seems to need a local-CLT level of control to establish a plain CLT — no arguments (in $d \geq 3$) that avoid heat-kernel estimates are known at present.

The reliance on heat-kernel bounds also suffers from another problem: (1.9) may actually *fail* once the conductance law has sufficiently heavy tails at zero. This was first noted to happen by Fontes and Mathieu [14] for the heat-kernel averaged over the environment; the quenched situation was analyzed recently by Berger, Biskup, Hoffman and Kozma [7]. The main conclusion of [7] is that the diagonal (i.e., $x = y$) bound in (1.9) holds in $d = 2, 3$ but the decay can be slower than any $o(n^{-2})$ sequence in $d \geq 5$. (The threshold sequence in $d = 4$ is presumably $o(n^{-2} \log n)$.) This is caused by the existence of *traps* that may capture the walk for a long time and thus, paradoxically, increase its chances to arrive back to the starting point.

The aforementioned facts lead to a natural question: In the absence of heat-kernel estimates, does the quenched CLT still hold? Our answer to this question is affirmative and constitutes the main result of this note. Another interesting question is what happens when the conductances are unbounded from above; this is currently being studied by Barlow and Deuschel [3].

Note: While this paper was in the process of writing, we received a preprint from Pierre Mathieu [24] in which he proves a result that is a continuous-time version of our main theorem. The strategy of [24] differs from ours by the consideration of a time-changed process (which we use only marginally) and proving that the “new” and “old” time scales are commensurate. Our approach is focused on proving the (pointwise) sublinearity of the corrector and it streamlines considerably the proof of [6] in $d \geq 3$ in that it limits the use of “heat-kernel technology” to a uniform bound on the heat-kernel decay (implied by isoperimetry) and a diffusive bound on the expected distance travelled by the walk (implied by regular volume growth).

2. MAIN RESULTS AND OUTLINE

Let $\Omega := [0, 1]^{\mathbb{B}_d}$ be the set of all admissible random environments and let \mathbb{P} be an i.i.d. law on Ω . Assuming (1.1), let \mathcal{C}_∞ denote the a.s. unique infinite connected component of edges with positive conductances and introduce the conditional measure

$$\mathbb{P}_0(-) := \mathbb{P}(-|0 \in \mathcal{C}_\infty). \quad (2.1)$$

For $T > 0$, let $(C[0, T], \mathscr{W}_T)$ be the space of continuous functions $f: [0, T] \rightarrow \mathbb{R}^d$ equipped with the Borel σ -algebra defined relative to the supremum topology.

Here is our main result:

Theorem 2.1 *Suppose $d \geq 2$ and $\mathbb{P}(\omega_b > 0) > p_c(d)$. For $\omega \in \{0 \in \mathcal{C}_\infty\}$, let $(X_n)_{n \geq 0}$ be the random walk with law $P_{\omega,0}$ and let*

$$B_n(t) := \frac{1}{\sqrt{n}}(X_{\lfloor tn \rfloor} + (tn - \lfloor tn \rfloor)(X_{\lfloor tn \rfloor + 1} - X_{\lfloor tn \rfloor})), \quad t \geq 0. \quad (2.2)$$

Then for all $T > 0$ and for \mathbb{P}_0 -almost every ω , the law of $(B_n(t): 0 \leq t \leq T)$ on $(C[0, T], \mathscr{W}_T)$ converges, as $n \rightarrow \infty$, weakly to the law of an isotropic Brownian motion $(B_t: 0 \leq t \leq T)$ with a positive and finite diffusion constant (which is independent of ω).

Using a variant of [6, Lemma 6.4], from here we can extract a corresponding conclusion for the “agile” version of our random walk (cf [6, Theorem 1.2]) by which we mean the walk that jumps from x to its neighbor y with probability $\omega_{xy}/\pi_\omega(x)$ where $\pi_\omega(x)$ is the sum of ω_{xz} over all of the neighbors z of x . Replacing discrete times by sums of i.i.d. exponential random variables, these invariance principles then extend also to the corresponding continuous-time processes. Theorem 2.1 of course implies also an annealed invariance principle, which is the above convergence for the walk sampled from the path measure integrated over the environment.

Remark 2.2 As we were reminded by Y. Peres, the above functional CLT automatically implies the “usual” lower bound on the heat-kernel. Indeed, the Markov property and reversibility of X yield

$$P_{\omega,0}(X_{2n} = 0) \geq \sum_{\substack{x \in \mathcal{C}_\infty \\ |x| \leq \sqrt{n}}} P_{\omega,0}(X_n = x)^2. \quad (2.3)$$

Cauchy-Schwarz then gives

$$P_{\omega,0}(X_{2n} = 0) \geq P_{\omega,0}(|X_n| \leq \sqrt{n})^2 \frac{1}{|\mathcal{C}_\infty \cap [-\sqrt{n}, \sqrt{n}]^d|}. \quad (2.4)$$

Now Theorem 2.1 implies that $P_{\omega,0}(|X_n| \leq \sqrt{n})$ is uniformly positive as $n \rightarrow \infty$ and the Spatial Ergodic Theorem shows that $|\mathcal{C}_\infty \cap [-\sqrt{n}, \sqrt{n}]^d|$ grows proportionally to $n^{d/2}$. Hence we get

$$P_{\omega,0}(X_{2n} = 0) \geq \frac{C(\omega)}{n^{d/2}}, \quad n \geq 1, \quad (2.5)$$

with $C(\omega) > 0$ a.s. on the set $\{0 \in \mathcal{C}_\infty\}$. Note that, in $d = 2, 3$, this complements nicely the “universal” upper bounds derived in [7].

The remainder of this paper is devoted to the proof of Theorem 2.1. The main line of attack is similar to the above 1D solution: We define a harmonic coordinate φ_ω — an analogue of (1.6) — and then prove an a.s. invariance principle for

$$t \mapsto \varphi_\omega(X_{\lfloor nt \rfloor})/\sqrt{n} \quad (2.6)$$

along the martingale argument sketched before. The difficulty comes with showing the sublinearity of the corrector,

$$\varphi_\omega(x) - x = o(x), \quad |x| \rightarrow \infty. \quad (2.7)$$

As in Berger and Biskup [6], sublinearity can be proved directly along coordinate directions by soft ergodic-theory arguments. The crux is to extend this to a bound throughout d -dimensional boxes.

Following the $d \geq 3$ proof of [6], the bound along coordinate axes readily implies *sublinearity on average*, meaning that the set of x where $|\varphi_\omega(x) - x|$ exceeds $\epsilon|x|$ has zero density. The extension of sublinearity on average to pointwise sublinearity is the main novel part of the proof which, unfortunately, still makes non-trivial use of the “heat-kernel technology.” A heat-kernel upper bound of the form (1.9) would do but, to minimize the extraneous input, we show that it suffices to have a diffusive bound for the expected displacement of the walk from its starting position. This step still requires detailed control of isoperimetry and volume growth as well as the comparison of the graph-theoretic and Euclidean distances, but it avoids many spurious calculations that are needed for the full-fledged heat-kernel estimates.

Of course, the required isoperimetric inequalities may not be true on \mathcal{C}_∞ because of the presence of “weak” bonds. As in [7] we circumvent this by observing the random walk on the set of sites that have a connection to infinity by bonds with *uniformly* positive conductances. Specifically we pick $\alpha > 0$ and let $\mathcal{C}_{\infty,\alpha}$ denote the set of sites in \mathbb{Z}^d that are connected to infinity by a path whose edges obey $\omega_b \geq \alpha$. Here we note:

Proposition 2.3 *Let $d \geq 2$ and $p = \mathbb{P}(\omega_b > 0) > p_c(d)$. Then there exists $c(p, d) > 0$ such that if α satisfies*

$$\mathbb{P}(\omega_b \geq \alpha) > p_c(d) \quad (2.8)$$

and

$$\mathbb{P}(0 < \omega_b < \alpha) < c(p, d) \quad (2.9)$$

then $\mathcal{C}_{\infty,\alpha}$ is nonempty and $\mathcal{C}_\infty \setminus \mathcal{C}_{\infty,\alpha}$ has only finite components a.s. In fact, if $\mathcal{F}(x)$ is the set of sites (possibly empty) in the finite component of $\mathcal{C}_\infty \setminus \mathcal{C}_{\infty,\alpha}$ containing x , then

$$\mathbb{P}(x \in \mathcal{C}_\infty \ \& \ \text{diam } \mathcal{F}(x) \geq n) \leq Ce^{-\eta n}, \quad n \geq 1, \quad (2.10)$$

for some $C < \infty$ and $\eta > 0$. Here “diam” is the diameter in the ℓ_∞ distance on \mathbb{Z}^d .

The restriction of φ_ω to $\mathcal{C}_{\infty,\alpha}$ is still harmonic, but with respect to a walk that can “jump the holes” of $\mathcal{C}_{\infty,\alpha}$. A discrete-time version of this walk was utilized heavily in [7]; for the purposes of this paper it will be more convenient to work with its continuous-time counterpart $Y = (Y_t)_{t \geq 0}$. Explicitly, sample a path of the random walk $X = (X_n)$ from $P_{\omega,0}$ and denote by T_1, T_2, \dots the time intervals between successive visits of X to $\mathcal{C}_{\infty,\alpha}$. These are defined recursively by

$$T_{j+1} := \inf \{ n \geq 1 : X_{T_1 + \dots + T_j + n} \in \mathcal{C}_{\infty,\alpha} \}, \quad (2.11)$$

with $T_0 = 0$. For each $x, y \in \mathcal{C}_{\infty,\alpha}$, let

$$\hat{\omega}_{xy} := P_{\omega,x}(X_{T_1} = y) \quad (2.12)$$

and define the operator

$$(\mathcal{L}_\omega^{(\alpha)} f)(x) := \sum_{y \in \mathcal{C}_{\infty, \alpha}} \hat{\omega}_{xy} [f(y) - f(x)]. \quad (2.13)$$

The continuous-time random walk Y is a Markov process with this generator; alternatively take the standard Poisson process $(N_t)_{t \geq 0}$ with jump-rate one and set

$$Y_t := X_{T_1 + \dots + T_{N_t}}. \quad (2.14)$$

Note that, while Y may jump “over the holes” of $\mathcal{C}_{\infty, \alpha}$, Proposition 2.3 ensures that all of its jumps are finite. The counting measure on $\mathcal{C}_{\infty, \alpha}$ is still invariant for this random walk, $\mathcal{L}_\omega^{(\alpha)}$ is self-adjoint on the corresponding space of square integrable functions and $\mathcal{L}_\omega^{(\alpha)} \varphi_\omega = 0$ on $\mathcal{C}_{\infty, \alpha}$ (see Lemma 5.2).

The skeleton of the proof is condensed into the following statement whose proof, and adaptation to the present situation, is the main novel part of this note:

Theorem 2.4 *Fix α as in (2.8–2.9) and suppose $\psi_\omega : \mathcal{C}_{\infty, \alpha} \rightarrow \mathbb{R}^d$ is a function and $\theta > 0$ is a number such that the following holds for a.e. ω :*

- (1) *(Harmonicity) If $\varphi_\omega(x) := x + \psi_\omega(x)$, then $\mathcal{L}_\omega^{(\alpha)} \varphi_\omega = 0$ on $\mathcal{C}_{\infty, \alpha}$.*
- (2) *(Sublinearity on average) For every $\epsilon > 0$,*

$$\lim_{n \rightarrow \infty} \frac{1}{n^d} \sum_{\substack{x \in \mathcal{C}_{\infty, \alpha} \\ |x| \leq n}} 1_{\{|\psi_\omega(x)| \geq \epsilon n\}} = 0. \quad (2.15)$$

- (3) *(Polynomial growth)*

$$\lim_{n \rightarrow \infty} \max_{\substack{x \in \mathcal{C}_{\infty, \alpha} \\ |x| \leq n}} \frac{|\psi_\omega(x)|}{n^\theta} = 0. \quad (2.16)$$

Let $Y = (Y_t)$ be the continuous-time random walk on $\mathcal{C}_{\infty, \alpha}$ with generator $\mathcal{L}_\omega^{(\alpha)}$ and suppose also:

- (4) *(Diffusive upper bounds) For a deterministic sequence $b_n = o(n^2)$ and a.e. ω ,*

$$\sup_{n \geq 1} \max_{\substack{x \in \mathcal{C}_{\infty, \alpha} \\ |x| \leq n}} \sup_{t \geq b_n} \frac{E_{\omega, x} |Y_t - x|}{\sqrt{t}} < \infty \quad (2.17)$$

and

$$\sup_{n \geq 1} \max_{\substack{x \in \mathcal{C}_{\infty, \alpha} \\ |x| \leq n}} \sup_{t \geq b_n} t^{d/2} P_{\omega, x}(Y_t = x) < \infty. \quad (2.18)$$

Then for almost every ω ,

$$\lim_{n \rightarrow \infty} \max_{\substack{x \in \mathcal{C}_{\infty, \alpha} \\ |x| \leq n}} \frac{|\psi_\omega(x)|}{n} = 0. \quad (2.19)$$

This result — with ψ_ω playing the role of the corrector — shows that $\varphi_\omega(x) - x = o(x)$ on $\mathcal{C}_{\infty,\kappa}$. This readily extends to sublinearity on \mathcal{C}_∞ by the maximum principle applied to φ_ω on the finite components of $\mathcal{C}_\infty \setminus \mathcal{C}_{\infty,\kappa}$ and using that the component sizes obey a polylogarithmic upper bound. The assumptions (1-3) are known to hold for the corrector of the supercritical bond-percolation cluster and the proof applies, with minor modifications, to the present case as well. The crux is to prove (2.17–2.18) which is where we need to borrow ideas from the “heat-kernel technology.” For our purposes it will suffice to take $b_n = n$ in part (4).

We remark that the outline strategy of proof extends rather seamlessly to other (translation-invariant, ergodic) conductance distributions with conductances bounded from above. Of course, one has to assume a number of specific properties for the “strong” component $\mathcal{C}_{\infty,\kappa}$ that, in the i.i.d. case, we are able to check explicitly.

The plan of the rest of this paper is as follows: Sect. 3 is devoted to some basic percolation estimates needed in the rest of the paper. In Sect. 4 we define and prove some properties of the corrector χ , which is a random function marking the difference between the harmonic coordinate $\varphi_\omega(x)$ and the geometric coordinate x . In Sect. 5 we establish the a.s. sublinearity of the corrector as stated in Theorem 2.4 subject to the diffusive bounds (2.17–2.18). Then we assemble all facts into the proof of Theorem 2.1. Finally, in Sect. 6 we adapt some arguments from Barlow [2] to prove (2.17–2.18); first in rather general Propositions 6.1 and 6.2 and then for the case at hand.

3. PERCOLATION ESTIMATES

In this section we provide a proof of Proposition 2.3 and also of a lemma dealing with the maximal distance the random walk Y can travel in a given number of steps. We will need to work with the “static” renormalization (cf Grimmett [17, Section 7.4]) whose salient features we will now recall. The underlying ideas go back to the work of Kesten and Zhang [19], Grimmett and Marstrand [18] and Antal and Pisztora [1].

We say that an edge b is occupied if $\omega_b > 0$. Consider the lattice cubes

$$B_L(x) := x + [0, L]^d \cap \mathbb{Z}^d \quad (3.1)$$

and

$$\tilde{B}_{3L}(x) := x + [-L, 2L]^d \cap \mathbb{Z}^d \quad (3.2)$$

and note that $\tilde{B}_{3L}(x)$ consists of 3^d copies of $B_L(x)$ that share only sites on their adjacent boundaries. Let $G_L(x)$ be the “good event” — whose occurrence designates $B_L(Lx)$ to be a “good block” — which is the set of configurations such that:

- (1) For each neighbor y of x , the side of the block $B_L(Ly)$ adjacent to $B_L(Lx)$ is connected to the opposite side of $B_L(Ly)$ by an occupied path.
- (2) Any two occupied paths connecting $B_L(Lx)$ to the boundary of $\tilde{B}_{3L}(Lx)$ are connected by an occupied path using only edges with both endpoints in $\tilde{B}_{3L}(Lx)$.

The sheer existence of infinite cluster implies that (1) occurs with high probability once L is large (see Grimmett [17, Theorem 8.97]) while the situation in (2) occurs with large

probability once there is percolation in half space (see Grimmett [17, Lemma 7.89]). It follows that

$$\mathbb{P}(G_L(x)) \xrightarrow{L \rightarrow \infty} 1 \quad (3.3)$$

whenever $\mathbb{P}(\omega_b > 0) > p_c(d)$. A crucial consequence of the above conditions is that, if $G_L(x)$ and $G_L(y)$ occur for neighboring sites $x, y \in \mathbb{Z}^d$, then the largest connected components in $\tilde{B}_{3L}(Lx)$ and $\tilde{B}_{3L}(Ly)$ — sometimes referred to as *spanning clusters* — are connected. Thus, if $G_L(x)$ occurs for all x along an infinite path on \mathbb{Z}^d , the corresponding spanning clusters are subgraphs of \mathcal{C}_∞ .

A minor complication is that the events $\{G_L(x) : x \in \mathbb{Z}^d\}$ are not independent. However, they are 4-dependent in the sense that if (x_i) and (y_j) are such that $|x_i - y_j| > 4$ for each i and j , then the families $\{G_L(x_i)\}$ and $\{G_L(y_j)\}$ are independent. By the main result of Liggett, Schonmann and Stacey [21, Theorem 0.0] (cf [17, Theorem 7.65]) the indicators $\{1_{G_L(x)} : x \in \mathbb{Z}^d\}$, regarded as a random process on \mathbb{Z}^d , stochastically dominate i.i.d. Bernoulli random variables whose density (of ones) tends to one as $L \rightarrow \infty$.

Proof of Proposition 2.3. In $d = 2$ the proof is actually very simple because it suffices to choose α such that (2.8) holds. Then $\mathcal{C}_\infty \setminus \mathcal{C}_{\infty, \alpha} \subset \mathbb{Z}^2 \setminus \mathcal{C}_{\infty, \alpha}$ has only finite (subcritical) components whose diameter has exponential tails (2.10) by, e.g., [17, Theorem 6.10].

To handle general dimensions we will have to invoke the above “static” renormalization. Let $G_L(x)$ be as above and consider the event $G_{L, \alpha}(x)$ where we in addition require that $\omega_b \notin (0, \alpha)$ for every edge with both endpoints in $\tilde{B}_{3L}(Lx)$. Clearly,

$$\lim_{L \rightarrow \infty} \lim_{\alpha \downarrow 0} \mathbb{P}(G_{L, \alpha}(x)) = 1. \quad (3.4)$$

Using the aforementioned domination by site percolation, and adjusting L and α we can thus ensure that, with probability one, the set

$$\{x \in \mathbb{Z}^d : G_{L, \alpha}(x) \text{ occurs}\} \quad (3.5)$$

has a unique infinite component \mathcal{C}_∞ , whose complement has only finite components. Moreover, if $\mathcal{G}(0)$ is the finite connected component of $\mathbb{Z}^d \setminus \mathcal{C}_\infty$ containing the origin, then a standard Peierls argument yields

$$\mathbb{P}(\text{diam } \mathcal{G}(0) \geq n) \leq e^{-\zeta n} \quad (3.6)$$

for some $\zeta > 0$. To prove (2.10), it suffices to show that

$$\mathcal{F}(0) \subset \bigcup_{x \in \mathcal{G}(0)} B_L(Lx) \quad (3.7)$$

once $\text{diam } \mathcal{F}(0) > 3L$. Indeed, then $\text{diam } \mathcal{F}(0) \leq L \text{diam } \mathcal{G}(0)$ and so (3.6) implies (2.10) with $\eta := \zeta/L$ and $C := e^{3L\eta}$.

To prove (3.7), pick $z \in \mathcal{F}(0)$ and let x be such that $z \in B_L(Lx)$. It suffices to show that if $G_{L, \alpha}(x)$ occurs, then x is not adjacent to an infinite component in (3.5). Assuming that x is adjacent to such a component, the fact that the spanning clusters in adjacent “good blocks” are connected and thus contained in $\mathcal{C}_{\infty, \alpha}$ implies

$$\mathcal{C}_{\infty, \alpha} \cap B_L(Lx) \neq \emptyset. \quad (3.8)$$

But then $B_L(Lx)$ is intersected by two “large” components, $\mathcal{C}_{\infty,\alpha}$ and $\mathcal{F}(0)$, of edges with $\omega_b \geq \alpha$. (This is where we need $\text{diam } \mathcal{F}(0) > 3L$.) If these components are joined by an occupied path — i.e., a path of edges with $\omega_b > 0$ — within $\tilde{B}_{3L}(Lx)$, then $\tilde{B}_{3L}(Lx)$ contains a “weak” bond and so $G_{L,\alpha}$ fails. In the absence of such a path the requirement (2) in the definition of $G_L(x)$ is not satisfied and so $G_{L,\alpha}(x)$ fails too. \square

Let $d(x, y)$ be the “Markov distance” on $V = \mathcal{C}_{\infty,\alpha}$, i.e., the minimal number of jumps the random walk $Y = (Y_t)$ needs to make to get from x to y . Note that $d(x, y)$ could be quite smaller than the graph-theoretic distance on $\mathcal{C}_{\infty,\alpha}$ and/or the Euclidean distance. (The latter distances are known to be comparable, see Antal and Pisztora [1].) To control the volume growth for the Markov graph of the random walk Y we will need to know that $d(x, y)$ is nevertheless comparable with the Euclidean distance $|x - y|$:

Lemma 3.1 *There exists $\varrho > 0$ and for each $\gamma > 0$ there is $\alpha > 0$ obeying (2.8–2.9) and $C < \infty$ such that*

$$\mathbb{P}(0, x \in \mathcal{C}_{\infty,\alpha} \ \& \ d(0, x) \leq \varrho|x|) \leq Ce^{-\gamma|x|}, \quad x \in \mathbb{Z}^d. \quad (3.9)$$

Proof. Suppose α is as in the proof of Proposition 2.3. Let (η_x) be independent Bernoulli that dominate the indicators $1_{G_{L,\alpha}}$ from below and let \mathcal{C}_∞ be the unique infinite component of the set $\{x \in \mathbb{Z}^d : \eta_x = 1\}$. We may “wire” the “holes” of \mathcal{C}_∞ by putting an edge between every pair of sites on the external boundary of each finite component of $\mathbb{Z}^d \setminus \mathcal{C}_\infty$; we use $d'(0, x)$ to denote the distance between 0 and x on the induced graph. The processes η and $(1_{G_{L,\alpha}(x)})$ can be coupled so that each connected component of $\mathcal{C}_\infty \setminus \mathcal{C}_{\infty,\alpha}$ with diameter exceeding $3L$ is “covered” by a finite component of $\mathbb{Z}^d \setminus \mathcal{C}_\infty$, cf (3.7). As is easy to check, this implies

$$d(0, x) \geq d'(0, x') \quad \text{and} \quad |x'| \geq \frac{1}{L}|x| - 1 \quad (3.10)$$

whenever $x \in B_L(Lx')$. It thus suffices to show the above bound for distance $d'(0, x')$.

Let $p = p_{L,\alpha}$ be the parameter of the Bernoulli distribution and recall that p can be made as close to one as desired by adjusting L and α . Let $z_0 = 0, z_1, \dots, z_n = x$ be a nearest-neighbor path on \mathbb{Z}^d . Let $\mathcal{G}(z_i)$ be the unique finite component of $\mathbb{Z}^d \setminus \mathcal{C}_\infty$ that contains z_i — if $z_i \in \mathcal{C}_\infty$, we have $\mathcal{G}(z_i) = \emptyset$. Define

$$\ell(z_0, \dots, z_n) := \sum_{i=0}^n \text{diam } \mathcal{G}(z_i) \left(\prod_{j<i} 1_{\{z_j \notin \mathcal{G}(z_i)\}} \right). \quad (3.11)$$

We claim that for each $\lambda > 0$ we can adjust p so that

$$\mathbb{E} e^{\lambda \ell(z_0, \dots, z_n)} \leq e^n \quad (3.12)$$

for all $n \geq 1$ and all paths as above. To verify this we note that the components contributing to $\ell(z_0, \dots, z_n)$ are distance at least one from one another. So conditioning on all but the last component, and the sites in the ultimate vicinity, we may use the Peierls argument to estimate the conditional expectation of $e^{\lambda \text{diam } \mathcal{G}(z_n)}$ by, say, e^1 . (We are using

also that $\text{diam } \mathcal{G}(z_n)$ is smaller than the boundary of $\mathcal{G}(z_n)$.) Proceeding by induction n times, (3.12) follows.

For any given $\gamma > 0$ we can adjust p so that (3.12) holds for

$$\lambda := 2(1 + \log(2d) + \gamma) \quad (3.13)$$

As the number of nearest-neighbor paths $(z_0 = 0, \dots, z_n = x)$ is bounded by $(2d)^n$, an exponential Chebyshev estimate then shows

$$\mathbb{P}\left(\exists(z_0 = 0, \dots, z_n = x): \ell(z_0, \dots, z_n) > \frac{n}{2}\right) \leq e^{-\gamma n}. \quad (3.14)$$

But if $(z_0 = 0, \dots, z_n = x)$ is the shortest nearest-neighbor interpolation of a path that achieves $d'(0, x)$, then

$$d'(0, x) \geq n - \ell(z_0, \dots, z_n). \quad (3.15)$$

Since, trivially, $|x| \leq n$, we deduce

$$\mathbb{P}(d'(0, x) \leq \frac{1}{2}|x|) \leq e^{-\gamma|x|} \quad (3.16)$$

as desired. \square

4. CORRECTOR

The purpose of this section is to define, and prove some properties of, the *corrector* $\chi(\omega, x) := \varphi_\omega(x) - x$. This object could be defined probabilistically by the limit

$$\chi(\omega, x) = \lim_{n \rightarrow \infty} (E_{\omega, x}(X_n) - E_{\omega, 0}(X_n)) - x, \quad (4.1)$$

unfortunately, at this moment we seem to have no direct (probabilistic) argument showing that the limit exists. The traditional definition of the corrector involves spectral calculus (Kipnis and Varadhan [20]); we will invoke a projection construction from Mathieu and Piatnitski [25] (see also Giacomin, Olla and Spohn [15]).

Let \mathbb{P} be an i.i.d. law on (Ω, \mathcal{F}) where $\Omega := [0, 1]^{\mathbb{B}^d}$ and \mathcal{F} is the natural product σ -algebra. Let $\tau_x: \Omega \rightarrow \Omega$ denote the shift by x , i.e.,

$$(\tau_x \omega)_{xy} := \omega_{x+z, y+z} \quad (4.2)$$

and note that $\mathbb{P} \circ \tau_x^{-1} = \mathbb{P}$ for all $x \in \mathbb{Z}^d$. Recall that \mathcal{C}_∞ is the infinite connected component of edges with $\omega_b > 0$ and, for $\alpha > 0$, let $\mathcal{C}_{\infty, \alpha}$ denote the set of sites connected to infinity by edges with $\omega_b \geq \alpha$. If $\mathbb{P}(0 \in \mathcal{C}_{\infty, \alpha}) > 0$, let

$$\mathbb{P}_\alpha(-) := \mathbb{P}(-|0 \in \mathcal{C}_{\infty, \alpha}) \quad (4.3)$$

and let \mathbb{E}_α be the corresponding expectation. Given $\omega \in \Omega$ and sites $x, y \in \mathcal{C}_{\infty, \alpha}(\omega)$, let $d_\omega^{(\alpha)}(x, y)$ denote the graph distance between x and y as measured on $\mathcal{C}_{\infty, \alpha}$. (Note this is distinct from the Markov distance $d(x, y)$ discussed, e.g., in Lemma 3.1.) We will also use \mathcal{L}_ω to denote the generator of the continuous-time version of the walk X , i.e.,

$$(\mathcal{L}_\omega f)(x) := \frac{1}{2d} \sum_{y: |y-x|=1} \omega_{xy} [f(y) - f(x)]. \quad (4.4)$$

The following theorem summarizes all relevant properties of the corrector:

Theorem 4.1 *Suppose $\mathbb{P}(0 \in \mathcal{C}_\infty) > 0$. There exists a function $\chi: \Omega \times \mathbb{Z}^d \rightarrow \mathbb{R}^d$ such that the following holds \mathbb{P}_0 -a.s.:*

- (1) (Gradient field) $\chi(0, \omega) = 0$ and, for all $x, y \in \mathcal{C}_\infty(\omega)$,

$$\chi(\omega, x) - \chi(\omega, y) = \chi(\tau_y \omega, x - y). \quad (4.5)$$

- (2) (Harmonicity) $\varphi_\omega(x) := x + \chi(\omega, x)$ obeys $\mathcal{L}_\omega \varphi_\omega = 0$.

- (3) (Square integrability) There is a constant $C = C(\alpha) < \infty$ such that for all $x, y \in \mathbb{Z}^d$ satisfying $|x - y| = 1$,

$$\mathbb{E}_\alpha(|\chi(\cdot, y) - \chi(\cdot, x)|^2 \omega_{xy} 1_{\{x \in \mathcal{C}_\infty\}}) < C \quad (4.6)$$

Let $\alpha > 0$ be such that $\mathbb{P}(0 \in \mathcal{C}_{\infty, \alpha}) > 0$. Then we also have:

- (4) (Polynomial growth) For every $\theta > d$, a.s.,

$$\lim_{n \rightarrow \infty} \max_{\substack{x \in \mathcal{C}_{\infty, \alpha} \\ |x| \leq n}} \frac{|\chi(\omega, x)|}{n^\theta} = 0. \quad (4.7)$$

- (5) (Zero mean under random shifts) Let $Z: \Omega \rightarrow \mathbb{Z}^d$ be a random variable such that

- (a) $Z(\omega) \in \mathcal{C}_{\infty, \alpha}(\omega)$,
- (b) \mathbb{P}_α is preserved by $\omega \mapsto \tau_{Z(\omega)}(\omega)$,
- (c) $\mathbb{E}_\alpha(d_\omega^{(\alpha)}(0, Z(\omega))^q) < \infty$ for some $q > 3d$.

Then $\chi(\cdot, Z(\cdot)) \in L^1(\Omega, \mathcal{F}, \mathbb{P}_\alpha)$ and

$$\mathbb{E}_\alpha[\chi(\cdot, Z(\cdot))] = 0. \quad (4.8)$$

As noted before, to construct the corrector we will invoke a projection argument. Abbreviate $L^2(\Omega) = L^2(\Omega, \mathcal{F}, \mathbb{P}_0)$ and let $B := \{\hat{e}_1, \dots, \hat{e}_d\}$ be the set of coordinate vectors. Consider the space $L^2(\Omega \times B)$ of square integrable functions $u: \Omega \times B \rightarrow \mathbb{R}^d$ equipped with the inner product

$$(u, v) := \mathbb{E}_0 \left(\sum_{b \in B} u(\omega, b) \cdot v(\omega, b) \omega_b \right). \quad (4.9)$$

We may interpret $u \in L^2(\Omega \times B)$ as a flow by putting $u(\omega, -b) = -u(\tau_{-b} \omega, b)$. Some, but not all, elements of $L^2(\Omega \times B)$ can be obtained as gradients of local functions, where the *gradient* ∇ is the map $L^2(\Omega) \rightarrow L^2(\Omega \times B)$ defined by

$$(\nabla \phi)(\omega, b) := \phi \circ \tau_b(\omega) - \phi(\omega). \quad (4.10)$$

Let L^2_∇ denote the closure of the set of gradients of all *local* functions — i.e., those depending only on the portion of ω in a finite subset of \mathbb{Z}^d — and note the following orthogonal decomposition $L^2(\Omega \times B) = L^2_\nabla \oplus (L^2_\nabla)^\perp$.

The elements of $(L_{\nabla}^2)^\perp$ can be characterized using the concept of *divergence*, which for $u: \Omega \times B \rightarrow \mathbb{R}^d$ is the function $\operatorname{div} u: \Omega \rightarrow \mathbb{R}^d$ defined by

$$\operatorname{div} u(\omega) := \sum_{b \in B} [\omega_b u(\omega, b) - \omega_{-b} u(\tau_{-b} \omega, b)]. \quad (4.11)$$

Using the interpretation of u as a flow, $\operatorname{div} u$ is simply the net flow out of the origin. The characterization of $(L_{\nabla}^2)^\perp$ is now as follows:

Lemma 4.2 $u \in (L_{\nabla}^2)^\perp$ if and only if $\operatorname{div} u(\omega) = 0$ for \mathbb{P}_0 -a.e. ω .

Proof. Let $u \in L^2(\Omega \times B)$ and let $\phi \in L^2(\Omega)$ be a local function. A direct calculation and the fact that $\omega_{-b} = (\tau_{-b} \omega)_b$ yield

$$(u, \nabla \phi) = -\mathbb{E}_0 \left(\phi(\omega) \operatorname{div} u(\omega) \right). \quad (4.12)$$

If $u \in (L_{\nabla}^2)^\perp$, then $\operatorname{div} u$ integrates to zero against all local functions. Since these are dense in $L^2(\Omega)$, we have $\operatorname{div} u = 0$ a.s. \square

It is easy to check that every $u \in L_{\nabla}^2$ is *curl-free* in the sense that for any oriented loop (x_0, x_1, \dots, x_n) on $\mathcal{C}_\infty(\omega)$ with $x_n = x_0$ we have

$$\sum_{j=0}^{n-1} u(\tau_{x_j} \omega, x_{j+1} - x_j) = 0. \quad (4.13)$$

On the other hand, every $u: \Omega \times B \rightarrow \mathbb{R}^d$ which is curl-free can be integrated into a unique function $\phi: \Omega \times \mathcal{C}_\infty(\cdot) \rightarrow \mathbb{R}^d$ such that

$$\phi(\omega, x) = \sum_{j=0}^{n-1} u(\tau_{x_j} \omega, x_{j+1} - x_j) \quad (4.14)$$

holds for any path (x_0, \dots, x_n) on $\mathcal{C}_\infty(\omega)$ with $x_0 = 0$ and $x_n = x$. This function will automatically satisfy the *shift-covariance* property

$$\phi(\omega, x) - \phi(\omega, y) = \phi(\tau_y \omega, x - y), \quad x, y \in \mathcal{C}_\infty(\omega). \quad (4.15)$$

We will denote the space of such functions $\mathcal{H}(\Omega \times \mathbb{Z}^d)$. To denote the fact that ϕ is assembled from the shifts of u , we will write

$$u = \operatorname{grad} \phi, \quad (4.16)$$

i.e., “grad” is a map from $\mathcal{H}(\Omega \times \mathbb{Z}^d)$ to functions $\Omega \times B \rightarrow \mathbb{R}^d$ that takes a function $\phi \in \mathcal{H}(\Omega \times \mathbb{Z}^d)$ and assigns to it the collection of values $\{\phi(\cdot, b) - \phi(\cdot, 0) : b \in B\}$.

Lemma 4.3 Let $\phi \in \mathcal{H}(\Omega \times \mathbb{Z}^d)$ be such that $\operatorname{grad} \phi \in (L_{\nabla}^2)^\perp$. Then ϕ is (discrete) harmonic for the random walk on \mathcal{C}_∞ , i.e., for \mathbb{P}_0 -a.e. ω and all $x \in \mathcal{C}_\infty(\omega)$,

$$(\mathcal{L}_\omega \phi)(\omega, x) = 0. \quad (4.17)$$

Proof. Our definition of divergence is such that “ $\operatorname{div} \operatorname{grad} = 2d \mathcal{L}_\omega$ ” holds. Lemma 4.2 implies that $u \in (L^2_{\nabla})^\perp$ if and only if $\operatorname{div} u = 0$, which is equivalent to $(\mathcal{L}_\omega \phi)(\omega, 0) = 0$. By translation covariance, this extends to all sites in \mathcal{C}_∞ . \square

Proof of Theorem 4.1(1-3). Consider the function $\phi(\omega, x) := x$ and let $u := \operatorname{grad} \phi$. Clearly, $u \in L^2(\Omega \times B)$. Let $G \in L^2_{\nabla}$ be the orthogonal projection of $-u$ onto L^2_{∇} and define $\chi \in \mathcal{H}(\Omega \times \mathbb{Z}^d)$ to be the unique function such that

$$G = \operatorname{grad} \chi \quad \text{and} \quad \chi(\cdot, 0) = 0. \quad (4.18)$$

This definition immediately implies (4.5), while the definition of the inner product on $L^2(\Omega \times B)$ directly yields (4.6). Since u projects to $-G$ on L^2_{∇} , we have $u + G \in (L^2_{\nabla})^\perp$. But $u + G = \operatorname{grad} [x + \chi(\omega, x)]$ and so, by Lemma 4.3, $x \mapsto x + \chi(\omega, x)$ is harmonic with respect to \mathcal{L}_ω . \square

Remark 4.4 We note that the corrector is actually uniquely determined by properties (1-3) of Theorem 4.1. In fact, $x + \chi$ spans the orthogonal complement of L^2_{∇} in the space of shift-covariant functions. See Biskup and Spohn [9].

For the remaining parts of Theorem 4.1 we will need to work on $\mathcal{C}_{\infty, \alpha}$. However, we do not yet need the full power of Proposition 2.3; it suffices to note that $\mathcal{C}_{\infty, \alpha}$ has the law of a supercritical percolation cluster.

Proof of Theorem 4.1(4). Let $\theta > d$ and abbreviate

$$R_n := \max_{\substack{x \in \mathcal{C}_{\infty, \alpha} \\ |x| \leq n}} |\chi(\omega, x)|. \quad (4.19)$$

By Theorem 1.1 of Antal and Pisztora [1],

$$\lambda(\omega) := \sup_{x \in \mathcal{C}_{\infty, \alpha}} \frac{d_\omega^{(\alpha)}(0, x)}{|x|} < \infty, \quad \mathbb{P}_\alpha\text{-a.s.}, \quad (4.20)$$

and so it suffices to show that $R_n/n^\theta \rightarrow 0$ on $\{\lambda(\omega) \leq \lambda\}$ for every $\lambda < \infty$. But on $\{\lambda(\omega) \leq \lambda\}$ every $x \in \mathcal{C}_{\infty, \alpha}$ with $|x| \leq n$ can be reached by a path on $\mathcal{C}_{\infty, \alpha}$ that does not leave $[-\lambda n, \lambda n]^d$ and so, on $\{\lambda(\omega) \leq \lambda\}$,

$$R_n \leq \sum_{\substack{x \in \mathcal{C}_{\infty, \alpha} \\ |x| \leq \lambda n}} \sum_{b \in B} \sqrt{\frac{\omega_{x, x+b}}{\alpha}} |\chi(\omega, x+b) - \chi(\omega, x)|. \quad (4.21)$$

Invoking the bound (4.6) we then get

$$\|R_n \mathbf{1}_{\{\lambda(\omega) \leq \lambda\}}\|_2 \leq Cn^d \quad (4.22)$$

for some constant $C = C(\alpha, \lambda, d) < \infty$. Applying Chebyshev’s inequality and summing n over powers of 2 then yields $R_n/n^\theta \rightarrow 0$ a.s. on $\{\lambda(\omega) \leq \lambda\}$. \square

Proof of Theorem 4.1(5). Let Z be a random variable satisfying the properties (a-c). By the fact that $G \in L^2_{\nabla}$, there exists a sequence $\psi_n \in L^2(\Omega)$ such that

$$\psi_n \circ \tau_x - \psi_n \xrightarrow[n \rightarrow \infty]{} \chi(\cdot, x) \quad \text{in } L^2(\Omega \times B). \quad (4.23)$$

Abbreviate $\chi_n(\omega, x) = \psi_n \circ \tau_x(\omega) - \psi_n(\omega)$ and without loss of generality assume that $\chi_n(\cdot, x) \rightarrow \chi(\cdot, x)$ almost surely.

By the fact that Z is \mathbb{P}_α -preserving we have $\mathbb{E}_\alpha(\chi_n(\cdot, Z(\cdot))) = 0$ as soon as we can show that $\chi_n(\cdot, Z(\cdot)) \in L^1(\Omega)$. It thus suffices to prove that

$$\chi_n(\cdot, Z(\cdot)) \xrightarrow[n \rightarrow \infty]{} \chi(\cdot, Z(\cdot)) \quad \text{in } L^1(\Omega). \quad (4.24)$$

Abbreviate $K(\omega) := d_\omega^{(\alpha)}(0, Z(\omega))$ and note that, as in part (4),

$$|\chi_n(\omega, Z(\omega))| \leq \sum_{\substack{x \in \mathcal{C}_{\infty, \alpha} \\ |x| \leq K(\omega)}} \sum_{b \in B} \sqrt{\frac{\omega_{x, x+b}}{\alpha}} |\chi_n(\omega, x+b) - \chi_n(\omega, x)|. \quad (4.25)$$

The quantities

$$\sqrt{\omega_{x, x+b}} |\chi_n(\omega, x+b) - \chi_n(\omega, x)| \mathbf{1}_{\{x \in \mathcal{C}_{\infty, \alpha}\}} \quad (4.26)$$

are bounded in L^2 , uniformly in x, b and n , and assumption (c) tells us that $K \in L^q$ for some $q > 3d$. Ordering the edges in \mathbb{B}_d according to their distance from the origin, Lemma 4.5 of Berger and Biskup [6] with the specific choices

$$p := 2, \quad s := q/d \quad \text{and} \quad N := d(2K + 1)^d \quad (4.27)$$

(note that $N \in L^s(\Omega)$) implies that for some $r > 1$,

$$\sup_{n \geq 1} \|\chi_n(\cdot, Z(\cdot))\|_r < \infty. \quad (4.28)$$

Hence, the family $\{\chi_n(\cdot, Z(\cdot))\}$ is uniformly integrable and (4.24) thus follows by the fact that $\chi_n(\cdot, Z(\cdot))$ converge almost surely. \square

Remark 4.5 It is worth pointing out that the proof of properties (1-3) extends nearly verbatim to the setting with arbitrary conductances and arbitrary long jumps (i.e., the case when B is simply all of \mathbb{Z}^d). One only needs that x is in $L^2(\Omega \times B)$, i.e.,

$$\mathbb{E} \left(\sum_{x \in \mathbb{Z}^d} \omega_{0,x} |x|^2 \right) < \infty. \quad (4.29)$$

The proof of (4-5) seems to require additional (and somewhat unwieldy) conditions.

5. CONVERGENCE TO BROWNIAN MOTION

Here we will prove Theorem 2.1. We commence by establishing the conclusion of Theorem 2.4 whose proof draws on an idea, suggested to us by Yuval Peres, that sublinearity

on average plus heat kernel upper bounds imply pointwise sublinearity. We have reduced the extraneous input from heat-kernel technology to the assumptions (2.17–2.18). These imply heat-kernel upper bounds but generally require less work to prove.

The main technical part of Theorem 2.1 is encapsulated into the following lemma:

Lemma 5.1 *Abusing the notation from (4.19) slightly, let*

$$R_n := \max_{\substack{x \in \mathcal{C}_{\infty, \alpha} \\ |x| \leq n}} |\psi_\omega(x)|. \quad (5.1)$$

Under the conditions (1,2,4) of Theorem 2.4, for each $\epsilon > 0$ and $\delta > 0$, there exists an a.s. finite random variable $n_0 = n_0(\omega, \epsilon, \delta)$ such that

$$R_n \leq \epsilon n + \delta R_{3n}. \quad n \geq n_0. \quad (5.2)$$

Before we prove this, let us see how this and (2.16) imply (2.19).

Proof of Theorem 2.4. Suppose that $R_n/n \not\rightarrow 0$ and pick c with $0 < c < \limsup_{n \rightarrow \infty} R_n/n$. Let θ be as in (2.16) and choose

$$\epsilon := \frac{c}{2} \quad \text{and} \quad \delta := \frac{1}{3^{\theta+1}}. \quad (5.3)$$

Note that then $c' - \epsilon \geq 3^\theta \delta c'$ for all $c' \geq c$. If $R_n \geq cn$ — which happens for infinitely many n 's — and $n \geq n_0$, then (5.2) implies

$$R_{3n} \geq \frac{c - \epsilon}{\delta} n \geq 3^\theta cn \quad (5.4)$$

and, inductively, $R_{3^k n} \geq 3^{k\theta} cn$. However, that contradicts (2.16) by which $R_{3^k n}/3^{k\theta} \rightarrow 0$ as $k \rightarrow \infty$ (with n fixed). \square

The idea underlying Lemma 5.1 is simple: We run a continuous-time random walk (Y_t) for time $t = o(n^2)$ starting from the maximizer of R_n and apply the harmonicity of $x \mapsto x + \psi_\omega(x)$ to derive an estimate on the expectation of $\psi(Y_t)$. The right-hand side of (5.2) expresses two characteristic situations that may occur at time t : Either we have $|\psi_\omega(Y_t)| \leq \epsilon n$ — which, by “sublinearity on average,” happens with overwhelming probability — or Y will not yet have left the box $[-3n, 3n]^d$ and so $\psi_\omega(Y_t) \leq R_{3n}$. The point is to show that these are the dominating strategies.

Proof of Lemma 5.1. Fix $\epsilon, \delta > 0$ and let $C_1 = C_1(\omega)$ and $C_2 = C_2(\omega)$ denote the suprema in (2.17) and (2.18), respectively. Let z be the site where the maximum R_n is achieved and denote

$$\mathcal{O}_n := \{x \in \mathcal{C}_{\infty, \alpha} : |x| \leq n, |\psi_\omega(x)| \geq \frac{1}{2}\epsilon n\}. \quad (5.5)$$

Let $Y = (Y_t)$ be a continuous-time random walk on $\mathcal{C}_{\infty, \alpha}$ with expectation for the walk started at z denoted by $E_{\omega, z}$. Define the stopping time

$$S_n := \inf\{t \geq 0 : |Y_t - z| \geq 2n\} \quad (5.6)$$

and note that, in light of Proposition 2.3, we have $|Y_{t \wedge S_n} - z| \leq 3n$ for all $t > 0$ provided $n \geq n_1(\omega)$ where $n_1(\omega) < \infty$ a.s. The harmonicity of $x \mapsto x + \psi_\omega(x)$ and the Optional Stopping Theorem yield

$$R_n \leq E_{\omega,z} |\psi_\omega(Y_{t \wedge S_n}) + Y_{t \wedge S_n} - z|. \quad (5.7)$$

Restricting to t satisfying

$$t \geq b_{4n}, \quad (5.8)$$

where $b_n = o(n^2)$ is the sequence in part (4) of Theorem 2.4, we will now estimate the expectation separately on $\{S_n < t\}$ and $\{S_n \geq t\}$.

On the event $\{S_n < t\}$, the absolute value in the expectation can simply be bounded by $R_{3n} + 3n$. To estimate the probability of $S_n < t$ we decompose according to whether $|Y_{2t} - z| \geq \frac{3}{2}n$ or not. For the former, (5.8) and (2.17) imply

$$P_{\omega,z}(|Y_{2t} - z| \geq \frac{3}{2}n) \leq \frac{E_{\omega,z}|Y_{2t} - z|}{\frac{3}{2}n} \leq \frac{2}{3}C_1 \frac{\sqrt{2t}}{n}. \quad (5.9)$$

For the latter we invoke the inclusion

$$\{|Y_{2t} - z| \leq \frac{3}{2}n\} \cap \{S_n < t\} \subset \{|Y_{2t} - Y_{S_n}| \geq \frac{1}{2}n\} \cap \{S_n < t\} \quad (5.10)$$

and note that $2t - S_n \in [t, 2t]$, (5.8) and (2.17) give us similarly

$$P_{\omega,x}(|Y_s - x| \geq n/2) \leq \frac{2}{n}C_1\sqrt{2t} \quad \text{when } x := Y_{S_n} \text{ and } s := 2t - S_n. \quad (5.11)$$

From the Strong Markov Property we thus conclude that this serves also as a bound for $P_{\omega,z}(S_n < t, |Y_{2t} - z| \geq \frac{3}{2}n)$. Combining both parts and using $\frac{8}{3}\sqrt{2} \leq 4$ we thus have

$$P_{\omega,z}(S_n < t) \leq \frac{4C_1\sqrt{t}}{n}. \quad (5.12)$$

The $S_n < t$ part of the expectation (5.7) is bounded by $R_{3n} + 3n$ times as much.

On the event $\{S_n \geq t\}$, the expectation in (5.7) is bounded by

$$E_{\omega,z}(|\psi_\omega(Y_t)| 1_{\{S_n \geq t\}}) + E_{\omega,z}|Y_t - z|. \quad (5.13)$$

The second term on the right-hand side is then less than $C_1\sqrt{t}$ provided $t \geq b_n$. The first term is estimated depending on whether $Y_t \notin \mathcal{O}_{2n}$ or not:

$$E_{\omega,z}(|\psi_\omega(Y_t)| 1_{\{S_n \geq t\}}) \leq \frac{1}{2}\epsilon n + R_{3n}P_{\omega,z}(Y_t \in \mathcal{O}_{2n}). \quad (5.14)$$

For the probability of $Y_t \in \mathcal{O}_{2n}$ we get

$$P_{\omega,z}(Y_t \in \mathcal{O}_{2n}) = \sum_{x \in \mathcal{O}_{2n}} P_{\omega,z}(Y_t = x) \quad (5.15)$$

which, in light of the Cauchy-Schwarz estimate

$$P_{\omega,z}(Y_t = x)^2 \leq P_{\omega,z}(Y_t = z)P_{\omega,x}(Y_t = x) \quad (5.16)$$

and the definition of C_2 , is further estimated by

$$P_{\omega,z}(Y_t \in \mathcal{O}_{2n}) \leq C_2 \frac{|\mathcal{O}_{2n}|}{t^{d/2}}. \quad (5.17)$$

From the above calculations we conclude that

$$R_n \leq (R_{3n} + 3n) \frac{4C_1 \sqrt{t}}{n} + C_1 \sqrt{t} + \frac{1}{2} \epsilon n + R_{3n} C_2 \frac{|\mathcal{O}_{2n}|}{t^{d/2}}. \quad (5.18)$$

Since $|\mathcal{O}_{2n}| = o(n^d)$ as $n \rightarrow \infty$, by (2.15) we can choose $t := \zeta n^2$ with $\zeta > 0$ so small that (5.8) applies and (5.2) holds for the given ϵ and δ once n is sufficiently large. \square

We now proceed to prove convergence of the random walk $X = (X_n)$ to Brownian motion. Most of the ideas are drawn directly from Berger and Biskup [6] so we stay rather brief. We will frequently work on the truncated infinite component $\mathcal{C}_{\infty,\alpha}$ and the corresponding restriction of the random walk; cf (2.11–2.13). We assume throughout that α is such that (2.8–2.9) hold.

Lemma 5.2 *Let χ be the corrector on \mathcal{C}_{∞} . Then $\varphi_{\omega}(x) := x + \chi(\omega, x)$ is harmonic for the random walk observed only on $\mathcal{C}_{\infty,\alpha}$, i.e.,*

$$\mathcal{L}_{\omega}^{(\alpha)} \varphi_{\omega}(x) = 0, \quad \forall x \in \mathcal{C}_{\infty,\alpha}. \quad (5.19)$$

Proof. We have

$$(\mathcal{L}_{\omega}^{(\alpha)} \varphi_{\omega})(x) = E_{\omega,x}(\varphi_{\omega}(X_{T_1})) - \varphi_{\omega}(x) \quad (5.20)$$

But X_n is confined to a finite component of $\mathcal{C}_{\infty} \setminus \mathcal{C}_{\infty,\alpha}$ for $n \in [0, T_1]$, and so $\varphi_{\omega}(X_n)$ is bounded. Since $(\varphi_{\omega}(X_n))$ is a martingale and T_1 is an a.s. finite stopping time, the Optional Stopping Theorem tells us $E_{\omega,x} \varphi_{\omega}(X_{T_1}) = \varphi_{\omega}(x)$. \square

Next we recall the proof of sublinearity of the corrector along coordinate directions:

Lemma 5.3 *For $\omega \in \{0 \in \mathcal{C}_{\infty,\alpha}\}$, let $(x_n(\omega))_{n \in \mathbb{Z}}$ mark the intersections of $\mathcal{C}_{\infty,\alpha}$ and one of the coordinate axis so that $x_0(\omega) = 0$. Then*

$$\lim_{n \rightarrow \infty} \frac{\chi(\omega, x_n(\omega))}{n} = 0, \quad \mathbb{P}_{\alpha}\text{-a.s.} \quad (5.21)$$

Proof. Let τ_x be the “shift by x ” on Ω and let $\sigma(\omega) := \tau_{x_1(\omega)}(\omega)$ denote the “induced” shift. Standard arguments (cf [6, Theorem 3.2]) prove that σ is \mathbb{P}_{α} preserving and ergodic. Moreover,

$$\mathbb{E}_{\alpha}(d_{\omega}^{(\alpha)}(0, x_1(\omega))^p) < \infty, \quad p < \infty, \quad (5.22)$$

by [6, Lemma 4.3] (based on Antal and Pisztora [1]). Define $\Psi(\omega) := \chi(\omega, x_1(\omega))$. Theorem 4.1(5) tells us that

$$\Psi \in L^1(\mathbb{P}_{\alpha}) \quad \text{and} \quad \mathbb{E}_{\alpha} \Psi(\omega) = 0. \quad (5.23)$$

But the gradient property of χ implies

$$\frac{\chi(\omega, x_n(\omega))}{n} = \frac{1}{n} \sum_{k=0}^{n-1} \Psi \circ \sigma^k(\omega) \quad (5.24)$$

and so the left-hand side tends to zero a.s. by the Pointwise Ergodic Theorem. \square

We will also need sublinearity of the corrector on average:

Lemma 5.4 *For each $\epsilon > 0$ and \mathbb{P}_α -a.e. ω :*

$$\lim_{n \rightarrow \infty} \frac{1}{n^d} \sum_{\substack{x \in \mathcal{C}_{\infty, \alpha} \\ |x| \leq n}} \mathbf{1}_{\{|\chi(\omega, x)| \geq \epsilon n\}} = 0. \quad (5.25)$$

Proof. This follows from Lemma 5.3 exactly as [6, Theorem 5.4]. \square

Remark 5.5 The proof of [6, Theorem 5.4] makes a convenient use of separate ergodicity (i.e., that with respect to shifts only in one of the coordinate directions). This is sufficient for i.i.d. environments as considered in the present situation. However, it is not hard to come up with a modification of the proof that covers general ergodic environments as well (Biskup and Deuschel [8]).

Finally, we will assert the validity of the bounds on the return probability and expected displacement of the walk from Theorem 2.4:

Lemma 5.6 *Let (Y_t) denote the continuous-time random walk on $\mathcal{C}_{\infty, \alpha}$. Then the diffusive bounds (2.17–2.18) hold for \mathbb{P}_α -a.e. ω .*

We will prove this lemma at the very end of Sect. 6.

Proof of Theorem 2.1. Let α be such that (2.8–2.9) hold and let χ denote the corrector on \mathcal{C}_∞ as constructed in Theorem 4.1. The crux of the proof is to show that χ grows sublinearly with x , i.e., $\chi(\omega, x) = o(|x|)$ a.s.

By Lemmas 5.2 and 5.4, Theorem 4.1(4) and Lemma 5.6, the corrector satisfies the conditions of Theorem 2.4. It follows that χ is sublinear on $\mathcal{C}_{\infty, \alpha}$ as stated in (2.19). However, by (2.10) the largest component of $\mathcal{C}_\infty \setminus \mathcal{C}_{\infty, \alpha}$ in a box $[-2n, 2n]$ is less than $C \log n$ in diameter, for some random but finite $C = C(\omega)$. Invoking the harmonicity of φ_ω on \mathcal{C}_∞ , the Optional Stopping Theorem gives

$$\max_{\substack{x \in \mathcal{C}_\infty \\ |x| \leq n}} |\chi(\omega, x)| \leq \max_{\substack{x \in \mathcal{C}_{\infty, \alpha} \\ |x| \leq n}} |\chi(\omega, x)| + 2C(\omega) \log(2n), \quad (5.26)$$

whereby we deduce that χ is sublinear on \mathcal{C}_∞ as well.

Having proved the sublinearity of χ on \mathcal{C}_∞ , we proceed as in the $d = 2$ proof of [6]. Let $\varphi_\omega(x) := x + \chi(\omega, x)$ and abbreviate $M_n := \varphi_\omega(X_n)$. Fix $\hat{v} \in \mathbb{R}^d$ and define

$$f_K(\omega) := E_{\omega, 0}((\hat{v} \cdot M_1)^2 \mathbf{1}_{\{|\hat{v} \cdot M_1| \geq K\}}). \quad (5.27)$$

By Theorem 4.1(3), $f_K \in L^1(\Omega, \mathcal{F}, \mathbb{P}_0)$ for all K . Since the Markov chain on environments, $n \mapsto \tau_{X_n}(\omega)$, is ergodic (cf [6, Section 3]), we thus have

$$\frac{1}{n} \sum_{k=0}^{n-1} f_K \circ \tau_{X_k}(\omega) \xrightarrow[n \rightarrow \infty]{} \mathbb{E}_0 f_K, \quad (5.28)$$

for \mathbb{P}_0 -a.e. ω and $P_{\omega,0}$ -a.e. path $X = (X_k)$ of the random walk. Using this for $K := 0$ and $K := \varepsilon\sqrt{n}$ along with the monotonicity of $K \mapsto f_K$ verifies the conditions of the Lindeberg-Feller Martingale Functional CLT (e.g., Durrett [13, Theorem 7.7.3]). Thereby we conclude that the random continuous function

$$t \mapsto \frac{1}{\sqrt{n}} (\hat{v} \cdot M_{\lfloor nt \rfloor} + (nt - \lfloor nt \rfloor) \hat{v} \cdot (M_{\lfloor nt \rfloor + 1} - M_{\lfloor nt \rfloor})) \quad (5.29)$$

converges weakly to Brownian motion with mean zero and covariance

$$\mathbb{E}_0 f_0 = \mathbb{E}_0 E_{\omega,0} ((\hat{v} \cdot M_1)^2). \quad (5.30)$$

This can be written as $\hat{v} \cdot D \hat{v}$ where D is the matrix with coefficients

$$D_{i,j} := \mathbb{E}_0 E_{\omega,0} ((\hat{e}_i \cdot M_1)(\hat{e}_j \cdot M_1)). \quad (5.31)$$

Invoking the Cramér-Wold device (e.g., Durrett [13, Theorem 2.9.5]) and the fact that continuity of a stochastic process in \mathbb{R}^d is implied by the continuity of its d one-dimensional projections we get that the linear interpolation of $t \mapsto M_{\lfloor nt \rfloor} / \sqrt{n}$ scales to d -dimensional Brownian motion with covariance matrix D . The sublinearity of the corrector then ensures, as in [6, (6.11–6.13)], that

$$X_n - M_n = \chi(\omega, X_n) = o(|X_n|) = o(|M_n|) = o(\sqrt{n}), \quad (5.32)$$

and so the same conclusion applies to $t \mapsto B_n(t)$ in (2.2).

The reflection symmetry of \mathbb{P}_0 forces D to be diagonal; the rotation symmetry then ensures that $D = (1/d)\sigma^2 1$ where

$$\sigma^2 := \mathbb{E}_0 E_{\omega,0} |M_1|^2 \quad (5.33)$$

To see that the limiting process is not degenerate to zero we note that if we had $\sigma = 0$ then $\chi(\cdot, x) = -x$ would hold a.s. for all $x \in \mathbb{Z}^d$. But that is impossible since, as we proved above, $x \mapsto \chi(\cdot, x)$ is sublinear a.s. \square

Remark 5.7 Note that, unlike the proofs in [29, 6, 25], the above line of argument does not require a separate proof of tightness. In our approach, this comes rather automatically for the deformed random walk $\varphi_\omega(X_n)$ — via the (soft) stationarity argument (5.28) and the Martingale Functional CLT. Sublinearity of the corrector then extends it readily to the original random walk.

Remark 5.8 We also wish to use the opportunity to correct an erroneous argument from [6]. There, at the end of the proof of Theorem 6.2 it is claimed that the expectation $\mathbb{E}_0 E_{\omega,0}(X_1 \cdot \chi(X_1, \omega))$ is zero. Unfortunately, this is false. In fact, we have

$$\mathbb{E}_0 E_{\omega,0}(X_1 \cdot \chi(X_1, \omega)) = -\mathbb{E}_0 E_{\omega,0} |\chi(X_1, \omega)|^2 < 0. \quad (5.34)$$

where the strict inequality assumes that \mathbb{P} is non-degenerate. This shows

$$\mathbb{E}_0 E_{\omega,0} |M_1|^2 = \mathbb{E}_0 E_{\omega,0} |X_1|^2 - \mathbb{E}_0 E_{\omega,0} |\chi(X_1, \omega)|^2 < \mathbb{E}_0 E_{\omega,0} |X_1|^2. \quad (5.35)$$

Thus, once \mathbb{P} is non-degenerate, the diffusion constant of the limiting Brownian motion is strictly *smaller* than the variance of the first step.

A consequence of the above error for the proof of Theorem 6.2 in [6] is that it invalidates one of the three listed arguments to prove that the limiting Brownian motion is non-degenerate. Fortunately, the remaining two arguments are correct.

6. HEAT KERNEL AND EXPECTED DISTANCE

Here we will derive the bounds (2.17–2.18) and thus establish Lemma 5.6. Most of the derivation will be done for a general countable-state Markov chain; we will specialize to random walk among i.i.d. conductances at the very end of this section. The general ideas underlying these derivations are fairly standard and exist, in some form, in the literature. A novel aspect is the way we control the non-uniformity of volume-growth caused by local irregularities of the underlying graph; cf (6.4) and Lemma 6.3(1). A well informed reader may nevertheless wish to read only the statements of Propositions 6.1 and 6.2 and then pass directly to the proof of Lemma 5.6.

Let V be a countable set and let $(a_{xy})_{x,y \in V}$ denote the collection of positive numbers with the following properties: For all $x, y \in V$,

$$a_{xy} = a_{yx} \quad \text{and} \quad \pi(x) := \sum_{y \in V} a_{xy} < \infty. \quad (6.1)$$

Consider a continuous time Markov chain (Y_t) on V with the generator

$$(\mathcal{L}f)(x) := \frac{1}{\pi(x)} \sum_{y \in V} a_{xy} [f(y) - f(x)]. \quad (6.2)$$

We use P^x to denote the law of the chain started from x , and E^x to denote the corresponding expectation. Consider a graph $G = (V, E)$ where E is the set of all pairs (x, y) such that $a_{xy} > 0$. Let $d(x, y)$ denote the distance between x and y as measured on G .

For each $x \in V$, let $B_n(x) := \{y \in V : d(x, y) \leq n\}$. If $\Lambda \subset V$, we use $Q(\Lambda, \Lambda^c)$ to denote the sum

$$Q(\Lambda, \Lambda^c) := \sum_{x \in \Lambda} \sum_{y \in \Lambda^c} a_{xy}. \quad (6.3)$$

Suppose that there are constants $d > 0$ and $\nu \in (0, 1/2)$ such that, for some $a > 0$,

$$C_{\text{vol}}(x, a) := \sup_{0 < s \leq a} \left[s^d \sum_{y \in V} \pi(y) e^{-sd(x,y)} \right] < \infty \quad (6.4)$$

and

$$C_{\text{iso}}(x) := \inf_{n \geq 1} \inf \left\{ \frac{Q(\Lambda, \Lambda^c)}{\pi(\Lambda)^{\frac{d-1}{d}}} : \Lambda \subset B_{2n}(x), \pi(\Lambda) \geq n^\nu, \Lambda \text{ connected} \right\} > 0. \quad (6.5)$$

Let $V(\epsilon) \subset V$ denote the set of all $x \in V$ that are connected to infinity by a self-avoiding path $(x_0 = x, x_1, \dots)$ with $a_{x_i, x_{i+1}} \geq \epsilon$ for all $i \geq 0$. Suppose that

$$a_* := \sup\{\epsilon > 0: V(\epsilon) = V\} > 0. \quad (6.6)$$

(Note that this does not require a_{xy} be bounded away from zero.)

The first observation is that the heat-kernel, defined by

$$q_t(x, y) := \frac{P^x(Y_t = y)}{\pi(y)}, \quad (6.7)$$

can be bounded in terms of the isoperimetry constant $C_{\text{iso}}(x)$. Bounds of this form are well known and have been derived by, e.g., Coulhon, Grigor'yan and Pittet [10] for heat-kernel on manifolds, and by Lovász and Kannan [22], Morris and Peres [23] and Goel, Montenegro and Tetali [16] in the context of countable-state Markov chains. We will use the formulation for infinite graphs developed in Morris and Peres [23].

Proposition 6.1 *There exists a constant $c_1 \in (1, \infty)$ depending only on d and a_* such that for $t(x) := c_1[\log(C_{\text{iso}}(x) \vee c_1)]^{\frac{1}{1-2v}}$ we have*

$$\sup_{z \in B_t(x)} \sup_{y \in V} q_t(z, y) \leq c_1 \frac{C_{\text{iso}}(x)^{-d}}{t^{d/2}}, \quad t \geq t(x). \quad (6.8)$$

Proof. We will first derive the corresponding bound for the discrete-time version of (Y_t) . Let $P(x, y) := a_{xy}/\pi(x)$ and define $\hat{P} := \frac{1}{2}(1 + P)$. Let

$$\hat{q}_n(x, y) := \frac{\hat{P}^n(x, y)}{\pi(y)} \quad (6.9)$$

We claim that, for some absolute constant c_1 and any $z \in B_n(x)$,

$$\hat{q}_n(z, y) \leq c_1 \frac{C_{\text{iso}}(x)^{-d}}{n^{d/2}}, \quad n \geq t(x). \quad (6.10)$$

To this end, let \hat{Q} be the object Q for the Markov chain \hat{P} and let

$$\phi(r) := \inf\left\{\frac{\hat{Q}(\Lambda, \Lambda^c)}{\pi(\Lambda)} : \pi(\Lambda) \leq r, \Lambda \subset B_{2n}(x)\right\}. \quad (6.11)$$

We claim that Theorem 2 of Morris and Peres [23] then implies that, for any ϵ that satisfies $0 < \epsilon < [\pi(x) \wedge \pi(y)]^{-1}$ and

$$n \geq 1 + \int_{4(\pi(z) \wedge \pi(y))}^{4/\epsilon} \frac{4dr}{r\phi(r)^2}, \quad (6.12)$$

we have $\hat{q}_n(z, y) \leq \epsilon$. To see this we have to check that the restriction $\Lambda \subset B_{2n}(x)$ in the definition of $\phi(r)$, which is absent from the corresponding object in [23], causes no harm. First note that the Markov chain started at $z \in B_n(x)$ will not leave $B_{2n}(x)$ by time n . Thus, we can modify the chain outside $B_{2n}(x)$ arbitrarily. It is easy to come up with a modification that will effectively reduce the infimum in (6.11) to sets inside $B_{2n}(x)$.

It is well known (and easy to check) that the infimum in (6.11) can be restricted to connected sets Λ . Then (6.5–6.6) give us

$$\phi(r) \geq \frac{1}{2} (C_{\text{iso}}(x) r^{-1/d} \wedge a_* n^{-\nu}) \quad (6.13)$$

where the extra half arises due the consideration of time-delayed chain $\hat{\mathbb{P}}$. The two regimes cross over at $r_n := (C_{\text{iso}}(x)/a_*)^d n^{d\nu}$; the integral is thus bounded by

$$\int_{4(\pi(z) \wedge \pi(y))}^{4/\epsilon} \frac{4dr}{r\phi(r)^2} \leq 4 \frac{n^{2\nu}}{a_*^2} \log\left(\frac{r_n}{4a_*}\right) + 2d C_{\text{iso}}(x)^{-2} \left(\frac{4}{\epsilon}\right)^{2/d}. \quad (6.14)$$

The first term splits into a harmless factor of order $n^{2\nu} \log n = o(n)$ and a term proportional to $n^{2\nu} \log C_{\text{iso}}(x)$. This is $O(n)$ by $n \geq t(x)$ where the (implicit) constant can be made as small as desired by choosing c_1 sufficiently large. Setting

$$\epsilon := c[C_{\text{iso}}(x)^2 n]^{-d/2} \quad (6.15)$$

we can thus adjust the constant c in such a way that (6.14) is less than $n - 1$ for all $n \geq t(x)$. Thereby (6.10) follows.

To extend the bound (6.10) to continuous time, we note that $\mathcal{L} = 2(\hat{\mathbb{P}} - 1)$. Thus if N_t is Poisson with parameter $2t$, then

$$q_t(z, y) = E\hat{q}_{N_t}(z, y). \quad (6.16)$$

But $P(N_t \leq \frac{3}{2}t \text{ or } N_t \geq 3t)$ is exponentially small in t and, in particular, much smaller than (6.8) for $t \geq c_1 \log C_{\text{iso}}(x)$ with c_1 sufficiently large. As $q_t \leq (a_*)^{-1}$, the $N_t \notin (\frac{3}{2}t, 3t)$ portion of the expectation in (6.16) is thus negligible. Once N_t is constrained to the interval $(t, 3t)$ the uniform bound (6.10) implies (6.8). \square

Our next item of business is a diffusive bound on the expected (graph-theoretical) distance traveled by the walk Y_t by time t . As was noted by Bass [4] and Nash [27], this can be derived from the above uniform bound on the heat-kernel assuming regularity of the volume growth. Our proof is an adaptation of an argument of Barlow [2].

Proposition 6.2 *There exist constants $c_2 = c_2(d)$ and $c_3 = c_3(d)$ such that the following holds: Let $x \in V$ and suppose $A > 0$ and $t(x) > 1$ are numbers for which*

$$\sup_{y \in V} q_t(x, y) \leq \frac{A}{t^{d/2}}, \quad t \geq t(x), \quad (6.17)$$

holds and let $T(x) := \frac{1}{d}(Aa_)^{-4/d} \vee [t(x) \log t(x)]$. Then*

$$E^x d(x, Y_t) \leq A'(x, t) \sqrt{t}, \quad t \geq T(x), \quad (6.18)$$

with $A'(x, t) := c_2 + c_3[\log A + C_{\text{vol}}(x, t^{-1/2})]$.

Much of the proof boils down to the derivation of rather inconspicuous but deep relations (discovered by Nash [27]) between the following quantities:

$$M(x, t) := E^x d(x, Y_t) = \sum_y \pi(y) q_t(x, y) d(x, y) \quad (6.19)$$

and

$$Q(x, t) := -E^x \log q_t(x, Y_t) = - \sum_y \pi(y) q_t(x, y) \log q_t(x, y). \quad (6.20)$$

Note that $q_t(x, \cdot) \leq (a_\star)^{-1}$ implies $Q(x, t) \geq \log a_\star$.

Lemma 6.3 For all $t \geq 0$ and all $x \in V$,

- (1) $M(x, t)^d \geq \exp\{-1 - C_{\text{vol}}(x, M(x, t)^{-1}) + Q(x, t)\}$,
- (2) $M'(x, t)^2 \leq Q'(x, t)$.

Proof. Part (2) is identical to the proof of Lemma 3.3 in Barlow [2] so we only need to prove part (1). We will follow the proof of [2, Lemma 3.3] except that we need to keep track of $C_{\text{vol}}(x)$ in the calculation. Pick two numbers $a > 0$ and $b \in \mathbb{R}$ and note that

$$u \log u + \lambda u \geq -e^{-\lambda-1}. \quad (6.21)$$

This implies

$$-Q(x, t) + aM(x, t) + b \geq - \sum_y \pi(y) e^{-b-1-ad(x,y)}. \quad (6.22)$$

Using the definition of $C_{\text{vol}}(x, a)$ and bounding $e^{-1} \leq 1$ we get

$$-Q(x, t) + aM(x, t) + b \geq -C_{\text{vol}}(x, a) e^{-b} a^{-d}. \quad (6.23)$$

Now set $e^{-b} := a^d$ with $a := M(x, t)^{-1}$ to get the result. \square

These bounds imply the desired diffusive estimate on $M(x, t)$:

Proof of Proposition 6.2. Suppose without loss of generality that $M(x, t) \geq \sqrt{t}$, because otherwise there is nothing to prove. We follow the proof of [2, Proposition 3.4]. The key input is provided by the inequalities in Lemma 6.3. Define the function

$$L(t) := \frac{1}{d} \left(Q(x, t) + \log A - \frac{d}{2} \log t \right) \quad (6.24)$$

and note that $L(t) \geq 0$ for $t \geq t(x)$. Let

$$t_0 := (Aa_\star)^{-2/d} \vee \sup\{t \geq 0: L(t) \leq 0\}. \quad (6.25)$$

We claim that $M(x, t_0) \leq \sqrt{dT(x)}$. Indeed, when $t_0 = (Aa_\star)^{-2/d}$ then this follows by

$$M(x, t_0) \leq t_0 = (Aa_\star)^{-2/d} \leq \sqrt{dT(x)} \quad (6.26)$$

due to our choice of $T(x)$. On the other hand, when $t_0 > (Aa_\star)^{-2/d}$ we use Lemma 6.3(2), the Fundamental Theorem of Calculus and the Cauchy-Schwarz inequality to derive

$$M(x, t_0) \leq \sqrt{t_0} [Q(x, t_0) - Q(x, 0)]^{1/2}. \quad (6.27)$$

Since $Q(x, 0) \geq \log a_\star$ and $L(t_0) = 0$ by continuity, we have

$$M(x, t_0) \leq \sqrt{t_0} \left(\frac{d}{2} \log t_0 - \log A - \log a_\star \right)^{1/2} \leq \sqrt{dt_0 \log t_0} \quad (6.28)$$

where we used that $t_0 \geq (Aa_\star)^{-2/d}$ implies $\log A + \log a_\star \geq -\frac{d}{2} \log t_0$. Since this implies $t_0 \geq 1$, the condition $t_0 \leq t(x)$ shows that the right-hand side is again less than $\sqrt{dT(x)}$.

For $t \geq t_0$ we have $L(t) \geq 0$. Lemma 6.3(2) yields

$$\begin{aligned} M(x, t) - M(x, t_0) &\leq \sqrt{d} \int_{t_0}^t \left(\frac{1}{2s} + L'(s) \right)^{1/2} ds \\ &\leq \sqrt{d} \int_{t_0}^t \left(\frac{1}{\sqrt{2s}} + L'(s) \sqrt{s/2} \right) ds \leq \sqrt{2dt} + L(t) \sqrt{dt}, \end{aligned} \quad (6.29)$$

where we used integration by parts and the positivity of L to derive the last inequality. Now put this together with $M(x, t_0) \leq \sqrt{dt}$ and apply Lemma 6.3(1), noting that $C_{\text{vol}}(z, M(x, t)^{-1}) \leq C_{\text{vol}}(z, t^{-1/2})$ by the assumption $M(x, t) \geq \sqrt{t}$. Dividing out an overall factor \sqrt{t} , we thus get

$$\left[A e^{C_{\text{vol}}(x, t^{-1/2})} \right]^{-1/d} e^{-1/d + L(t)} \leq 3\sqrt{d} + \sqrt{d} L(t). \quad (6.30)$$

This implies

$$L(t) \leq \tilde{c}_2 + \tilde{c}_3 [\log A + C_{\text{vol}}(x, t^{-1/2})] \quad (6.31)$$

for some constants \tilde{c}_2 and \tilde{c}_3 depending only on d . Plugging this in (6.29), we get the desired claim. \square

We are now finally ready to complete the proof of our main theorem:

Proof of Lemma 5.6. We will apply the above estimates to obtain the proof of the bounds (2.17–2.18). We use the following specific choices

$$V := \mathcal{C}_{\infty, \alpha}, \quad a_{xy} := \hat{\omega}_{xy}, \quad \pi(x) := 2d, \quad \text{and} \quad b_n := n. \quad (6.32)$$

As $a_\star \geq \alpha$, all required assumptions are satisfied.

To prove (2.18), we note that by Lemma 3.4 of Berger, Biskup, Hoffman and Kozma [7] (based on the isoperimetric inequality for the supercritical bond-percolation cluster, cf Benjamini and Mossel [5], Rau [28, Proposition 1.2] or [7, Section 5]) we have $C_{\text{iso}}(0) > 0$ a.s. Hence, Proposition 6.1 ensures that, for all $z \in \mathcal{C}_{\infty, \alpha}$ with $|z| \leq t$,

$$t^{d/2} P_{\omega, z}(Y_t = z) \leq 2dc_1 C_{\text{iso}}(0)^{-d} \quad (6.33)$$

provided t exceeds some t_1 depending on $C_{\text{iso}}(0)$. From here (2.18) immediately follows.

To prove (2.17), we have to show that, a.s.,

$$\sup_{n \geq 1} \max_{\substack{z \in \mathcal{C}_{\infty, \alpha} \\ |z| \leq n}} \sup_{t \geq n} C_{\text{vol}}(z, t^{-1/2}) < \infty. \quad (6.34)$$

To this end we note that Lemma 3.1 implies that there is a.s. finite $C = C(\omega)$ such that for all $z, y \in \mathcal{C}_{\infty, \alpha}$ with $|z| \leq n$ and $|z - y| \geq C \log n$,

$$d(z, y) \geq \varrho |z - y|. \quad (6.35)$$

It follows that, once $1/a > C \log n$, for every $z \in \mathcal{C}_{\infty, \kappa}$ with $|z| \leq n$ we have

$$\sum_{y \in \mathcal{C}_{\infty, \kappa}} e^{-ad(z,y)} \leq c_4 a^{-d} + \sum_{\substack{y \in \mathcal{C}_{\infty, \kappa} \\ |y-z| \geq 1/a}} e^{-a\varrho|z-y|} \leq c_5 a^{-d}, \quad (6.36)$$

where c_4 and c_5 are constants depending on d and ϱ . Since $1/a = t^{1/2} \geq \sqrt{n} \gg \log n$, (6.34) follows.

Once we have the uniform bound (6.34), as well as the uniform bound (6.17), Proposition 6.2 yields the a.s. inequality

$$\sup_{n \geq 1} \max_{\substack{z \in \mathcal{C}_{\infty, \kappa} \\ |z| \leq n}} \sup_{t \geq n} \frac{E_{\omega, z} d(z, Y_t)}{\sqrt{t}} < \infty. \quad (6.37)$$

To convert $d(z, Y_t)$ into $|z - Y_t|$ in the expectation, we invoke (6.35) one more time. \square

ACKNOWLEDGMENTS

The research of M.B. was supported by the NSF grant DMS-0505356. The authors wish to thank N. Berger, M. Barlow, J.-D. Deuschel, C. Hoffman, G. Kozma, Y. Peres and G. Pete for discussions on this problem.

REFERENCES

- [1] P. Antal and A. Pisztora (1996). On the chemical distance for supercritical Bernoulli percolation. *Ann. Probab.* **24**, no. 2, 1036–1048.
- [2] M.T. Barlow (2004). Random walks on supercritical percolation clusters. *Ann. Probab.* **32**, no. 4, 3024–3084.
- [3] M.T. Barlow and J.-D. Deuschel (2007). Quenched invariance principle for the random conductance model with unbounded conductances. In preparation.
- [4] R.F. Bass (2002). On Aronsen’s upper bounds for heat kernels. *Bull. London Math. Soc.* **34** 415–419.
- [5] I. Benjamini and E. Mossel (2003). On the mixing time of a simple random walk on the super critical percolation cluster. *Probab. Theory Rel. Fields* **125**, no. 3, 408–420.
- [6] N. Berger and M. Biskup (2007). Quenched invariance principle for simple random walk on percolation clusters. *Probab. Theory Rel. Fields* **137**, no. 1-2, 83–120.
- [7] N. Berger, M. Biskup, C.E. Hoffman and G. Kozma (2006). Anomalous heat-kernel decay for random walk among bounded random conductances. *Ann. Inst. Henri Poincaré* (to appear).
- [8] M. Biskup and J.-D. Deuschel, in preparation.
- [9] M. Biskup and H. Spohn (2007). Scaling limit for a class of gradient fields with non-convex potentials. Preprint (arxiv:0704.3086).
- [10] T. Coulhon, A. Grigor’yan and C. Pittet (2001). A geometric approach to on-diagonal heat kernel lower bound on groups. *Ann. Inst. Fourier* **51** 1763–1827.
- [11] A. De Masi, P.A. Ferrari, S. Goldstein and W.D. Wick (1985). Invariance principle for reversible Markov processes with application to diffusion in the percolation regime. In: *Particle Systems, Random Media and Large Deviations (Brunswick, Maine)*, pp. 71–85, *Contemp. Math.*, **41**, AMS, Providence, RI.
- [12] A. De Masi, P.A. Ferrari, S. Goldstein and W.D. Wick (1989). An invariance principle for reversible Markov processes. Applications to random motions in random environments. *J. Statist. Phys.* **55**, no. 3–4, 787–855.
- [13] R. Durrett (1996). *Probability Theory and Examples* (Third edition), Duxbury Press, Belmont, CA.

- [14] L.R.G. Fontes and P. Mathieu (2006). On symmetric random walks with random conductances on \mathbb{Z}^d . *Probab. Theory Rel. Fields* **134**, no. 4, 565–602.
- [15] G. Giacomin, S. Olla and H. Spohn (2001). Equilibrium fluctuations for $\nabla\phi$ interface model, *Ann. Probab.* **29**, no. 3, 1138–1172.
- [16] S. Goel, R. Montenegro and P. Tetali (2006). Mixing time bounds via the spectral profile. *Electron. J. Probab.* **11**, paper no. 1, 1–26.
- [17] G.R. Grimmett (1999). *Percolation* (Second edition), Grundlehren der Mathematischen Wissenschaften, vol. 321. Springer-Verlag, Berlin.
- [18] G.R. Grimmett and J.M. Marstrand (1990). The supercritical phase of percolation is well behaved, *Proc. Roy. Soc. London Ser. A* **430**, no. 1879, 439–457.
- [19] H. Kesten and Y. Zhang (1990). The probability of a large finite cluster in supercritical Bernoulli percolation. *Ann. Probab.* **18**, no. 2, 537–555.
- [20] C. Kipnis, and S.R.S. Varadhan (1986). A central limit theorem for additive functionals of reversible Markov processes and applications to simple exclusions. *Commun. Math. Phys.* **104**, no. 1, 1–19.
- [21] T.M. Liggett, R.H. Schonmann and A.M. Stacey (1997). Domination by product measures. *Ann. Probab.* **25**, no. 1, 71–95.
- [22] L. Lovász and R. Kannan (1999). Faster mixing via average conductance. *Proc. of the 31st Annual ACM Symp. on Theory of Computing*, 282–287.
- [23] B. Morris and Y. Peres (2005). Evolving sets, mixing and heat kernel bounds. *Probab. Theory Rel. Fields* **133**, no. 2, 245–266.
- [24] P. Mathieu (2006). Quenched invariance principles for random walks with random conductances, *J. Statist. Phys.* (to appear).
- [25] P. Mathieu and A.L. Piatnitski (2007). Quenched invariance principles for random walks on percolation clusters. *Proc. Roy. Soc. A* **463**, 2287–2307.
- [26] P. Mathieu and E. Remy (2004). Isoperimetry and heat kernel decay on percolation clusters. *Ann. Probab.* **32**, no. 1A, 100–128.
- [27] J. Nash (1958). Continuity of solutions of parabolic and elliptic equations. *Amer. J. Math.* **80** 931–954.
- [28] C. Rau (2006). Sur le nombre de points visités par une marche aléatoire sur un amas infini de percolation, *Bull. Soc. Math. France* (to appear).
- [29] V. Sidoravicius and A.-S. Sznitman (2004). Quenched invariance principles for walks on clusters of percolation or among random conductances. *Probab. Theory Rel. Fields* **129**, no. 2, 219–244.