

On the absence of ferromagnetism in typical 2D ferromagnets

Marek Biskup¹, Lincoln Chayes¹, Steven A. Kivelson²

¹ Department of Mathematics, UCLA, Los Angeles, CA 90095-1555, U.S.A.

² Department of Physics, Stanford University, Stanford, CA 94305-4045, U.S.A.

Abstract: We consider the Ising systems in d dimensions with nearest-neighbor ferromagnetic interactions and long-range repulsive (antiferromagnetic) interactions that decay with power s of the distance. The physical context of such models is discussed; primarily this is $d = 2$ and $s = 3$ where, at long distances, genuine magnetic interactions between genuine magnetic dipoles are of this form. We prove that when the power of decay lies above d and does not exceed $d + 1$, then for all temperatures the spontaneous magnetization is zero. In contrast, we also show that for powers exceeding $d + 1$ (with $d \geq 2$) magnetic order can occur.

1. Introduction

While most of our knowledge of statistical mechanics is derived from studies of model problems with short-range forces, in nature interactions more often fall off only in proportion to an inverse power of the distance, $U(r) \sim 1/r^s$. This includes systems interacting via Coulomb forces ($s = 1$), dipolar interactions ($s = 3$) as well as interactions caused by collective effects such as strain induced interactions in solids or the effective entropic interactions (analogous to Casimir forces) in lipid films. When the interactions are sufficiently long-range, i.e., when $s \leq d$ where d is the spatial dimension, the very definition of the thermodynamic limit is different than for short-ranged models. However, even when $s > d$ there can be qualitatively new, or at least unexpected, phenomena, cf, e.g., [2–4, 29].

In the present paper we study a class of systems with long-range forces; namely, the Ising models on \mathbb{Z}^d , $d \geq 1$, which are defined by the (formal) Hamiltonians

$$H = - \sum_{(i,j)} J \sigma_i \sigma_j + \frac{1}{2} \sum_{i,j} K_{i,j} \sigma_i \sigma_j. \quad (1.1)$$

Here $\sigma_i \in \{+1, -1\}$, i and j index sites in \mathbb{Z}^d and $\langle i, j \rangle$ denotes a nearest neighbor pair. The above notation expresses the relevant signs of all the couplings: $J > 0$ is the short-range *ferromagnetic* interaction while $K_{i,j} \geq 0$ represents the *antiferromagnetic* long range interaction which we assume decays with power s of the distance between i and j . We investigate the question of presence, and absence, of spontaneous magnetization in such models.

The motivation for this work was provided by a paper of Spivak and one of us [27] where it was conjectured that, in the presence (or absence) of an external field, discontinuous transitions permitting coexisting states of different magnetization are forbidden for antiferromagnetic power law interactions with range $d < s \leq d + 1$. A heuristic proof by contradiction was presented based on the explicit construction of a “micro-emulsion” phase which has a lower free energy than the state of macroscopic two-phase coexistence. Simply put, the anticipated surface tension between the two pure phases would be negative—and divergent. The proof is heuristic in the sense that it makes the physically plausible assumption that correlations in the putative coexisting phases have reasonable decay and that there is a well defined interface.

As it turns out, versions of the above conjecture are actually more than 20 years old. For example, on the physics side, modulated phases in 2D dipolar ferromagnets were analyzed in [1, 11, 18]. On the mathematics side, in [8], models with extreme anisotropic repulsive interactions which have *very* slow decay, but only among a sparse set of spins, were considered and absence of spontaneous magnetism was proved. The isotropic case, $U(r) \sim 1/r^s$, was also mentioned in [8] and the significance of the interval $d < s < d + 1$ for the absence of magnetization was highlighted (with no mention of $s = d + 1$). Related problems were described in [17] for systems with longer range, e.g., Coulomb, interactions and in [8, 9] for the current setup with $O(n)$ -spins. Furthermore, general theorems demonstrating instability of phase coexistence under the addition of generic long-range interactions have been proved in [7, 16, 26]. In the present paper we provide a full proof of the absence of ferromagnetism in the model (1.1) with $d < s \leq d + 1$ thereby vindicating completely the arguments of [27]—at least for $h = 0$.

The mathematical result presented in this note has the following consequence for 2D physics: Two-dimensional magnetic systems often have strong “crystal field” effects which orient the electron spins (largely or entirely) in the z direction, perpendicular to the plane in which they reside. This gives the problem of magnetic ordering an Ising character. Interactions between nearby spins—quantum mechanical and somewhat complicated—are, often enough, of the ferromagnetic type and considerably stronger than the direct magnetic dipolar interactions (which are a relativistic effect). Thus, it seems reasonable to study Ising ferromagnets in 2D contexts and conclude that there is a definitive possibility for ferromagnetism. However, while possibly weak, there is *always* the long-range $1/r^3$ repulsive interaction. The conclusion of this note is that, no matter how small its relative strength may be, this interaction will preclude the possibility of ferromagnetism among the z -components.

We remark that the absence of magnetization certainly does not disallow other types of ordering. Indeed, a large body of physics literature [1, 5, 6, 10, 11, 14, 18, 19, 21–23, 27], points in the direction of modulated (striped and/or bubble) states in this and related systems. (For an extremely insightful review of the phases produced by models of this sort and many experimentally clear realizations of the corresponding physics, see [24].) From the perspective of mathematics, recent rigorous estimates on ground-state energies [13], which are asymptotic in $d \geq 2$ and exact in $d = 1$, also indicate

striped order in the ground state. In fact, for certain special cases of the 1D ground-state problem, this has been established completely.

The organization of the rest of this paper is as follows: In the next section we define all necessary background and state the main results. In Sect. 3 we derive some estimates on the strength of the long-range interaction between a box and its complement. These are assembled into the proof of the main result in Sect. 4. Sect. 5 contains some open problems and further discussion.

2. Statement of main results

As mentioned, for the problem of central interest we have $K_{i,j} \sim |i - j|^{-3}$ in $d = 2$, where $|i - j|$ is the Euclidean distance, but we may as well treat all powers for which the interaction is absolutely summable. To be definitive we will simply take, for $s > d$,

$$K_{i,j} = \frac{1}{|i - j|^s} \quad (2.1)$$

with the proviso $K_{i,i} = 0$. We remark that more generality than (2.1) is manifestly possible as is also the case with the ferromagnetic portion of the interaction in (1.1). However, these generalities would tend to obscure the mechanics of the proofs and so we omit them.

In order to define the corresponding Gibbs measures, let $\Lambda \subset \mathbb{Z}^d$ be a finite set and, given a configuration $\sigma \in \{+1, -1\}^{\mathbb{Z}^d}$, let $H_\Lambda = H_\Lambda(\sigma_\Lambda, \sigma_{\Lambda^c})$ denote the Hamiltonian in Λ which is obtained from (1.1) by pitching out all terms with *both* i and j outside Λ . Since $s > d$, the corresponding object is bounded uniformly in σ . Then the DLR formalism tells us that a probability measure on $\{+1, -1\}^{\mathbb{Z}^d}$ —equipped with the product σ -algebra—is a Gibbs measure if the regular conditional distribution of $\sigma_\Lambda = (\sigma_i)_{i \in \Lambda}$ given a configuration $\sigma_{\Lambda^c} = (\sigma_i)_{i \in \Lambda^c}$ in the complement $\Lambda^c = \mathbb{Z}^d \setminus \Lambda$ is of the form

$$Z_\Lambda(\sigma_{\Lambda^c})^{-1} e^{-\beta H_\Lambda(\sigma_\Lambda, \sigma_{\Lambda^c})}, \quad (2.2)$$

where

$$Z_\Lambda(\sigma_{\Lambda^c}) = \sum_{\sigma_\Lambda} e^{-\beta H_\Lambda(\sigma_\Lambda, \sigma_{\Lambda^c})} \quad (2.3)$$

is the partition function. We will use the notation $\langle - \rangle$ to denote expectations with respect to Gibbs measures (which may often stay implicit).

We wish to establish that all Gibbs measures corresponding to the above Hamiltonian have zero average magnetization once $s \in (d, d + 1]$. We will employ some thermodynamic arguments based, ultimately, on the notion of the free energy. To define this quantity, let $Z_{\Lambda,h}(\sigma_{\Lambda^c})$ denote the partition function in Λ with the Hamiltonian

$$H_\Lambda(\sigma_\Lambda, \sigma_{\Lambda^c}) - h \sum_{i \in \Lambda} \sigma_i, \quad (2.4)$$

i.e., for the model in homogeneous external field h . Let

$$\Lambda_L = [-L, L]^d \cap \mathbb{Z}^d. \quad (2.5)$$

Then there exists $\epsilon_L = o(|\Lambda_L|)$ —with little- o uniform in h —such that for all $\sigma, \tilde{\sigma} \in \{-1, 1\}^{\mathbb{Z}^d}$,

$$\left| \log \frac{Z_{\Lambda_L, h}(\sigma_{\Lambda_L^c})}{Z_{\Lambda_L, h}(\tilde{\sigma}_{\Lambda_L^c})} \right| \leq \epsilon_L. \quad (2.6)$$

In particular, the limit

$$f(\beta, h) = -\frac{1}{\beta} \lim_{L \rightarrow \infty} \frac{1}{|\Lambda_L|} \log Z_{\Lambda_L, h}(\sigma_{\Lambda_L^c}) \quad (2.7)$$

exists and is independent of the boundary condition. Furthermore, the function $h \mapsto f(\beta, h)$ is concave for all h .

The independence of the free energy on the boundary condition is standard and follows from the uniform bound on energy per site; cf [25, Theorem II.3.1]. In Sect. 3 we will show that, perhaps not surprisingly, ϵ_L is order $L^{\max\{2d-s, d-1\}}$ with a logarithmic correction at $s = d + 1$.

The concavity of the free energy now permits us to define the *spontaneous magnetization* $m_\star = m_\star(\beta)$ via the right-derivative of $h \mapsto f(\beta, h)$ at $h = 0$:

$$m_\star = -\left. \frac{\partial f}{\partial h^+} \right|_{h=0}. \quad (2.8)$$

It is clear that, by the plus-minus symmetry built into the model, the corresponding left derivative equals $-m_\star$.

The statement of our main result is as follows:

Theorem 2.1. *Consider the interaction described by the Hamiltonian in (1.1-2.1). Then for all $s \in (d, d + 1]$ and all $\beta \in (0, \infty)$, the spontaneous magnetization, m_\star , is zero.*

The regime $d < s < d + 1$ of exponents for the vanishing of the spontaneous magnetization was surmised already in [8]; the present work covers this and, in addition, the somewhat subtle borderline case $s = d + 1$. The above is about as strong a statement as possible concerning the absence of magnetic order from a thermodynamic perspective; the implications for statistical mechanics are similar in their finality. Indeed, the following standard conclusions are implied for the properties of equilibrium states:

Corollary 2.2. *Let $s \in (d, d + 1]$ and let μ be any infinite-volume Gibbs measure for the Hamiltonian in (1.1-2.1) at inverse temperature $\beta \in (0, \infty)$. Let Λ_L be as in (2.5). For each $\epsilon > 0$ there exists $\delta > 0$ such that for all L sufficiently large and μ -almost every boundary condition $\sigma_{\Lambda_L^c}$,*

$$\mu \left(\left| \sum_{i \in \Lambda_L} \sigma_i \right| > \epsilon |\Lambda_L| \mid \sigma_{\Lambda_L^c} \right) \leq e^{-\delta |\Lambda_L|}. \quad (2.9)$$

In particular, μ -almost all configurations σ have zero block-average magnetization,

$$\lim_{L \rightarrow \infty} \frac{1}{|\Lambda_L|} \sum_{i \in \Lambda_L} \sigma_i = 0. \quad (2.10)$$

Finally, in any translation-invariant (infinite volume) Gibbs state, the expectation of the spin at the origin is zero.

The last statement should not be interpreted as a claim that the state is disordered. In fact, as already mentioned, one expects the occurrence of “striped states” at sufficiently low temperatures; see our discussion in Sect. 1 and also Sect. 5. Note that no restrictions are put on the nearest-neighbor coupling J ; the theorem works for all $J \in \mathbb{R}$.

To complement our “no-go” Theorem 2.1, we note that for exponents $s > d + 1$, spontaneous magnetization *will* occur under the “usual” conditions:

Theorem 2.3. *Let $d \geq 2$, pick $s > d + 1$ and consider the interaction as described in (1.1-2.1). Then there exist $J_0 = J_0(s, d) \in (0, \infty)$ and $C_0 = C_0(d) \in (0, \infty)$ such that for all $\beta(J - J_0) \geq C_0$,*

$$m_* > 0. \quad (2.11)$$

In particular, under such conditions, there exist two distinct, translation-invariant extremal Gibbs states $\langle - \rangle^+$ and $\langle - \rangle^-$ such that

$$\langle \sigma_0 \rangle^+ = -\langle \sigma_0 \rangle^- > 0. \quad (2.12)$$

Strictly speaking, this result could be proved by directly plugging in a theorem from [12, Section 3], which is based on an enhanced Peierls estimate. Instead, we provide an independent way to estimate the contour-flip energy which is technically no more demanding and permits the use of sharp contour-counting arguments [20] to derive good estimates on J_0 and the critical value of β at which the transition occurs. As a result, the corresponding constants can be bounded as follows:

$$J_0 \leq C \frac{\pi_d}{d + 1 - s} \quad \text{and} \quad C_0 \leq C \frac{\log d}{d}, \quad (2.13)$$

where π_d is the “surface” measure of the unit sphere in \mathbb{R}^d , and C is a constant of order unity.

3. Estimates on interaction strength

In this section we will perform some elementary but in places tedious calculations that are needed for the proof of our main results. We begin by an estimate on the energy cost of turning large magnetized blocks to opposite magnetization:

Proposition 3.1. *Let Λ_L be as above and, for the couplings $K_{i,j}$ described in (2.1), consider the discrete sum*

$$T_L = \sum_{\substack{i \in \Lambda_L \\ j \in \Lambda_L^c}} K_{i,j}. \quad (3.1)$$

Then, as L tends to infinity:

(i) *For $d < s < d + 1$,*

$$T_L \sim L^{2d-s} Q \quad (3.2)$$

where $Q \in (0, \infty)$ is the integral

$$Q = \int_{\substack{x \in S_1 \\ y \in S_1^c}} \frac{dx dy}{|x - y|^s}. \quad (3.3)$$

with $S_1 = \{x \in \mathbb{R}^d : |x|_1 \leq 1\}$.

(ii) For $s = d + 1$, there exists a constant $A = A(d) \in (0, \infty)$ such that

$$T_L \sim AL^{d-1} \log L. \quad (3.4)$$

In both (i) and (ii) the symbol \sim is interpreted to mean that the ratio of the two sides tends to unity in the stated limit.

To prove this claim, we will instead consider the quantity $T_{L,a}$ which is defined in the same fashion as T_L except that the “inside sum” now ranges over Λ_{L-a} instead of Λ_L , providing us with a cutoff scale a . Of course we must allow $a \rightarrow \infty$ and, for $s \in (d, d + 1)$, not much more is actually required but, to save work, we shall insist that $a/L^{d+1-s} \rightarrow 0$. (Indeed, we remark that while most of the up and coming is not *strictly* necessary for these cases, it will allow for a unified treatment later.) For the the marginal case of $s = d + 1$ we need to implement the stronger requirement that $a/\log L \rightarrow 0$.

Our claim is that the augmented quantities have the asymptotics that was stated for their unadorned counterparts. This is sufficient since, keeping in mind the above requirements,

$$T_{L,a} \leq T_L \leq T_{L,a} + 2d\mathcal{E}_s L^{d-1}a \quad (3.5)$$

where

$$\mathcal{E}_s = \sum_j K_{0,j} < \infty \quad (3.6)$$

denotes the maximum antiferromagnetic energy associated with a single spin flip.

For the purposes of explicit calculations, it will be convenient to replace $K_{i,j}$ with the quantities $\tilde{K}_{i,j}$ obtained by “smearing” the interaction about the unit cells surrounding the sites i and j :

$$\tilde{K}_{i,j} = \int_{\substack{|x-i|_\infty \leq 1/2 \\ |y-j|_\infty \leq 1/2}} \frac{dx dy}{|x-y|^s}. \quad (3.7)$$

It is noted that since all distances exceed (the large quantity) a , the approximation is not severe:

$$\frac{K_{i,j}}{(1 + \theta a^{-1})^s} \leq \tilde{K}_{i,j} \leq \frac{K_{i,j}}{(1 - \theta a^{-1})^s} \quad (3.8)$$

where θ is a number of order unity. Thus, to prove the asymptotics for the $T_{L,a}$, we may insert the $\tilde{K}_{i,j}$ and then perform blatant continuum integration.

As a technical step, for the proof we will need to calculate the total (long-range) interaction between the line segment $(-L, -a)$ on the x -axis and the half-space in \mathbb{R}^d containing all points with positive x -coordinate:

Lemma 3.2. *Consider the integral*

$$I_1(L, a) = \int_a^L dx \int_0^\infty dy \int_{\mathbb{R}^{d-1}} dz \frac{1}{[(x+y)^2 + |z|^2]^{s/2}}. \quad (3.9)$$

In the limit when $a/L \rightarrow 0$ (with $L \geq 1$) when $s < d + 1$ and $|\log a|/\log L \rightarrow 0$ when $s = d + 1$,

$$I_1(L, a) \sim \begin{cases} C_1 L^{d+1-s}, & \text{if } d < s < d + 1, \\ C_1 \log L, & \text{if } s = d + 1, \end{cases} \quad (3.10)$$

where $C_1 = C_1(d, s) \in (0, \infty)$.

Proof. Scaling z by $x + y$ yields

$$I_1(L, a) = \tilde{C}_1 \int_a^L dx \int_0^\infty dy (x + y)^{d-1-s} \quad (3.11)$$

where

$$\tilde{C}_1 = \int_{\mathbb{R}^{d-1}} \frac{dz}{[1 + |z|^2]^{s/2}}. \quad (3.12)$$

From here the result follows by direct integration. \square

Now we are ready to prove the $s < d + 1$ part of Proposition 3.1:

Proof of Proposition 3.1(i). For $r < 1$ let \mathcal{Q}_r denote the integral (3.3) with x restricted to a cube S_r instead of S_1 . Let $\tilde{T}_{L,a}$ denote the quantity $T_{L,a}$ with $K_{i,j}$ replaced by $\tilde{K}_{i,j}$. A simple scaling yields

$$\tilde{T}_{L,a} = L^{2d-s} \mathcal{Q}_{1-a/L}. \quad (3.13)$$

Hence, all we need to show is that \mathcal{Q}_r remains finite as $r \uparrow 1$. This in turn boils down to the absolute convergence of the integral defining \mathcal{Q} .

To show that $\mathcal{Q} < \infty$ we note that the quantity $(L-a)^{d-1} I_1(L-a, a)$ in Lemma 3.2 may be interpreted as the integral of $|x - y|^{-s}$ over $x \in \Lambda_{L-a}$ and over y ranging through the half-space marked by the hyperplane passing through a given side of the cube Λ_L . This implies

$$\tilde{T}_{L,a} \leq 2dL^{d-1} I_1(L-a, a) \quad (3.14)$$

and, more importantly,

$$\mathcal{Q} \leq 2dI_1(1, 0). \quad (3.15)$$

By Lemma 3.2 and the Monotone Convergence Theorem, $I_1(1, 0) < \infty$ when $s < d + 1$. \square

The proof of the critical case, $s = d + 1$, is more subtle. The following lemma encapsulates the calculations that are needed on top of those in Lemma 3.2:

Lemma 3.3. *Let $s \in (d, d + 1]$ and consider the integral*

$$I_2(L, a) = \int_a^L dx \int_a^L dy \int_0^\infty d\tilde{x} \int_0^\infty d\tilde{y} \int_{\mathbb{R}^{d-2}} dz \frac{1}{[(x + \tilde{x})^2 + (y + \tilde{y})^2 + |z|^2]^{s/2}}. \quad (3.16)$$

There exists $C_2 = C_2(d, s) < \infty$ such that for $L \gg a \gg 1$,

$$I_2(L, a) \leq C_2 L^{d+2-s}. \quad (3.17)$$

Similarly to the quantity $I_1(L, a)$ in Lemma 3.2, the integral $I_2(L, a)$ may be interpreted as the total interaction between the square $(-L, -a) \times (-L, -a)$ in the (x, y) -plane and the quarter-space in \mathbb{R}^d containing all points with positive x and y coordinates.

Proof of Lemma 3.3. Applying the bound

$$(x + \tilde{x})^2 + (y + \tilde{y})^2 + |z|^2 \geq x^2 + \tilde{x}^2 + y^2 + \tilde{y}^2 + |z|^2 \quad (3.18)$$

and scaling z by the root of $x^2 + \tilde{x}^2 + y^2 + \tilde{y}^2$ we get

$$I_2(L, a) \leq O(1) \int_a^L dx \int_a^L dy \int_0^\infty d\tilde{x} \int_0^\infty d\tilde{y} [x^2 + \tilde{x}^2 + y^2 + \tilde{y}^2]^{\frac{d-2-s}{2}} \quad (3.19)$$

Writing

$$r^2 = x^2 + y^2 \quad \text{and} \quad \rho^2 = \tilde{x}^2 + \tilde{y}^2 \quad (3.20)$$

we pass to the polar coordinates in both pairs of variables—with $\rho \in (0, \infty)$ and, as an upper bound, $r \in (a/2, 2L)$ —yielding the result

$$I_2(L, a) \leq O(1) \int_{a/2}^{2L} dr r \int_0^\infty d\rho \rho [r^2 + \rho^2]^{\frac{d-2-s}{2}} = O(1) \int_{a/2}^{2L} dr r^{d+1-s}. \quad (3.21)$$

Here we scaled ρ by r and integrated ρ out to get the last integral. Since $s \leq d + 1$, the integral over r is order L^{d+2-s} . \square

Proof of Proposition 3.1(ii). In this case we cannot simply set $a = 0$ and apply scaling. Notwithstanding, we still have the bound

$$\tilde{T}_{L,a} \leq 2dL^{d-1} I_1(L - a, a). \quad (3.22)$$

By Lemma 3.2, we have

$$\tilde{T}_{L,a} \leq 2dC_1 L^{d-1} \log L [1 + o(1)], \quad L \rightarrow \infty. \quad (3.23)$$

We claim that this bound is asymptotically sharp. Indeed, (3.23) overcounts by including (the integral over y in) the intersection of two halfspaces—marked by two neighboring sides of Λ_L —multiple times. In light of the aforementioned interpretation of $I_2(L, a)$, the contribution from each such intersection is bounded by $L^{d-2} I_2(L, a)$. By Lemma 3.3, this is at most order L^{d-1} . Hence we have (3.4) with $A = 2dC_1$. \square

Theorem 2.3 will require us to show that, for $s > d + 1$, the total strength of the long-range interaction through the boundary of a finite set is of order boundary:

Proposition 3.4. *Let $s > d + 1$. Then there is a constant $C_3 = C_3(d, s) < \infty$ such that if $\Lambda \subset \mathbb{Z}^d$ is finite and connected, then*

$$\sum_{i \in \Lambda} \sum_{j \in \Lambda^c} K_{i,j} \leq C_3 |\partial\Lambda| \quad (3.24)$$

where $|\partial\Lambda|$ denotes the number of bonds with one endpoint in Λ and the other in Λ^c .

Proof. Let $V \subset \mathbb{R}^d$ denote the union of unit cubes centered at the sites of Λ . Let

$$W = \{y \in V^c : \text{dist}(y, V) \geq 1\}. \quad (3.25)$$

In light of (3.8), it suffices to show that, for some $C < \infty$,

$$\int_W dy \int_V dx \frac{1}{|x - y|^s} \leq C \Sigma(\partial V), \quad (3.26)$$

where Σ denotes the surface measure on ∂V . (Indeed, we have $\Sigma(\partial V) = |\partial\Lambda|$.) To this end we note that the function $x \mapsto (d - s)|x|^{-s}$ is the divergence of the vector field $x \mapsto x/|x|^s$. The Gauss-Green formula thus tells us that for all $y \in (V^c)^\circ$,

$$\int_V dx \frac{1}{|x - y|^s} = \frac{1}{d - s} \int_{\partial\Lambda} \frac{\tau(x) \cdot (x - y)}{|x - y|^s} \Sigma(dx), \quad (3.27)$$

where $\tau(x)$ is the unit outer normal to the surface at point x (which is well defined Σ -a.e. because ∂V is piecewise smooth). But $|\tau(x) \cdot (x - y)| \leq |x - y|$ and so

$$\int_V dx \frac{1}{|x - y|^s} \leq \frac{1}{s - d} \int_{\partial\Lambda} \frac{1}{|x - y|^{s-1}} \Sigma(dx) \quad (3.28)$$

But $s > d + 1$ ensures that $y \mapsto |x - y|^{-(s-1)}$ is integrable over $\{y \in \mathbb{R}^d : |y - x| \geq 1\}$ and so integrating over y , applying Fubini's theorem, extending the y -integral from $y \in W$ to $\{y : |y - x| \geq 1\}$, and setting

$$C = (s - d)^{-1} \int_{\mathbb{R}^d} |z|^{1-s} 1_{\{|z| \geq 1\}} dz, \quad (3.29)$$

we get (3.26). \square

4. Proofs of main results

Here we will prove the results from Sect. 2; we begin with Theorem 2.1. In our efforts to rule out that $m_\star > 0$, it is useful to have a definite state that exhibits the magnetization. Our choice will be the limit of states at positive external field that are constructed on the torus.

Definition 4.1. Let $h > 0$ and let $\langle - \rangle_{\mathbb{T};h}$ denote an infinite volume state for the interaction described in (1.1-2.1) at inverse temperature β and external field h that is constructed as a limit of finite volume states with toroidal boundary conditions. We define $\langle - \rangle_{\mathbb{T}}$ to be any $h \downarrow 0$ weak limit of the states $\langle - \rangle_{\mathbb{T};h}$. When the occasion arises, we will denote the measure associated with this state by $w_{\mathbb{T}}$.

Lemma 4.2. The measure $w_{\mathbb{T}}$ is a Gibbs measure for the interaction described in (1.1-2.1) at inverse temperature β . Moreover, $w_{\mathbb{T}}$ is translation invariant, it satisfies $\langle \sigma_0 \rangle_{\mathbb{T}} = m_\star$ and if m_L denotes the block magnetizations,

$$m_L = \frac{1}{|\Lambda_L|} \sum_{i \in \Lambda_L} \sigma_i, \quad (4.1)$$

then for any μ with $0 < \mu < m_\star$,

$$\lim_{L \rightarrow \infty} w_{\mathbb{T}}(m_L > \mu) = 1. \quad (4.2)$$

Proof. These are standard results from the general theory of Gibbs states. Indeed, translation invariance follows by construction while the fact that $w_{\mathbb{T}}$ is Gibbs is a result of the absolute summability of interactions; cf. [25, Corollary III.2.3]. To compute the expectation $\langle \sigma_0 \rangle_{\mathbb{T}}$ we recall that concavity of the free energy ensures that for any $h' < h < h''$ and any translation-invariant Gibbs state $\langle - \rangle_h$ at external field h ,

$$-\frac{\partial f}{\partial h^+}(\beta, h') \leq \langle \sigma_0 \rangle_h \leq -\frac{\partial f}{\partial h^-}(\beta, h''). \quad (4.3)$$

The definition of m_\star —and the construction of $\langle - \rangle_{\mathbb{T}}$ —then implies $\langle \sigma_0 \rangle_{\mathbb{T}} = m_\star$. Finally, we claim that $m_L \rightarrow m_\star$ in $w_{\mathbb{T}}$ -probability, implying (4.2). Indeed, if the random variable m_L were not asymptotically concentrated, then

$$c_L := w_{\mathbb{T}}(m_L > m_\star + \epsilon) \quad (4.4)$$

would be uniformly positive (at least along a subsequence) for some $\epsilon > 0$. But then the DLR conditions and (2.6-2.7) would imply that, for any $h > 0$,

$$c_L e^{h|\Lambda_L|(m_* + \epsilon)} \leq \langle e^{h|\Lambda_L| m_L} \rangle_{\mathbb{T}} = \left\langle \frac{Z_{\Lambda_L, h}}{Z_{\Lambda_L, 0}} \right\rangle_{\mathbb{T}} = e^{-|\Lambda_L| [f(\beta, h) - f(\beta, 0) + o(1)]}. \quad (4.5)$$

Hence we would conclude

$$f(\beta, h) - f(\beta, 0) \leq -(m_* + \epsilon)h, \quad (4.6)$$

in contradiction with (2.8).

We now define the random analogue of the quantity T_L denoted by \mathbf{T}_L . In each configuration this quantity measures the antiferromagnetic interaction between the inside and outside of a box of scale L :

$$\mathbf{T}_L = \sum_{\substack{i \in \Lambda_L \\ j \in \Lambda_L^c}} K_{i,j} \sigma_i \sigma_j. \quad (4.7)$$

The central estimate—from which Theorem 2.1 will be readily proved—is as follows:

Proposition 4.3. *Consider the interaction described by (1.1-2.1) with $s \in (d, d + 1]$ and $\beta \in (0, \infty)$ and let m_* denote the spontaneous magnetization corresponding to these parameters. For each $\lambda \in (0, 1)$ there is $L_0 < \infty$ such that for $L \geq L_0$,*

$$\langle \mathbf{T}_L \rangle_{\mathbb{T}} \geq \lambda m_*^2 T_L. \quad (4.8)$$

To facilitate the proof we will state and prove a small lemma concerning the averaging behavior of the $K_{i,j}$'s:

Lemma 4.4. *Let ℓ and a be such that $a \gg \ell$ and let V_1 and V_2 be two translates of Λ_ℓ such that $\text{dist}(V_1, V_2) \geq a$. Then for any $\sigma \in \{+1, -1\}^{\mathbb{Z}^d}$ and any $i_0 \in V_1$ and $j_0 \in V_2$,*

$$\left| \sum_{\substack{i \in V_1 \\ j \in V_2}} K_{i,j} \sigma_i \sigma_j - K_{i_0, j_0} \left(\sum_{i \in V_1} \sigma_i \right) \left(\sum_{j \in V_2} \sigma_j \right) \right| \leq C \frac{\ell}{a} K_{i_0, j_0} |\Lambda_\ell|^2. \quad (4.9)$$

Here C is a constant independent of a , ℓ , σ , i_0 or j_0 .

Proof. This is a simple consequence of the bound

$$|K_{i,j} - K_{i_0, j_0}| \leq C \frac{\ell}{a} K_{i_0, j_0} \quad (4.10)$$

which follows by (discrete) differentiation of the formula (2.1) and using the fact that the distance between V_1 and V_2 is at least a , while the difference between the minimum and maximum separation of V_1 and V_2 is a number of order ℓ and $\ell \ll a$. \square

Proof of Proposition 4.3. For $a = a(L)$ tending to infinity in the fashion described in the proof of Lemma 3.1, it is sufficient to establish the inequality in (4.8) with T_L replaced by $T_{L,a}$ and \mathbf{T}_L replaced by its random analogue, $\mathbf{T}_{L,a}$, defined by the corresponding modification of (4.7). We will need to introduce one more length scale, namely $\ell = \ell(L)$ which will also tend to infinity but in such a way that $\ell/a \rightarrow 0$. We will assume

that L , a and ℓ are such that both Λ_{L-a} and Λ_L^c may be tiled by disjoint copies of Λ_ℓ . (Technically this only proves the result for a subsequence but the extension is trivial.)

Let V_1 and V_2 denote translates of Λ_ℓ with $V_1 \subset \Lambda_{L-a}$ and $V_2 \subset \Lambda_L^c$ and let us pick $i_0 \in V_1$ and $j_0 \in V_2$. Let

$$q_\ell = w_{\mathbb{T}}(m_\ell > \mu). \quad (4.11)$$

The following is now easily derived using Lemma 4.4: On the event that the average magnetization in both V_1 and V_2 exceeds μ (which has probability at least $2q_\ell - 1$) the contribution of $i \in V_1$ and $j \in V_2$ to the random variable $\mathbf{T}_{L,a}$ is at least

$$[1 + O(\ell/a)]K_{i_0, j_0}|\Lambda_\ell|^2\mu^2. \quad (4.12)$$

On the other hand, on the complementary event (which has probability $1 - q_\ell$) the contribution can be as small as

$$-[1 + O(\ell/a)]K_{i_0, j_0}|\Lambda_\ell|^2. \quad (4.13)$$

This means that the blocks V_1 and V_2 contribute to $\langle \mathbf{T}_{L,a} \rangle_{\mathbb{T}}$ at least

$$[1 + O(\ell/a)]K_{i_0, j_0}|\Lambda_\ell|^2[\mu^2(2q_\ell - 1) - (1 - q_\ell)]. \quad (4.14)$$

Finally, Lemma 4.4 also gives

$$K_{i_0, j_0}|\Lambda_\ell|^2 = [1 + O(\ell/a)] \sum_{i \in V_1} \sum_{j \in V_2} K_{i, j}. \quad (4.15)$$

Noting that the error $O(\ell/a)$ holds uniformly in the position of V_1 and V_2 , we may now sum over all (disjoint) translates of V_1 and V_2 in Λ_{L-a} and Λ_L^c , respectively, to get

$$\langle \mathbf{T}_{L,a} \rangle_{\mathbb{T}} \geq [1 + O(\ell/a)](\mu^2(2q_\ell - 1) - (1 - q_\ell))T_{L,a}. \quad (4.16)$$

Since we assumed $\ell/a \rightarrow 0$ and $q_\ell \rightarrow 1$ as $L \rightarrow \infty$, the right-hand side exceeds $\lambda\mu^2T_L$ once $L \gg 1$. \square

Proof of Theorem 2.1. By the inherent spin-reversal symmetry, an enhancement of the standard Peierls contour (de)erasure procedure yields, for any $\kappa > 0$,

$$w_{\mathbb{T}}(\mathbf{T}_L \geq \kappa T_L) \leq e^{-2\beta[\kappa T_L - 2dJL^{d-1}]}. \quad (4.17)$$

Indeed, considering the probability conditioned on the configuration outside Λ_L , we may split the energy into two parts: the energy inside $E_{\text{in}}(\sigma)$ and the energy $E_{\text{bdry}}(\sigma)$ across the boundary of Λ_L . The important difference between these objects is that E_{in} is invariant under the (joint) reversal of all spins in Λ_L , while E_{bdry} changes sign. Using the fact that the conditional measure has the Gibbs-Boltzmann form, and restricting the partition function in the denominator to configurations obeying $\mathbf{T}_L \leq -\kappa T_L$, we get

$$w_{\mathbb{T}}(\mathbf{T}_L \geq \kappa T_L | \sigma_{\Lambda^c}) \leq \frac{\sum_{\sigma: \mathbf{T}_L \geq \kappa T_L} e^{-\beta[E_{\text{in}}(\sigma) + E_{\text{bdry}}(\sigma)]}}{\sum_{\sigma: \mathbf{T}_L \leq -\kappa T_L} e^{-\beta[E_{\text{in}}(\sigma) + E_{\text{bdry}}(\sigma)]}}. \quad (4.18)$$

Now let us reverse all spins in Λ_L in the lower sum; this yields

$$w_{\mathbb{T}}(\mathbf{T}_L \geq \kappa T_L | \sigma_{\Lambda^c}) \leq \frac{\sum_{\sigma: \mathbf{T}_L \geq \kappa T_L} e^{-\beta[E_{\text{in}}(\sigma) + E_{\text{bdry}}(\sigma)]}}{\sum_{\sigma: \mathbf{T}_L \geq \kappa T_L} e^{-\beta[E_{\text{in}}(\sigma) - E_{\text{bdry}}(\sigma)]}}. \quad (4.19)$$

But

$$E_{\text{bdry}}(\sigma) \geq \kappa T_L - 2dJL^{d-1} \quad (4.20)$$

for every σ in these sums and so (4.17) holds pointwise for $w_{\mathbb{T}}(\mathbf{T}_L \geq \kappa T_L | \sigma_{\Lambda^c})$. Integrating over the boundary condition, we get (4.17).

To finish the proof, we now note

$$\langle \mathbf{T}_L \rangle_{\mathbb{T}} \leq T_L w_{\mathbb{T}}(\mathbf{T}_L \geq \kappa T_L) + \kappa T_L w_{\mathbb{T}}(\mathbf{T}_L < \kappa T_L). \quad (4.21)$$

We learned in Proposition 4.3 that for any $\lambda < 1$ the left hand side is bounded below by $\lambda m_{\star}^2 T_L$ for all L large enough. Thus we have, $\forall \kappa \in (0, 1)$ and $\forall \lambda \in (0, 1)$

$$\frac{\lambda m_{\star}^2 - \kappa}{(1 - \kappa)} \leq w_{\mathbb{T}}(\mathbf{T}_L \geq \kappa T_L) \quad (4.22)$$

once $L \gg 1$. But Proposition 3.1 tells us $T_L \gg L^{d-1}$ and so, in light of (4.17), the $L \rightarrow \infty$ limit forces $\lambda m_{\star}^2 \leq \kappa$. Taking $\kappa \downarrow 0$ yields $m_{\star} = 0$ as claimed. \square

Proof of Corollary 2.2. Let μ be an arbitrary Gibbs state. A variant of the inequality in (4.5) tells us that, for any $h > 0$,

$$\mu(m_L > \epsilon | \sigma_{\Lambda_L^c}) \leq e^{-h|\Lambda_L|\epsilon} \frac{Z_{\Lambda_L, h}(\sigma_{\Lambda_L^c})}{Z_{\Lambda_L, 0}(\sigma_{\Lambda_L^c})} \quad (4.23)$$

Since $m_{\star} = 0$, the ratio of the partition functions behaves like

$$\frac{Z_{\Lambda_L, h}(\sigma_{\Lambda_L^c})}{Z_{\Lambda_L, 0}(\sigma_{\Lambda_L^c})} = \exp\{|\Lambda_L|[o(h) + o(1)]\} \quad (4.24)$$

and so, choosing $0 < h \ll 1$, the right-hand side decays exponentially in $|\Lambda_L|$. An analogous derivation (involving $h < 0$) shows a bound on $\mu(m_L < -\epsilon)$. The second part of the claim now follows by the Borel-Cantelli lemma. \square

We will also finish the proof of the existence of magnetic order for $s > d + 1$:

Proof of Theorem 2.3. The proof is a simple modification of the standard Peierls argument. Consider the box Λ_L and let μ_L^+ denote the Gibbs measure in Λ_L with plus boundary condition in Λ_L^c . We claim that $\mu_L^+(\sigma_0 = -1) \ll 1$ once J and β are sufficiently large (in $d \geq 2$). Indeed, given a connected set $\Lambda \subset \Lambda_L$ whose component is connected and which contains the origin, let \mathcal{A}_{Λ} denote the event that $\sigma_0 = -1$ and that $\partial\Lambda$ is the outer boundary of the connected component of -1 's containing the origin. (In other words, Λ^c is the unique infinite connected component in the complement of the connected component of -1 's containing 0.)

Given $\sigma \in \mathcal{A}_{\Lambda}$, let σ' be the result of flipping *all* spins in Λ (including the $+1$'s). We have

$$H_{\Lambda_L}(\sigma) - H_{\Lambda_L}(\sigma') \geq 2J|\partial\Lambda| - 2 \sum_{i \in \Lambda} \sum_{j \in \Lambda^c} K_{i,j}. \quad (4.25)$$

By Proposition 3.4 the second term in the exponent is bounded by $C_3|\partial\Lambda|$. Letting $J_0 = C_3$ and applying the argument in (4.18-4.19), we thus get

$$\mu_L^+(\mathcal{A}_{\Lambda}) \leq e^{-2\beta(J-J_0)|\partial\Lambda|}. \quad (4.26)$$

But $\mu_L^+(\sigma_0 = -1)$ can be written as the sum of $\mu_L^+(\mathcal{A}_\Lambda)$ over all connected $\Lambda \subset \Lambda_L$ (with connected complement) containing the origin. The standard Peierls argument shows that this sum is dominated by the $\Lambda = \{0\}$ term once $e^{+2\beta(J-J_0)}$ exceeds the connectivity constant for the so-called Peierls contours. It follows that $\mu_L^+(\sigma_0 = -1) \ll 1$ for $J > J_0$ and β sufficiently large, uniformly in L . Taking the weak limit $L \rightarrow \infty$ produces a magnetized infinite volume Gibbs measure μ^+ and, by symmetry, a counterpart negatively-magnetized state μ^- . \square

5. Open problems

We finish by some comments and a few open problems. First, the present paper shows the absence of magnetization at $h = 0$. A natural question is now as follows:

Problem 1. *Let $s \in (d, d + 1]$. Characterize the values $h \neq 0$ at which the free energy is continuously differentiable in homogeneous external field h .*

An answer to this question depends strongly on the precise structure of low-temperature states. In particular, if there is a rigid stripe order (see Problem 3) it is possible that, for some particular values of h , there will be phase coexistence between different arrangements of stripes. Whether that has an effect on the continuity of the magnetization is not clear.

To move to our next problem, let us recall the main reason why the exponent $s = d + 1$ is critical for the disappearance of magnetic order: For $s \leq d + 1$, the gain to be obtained from the antiferromagnetic interaction “through” the boundary of a volume of scale L is order L^{2d-s} which—including the $\log L$ correction when $s = d + 1$ —overpowers the short-range surface cost of order L^{d-1} . However the short-range calculation only applies under the conditions where one envisions a surface tension, e.g., discrete spins. If we replace the Ising spins by, say, plane rotors, the cost due to local interactions for turning over a block now scales as L^{d-2} . Various exponents will readjust accordingly. Thus we pose:

Problem 2. *For the Ising spins replaced by $O(n)$ -spins, and the spin-spin interactions given by the dot product, find the range of exponents s for which the spontaneous magnetization vanishes.*

The problem is interesting due to competing effects in the vicinity of the (purported) interfaces. It has been stipulated in [8] that, in these cases, magnetism will not occur for $d < s < d + 2$. See [9] for some relevant calculations.

As for our next problem we note that, as already mentioned, absence of magnetism is far from ruling out other types of order, with striped states being a prime candidate. Thus we ask:

Problem 3. *Prove the existence of striped states at low temperatures for interactions of the type discussed in this note.*

Some mathematical progress [13] and a great deal of physical progress [1, 5, 6, 10, 11, 14, 18, 19, 21–23, 27] in this direction has been made for the ground state problem. But, at present, the positive-temperature case is far from resolved.

Finally, we recall that much of our proof was based on thermodynamic arguments which, to begin with, require the existence of thermodynamics. Notwithstanding, analogous results should hold even for interactions that decay so slowly that the standard

techniques ensuring the existence of the free energy fail. An instance of some genuine interest arises from Ref. [17]: Consider the model with the Hamiltonian as in (1.1) but with the long-range interaction term modified into

$$\sum_{i,j} K_{i,j}(\sigma_i - \rho)(\sigma_j - \rho). \quad (5.1)$$

The quantity ρ plays the role of “background charge” density; the spin configurations are restricted to have average ρ (otherwise their energy diverges).

Problem 4. Suppose $K_{i,j} \sim |i - j|^{-1}$ in $d = 2, 3$ (and, in general, $K_{i,j} \sim |i - j|^{-s}$ with $s_d < s \leq d$ and $d \geq 2$). Prove that the free energy is differentiable in ρ , at $\rho = 0$.

On the basis of [28] one can infer that the lower bound, s_d , on the region of exponents in the previous open problem satisfies $s_d \leq d - 1$. However, it is noted that, for $s = d - 2$, there is a (complicated) counterexample to differentiability [15] so, presumably, $s_d \geq d - 2$.

Acknowledgments

The authors wish to thank Aernout van Enter for many useful comments on the content and literature. This research was partially supported by the grants NSF DMS-0505356 (M.B.), NSF DMS-0306167 (L.C.) and DOE DE-FG03-00ER45798 (S.K.).

References

1. Ar. Abanov, V. Kalatsky, V.L. Pokrovsky and W.M. Saslow, Phase diagram of ultrathin ferromagnetic films with perpendicular anisotropy, *Phys. Rev. B* **51** (1995), no. 2, 1023–1038.
2. M. Aizenman, J.T. Chayes, L. Chayes and C.M. Newman, Discontinuity of the magnetization in one-dimensional $1/|x - y|^2$ Ising and Potts models, *J. Statist. Phys.* **50** (1988), no. 1-2, 1–40.
3. M. Aizenman and C. Newman, Discontinuity of the percolation density in one-dimensional $1/|x - y|^2$ percolation models, *Commun. Math. Phys.* **107** (1986), no. 4, 611–647.
4. P.W. Anderson, G. Yuval and D.R. Hamann, Exact results in the Kondo problem. II. Scaling theory, qualitatively correct solution, and some new results on one-dimensional classical statistical models, *Phys. Rev. B* **1** (1970), no. 1, 4464–4473.
5. P. Bak and R. Bruinsma, One-dimensional Ising model and the complete devil’s staircase, *Phys. Rev. Lett.* **49** (1982), no. 4, 249–251.
6. L. Chayes, V.J. Emery, S.A. Kivelson, Z. Nussinov and G. Tarjus, Avoided critical behavior in a uniformly frustrated system, *Physica A* **225** (1996), no. 1, 129–153.
7. H.A.M. Daniëls and A.C.D. van Enter, Differentiability properties of the pressure in lattice systems, *Commun. Math. Phys.* **71** (1980), no. 1, 65–76.
8. A.C.D. van Enter, A note on the stability of phase diagrams in lattice systems, *Commun. Math. Phys.* **79** (1981), no. 1, 25–32.
9. A.C.D. van Enter, Instability of phase diagrams for a class of “irrelevant” perturbations, *Phys. Rev. B* **26** (1982), no. 3, 1336–1339.
10. M.M. Fogler, A.A. Koulakov and B.I. Shklovskii, Ground state of a two-dimensional electron liquid in a weak magnetic field, *Phys. Rev. B* **54** (1996), no. 3, 1853–1871.
11. T. Garel and S. Doniach, Phase-transitions with spontaneous modulation: the dipolar ferromagnet, *Phys. Rev. B* **26** (1982), no. 1, 325–329.
12. J. Ginibre, A. Grossmann and D. Ruelle, Condensation of lattice gases, *Commun. Math. Phys.* **3** (1966), no. 3, 187–193.
13. A. Giuliani, J.L. Lebowitz and E.H. Lieb, Ising models with long-range dipolar and short range ferromagnetic interactions, *Phys. Rev. B* **74** (2006), no. 6, 064420.
14. M. Grousson, G. Tarjus and P. Viot, Evidence for “fragile” glass-forming behavior in the relaxation of Coulomb frustrated three-dimensional systems, *Phys. Rev. E* **65** (2002), no. 6, 065103.

15. D. Huse, private communication.
16. R.B. Israel, Generic triviality of phase diagrams in spaces of long-range interactions, *Commun. Math. Phys.* **106** (1986), no. 3, 459–466.
17. R. Jamei, S. Kivelson and B. Spivak, Universal aspects of Coulomb-frustrated phase separation, *Phys. Rev. Lett.* **94** (2005), no. 5, 056805.
18. A.B. Kashuba and V.L. Pokrovsky, Stripe domain structures in a thin ferromagnetic film, *Phys. Rev. B* **48** (1993), no. 14, 10335–10344.
19. A.A. Koulakov, M.M. Fogler and B.I. Shklovskii, Charge density wave in two-dimensional electron liquid in weak magnetic field, *Phys. Rev. Lett.* **76** (1996), no. 3, 499–502.
20. J.L. Lebowitz and A.E. Mazel, Improved Peierls argument for high-dimensional Ising models, *J. Statist. Phys.* **90** (1998), no. 3–4, 1051–1059.
21. U. Löw, V.J. Emery, K. Fabricius and S.A. Kivelson, Study of an Ising model with competing long- and short-range interactions, *Phys. Rev. Lett.* **72** (1994), no. 12, 1918–1921.
22. K.-O. Ng and D. Vanderbilt, Stability of periodic domain structures in a two-dimensional dipolar model, *Phys. Rev. B* **52** (1995), no. 3, 2177–2183.
23. C. Ortix, J. Lorenzana and C. Di Castro, Frustrated phase separation in two-dimensional charged systems, *Phys. Rev. B* **73** (2006), no. 24, 245117.
24. M. Seul and D. Andelman, Domain shapes and patterns: The phenomenology of modulated phases, *Science* **267** (1995), no. 5197, 476–483.
25. B. Simon, *The Statistical Mechanics of Lattice Gases*, Vol. I., Princeton Series in Physics, Princeton University Press, Princeton, NJ, 1993.
26. A.D. Sokal, More surprises in the general theory of lattice systems, *Commun. Math. Phys.* **86** (1982), no. 3, 327–336.
27. B. Spivak and S. Kivelson, Phases intermediate between a two-dimensional electron liquid and Wigner crystal, *Phys. Rev. B* **70** (2004), no. 15, 155114.
28. B. Spivak and S. Kivelson, Transport in two dimensional electronic micro-emulsions, *Ann. Phys.* **321** (2006), no. 9, 2071–2115.
29. D. J. Thouless, Critical region for the Ising model with a long-range interaction, *Phys. Rev.* **181** (1969), no. 2, 954–968.