

# LOCAL INDECOMPOSABILITY OF TATE MODULES OF NON CM ABELIAN VARIETIES WITH REAL MULTIPLICATION

HARUZO HIDA

ABSTRACT. Indecomposability of  $p$ -adic Tate modules over the  $p$ -inertia group for non CM (partially  $p$ -ordinary) abelian varieties with real multiplication is proven under unramifiedness of  $p$  in the base field and in the multiplication field.

## CONTENTS

1. Deformation space of an AVRМ	3
2. Hilbert modular Shimura variety	5
3. Eigen differentials	7
4. Eigen coordinates	8
5. Local indecomposability of an AVRМ	14
6. CM periods and proportionality constants	16
7. Density of partially ordinary primes	18
References	18

As quoted by Ghate–Vatsal in [GV] Question 1 (and answered there affirmatively to a good extent), it is a fundamental problem posed by R. Greenberg to decide indecomposability of an elliptic modular  $p$ -adic Galois representation restricted to the decomposition group at  $p$  in the ordinary case (as long as the representation is not of CM type). Jointly with B. Balasubramanyam, they have generalized the result to the cases of Hilbert modular forms for primes  $p$  splitting completely in the totally real base field (see [GV1]). There are some good applications of their results (see for example [E] and [G]). We can ask the same question for  $p$ -adic Galois representations arising from the Tate module of an abelian variety  $A$  with real multiplication defined over a number field  $k$ ; so,  $\text{End}(A/k)$  is the integer ring  $O$  of a totally real field of degree  $\dim A$ . This concerns the  $\mathfrak{p}$ -adic Tate module  $T_{\mathfrak{p}}A$  for a prime  $\mathfrak{p} | (p)$  of  $O$  as a module over the decomposition group  $D_p$  at a place  $\mathfrak{P}$  over  $p$  of  $k$ . Suppose that  $A$  has good reduction modulo  $\mathfrak{P}$ . If the finite flat group scheme  $A[\mathfrak{p}]$  of  $\mathfrak{p}$ -torsion points is local-local at  $\mathfrak{P}$ , as is well known, the  $D_p$ -module  $A[\mathfrak{p}](\overline{\mathbb{Q}}_p)$  or  $T_{\mathfrak{p}}A \otimes_{O_{\mathfrak{p}}} F_{\mathfrak{p}}$  is often irreducible (as proven in [ALR] IV-38, §A.2.2 for elliptic curves, see also the generalization of Serre’s modulo  $p$  modularity conjecture for totally real  $k$ ; in particular, §3.1 of [BDJ]). Anyway in this paper, we limit ourselves to the case where  $A[\mathfrak{p}](\overline{\mathbb{F}}_p) \neq \{0\}$ . This assumption does not necessarily mean that  $A$  has ordinary good reduction at  $p$ .

We answer this question affirmatively in Theorem 5.1 for an abelian variety with sufficiently many real multiplication but without complex multiplication (over  $\overline{\mathbb{Q}}$ ) if the endomorphism totally real field  $F$  and the field  $k$  of definition are unramified at  $p$  over  $\mathbb{Q}$ . Since simple factors of the jacobian of a Shimura curve are often of this type, we should be able to answer the original question by Greenberg. Indeed, Bin Zhao [Z] has just removed our assumption of unramifiedness of  $p$  in the base field  $k$  and in the multiplication field  $F$  and has proved unconditionally indecomposability for the

---

*Date:* December 30, 2012.

*2010 Mathematics Subject Classification.* primary 14G35, 11G15, 11G18, 11F80 secondary 11G10, 14L05.

*Key words and phrases.* abelian variety, Tate module, real/complex multiplication, Galois representation, deformation.

The author is partially supported by the NSF grants DMS 0753991 and DMS 0854949.

nearly ordinary Galois representation of each weight 2 Hilbert cusp form at least over an odd degree totally real base field. If the base field has even degree, he had to assume square-integrability at a finite place or the existence of an abelian variety associated to the cusp form.

Since a CM abelian variety with ordinary good reduction at  $p$  has semi-simple Tate module over  $D_p$ , we may regard the  $p$ -local indecomposability of the  $p$ -adic Tate module as a characterization of non CM/CM abelian varieties via local  $p$ -inertia action. The importance of such characterization is emphasized in [H11a], and other examples of characterization are listed there.

Throughout this paper, we fix an algebraic closure  $\overline{\mathbb{Q}}$  in  $\mathbb{C}$  and an algebraic closure  $\overline{\mathbb{Q}}_v$  of  $\mathbb{Q}_v$  (the  $v$ -adic field or  $\mathbb{R}$  for  $v = \infty$ ) for each rational place  $v$ . We write  $\mathbb{C}_v$  for the completion of  $\overline{\mathbb{Q}}_v$  under  $v$ . Therefore, for each  $v$ , we fix a field embedding  $i_v : \overline{\mathbb{Q}} \hookrightarrow \mathbb{C}_v$  (here  $i_\infty$  coincides with the original inclusion  $\overline{\mathbb{Q}} \subset \mathbb{C}$ ) and identify all  $\mathbb{C}_v$  with  $\mathbb{C}$  so that we have the following commutative diagram for each  $v$ :

$$\begin{array}{ccc} \overline{\mathbb{Q}} & \xlongequal{\quad} & \overline{\mathbb{Q}} \\ i_v \downarrow \cap & & \cap \downarrow i_\infty \\ \mathbb{C}_v & \xlongequal{\quad} & \mathbb{C}. \end{array}$$

Here is a sketch of our proof of indecomposability (and an outline of the paper), assuming for simplicity that the abelian variety  $A$  over a number field  $k \subset \overline{\mathbb{Q}}$  has multiplication by the integer ring  $O$  of a real quadratic field  $F$ ; so,  $2 = [F : \mathbb{Q}] = \dim A$  and  $\text{End}_F^0(A/\overline{\mathbb{Q}}) = \text{End}_F^0(A/k \times_k \overline{\mathbb{Q}}/\overline{\mathbb{Q}}) = F$  (this means that  $A$  does not have complex multiplication), where “ $\text{End}_F^0(?)$ ” denotes the total quotient ring of the  $O$ -linear endomorphism algebra  $\text{End}_O(?)$  of the object “?” inside. Such abelian varieties are parameterized by Hilbert modular varieties whose theory we recall in Section 2 to the extent we need (explicitly or implicitly).

Let  $\mathfrak{O}$  denote the integer ring of  $k$ . We denote primes of  $k$  by upper case Gothic letters and those of  $F$  by lower case Gothic letters, and the corresponding Roman (lower case) character is the residual characteristic of a prime denoted by a Gothic character. Suppose further for simplicity that  $A$  has ordinary good reduction at primes  $\mathfrak{P}$  and  $\mathfrak{L}$  with  $p \neq l$ . On the contrary to what we want to prove, suppose that  $A[\mathfrak{p}^\infty]$  for a prime  $\mathfrak{p}$  above  $p$  has a semi-simple Tate module over  $D_p = \text{Gal}(\overline{\mathbb{Q}}_p/k_{\mathfrak{p}})$ . Then the  $\mathfrak{p}$ -adic Serre–Tate coordinate  $t_{\mathfrak{p}}(A)$  of  $A$  is equal to 1. If  $\mathfrak{p}$  is a unique prime over  $(p)$  in  $O$ , this identity  $t_{\mathfrak{p}}(A) = 1$  forces  $A$  to be a canonical lift having complex multiplication. Thus we assume  $(p) = \mathfrak{p}\mathfrak{p}'$  in  $F$  with  $\mathfrak{p} \neq \mathfrak{p}'$ . By the Serre–Tate theory, for the Witt vector ring  $W_p$  with coefficients in an algebraic closure  $\overline{\mathbb{F}}_p$  of  $\mathbb{F}_p$ , the deformation space of the reduction  $A_{\mathfrak{p}} = A \otimes_{\mathfrak{O}_{\mathfrak{p}}} \overline{\mathbb{F}}_p$  (for  $\mathbb{F}_{\mathfrak{p}} := \mathfrak{O}/\mathfrak{p}$  regarded as sitting in  $\overline{\mathbb{F}}_p$ ) is isomorphic to  $\widehat{S}_{/W_p} = \widehat{S}_{p/W_p} := \widehat{\mathbb{G}}_m \otimes_{\mathbb{Z}_p} O_p = \widehat{S}_{\mathfrak{p}} \times \widehat{S}_{\mathfrak{p}'}$  with  $\widehat{S}_{\mathfrak{p}} := \widehat{\mathbb{G}}_m \otimes_{\mathbb{Z}_p} O_{\mathfrak{p}}$ , and  $t_{\mathfrak{p}}$  is the coordinate of the multiplicative formal group  $\widehat{\mathbb{G}}_m \otimes_{\mathbb{Z}_p} O_{\mathfrak{p}} = \widehat{\mathbb{G}}_m$  (as  $O_{\mathfrak{p}} = \mathbb{Z}_p$ ). In other words,  $\widehat{\mathbb{G}}_{m/W_p}$  is the formal completion of  $\mathbb{G}_m = \text{Spec}(W_p[t_p, t_p^{-1}])$  along the origin (i.e., the identity) of its reduction modulo  $p$ . We recall in Section 1 some details of the construction of Serre–Tate coordinates.

The field  $M = \text{End}_F^0(A_{\mathfrak{p}/\overline{\mathbb{F}}_p})$  is a CM quadratic extension of  $F$ . We have two Serre–Tate coordinates  $t_{\mathfrak{p}}$  and  $t_{\mathfrak{p}'}$ . Since  $A$  does not have CM, we must have  $t_{\mathfrak{p}'}(A) \neq 1$  (as the origin of  $\widehat{S}$  corresponds to the canonical lift of  $A_{\mathfrak{p}}$  which has complex multiplication by the CM field  $M$ ). By the universality of  $\widehat{S}$ ,  $M^\times$  acts on  $\widehat{S}$  naturally by an isogeny action. Then by the reciprocity law of  $\widehat{S}$  in [H10] §3.3, we have  $t_{\mathfrak{p}} \circ \alpha = t_{\mathfrak{p}}^{\alpha \sigma(1-c)}$  for an embedding  $\sigma : M \hookrightarrow \overline{\mathbb{Q}}$  inducing the place  $\mathfrak{p}$  of  $F$  after composing with  $i_p$ . Here  $c$  denotes the generator of  $\text{Gal}(M/F)$ . For the  $l$ -adic place  $\mathfrak{l}$  induced by  $i_l \circ \sigma$ , we have the Serre–Tate coordinate  $t_{\mathfrak{l}}$  of the deformation space  $\widehat{S}_{\mathfrak{l}} = \widehat{\mathbb{G}}_m \otimes_{\mathbb{Z}_p} O_{\mathfrak{l}}$  of  $A_{\mathfrak{L}}$ . Now the abelian variety  $A$  gives rise to a  $k$ -point  $x$  of the Hilbert modular Shimura variety  $Sh_{/\mathbb{Q}}$  for  $GL(2)_{/F}$ . Looking at the  $k$ -tangent space  $Tan_x$  at  $x \in Sh_K(k)$  (for a neat open compact subgroup  $K \subset GL_2(F_{\mathbb{A}}^{(\infty)})$  maximal at  $p$  and  $l$ ), the Lie algebras  $Lie(\widehat{S}_{\mathfrak{p}})$  and  $Lie(\widehat{S}_{\mathfrak{l}})$  are related to (the  $\sigma$ -part of)  $Tan_x$  via appropriate scalar extension; therefore, the action of  $M^\times$  on  $Lie(\widehat{S}_{\mathfrak{p}})$  can be transferred to  $Lie(\widehat{S}_{\mathfrak{l}})$  via  $Tan_x$ . We will see in Section 4 that  $t_{\mathfrak{p}}$  (resp.  $t_{\mathfrak{l}}$ ) is the exponential of the corresponding linear coordinate  $\tau_{\mathfrak{p}}$  (resp.  $\tau_{\mathfrak{l}}$ ) of  $Lie(\widehat{S}_{\mathfrak{p}})$  (resp.  $Lie(\widehat{S}_{\mathfrak{l}})$ ). From this, we conclude in Theorem 4.5 that  $t_{\mathfrak{l}}$  also satisfies  $t_{\mathfrak{l}} \circ \alpha = t_{\mathfrak{l}}^{\alpha \sigma(1-c)}$  for  $\alpha \in M^\times$ ; so,  $\text{End}_F^0(A_{\mathfrak{L}/\overline{\mathbb{F}}_l})$  must be isomorphic to  $M$ . In Section 3, we describe

the canonical  $O$ -action on  $\Omega_{Sh/k}$ ,  $Tan_x$  and on  $Lie(\widehat{S})$ , and in Section 4, we study the  $M$ -action (extending the  $O$ -action) on the Serre–Tate coordinates (and its linear version).

Because of  $x \in Sh(k) \subset Sh(\mathbb{C}_p) = Sh(\mathbb{C}_l)$  (under our identification:  $\mathbb{C}_p = \mathbb{C}_l$ ), the formal completion  $\widehat{O}_{Sh,x}$  along the points  $x \in Sh(\mathbb{C}_p) = Sh(\mathbb{C}_l)$  canonically contains  $\mathcal{O}_{\widehat{S}_p}$  and  $\mathcal{O}_{\widehat{S}_l}$ . Thus in  $\widehat{O}_{Sh,x}$ , we can compare  $\tau_p = \log_p \circ t_p$  and  $\tau_l = \log_l \circ t_l$  for the  $v$ -adic logarithm  $\log_v$ . Since they both satisfy  $\tau_\gamma \circ \alpha = \alpha^{\sigma(1-c)} \tau_\gamma$ , they are proportional; so,  $\tau_l(A) = 0$  and hence  $t_l(A) = 1$ , which implies the Tate module  $T_l A$  of  $A[\mathfrak{l}^\infty]$  is also semi-simple under  $D_l$ . As is well known (cf. [O] 2.7 and Section 7 in the text), split ordinary primes for an abelian surface have Dirichlet density 1 for  $k$  (replacing  $k$  by its finite extension if necessary), and using non-CM property of the  $\mathfrak{p}$ -adic Galois representation on the Tate module  $T_p A$  (combined with Chebotarev density), we can find  $\mathfrak{l}$  such that  $M \not\cong \text{End}_F^0(A_{\mathfrak{D}/\overline{\mathbb{F}}_l})$ , getting a contradiction (whose details will be given in Section 5). In the text, we fill details of this argument and generalize this to totally real fields  $F$  with  $[F : \mathbb{Q}] \geq 2$ , to primes  $\mathfrak{p}$  with residual degree  $\geq 1$  and to abelian varieties  $A$  which may not be ordinary at  $\mathfrak{P}$  (i.e., only partially ordinary at  $\mathfrak{P}$ ).

As a byproduct of our proof, we get (in Theorem 4.5) the proportionality of  $\tau_p$  and  $\tau_l$ . Though it is irrelevant to the proof of the main theorem, we will compute in Section 6 explicitly the proportionality constant as the ratio of the square of the Katz  $p$ -adic and  $l$ -adic CM periods. In the above sketch of the proof, we used heavily existence of many primes of  $k$  at which the given abelian variety  $A/k$  has good (partially) ordinary reduction. We give a brief account of the fact that such primes has density 1 among primes of (the Galois closure of)  $k$  in Section 7.

As is clear from this sketch, there is much room for generalization of our argument to abelian varieties with less multiplication. We hope to come back to the study of local indecomposability for general abelian varieties in near future.

The author thanks Brian Conrad for some help of identifying an optimal reference in rigid analytic geometry for quotes here.

## 1. DEFORMATION SPACE OF AN AVRMS

Pick a rational primes  $p$ . Let  $F$  be a totally real field with integer ring  $O$  in which  $p$  is unramified. Write  $d = [F : \mathbb{Q}]$ . Let  $(A, \lambda, \theta)_{/k}$  be an abelian variety with real multiplication by  $O$  (in short an AVRMS by  $O$ ) over a number field  $k \subset \overline{\mathbb{Q}}$  with integer ring  $\mathfrak{D}$  in which  $p$  is unramified. Here  $\theta : O \hookrightarrow \text{End}(A_{/k})$  is a ring homomorphism sending the identity to the identity making  $Lie(A)$  a free  $O \otimes_{\mathbb{Z}} k$ -module of rank 1, and  $\lambda : A \rightarrow {}^t A$  for the dual abelian scheme  ${}^t A = \text{Pic}_{A/k}^0$  is an  $O$ -linear polarization of degree prime to  $p$ . Thus the  $p$ -adic Tate module  $T_p A$  is free of rank 2 over  $O_p = O \otimes_{\mathbb{Z}} \mathbb{Z}_p$ , and  $\dim A = [F : \mathbb{Q}]$ . Let  $\mathfrak{P} = \{x \in \mathfrak{D} : |i_p(x)|_p < 1\}$  be the prime of  $k$  induced by  $i_p : \overline{\mathbb{Q}} \hookrightarrow \mathbb{C}_p$ . Let  $\overline{\mathbb{F}}_p = \overline{\mathbb{F}}_{\mathfrak{P}}$  be an algebraic closure of  $\mathbb{F}_{\mathfrak{P}} = \mathfrak{D}/\mathfrak{P}$ , and write  $W = W_p$  for the ring of Witt vectors with coefficients in  $\overline{\mathbb{F}}_p$ . We identify  $W$  with the subring of  $\mathbb{C}_p$  which is the  $p$ -adic completion of the ring of integers in the maximal unramified extension of  $\mathbb{Q}_p$  in  $\overline{\mathbb{Q}}_p \subset \mathbb{C}_p$ . Thus  $i_p$  embeds  $\mathfrak{D}$  in  $W$ , and we regard  $\mathfrak{D}_{\mathfrak{P}} \subset W$ . We put  $\mathcal{W} = \mathcal{W}_p = i_p^{-1}(W_p) \subset \overline{\mathbb{Q}}$  which is a discrete valuation ring with residue field  $\overline{\mathbb{F}}_p$  (and is a strict henselization of  $\mathbb{Z}_{(p)} = \mathbb{Q} \cap \mathbb{Z}_p$ ).

Let  $\mathfrak{D}_{\mathfrak{P}}$  be the  $\mathfrak{P}$ -adic completion of  $\mathfrak{D}$ , and suppose that  $(A, \lambda, \theta)_{/k}$  extends to the triple  $(A, \lambda, \theta)_{/\mathfrak{D}_{\mathfrak{P}}}$  for an abelian scheme  $A_{/\mathfrak{D}_{\mathfrak{P}}}$ . Thus we can think of the special fiber  $(A_0, \lambda_0, \theta_0)_{/\overline{\mathbb{F}}_p} = (A, \lambda, \theta) \times_{\mathfrak{D}_{\mathfrak{P}}} \overline{\mathbb{F}}_p$ . Let  $CL_W$  denote the category of complete local  $W$ -algebras with residue field  $\overline{\mathbb{F}}_p$ . We consider the fiber category  $\mathcal{D}_{/W}$  (over  $CL_W$ ) of deformations over  $R$  of  $(A_0, \lambda_0, \theta_0)_{/\overline{\mathbb{F}}_p}$  over  $R \in CL_W$ . Objects of  $\mathcal{D}_{/W}$  are triples  $(A', \lambda', \theta')_{/R}$  satisfying  $\iota_{A'} : (A', \lambda', \theta') \times_R \overline{\mathbb{F}}_p \cong (A_0, \lambda_0, \theta_0)_{/\overline{\mathbb{F}}_p}$ . A morphism  $\phi : (A', \lambda', \theta')_{/R} \rightarrow (A'', \lambda'', \theta'')_{/R}$  is a morphism of AVRMS's  $\phi : A' \rightarrow A''$  such that

- (1) the special fiber  $\phi_0$  of  $\phi$  satisfies  $\iota_{A''} \circ \phi_0 \circ \iota_{A'}^{-1} = \text{id}_{A_0}$  for the identity map  $\text{id}_{A_0}$  of  $A_0$ ;
- (2)  $\lambda' = {}^t \phi \circ \lambda'' \circ \phi$ ;
- (3)  $\phi \circ \theta'(a) = \theta''(a) \circ \phi$  for all  $a \in O$ .

We suppose that

(NLL) the reduction  $A_0 := A_{\mathfrak{P}} = A \otimes_{\mathfrak{D}_{\mathfrak{P}}} \overline{\mathbb{F}}_p$  has nontrivial  $p$ -torsion  $\overline{\mathbb{F}}_p$ -points.

So, we have a prime ideal  $\mathfrak{p}|p$  of  $O$  such that the  $\mathfrak{p}$ -adic Barsotti–Tate group  $A[\mathfrak{p}^\infty]$  is ordinary, fitting into the following connected-étale exact sequence of Barsotti–Tate  $O_{\mathfrak{p}}$ -modules over  $W$ :

$$0 \rightarrow \mu_{p^\infty} \otimes_{\mathbb{Z}_p} O_{\mathfrak{p}} \rightarrow A[\mathfrak{p}^\infty]_{/W} \rightarrow F_{\mathfrak{p}}/O_{\mathfrak{p}} \rightarrow 0.$$

Let  $\Sigma^{ord} = \Sigma_p^{ord}$  be the set of primes  $\mathfrak{p}$  in  $O$  over  $p$  such that  $A[\mathfrak{p}^\infty]_{/W}$  is not connected. Then we put  $\Sigma^{ll} = \Sigma_p^{ll} = \{\mathfrak{p}|p : \mathfrak{p} \notin \Sigma^{ord}\}$ . We may regard  $\Sigma^?$  as a set of field embeddings  $\sigma : F \hookrightarrow \overline{\mathbb{Q}}$  made up of  $\sigma$  with  $\{\alpha \in O : |i_p(\sigma(\alpha))|_p < 1\} = \mathfrak{p} \in \Sigma^?$ .

We now want to relate  $\mathcal{D}_{/W}$  with the following fiber category  $DEF$  over  $CL_W$  of quadruples  $(A_0, D, \Lambda, \epsilon)_{/R}$ , where  $D$  is a Barsotti–Tate  $O$ -module over  $R$ ,  $\Lambda : D \rightarrow {}^tD$  is an  $O$ -linear isomorphism of Barsotti–Tate groups into the Cartier dual  ${}^tD$  of  $D$  and  $\epsilon : D_0 \cong A_0[p^\infty]$  is an  $O$ -linear isomorphism of Barsotti–Tate  $O$ -modules for the special fiber  $D_0 = D \otimes_{\text{Spec}(W)} \text{Spec}(\overline{\mathbb{F}}_p)$ . We require  $\epsilon$  to satisfy the following commutativity:

$$\begin{array}{ccc} D_0 & \xrightarrow{\epsilon} & A_0[p^\infty] \\ \Lambda_0 \downarrow & & \downarrow \lambda_0 \\ {}^tD_0 & \xleftarrow[{}^t\epsilon]{} & {}^tA_0[p^\infty], \end{array}$$

where  $\Lambda_0$  is the special fiber of  $\Lambda$ . A morphism

$$\phi : (A_0, D, \Lambda, \epsilon) \rightarrow (A'_0, D', \Lambda', \epsilon')$$

of  $DEF$  is a pair of  $O$ -linear morphisms  $\phi_{BT} : D \rightarrow D'$  in  $\text{Hom}_{BT_{/W}}(D, D')$  and  $\phi_{\mathcal{D}} : A_0 \rightarrow A'_0$  in  $\text{Hom}_{GSC\mathcal{H}_{/\overline{\mathbb{F}}_p}}(A_0, A'_0)$  making the following two diagrams commutative:

$$\begin{array}{ccc} A_0[p^\infty] & \xrightarrow{\phi_{\mathcal{D}}} & A'_0[p^\infty] \\ \epsilon \uparrow & & \uparrow \epsilon' \\ D_0 & \xrightarrow{\phi_{BT}} & D'_0 \end{array}$$

and

$$\begin{array}{ccc} D_0 & \xrightarrow{\phi_{BT}} & D'_0 \\ \Lambda \downarrow & & \downarrow \Lambda' \\ {}^tD_0 & \xleftarrow[{}^t\phi_{BT}]{} & {}^tD'_0. \end{array}$$

Here  $GSC\mathcal{H}$  (resp.  $BT$ ) stands for the fiber category of group schemes (resp. Barsotti–Tate groups) over  $CL_W$ . We have a natural functor:  $\mathcal{D}_{/W} \rightarrow DEF$  given by

$$A \mapsto (A_0 = A \otimes_W \overline{\mathbb{F}}_p, A[p^\infty], \lambda|_{A[p^\infty]}, \iota)$$

for the canonical isomorphism  $\iota : A[p^\infty] \otimes_W \overline{\mathbb{F}}_p \cong A_0[p^\infty]$ .

**Theorem 1.1** (Serre–Tate). *The above functor:  $\mathcal{D}_{/W} \rightarrow DEF$  is a canonical equivalence of categories.*

A proof of this is given in [K] though the input of  $\theta$  and  $\lambda$  is not there (see [K1] Lemma 1.11.6 or [Ra] §1.7 for how to modify the argument incorporating endomorphisms and a polarization).

We can split  $DEF$  into the product of the following fiber categories:  $DEF^{ord}$  and  $DEF^{ll}$  over  $CL_W$ . Here “*ord*” stands for ordinary Barsotti–Tate groups and “*ll*” stands for local-local Barsotti–Tate groups. The objects of  $DEF^?$  is given by  $(A_0, D^?, \Lambda^?, \epsilon^?)_{/R}$  such that  $D^?$  is a Barsotti–Tate  $O$ -module deforming  $A_0[p^\infty]^? = \prod_{\mathfrak{p} \in \Sigma^?} A_0[\mathfrak{p}^\infty]$  for  $? = ord, ll$  and  $\Lambda^? : D^? \cong {}^tD^?$  is an  $O$ -linear isomorphism and  $\epsilon^? : D_0^? \cong A_0[p^\infty]^?$ . Thus  $D^{ll}$  is local-local (i.e.,  $D^{ll}$  and  ${}^tD^{ll}$  both connected). The quadruple  $(A_0, D^?, \Lambda^?, \epsilon^?)$  is supposed to satisfy the same compatibility of diagrams defining  $DEF$ . Since by  $O$ -linearity, we have a canonical decomposition  $D = D^{ord} \times D^{ll}$  for an object

$(A_0, D, \Lambda, \epsilon)_{/R}$  of  $DEF$  with the maximal ordinary Barsotti–Tate  $O$ -module  $D^{ord}$  in  $D$  and the local-local complement  $D^{ll}$  of  $D^{ord}$  in  $D$  so that  $D = D^{ord} \times D^{ll}$ , we get a decomposition

$$DEF \ni (A_0, D, \Lambda, \epsilon) \mapsto \{(A_0, D^{ord}, \Lambda|_{D^{ord}}, \epsilon|_{D^{ord}}), (A_0, D^{ll}, \Lambda|_{D^{ll}}, \epsilon^{ll})\} \in DEF^{ord} \times DEF^{ll}$$

which induces the equivalence between  $DEF$  and  $DEF^{ord} \times DEF^{ll}$ .

We consider the following deformation functor  $\widehat{D} : CL_{/W} \rightarrow SETS$  given by

$$\widehat{D}(R) = \{(A, \lambda, \theta)_{/R} | (A, \lambda, \theta)_{/R} \in \mathcal{D}\} / \cong.$$

To describe the deformation space representing  $\widehat{D}$ , we write  $O_{\Xi} = \prod_{\mathfrak{l} \in \Xi} O_{\mathfrak{l}}$  for the  $\Xi$ -adic completion of  $O$  for any finite set of prime ideals  $\Xi$  of  $O$ . Then we put  $F_{\Xi} = O_{\Xi} \otimes_O F$  and  $O_{(\Xi)} = F \cap O_{\Xi}$  in  $F_{\Xi}$ .

**Proposition 1.2.** *Let the notation be as above. Suppose that  $p$  is unramified in  $F/\mathbb{Q}$  and in  $k/\mathbb{Q}$ . Then the deformation functor  $\widehat{D}$  is representable by the smooth formal scheme  $\widehat{S}_{/W}$  which is a product of two smooth formal schemes:  $\widehat{S}^{ord}$  isomorphic to  $\widehat{\mathbb{G}}_m \otimes_{\mathbb{Z}_p} O_{\Sigma^{ord}} = \prod_{\mathfrak{p} \in \Sigma^{ord}} \widehat{\mathbb{G}}_m \otimes_{\mathbb{Z}_p} O_{\mathfrak{p}}$  and  $\widehat{S}^{ll}$ , where  $\widehat{S}^{ll} \cong \mathrm{Spf}(W[[T_1, \dots, T_m]])$  (non-canonically) for  $m = \mathrm{rank}_{\mathbb{Z}_p} O_{\Sigma^{ll}}$ , and each factor  $\widehat{S}^?$  represents the formal stack  $DEF^?$ .*

*Proof.* As is well known, the formal stack  $DEF$  over  $W$  is represented by a formal scheme  $\widehat{S}$  smooth over  $W$  of relative dimension equal to  $\mathrm{rank}_{\mathbb{Z}_p} O_p$ . Indeed, adding an  $O$ -linear prime-to- $p$  adelic level structure  $\eta^{(p)} : (\widehat{O}^{(p)})^2 \cong T^{(p)}A = \varprojlim_{p|N} A_0[N](\overline{\mathbb{F}}_p)$  to  $(A_0, \lambda_0, \theta_0)$ , the quadruple  $(A_0, \lambda_0, \theta_0, \eta^{(p)})$  (called a test object) gives rise to an  $\overline{\mathbb{F}}_p$ -point  $x_0$  of the Hilbert modular Shimura variety  $Sh_{/W}^{(p)} = Sh_{/\mathbb{Z}_{(p)}}^{(p)} \times W$  of prime to  $p$  level. We recall in Section 2 the definition of the Shimura variety. Then  $\widehat{S}_{/W}$  can be identified with the formal completion of  $Sh_{/W}^{(p)}$  along  $x_0$  (and the Shimura variety is smooth over  $\mathbb{Z}_{(p)}$  of relative dimension equal to  $\mathrm{rank}_{\mathbb{Z}_p} O_p = [F : \mathbb{Q}]$ ). For the part  $DEF^{ord}$ , by the theory of Serre–Tate, once we identify  $A_0[p^{\infty}]^{ord}$  with  $(\mu_{p^{\infty}} \otimes_{\mathbb{Z}_p} O_{\Sigma^{ord}}) \times (F_{\Sigma^{ord}}/O_{\Sigma^{ord}})$  by a level  $p$ -structure

$$\eta_p : (\mu_{p^{\infty}} \otimes_{\mathbb{Z}_p} O_{\Sigma^{ord}}) \times (F_{\Sigma^{ord}}/O_{\Sigma^{ord}}) \cong A_0[p^{\infty}]^{ord},$$

the space  $\widehat{S}^{ord}$  is isomorphic to  $\widehat{\mathbb{G}}_m \otimes_{\mathbb{Z}_p} O_{\Sigma^{ord}}$  whose origin corresponds to the unique canonical lift of  $A_0[p^{\infty}]^{ord}$  with complex multiplication (which may be different from the starting  $A[p^{\infty}]^{ord}$ ). Since  $DEF \approx DEF^{ord} \times DEF^{ll}$  (equivalence of categories), we have a product decomposition  $\widehat{S} = \widehat{S}^{ord} \times \widehat{S}^{ll}$ . Since  $\widehat{S}$  and  $\widehat{S}^{ord}$  are formally smooth,  $\widehat{S}^{ll}$  must be formally smooth. Then by dimension computation, we have  $\dim_W \widehat{S}^{ll} = m$  as above.  $\square$

We have  $\widehat{S}_p^{ord} \cong \widehat{\mathbb{G}}_m \otimes_{\mathbb{Z}_p} O_{\Sigma^{ord}} = \prod_{\mathfrak{p} \in \Sigma^{ord}} \widehat{S}_p^{ord}$  for  $\widehat{S}_p^{ord} = \widehat{\mathbb{G}}_m \otimes_{\mathbb{Z}_p} O_{\mathfrak{p}}$ . Thus  $\widehat{S}_p^{ord}$  is the  $O_{\mathfrak{p}}$ -factor of the formal completion of  $\mathbb{G}_m \otimes_{\mathbb{Z}} O = \mathrm{Spec}(W[t^{\xi}]_{\xi \in O})$  along the origin corresponding to the maximal ideal  $(p, t-1)$ . Here we used the fact that for the cocharacter group  $L$  of a split torus  $T_{/W}$ , we have a canonical isomorphism:  $T \cong \mathrm{Spec}(W[t^l]_{l \in L}) = \mathbb{G}_m \otimes_{\mathbb{Z}} L$  for the group algebra  $W[L] \cong W[t^l]_{l \in L}$  with the canonical variable  $t$  of  $\mathbb{G}_m$ . Since  $\widehat{\mathbb{G}}_m \otimes_{\mathbb{Z}} O = \widehat{\mathbb{G}}_m \otimes_{\mathbb{Z}} O_p$  ( $O_p = \varprojlim_n O/p^n O$ ),  $\widehat{\mathbb{G}}_m \otimes_{\mathbb{Z}_p} O_{\mathfrak{p}}$  has variable  $t_{\mathfrak{p}} = t \otimes 1_{\mathfrak{p}}$  for the idempotent  $1_{\mathfrak{p}}$  of  $O_{\mathfrak{p}}$  in  $O_p = \prod_{\mathfrak{p}'|p} O_{\mathfrak{p}'}$ . We call  $t_{\mathfrak{p}}$  the Serre–Tate (multiplicative) coordinate of  $\widehat{S}_p^{ord}$ .

## 2. HILBERT MODULAR SHIMURA VARIETY

Here is a more detailed definition of AVRMS by  $O$ . Write  $O^* = \mathrm{Hom}_{\mathbb{Z}}(O, \mathbb{Z})$  (the  $\mathbb{Z}$ -linear dual), which can be identified with the different inverse of  $F_{/\mathbb{Q}}$  by the trace pairing. A triple  $(A, \lambda, \theta)_{/S}$  with an abelian scheme  $A$  over a scheme  $S$  with real multiplication is called an AVRMS by  $O$  if it satisfies the following four conditions:

- (rm1)  $\theta = \theta_A : O \hookrightarrow \mathrm{End}(A_{/S})$  is an embedding of algebras taking the identity to the identity.
- (rm2)  $\lambda$  is an  $O$ -linear symmetric isogeny  $\lambda : A \rightarrow {}^t A$  induced by an ample line bundle fiber-by-fiber geometrically (see [GIT] 6.2). Identifying  ${}^t A$  with  $A \otimes_O \mathfrak{c}$  for a fractional ideal  $\mathfrak{c}$  of  $F$ ,  $\lambda$  is called a  $\mathfrak{c}$ -polarization of  $A$ . Here  $\lambda$  is called *symmetric* if  $\lambda = {}^t \lambda$ .

- (rm3) The image of  $\theta_A$  is stable under the Rosati involution on the endomorphism algebra  $\text{End}^0(A)$ :  $\alpha \mapsto \alpha^* = \lambda^{-1} \circ {}^t\alpha \circ \lambda$ .
- (rm4) As  $O \otimes_{\mathbb{Z}} \mathcal{O}_S$ -modules, we have an isomorphism:  $\text{Lie}(A) \cong O \otimes_{\mathbb{Z}} \mathcal{O}_S (\Leftrightarrow \pi_*(\Omega_{A/S}) \cong O^* \otimes_{\mathbb{Z}} \mathcal{O}_S$  for  $\pi : A \rightarrow S$ ) locally under the Zariski topology of  $S$ , where the sheaf  $\text{Lie}(A)$  of Lie algebras of  $A$  (i.e., the pull-back of the tangent bundle over  $A/S$  by the 0-section) is an  $O$ -module by the action induced from  $\theta$ .

An AVRМ  $A$  defined over a field  $\kappa$  (potentially) has complex multiplication (or has CM) if the algebra  $\text{End}_F^0(A \times_{\kappa} \bar{\kappa}) = \text{End}_O(A \times_{\kappa} \bar{\kappa}) \otimes_O \mathbb{Q}$  (of  $F$ -linear endomorphisms of  $A \times_{\kappa} \bar{\kappa}$  defined over  $\bar{\kappa}$ ) for an algebraic closure  $\bar{\kappa}$  of  $\kappa$  is not equal to  $F$  (and if  $\kappa \subset \bar{\mathbb{Q}}$  and  $A$  has CM,  $\text{End}_F^0(A \times_{\kappa} \bar{\kappa})$  is a quadratic extension of  $F$ ). An AVRМ  $A$  over an integral domain  $\mathfrak{r}$  has CM if  $A \times_{\mathfrak{r}} \kappa$  has CM for the quotient field  $\kappa$  of  $\mathfrak{r}$ . An AVRМ  $A_{/\mathfrak{r}}$  is said to be *non CM* if  $\text{End}_F^0(A \times_{\mathfrak{r}} \bar{\kappa}) = F$ .

In the rest of the paper, we denote by  $\Xi$  a finite set of rational primes unramified in  $F$  containing the fixed prime  $p$ , and we put  $\mathbb{Z}_{(\Xi)} = \bigcap_{l \in \Xi} (\mathbb{Q} \cap \mathbb{Z}_l) \subset \mathbb{Q}$  (the localization at  $\Xi$  of  $\mathbb{Z}$ ). We write  $O_{(\Xi)}$  for the ring  $O \otimes_{\mathbb{Z}} \mathbb{Z}_{(\Xi)} \subset F$  and let  $O_{(\Xi)+}^{\times}$  be the group of totally positive units in  $O_{(\Xi)}$ . Let  $\mathcal{A}^{(\Xi)}$  be the fiber category over  $\mathbb{Z}_{(\Xi)}$ -schemes made of abelian schemes with real multiplication by  $O$  up to prime-to- $\Xi$  isogenies. Thus the objects of  $\mathcal{A}^{(\Xi)}$  are triples  $(A, \bar{\lambda}, \theta_A)_{/S}$  over a  $\mathbb{Z}_{(\Xi)}$ -scheme  $S$ , and  $\bar{\lambda}$  is the set  $\{a\lambda := \lambda \circ \theta_A(a)\}_{a \in O_{(\Xi)+}^{\times}}$  for a polarization  $\lambda$  with polarization ideal prime to  $\Xi$ . The morphisms:  $(A, \bar{\lambda}, \theta_A)_{/S} \rightarrow (A', \bar{\lambda}', \theta_{A'})_{/S}$  of  $\mathcal{A}^{(\Xi)}$  are given by the elements  $\phi : A \rightarrow A'$  in  $\text{Hom}_{G\text{SCH}/\mathbb{Z}_{(\Xi)}}(A, A') \otimes_{\mathbb{Z}} \mathbb{Z}_{(\Xi)}$  compatible with the data  $(\bar{\lambda}, \theta_A)$  and  $(\bar{\lambda}', \theta_{A'})$ . Here  $G\text{SCH}/R$  stands for the category of group schemes over a ring  $R$ . We consider the Hilbert modular Shimura variety  $Sh_{/\mathbb{Z}_{(\Xi)}}^{(\Xi)}$  which is known to represent the stack made out of  $(A, \bar{\lambda}, \theta_A, \eta^{(\Xi)})$  for a prime-to- $\Xi$   $O$ -linear level structure  $\eta^{(\Xi)} : (F_{\mathbb{A}}^{(\Xi, \infty)})^2 \cong T^{(\Xi)} A \otimes_{\mathbb{Z}} \mathbb{Z}_{(\Xi)}$  sending the determinant pairing  $(a, b) \mapsto \det(a, b)$  for  $a, b \in (F_{\mathbb{A}}^{(\Xi, \infty)})^2$  to the pairing on  $T^{(\Xi)} A$  induced by the polarization up to similitude factors in  $(F_{\mathbb{A}}^{(\Xi, \infty)})^{\times}$ , where  $T^{(\Xi)} A$  is the prime-to- $\Xi$  Tate module of  $A$ ,  $F_{\mathbb{A}}$  is the adèle ring of  $F$  and  $F_{\mathbb{A}}^{(\Xi, \infty)}$  is its prime-to- $\Xi$  part given by  $\{x \in F_{\mathbb{A}} | x_l = x_{\infty} = 0 \text{ for all } l \in \Xi\}$ . Strictly speaking, to get  $Sh_{/\mathbb{Z}_{(l)}}^{(\Xi)}$  ( $l \in \Xi$ ), for  $\Xi^{(l)} := \Xi \setminus \{l\}$ , we need to actually identify the  $\Xi^{(l)}$ -part of the Tate module  $T_{\Xi^{(l)}} A$  with  $O_{\Xi^{(l)}}^2$  (without specifying an isomorphism; see [H06] §1 for more details). Then the schemes  $\{Sh_{/\mathbb{Z}_{(l)}}^{(\Xi)}\}_{l \in \Xi}$  glue into  $Sh_{/\mathbb{Z}_{(\Xi)}}^{(\Xi)}$ . The pro-scheme  $Sh^{(\Xi)}$  is smooth over  $\mathbb{Z}_{(\Xi)}$  (cf. [PAF] Section 4.2).

Let  $G = \text{Res}_{F/\mathbb{Q}} GL(2)$  with center  $Z_{/\mathbb{Q}}$ . Under the action  $\eta^{(\Xi)} \mapsto \eta^{(\Xi)} \circ g$ , each  $g \in G(\mathbb{A}^{(\Xi, \infty)})$  acts on  $Sh^{(\Xi)}$  as an automorphism. Let  $\mathcal{W}_{\Xi} = \bigcap_{l \in \Xi} \mathcal{W}_l$  (inside  $\bar{\mathbb{Q}}$ ) which is a henselian semi-local Dedekind domain with localization at each maximal ideal  $\mathfrak{m}_l$  of residual characteristic  $l$  equal to  $\mathcal{W}_l$ . We fix a non CM test object  $(A, \bar{\lambda}, \theta, \eta^{(\Xi)})_{/\mathcal{W}_{\Xi}}$  satisfying (NLL). Let  $x \in Sh^{(\Xi)}(\mathcal{W}_{\Xi})$  be the point representing this test object.

Suppose  $p \in \Xi$ . By a theorem of Tate (cf. [ABV] Appendix I), the reduction  $(A_0, \bar{\lambda}_0, \theta_0, \eta_0^{(\Xi)})_{/\bar{\mathbb{F}}_p} = (A, \lambda, \theta, \eta^{(\Xi)})_{/\mathcal{W}_{\Xi}} \times_{\mathcal{W}_{\Xi}} \bar{\mathbb{F}}_p$  has complex multiplication by a field  $M = \text{End}_F^0(A_0/\bar{\mathbb{F}}_p)$  ( $F$ -linear endomorphism algebra). Regarding  $M \subset \bar{\mathbb{Q}}$  as an  $F$ -algebra, we write  $\tilde{\theta}_{\mathfrak{p}} : M \cong \text{End}_F^0(A_0/\bar{\mathbb{F}}_p)$  for the identification (so,  $\tilde{\theta}_{\mathfrak{p}}|_F = \theta_0$ ). Let  $R = \text{End}_O(A_0/\bar{\mathbb{F}}_p)$  which is an order of  $M$ . The quadruple  $(A_0, \bar{\lambda}_0, \theta_0, \eta_0^{(\Xi)})_{/\bar{\mathbb{F}}_p}$  gives rise to a point  $x_0 \in Sh^{(\Xi)}(\bar{\mathbb{F}}_p)$ . Let  $V$  be the geometrically irreducible component of  $Sh_{/\mathcal{W}_{\Xi}}^{(\Xi)}$  containing  $x$ ; so,  $V_{/\bar{\mathbb{F}}_p} = V \times_{\mathcal{W}_{\Xi}} \bar{\mathbb{F}}_p$  is geometrically irreducible and contains  $x_0$ . By the smoothness of  $Sh^{(\Xi)}$  over  $\mathbb{Z}_{(\Xi)}$  and the existence of the smooth projective toroidal compactification of  $Sh^{(\Xi)}$ ,  $V_{/\bar{\mathbb{F}}_p}$  is geometrically connected.

By the global reciprocity law of Shimura (cf. [Sh] II), we know

**Theorem 2.1.** *Let the notation and the assumption be as above. Suppose  $p \in \Xi$  which is unramified in  $F/\mathbb{Q}$ . Let  $\kappa = \bar{\mathbb{F}}_p$  or  $\bar{\mathbb{Q}}$ . For the geometrically connected component  $V_{/\kappa} \subset Sh_{/\mathbb{Z}_{(\Xi)}}^{(\Xi)} \times_{\mathbb{Z}_{(\Xi)}} \kappa$ , the scheme automorphism group  $\text{Aut}(V_{/\kappa}) \subset G(\mathbb{A}^{(\Xi, \infty)})$  is given by the semi-direct product of the field*

automorphism group  $\text{Aut}(F/\mathbb{Q})$  and

$$\bar{\mathcal{E}}^{(\Xi)} = \frac{\{g \in G(\mathbb{A}^{(\Xi)}) \mid \det(g) \in \overline{O_{(\Xi)+}^{\times} F_{\infty+}^{\times}}\}}{Z(\mathbb{Z}(\Xi))G(\mathbb{R})_+} = \frac{\{g \in G(\mathbb{A}^{(\Xi),\infty}) \mid \det(g) \in \overline{O_{(\Xi)+}^{\times}}\}}{Z(\mathbb{Z}(\Xi))},$$

where  $G(\mathbb{R})_+$  is the identity connected component of  $G(\mathbb{R})$  and “ $\overline{\phantom{x}}$ ” indicates closure under adelic topology. Here  $\text{Aut}(F/\mathbb{Q})$  acts on  $\bar{\mathcal{E}}^{(\Xi)}$  through its action on  $G(\mathbb{A})$ . For the above point  $x_0 \in \text{Sh}^{(\Xi)}(\overline{\mathbb{F}}_p)$ , the stabilizer of  $x_0$  in  $\bar{\mathcal{E}}^{(\Xi)}$  is the image of the morphism  $\hat{\rho}: R_{(\Xi)}^{\times} \rightarrow G(\mathbb{A}^{(\Xi),\infty})$  given by  $\tilde{\theta}_{\mathfrak{p}}(\alpha) \circ \eta_0^{(\Xi)} = \eta_0^{(\Xi)} \circ \hat{\rho}(\alpha)$  for  $\alpha \in R_{(\Xi)}^{\times} = (R \otimes_{\mathbb{Z}} \mathbb{Z}(\Xi))^{\times} \subset M^{\times}$ .

By this theorem, the stabilizer  $\mathcal{T}$  in  $\bar{\mathcal{E}} = \text{Aut}(V/\overline{\mathbb{F}}_p)$  of  $x_0$  is the image of  $R_{(\Xi)}^{\times}$  for the order  $R$  in the integer ring  $\mathfrak{o}$  of  $M$  under the morphism  $\hat{\rho}: T_M = \text{Res}_{R/\mathbb{Z}} \mathbb{G}_m \rightarrow G$  of group schemes defined over  $\mathbb{A}^{(\Xi),\infty}$  (so, it is a conjugate of an anisotropic torus in  $G$  under an element in  $G(\mathbb{A}^{(\Xi),\infty})$ ). It would appear later that we do not use this theorem, but it will be implicitly used to assure that our action of  $T_M(\mathbb{Z}(\Xi))$  preserves each geometrically irreducible component of  $\text{Sh}^{(\Xi)}$  as its action factors through  $\bar{\mathcal{E}}^{(\Xi)}$ .

*Proof.* The first assertion is proven by T. Miyake in [M] for  $\kappa = \overline{\mathbb{Q}}$  (following the method of Shimura in [Sh] II) via the well known identification of the automorphism group of a bounded symmetric domain with the corresponding Lie group (e.g., [DLS] Chapter IV); in our Hilbert modular case,  $\text{Aut}(\mathfrak{H}^I)$  ( $\mathfrak{H} = \{z \in \mathbb{C} \mid \text{Im}(z) > 0\}$ ) has identity connected component isomorphic to  $PSL_2(\mathbb{R})^I$  for  $I = \text{Hom}_{\text{field}}(F, \overline{\mathbb{Q}})$ . See [PAF] Theorem 4.14 for another algebraic proof for  $\kappa = \overline{\mathbb{Q}}$ , and the case where  $\kappa = \overline{\mathbb{F}}_p$  is treated in [H06]. If  $g \in \bar{\mathcal{E}}^{(\Xi)}$  stabilizes  $x_0$ , by universality, we have a prime-to- $\Xi$   $F$ -linear endo-isogeny

$$\alpha: (A_0, \bar{\lambda}_0, \theta_0, \eta_0^{(\Xi)})_{/\overline{\mathbb{F}}_p} \cong (A_0, \bar{\lambda}_0, \theta_0, \eta_0^{(\Xi)} \circ g)_{/\overline{\mathbb{F}}_p}.$$

In other words,  $\alpha$  can be regarded as an element of  $R_{(\Xi)}^{\times}$ . Since  $\tilde{\theta}_{\mathfrak{p}}(\alpha) \circ \eta^{(\Xi)} = \eta^{(\Xi)} \circ \hat{\rho}(\alpha)$ , the matrix  $\hat{\rho}(\alpha)$  coincides with  $g$  in the quotient  $\bar{\mathcal{E}}^{(\Xi)}$ .  $\square$

**Lemma 2.2.** *Suppose that the AVRMS  $(A_0, \bar{\lambda})_{/\overline{\mathbb{F}}_p}$  is defined over a finite field  $\mathbb{F}_q \subset \overline{\mathbb{F}}_p$  and  $\mathfrak{p} \in \Sigma^{\text{ord}}$ . Then the algebra  $M = \text{End}_F^0(A_0/\overline{\mathbb{F}}_p)$  is a CM quadratic extension  $M$  over  $F$  generated by the Frobenius endomorphism over  $\mathbb{F}_q$ . In particular, the prime  $\mathfrak{p}$  in  $F$  splits into  $\mathcal{P}\bar{\mathcal{P}}$  in  $R$  for primes  $\mathcal{P} \neq \bar{\mathcal{P}}$ .*

By this lemma, we have  $R_{\mathfrak{p}} = \mathfrak{o}_{\mathfrak{p}} = R_{\mathcal{P}} \times R_{\bar{\mathcal{P}}} \cong O_{\mathfrak{p}} \times O_{\mathfrak{p}}$  for the integer ring  $\mathfrak{o}$  of  $M$ .

*Proof.* By definition,  $M_{\mathfrak{p}}$  is isomorphic to  $F$ -linear endomorphism algebra  $\text{End}_F^0(A_0[\mathfrak{p}^{\infty}]_{/\mathbb{F}_q})$ , which is therefore a semi-simple quadratic extension of  $F_{\mathfrak{p}}$ . Thus  $M$  is commutative, and the embedding restricted to  $M \subset M_{\mathfrak{p}}$  factors through  $\text{End}_F(A_0/\mathbb{F}_q)$  and its image contains the  $q$ -th power Frobenius map  $\phi$ . Since  $F[\phi]/F$  is a CM field because of the positivity and non-triviality of Rosati involution  $c$  of  $\lambda_0$ , we must have  $M = F[\phi]$  which is a CM quadratic extension of  $F$ .

Note that  $\phi$  and its dual  $\phi^c = \lambda^{-1} \circ {}^t\phi \circ \lambda$  satisfy  $\phi\phi^c = \phi^c\phi = q$ . Since  $\phi$  is invertible on  $A_0(\overline{\mathbb{F}}_p) \supset A_0[\mathfrak{p}^{\infty}] \neq 0$ ,  $a = \phi + \phi^c \in O$  is prime to  $\mathfrak{p}$ . Thus the minimal polynomial  $X^2 - aX + q$  of  $\phi$  over  $F$  satisfies  $X^2 - aX + q \equiv X(X - a) \pmod{\mathfrak{p}}$  with  $(a \pmod{\mathfrak{p}}) \neq 0$ ; so,  $\mathfrak{p}$  must split in  $R \supset O[\phi]$ .  $\square$

We often choose  $\mathcal{P}$  so that  $\mathcal{P}$  contains  $\phi$ ; so,  $A_0[\mathcal{P}]$  is connected.

### 3. EIGEN DIFFERENTIALS

We keep the notation introduced in Theorem 2.1 in the rest of the paper. For an open compact subgroup  $K$  of  $G(\mathbb{A}^{(\Xi),\infty})$ , we write  $V_K$  for the image of  $V$  under  $V \subset \text{Sh}^{(\Xi)} \rightarrow \text{Sh}_K^{(\Xi)}$  for the

geometric quotient  $Sh_K^{(\Xi)} := Sh^{(\Xi)}/K$  (thus  $V$  is the pro-variety  $V = \varprojlim_K V_K$  in the sense of Grothendieck [EGA] IV.8 or [EAI] §5.1.4). We take  $K$  sufficiently small so that  $K$  is neat; i.e.,

$$\Gamma_{gKg^{-1}} = \{\alpha \in gKg^{-1} \cap G(\mathbb{Q}) \mid \det(\alpha) \text{ is totally positive}\}$$

does not have torsion modulo its center (this holds true if  $K$  is inside the principal congruence subgroup of level  $N \geq 3$ ). Under neatness,  $V_K$  is smooth over  $\mathcal{W}_\Xi$ .

Let  $\pi : (\mathbf{A}, \theta_{\mathbf{A}}) \rightarrow V_K$  be the universal AVRМ over  $V_K$ . Then  $\underline{\omega} = \pi_* \Omega_{\mathbf{A}/V_K}$  has an  $\mathcal{O}$ -action via  $\theta_{\mathbf{A}}$  and by (rm4), is locally free of rank 1 over  $\mathcal{O} \otimes_{\mathbb{Z}} \mathcal{O}_{V_K}$  (which is a module over  $\mathcal{O} \otimes_{\mathbb{Z}} \mathcal{W}_\Xi$ ). Since  $\mathcal{O} \otimes_{\mathbb{Z}} \mathcal{W}_\Xi = \mathcal{W}_\Xi^I$  by sending  $a \otimes b$  to  $\varphi(a)b$  for  $\varphi \in I = \text{Hom}_{\text{field}}(F, \mathbb{Q})$  (as primes in  $\Xi$  are unramified in  $\mathcal{O}$ ), we can accordingly split  $\underline{\omega} = \bigoplus_{\varphi \in I} \underline{\omega}^\varphi$  for a line bundle  $\underline{\omega}^\varphi$  on which the  $\mathcal{O}$ -action factors through  $\varphi$ . By the Kodaira-Spencer map (see [K] §3.6),  $\Omega_{V_K/\mathcal{W}_\Xi}$  is canonically isomorphic to  $\bigoplus_{\varphi} \underline{\omega}^{2\varphi}$  for  $\underline{\omega}^{2\varphi} = \underline{\omega}^\varphi \otimes_{\mathcal{O}_{V_K}} \underline{\omega}^\varphi$  (see [K1] 1.0.21). Thus the cotangent space  $\Omega_{V_K/\mathcal{W}}$  has a canonical  $\mathcal{O}_{(\Xi)}^\times$ -action. A section  $T$  of the structure sheaf of a Zariski (resp. complex analytic,  $p$ -adic rigid analytic) neighborhood  $D_x$  (of  $x$ ) is called “belonging to  $\varphi$ ” if  $dT \in H^0(D_x, (\underline{\omega}^{2\varphi})^*)$  for the line bundle  $(\underline{\omega}^{2\varphi})^*$  corresponding to the algebraic bundle  $\underline{\omega}^{2\varphi}$  (by GAGA theorems; see Example 2.4.6 in [C]). Here we use  $*$  =  $an$  in the complex analytic case and in the rigid analytic case (and we place nothing in the algebraic case; if we need to specify the place  $v$ , we write  $*$  =  $v$ -an). If  $T$  is a part of the coordinate system belonging to  $\varphi$  (at  $v \in \Xi \cup \{\infty\}$ ), we call  $T$  a  $\varphi$ -coordinate (at  $v$ ).

Recall  $M = \text{End}_F^0(A_0) \supset \text{End}_{\mathcal{O}}(A_0) = R$ . The field  $M$  is a CM quadratic extension of  $F$ , and let  $T_M = \text{Res}_{R/\mathbb{Z}} \mathbb{G}_m$ . Recall the morphism  $\hat{\rho} : T_M \rightarrow G(\mathbb{A}^{(\Xi), \infty})$  of algebraic groups over  $\mathbb{A}^{(\Xi), \infty}$  given by  $\hat{\theta}_{\mathfrak{q}}(\alpha) \circ \eta_0^{(\Xi)} = \eta_0^{(\Xi)} \circ \hat{\rho}(\alpha)$ . The point  $x_0 \in V(\overline{\mathbb{F}}_p)$  is a fixed point of  $\hat{\rho}(T_M(\mathbb{Z}(\Xi)))$ , and let  $U$  be a Zariski open neighborhood of  $x_0$ . Suppose that we have a non-constant  $f \in \overline{\mathbb{F}}_p(V)$  vanishing at  $x_0$  and a  $p$ -adically open subgroup  $\mathcal{T}$  of  $T_M(\mathbb{Z}(\Xi))$  such that  $f \circ \hat{\rho}(\alpha) = \lambda_\alpha f$  with a constant  $\lambda_\alpha \in \overline{\mathbb{F}}_p$  for all  $\alpha \in \mathcal{T}$ . Let  $X^\circ = \{u \in U \mid f(u) = 0\}$  and  $X$  be the Zariski closure of  $X^\circ$ . Since  $f$  is non-constant,  $\dim X = d - 1$ . Then, by definition,  $X$  is stable under the action of the image  $\mathcal{T} \subset \overline{\mathcal{E}}^{(\Xi)}$  of  $\hat{\rho}(T_M(\mathbb{Z}(\Xi)))$ . Supposing that  $x_0$  is in the ordinary locus, by [H10] Proposition 3.8,  $X = V$  if  $\dim X = d - 1 > 0$ . This is impossible, so such a nontrivial  $f$  does not exist.

Suppose that we have a flat integral covering space  $\tilde{U}/U$  whose image contains  $x_0$  on which the action of  $\mathcal{T}$  extends. If  $f \circ \hat{\rho}(\alpha) = \lambda_\alpha f$  with a constant  $\lambda_\alpha \in \overline{\mathbb{F}}_p$  for all  $\alpha \in \mathcal{T}$  for  $f \in \overline{\mathbb{F}}_p(\tilde{U})$ , we consider the image  $X^\circ \subset U$  of  $\{u \in \tilde{U} \mid f(u) = 0\}$  and its Zariski closure  $X$ . By the same argument, we have  $X = V$  if  $d - 1 > 0$ . Thus even in  $\overline{\mathbb{F}}_p(\tilde{U})$ , such an  $f \neq 0$  does not exist. Thus if  $0 \neq f \in \hat{\mathcal{O}}_{V, x_0}$  satisfies this property,  $f$  is transcendental over  $\mathcal{O}_{V, x_0}$  if  $[F : \mathbb{Q}] \geq 2$ . We will not use this transcendence result in the rest of the paper.

#### 4. EIGEN COORDINATES

Let  $V$  be the geometrically irreducible component of  $Sh_{/\mathcal{W}_\Xi}^{(\Xi)}$  as in Theorem 2.1 containing the point  $x \in V(\mathcal{W}_\Xi)$ , and take a neat open compact subgroup  $K$  of  $G(\mathbb{A}^{(\Xi), \infty})$ . As before, let  $x_0$  be the image of  $x$  in  $V(\overline{\mathbb{F}}_p)$  (assuming  $p \in \Xi$ ). Recall  $V_K$  defined in Section 3 which is the geometrically irreducible component of  $Sh_K^{(\Xi)}$  containing the projection of  $x$ . Let  $\mathcal{O}$  be the stalk  $\mathcal{O}_{V_K/\mathcal{W}_\Xi, x} \subset \mathcal{O}_{V/\mathcal{W}_\Xi, x} = \mathcal{O}_{Sh^{(\Xi)}, x}$  at the  $\mathcal{W}_\Xi$ -point  $x \in V(\mathcal{W}_\Xi)$ . The  $\mathcal{W}_\Xi$ -algebra  $\mathcal{O}$  is semi-local (of finite type), and its maximal ideal  $\mathfrak{m}_{l, x}$  over  $(l) \subset \mathcal{W}_\Xi$  for  $l \in \Xi$  is unique. Since  $V/V_K$  is étale, we may identify the  $P$ -adic localization-completion of  $\mathcal{O}$  and that of  $\mathcal{O}_{V, x}$  for any prime ideal  $P$  of  $\mathcal{O}_{V, x}$ . We consider the prime  $P = P_x$  given by the kernel of  $\mathcal{O} \rightarrow \mathcal{W}_\Xi$  induced by evaluation at  $x$ . Then  $P$  is a prime ideal inside the Jacobson radical of  $\mathcal{O}$  (i.e., the intersection of all the maximal ideals  $\{\mathfrak{m}_{l, x}\}_{l \in \Xi}$  of  $\mathcal{O}$  indexed by the residual characteristic  $l \in \Xi$  of  $\mathfrak{m}_{l, x}$ ). Let  $\mathcal{K}$  be the field of fraction of  $\mathcal{W}_\Xi$ , and consider the localization completion  $\hat{\mathcal{O}}_P \cong \mathcal{K}[[T_1, \dots, T_d]]$  for  $d$  variables  $T_j$ . We can extend scalar to  $\mathbb{C}_v$  for  $v = l, \infty$  by  $\mathcal{K} \subset \overline{\mathbb{Q}} \xrightarrow{i_v} \mathbb{C}_v$  and consider  $\mathcal{O}_v = \mathcal{O} \otimes_{\mathcal{W}_\Xi} \mathbb{C}_v$ . Then the evaluation at  $x \in V(\mathbb{C}_v)$  gives rise to a  $\mathbb{C}_v$ -algebra homomorphism  $\mathcal{O}_v \rightarrow \mathbb{C}_v$  whose kernel, we denote by  $P_v$ ; so,  $P = P_v \cap \mathcal{O}$ . Then we localize  $\mathcal{O}_v$  at  $P_v$  and write the localization as  $\mathcal{O}_{v, P_v}$ . We then complete it  $P_v$ -adically,

getting  $\widehat{\mathcal{O}}_{v, P_v} = \varprojlim_n \mathcal{O}_{v, P_v} / P_v^n \mathcal{O}_{v, P_v}$ . We then have a commutative diagram of inclusions

$$\begin{array}{ccc} \mathcal{O} & \xrightarrow{\quad \subset \quad} & \widehat{\mathcal{O}}_P = \mathcal{K}[[T_1, \dots, T_d]] \\ \cap \downarrow & & i_v \downarrow \cap \\ \widehat{\mathcal{O}}_{v, P_v} & \xlongequal{\quad} & \mathbb{C}_v[[T_1, \dots, T_d]]. \end{array}$$

As described above, we study coordinates around a point on  $\widehat{S}^{ord}$ . The coordinate can be a formal function on  $\widehat{S}^{ord}$  or a rigid analytic function defined on a rigid open neighborhood of the point in the associated  $p$ -adic rigid analytic space denoted by  $(\widehat{S}^{ord})^{p\text{-an}}$ . Here is what we mean by the rigid analytic space associated to  $\widehat{S}^{ord}$ . A functor associating to a formal scheme  $\text{Spf}(A)$  a rigid analytic space is studied by Raynaud and Berthelot. The theory of Raynaud is explained in [BL], and one can find a good account of that of Berthelot in [dJ] Section 7. As a special case of his theory, Berthelot associates, to  $\text{Spf}(W_p[[ (T) ]]/(f_1(T), \dots, f_m(T)))$  for a finite set of variable  $(T) = (T_1, \dots, T_n)$  and a finite set of power series  $f_j(T) \in W[[ (T) ]]$  :=  $W[[T_1, T_2, \dots, T_n]]$ , the rigid analytic space  $X^{rig}$  given by the zero set of  $\{f_j\}_j$  in the open unit disk  $D^n$  in the  $n$ -dimensional affine space over the field of fraction  $\text{Frac}(W)$  of  $W$ . Here we take the base-change of  $X^{rig}$  to  $\mathbb{C}_p$  (see [NAA] §9.3.6), and write it as  $X_{\mathbb{C}_p}^{p\text{-an}}$ . This construction appears very naive, but as exposed in [dJ], this functor has intrinsic meaning and functorial construction, and it extends naturally the results of Raynaud to formal schemes not necessarily of finite type over  $W$ . For  $\mathfrak{p}_\varphi \in \Sigma^{ord}$  associated to  $i_p \circ \varphi : F \rightarrow \mathbb{C}_p$ , we extend  $i_p \circ \varphi$  to a projection  $W_p \otimes_{\mathbb{Z}} \mathcal{O} = W_p^I \rightarrow \mathbb{C}_p$  (which is still denoted by  $i_p \circ \varphi$ ), and we put  $\tau_\varphi = \tau_{\varphi, p} = \log_p \circ i_p \circ \varphi \circ t_{\mathfrak{p}_\varphi}$  for the Serre–Tate coordinate  $t_{\mathfrak{p}_\varphi}$  on  $\widehat{S}_{\mathfrak{p}_\varphi}^{ord} = \widehat{\mathbb{G}}_m \otimes_{\mathbb{Z}_p} \mathcal{O}_{\mathfrak{p}_\varphi}$ . Then a convergent power series in  $\mathbb{C}_p\{\{\tau_\varphi\}\}_{\varphi \in \Sigma^{ord}}$  is a rigid analytic function well defined on  $(\widehat{S}^{ord})^{p\text{-an}}$ , where  $\mathbb{C}_p\{\{\tau_\varphi\}\}$  is made up of power series  $\sum_{n=(n_\varphi)_{\geq 0}} a_n \tau^n$  in  $\mathbb{C}_p[[\tau_\varphi]]_{\varphi \in \Sigma^{ord}}$  with  $\lim_{|n| \rightarrow \infty} |a_n|_p = 0$  for  $|n| = \sum_\varphi n_\varphi$ . The GAGA type theorems associating an analytic sheaf  $\mathfrak{F}^{p\text{-an}}$  (again  $\mathfrak{F}^{p\text{-an}}$  is the base change to  $\mathbb{C}_p$  of  $\mathfrak{F}^{rig}$  in [dJ]) to a formal coherent sheaf  $\mathfrak{F}$  are given in this setting in [dJ] §7.1.11, and the analytic/formal sheaf of Kähler differential is discussed in [dJ] §7.1.12. In particular, for a rigid point  $x \in X^{p\text{-an}}$  coming from a smooth formal point  $x \in X(W)$  in the generic fiber, the formal cotangent space  $\Omega_{X/W}(x) \otimes_W \mathbb{C}_p$  and the analytic cotangent space  $\Omega_{X^{p\text{-an}}/\mathbb{C}_p}(x)$  at  $x$  are identical.

For an  $l$ -adic Barsotti–Tate  $\mathcal{O}$ -module  $B$  defined over a ring  $C$ , we write  $\text{End}_{\mathcal{O}}(B/C)$  for the ring of  $\mathcal{O}$ -linear endomorphisms of  $B$  defined over a ring  $C$  and put  $\text{End}_F^0(B/C) = \text{End}_{\mathcal{O}}(B/C) \otimes_{\mathbb{Z}} \mathbb{Q}$ . For a finite extension  $k$  of  $\mathbb{Q}$  in  $\overline{\mathbb{Q}}$  with integer ring  $\mathfrak{D}$ , put  $\mathcal{W}_k = \mathcal{W}_{\Xi} \cap k$ .

**Lemma 4.1.** *Let  $l \in \Xi$  be a prime. Let  $E/F$  be a quadratic extension with  $T_E = \text{Res}_{E/\mathbb{Q}} \mathbb{G}_m$ . Suppose that  $(A, \overline{\lambda}, \theta, \eta^{(\Xi)})$  is the test object sitting over  $x \in V(\mathcal{W}_{\Xi})$  such that the image of  $x$  in  $V_K$  (for a neat open compact subgroup  $K$  of  $G(\mathbb{A}^{(\Xi), \infty})$ ) falls in  $V_K(\mathcal{W}_k)$  for a finite extension  $k/\mathbb{Q}$ . For  $\mathfrak{L} = \mathcal{W}_k \cap \mathfrak{m}_{l, x}$ , let  $(A_{\mathfrak{L}}, \overline{\lambda}_{\mathfrak{L}}, \theta_{\mathfrak{L}}, \eta_{\mathfrak{L}}^{(\Xi)})$  be its reduction modulo  $\mathfrak{m}_{l, x}$  sitting over  $x_{\mathfrak{L}} \in V(\overline{\mathbb{F}}_l)$  (so  $A_{\mathfrak{L}}$  is defined over  $\mathbb{F}_{\mathfrak{L}} = \mathfrak{D}/\mathfrak{L}$ ). Suppose that we have a level  $l$ -structure  $\eta_l : (\mu_{l^\infty} \otimes_{\mathbb{Z}_l} \mathcal{O}_l) \cong A[l^\infty]^\circ$  defined over  $\mathcal{W}_l$  for a prime  $l \in \Sigma_l^{ord}$  such that  $\eta_l \oplus {}^t\eta_l^{-1} : (\mu_{l^\infty} \otimes_{\mathbb{Z}_l} \mathcal{O}_l) \oplus (F_l/\mathcal{O}_l) \cong A[l^\infty]$  over  $\mathcal{W}_l$  as Barsotti–Tate  $\mathcal{O}$ -modules, where  $A[l^\infty]^\circ$  is the connected component of the Barsotti–Tate group  $A[l^\infty]_{/\mathcal{W}_l}$  and  ${}^t\eta_l^{-1}$  is the dual inverse of  $\eta_l$  under the Cartier duality induced by the polarization  $\lambda$ . If we have embeddings  $\widehat{\rho} : T_E(\mathbb{Q}) \rightarrow G(\mathbb{A}^{(\Xi), \infty})$  and  $E \ni \alpha \mapsto \alpha_q \in \text{End}_F^0(A_{\mathfrak{L}}[q^\infty]_{/\mathbb{F}_{\mathfrak{L}}})$  for all primes  $q$  outside  $\Xi$  satisfying  $\eta_{\mathfrak{L}, q}^{(\Xi)} \circ \widehat{\rho}(\alpha) = \alpha_q \circ \eta_{\mathfrak{L}, q}^{(\Xi)}$  for all  $q \notin \Xi$  for the  $q$ -component  $\eta_q^{(\Xi)}$  of the level structure  $\eta^{(\Xi)}$ , then*

- (1)  $E$  is a CM field isomorphic to  $\text{End}_F^0(A_{\mathfrak{L}}/\overline{\mathbb{F}}_l)$ ,
- (2)  $\widehat{\rho}(\mathfrak{r}_{(\Xi)}^\times)$  ( $\mathfrak{r}_{(\Xi)} = \mathfrak{r} \otimes_{\mathbb{Z}} \mathbb{Z}_{(\Xi)}$ ) fixes  $x_{\mathfrak{L}} \in V(\overline{\mathbb{F}}_l)$  for an order  $\mathfrak{r}$  of  $E$  maximal at  $l$ ,
- (3) we have an  $F$ -linear embedding  $\widetilde{\theta}_{\mathfrak{L}} : E \hookrightarrow \text{End}_F^0(A_{\mathfrak{L}}/\overline{\mathbb{F}}_l)$  such that  $\widetilde{\theta}_{\mathfrak{L}}(\alpha)$  induces  $\alpha_q$  in the endomorphism algebra  $\text{End}_F^0(A_{\mathfrak{L}}[q^\infty])$  for all  $q \notin \Xi$ ,
- (4) there exist an extension of  $\sigma : F \hookrightarrow \overline{\mathbb{Q}}$  to  $E$  (still denoted by  $\sigma$ ) and a  $l$ -adic rigid analytic  $\sigma$ -coordinate  $\tau_\sigma = \tau_{\sigma, l}$  of the rigid analytic space associated to  $\widehat{S}_l^{ord}$  centered at  $x$  in  $\widehat{S}(W_l)$

such that  $\tau_\sigma(x) = 0$  and  $\tau_\sigma \circ \widehat{\rho}(\alpha) = i_l(\alpha^{\sigma(1-c)})\tau_\sigma$  for each  $\sigma$  inducing the place  $l$  on  $F$  after composing  $i_l$ .

Moreover the coordinate  $\tau_\sigma$  as in (4) above is unique up to scalar multiples.

We call a  $\sigma$ -coordinate satisfying (4) above a  $\widehat{\rho}$ -eigen  $\sigma$ -coordinate.

*Proof.* The test object  $(A_\mathfrak{L}, \overline{\lambda}_\mathfrak{L}, \theta_\mathfrak{L})$  is defined over the finite field  $\mathbb{F}_\mathfrak{L} = \mathfrak{D}/\mathfrak{L}$ . Let  $R' = \text{End}_O(A_\mathfrak{L}/\mathbb{F}_\mathfrak{L})$ . Since  $\Sigma_l^{\text{ord}} \neq \emptyset$ , by a theorem of Tate (see [ABV] Appendix I, Theorem 3),  $R'$  is an order of a semi-simple quadratic extension  $M'$  of  $F$  (generated by the Frobenius endomorphism over  $\mathbb{F}_\mathfrak{L}$ ) whose simple components are all CM fields; so,  $M'$  must be a CM quadratic extension (as it cannot be isomorphic to  $F \times F$ ). By another theorem of Tate ([ABV] Appendix I, Theorem 2), we have

$$\text{End}_O(A_\mathfrak{L}[q^\infty]_{/\mathbb{F}_\mathfrak{L}}) = \text{End}_{O[\text{Gal}(\overline{\mathbb{F}}_l/\mathbb{F}_\mathfrak{L})]}(T_q A_\mathfrak{L}) = R' \otimes_{\mathbb{Z}} \mathbb{Z}_q =: R'_q.$$

Since  $E \hookrightarrow \text{End}_F^0(A_\mathfrak{L}[q^\infty]_{/\mathbb{F}_\mathfrak{L}})$  by  $\alpha \mapsto \alpha_q$ , we have  $E_q = E \otimes_{\mathbb{Z}} \mathbb{Z}_q \cong M'_q$  for all primes  $q$  outside  $\Xi$ , and we conclude to have  $\theta_\mathfrak{L} : E \cong M'$ ; so,  $E$  is a CM field. Put  $\mathfrak{r} = \widetilde{\theta}_\mathfrak{L}^{-1}(R') \subset E$ , which is an order of  $E$  maximal at  $l$  by Lemma 2.2 applied to  $l$ . Note that  $E_l \cong F_l \times F_l$ . Since  $\widetilde{\theta}_\mathfrak{L}(\alpha)$  for  $\alpha \in \mathfrak{r}_{(\Xi)}^\times$  is a prime-to- $\Xi$  isogeny sending  $(A_\mathfrak{L}, \overline{\lambda}_\mathfrak{L}, \eta_\mathfrak{L}^{(\Xi)})_{/\overline{\mathbb{F}}_l}$  to  $(A_\mathfrak{L}, \overline{\lambda}_\mathfrak{L}, \widetilde{\theta}_\mathfrak{L}(\alpha) \circ \eta_\mathfrak{L}^{(\Xi)})_{/\overline{\mathbb{F}}_l} = (A_\mathfrak{L}, \overline{\lambda}_\mathfrak{L}, \eta_\mathfrak{L}^{(\Xi)} \circ \widehat{\rho}(\alpha))_{/\overline{\mathbb{F}}_l}$ ,  $\widehat{\rho}(\alpha)$  fixes  $x_\mathfrak{L}$  by the universality of  $Sh_{/\overline{\mathbb{F}}_l}^{(\Xi)}$ .

By Serre–Tate theory, we have the Serre–Tate coordinate  $t := t_l$  of the  $l$ -component  $\widehat{\mathbb{G}}_m \otimes_{\mathbb{Z}_l} O_l$  of  $\widehat{S}_l^{\text{ord}}$  such that an  $O$ -linear automorphism  $a \in \text{Aut}_O(\widehat{\mathbb{G}}_m \otimes O_{l/W_l}) = O_l^\times$  acts by  $t \mapsto t^a$ . As seen in [H10] Lemma 3.3,  $\alpha \in \text{End}_F^0(A_\mathfrak{L}) = E$  acts on  $\widehat{S}_l^{\text{ord}}$  by universality, and this action induces a homomorphism  $\mathfrak{r}_{(\Xi)}^\times \rightarrow \text{Aut}_O(\widehat{\mathbb{G}}_m \otimes O_{l/W_l}) = O_l^\times$  sending  $\alpha \in E^\times$  to  $\alpha^{\sigma(1-c)} \in F_l$  under an extension  $\sigma : E \hookrightarrow \overline{\mathbb{Q}}$  of  $\sigma : F \hookrightarrow \overline{\mathbb{Q}}$  inducing  $l$  and the projection of  $E_l$  to one of its simple components isomorphic to  $F_l$ . Though the proof of [H10] Lemma 3.3 is given for ordinary  $A_\mathfrak{L}$ , the same argument applied to  $A_\mathfrak{L}[l^\infty]$  (in place of  $A_\mathfrak{L}[l^\infty]$  there) works equally well.

Let  $\log_l : 1 + lW_l \rightarrow W_l$  be the  $l$ -adic logarithm map. Write  $\sigma_l$  for the projection of  $W_l \otimes_{\mathbb{Z}} O \subset W_l^I$  to  $W_l$  induced by  $\sigma : O \hookrightarrow \overline{\mathbb{Q}}$ . Then we get a standard  $\sigma$ -coordinate  $\tau_\sigma$  as the following composition:

$$\widehat{\mathbb{G}}_m \otimes_{\mathbb{Z}_l} O_l(W_l) = (1 + lW_l) \otimes_{\mathbb{Z}_l} O_l \xrightarrow{\log_l \otimes 1} W_l \otimes_{\mathbb{Z}_l} O_l \xrightarrow{\sigma_l} W_l.$$

We can perform the same procedure for other primes  $l_\varphi \in \Sigma_l^{\text{ord}}$ , and get a coordinate  $\tau_\varphi$  for  $\varphi \in I$  associated to  $l_\varphi$  satisfying the same property above replacing  $\sigma$  by  $\varphi$ . The associated rigid analytic space  $(\widehat{S}_l^{\text{ord}})^{l\text{-an}}$  has by definition the coordinates  $\{\tau_\varphi\}_{\varphi \in \Sigma_l^{\text{ord}}}$  (here  $\Sigma_l^{\text{ord}}$  is regarded as a set of field embeddings of  $F$  into  $\mathbb{C}_l$ ). Note here  $t_\sigma(x) = 1$  and  $\tau_\sigma(x) = 0$  (by the splitting  $A[l^\infty] \cong (\mu_{l^\infty} \otimes_{\mathbb{Z}_l} O_l) \oplus (F_l/O_l)$  over  $W_l$ ) but for  $\varphi$  not associated to  $l$ ,  $\tau_\varphi(x)$  may not be equal to 0. Then any other rigid analytic coordinate  $f$  centered at  $x$  (i.e.,  $f(x) = 0$ ) is anyway a power series convergent on an open subset of the rigid analytic space  $(\widehat{S}_l^{\text{ord}})^{l\text{-an}}$  associated to  $\widehat{S}_l^{\text{ord}}$ . Therefore it has power series expansion  $f(\tau) = \sum_\nu a_\nu \tau^\nu$  for  $\tau^\nu = \prod_{\varphi \in \Sigma_l^{\text{ord}}} \tau_\varphi^\nu$ . If  $f \circ \widehat{\rho}(\alpha) = i_l(\alpha^{\sigma(1-c)})f$ , by equating the power series expansion, we find that  $f$  is a constant multiple of  $\tau_\sigma$ .  $\square$

If  $v = \infty$ , we put  $G_\infty = \text{Aut}_F(\mathfrak{H}^I)$  (the complex analytic automorphism group inducing  $O$ -linear map on  $\Omega_{\mathfrak{H}^I/\mathbb{C}}$ ); so,  $G_\infty = PSL_2(F_\mathbb{R}) = \prod_{\varphi \in I} PSL_2(\mathbb{R})$  according to the decomposition  $F_\mathbb{R} = F \otimes_{\mathbb{Q}} \mathbb{R} = \prod_{\varphi \in I} \mathbb{R}$  for which the projection to the  $\varphi$ -factor induces  $i_\infty \circ \varphi : F \rightarrow \mathbb{R}$  (e.g., [DLS] Chapter IV). As is well known, the stabilizer of  $x \in V(W_\Xi)$  in  $G_\infty$  is given by an anisotropic torus in  $G_\infty$ . Write the torus as  $\mathcal{T}_{\infty, x} = \mathcal{T}_x = \prod_{\varphi \in I} \mathcal{T}_{\varphi, x}$  for  $\mathcal{T}_{\varphi, x} \cong \mathbb{C}^\times / \mathbb{R}^\times$ . We put  $\Sigma_\infty^{\text{ord}} := I$ .

If  $v = p$  and  $\Sigma_p^{\text{ord}} \neq \emptyset$ , we write  $G_p = \text{Aut}_F(\widehat{S}^{\text{ord}})$  (the group of formal automorphisms over  $W_p$  compatible with  $O_{(\Xi)}^\times$ -action on  $\Omega_{\widehat{S}^{\text{ord}}/W_p}$ ). Note that  $\widehat{S}^{\text{ord}} \cong \widehat{\mathbb{G}}_m \otimes_{\mathbb{Z}_p} O_{\Sigma^{\text{ord}}}$  as seen in Section 1. We fix once and for all the identification  $\widehat{S}^{\text{ord}} = \widehat{\mathbb{G}}_m \otimes_{\mathbb{Z}_p} O_{\Sigma^{\text{ord}}}$ . This is tantamount to fixing an isomorphism  $\mu_{p^\infty} \otimes_{\mathbb{Z}_p} O_p \cong A_0[\mathfrak{p}^\infty]^\circ$  for each  $\mathfrak{p} \in \Sigma_p^{\text{ord}}$ ; so, as before, we write  $t_\mathfrak{p}$  for the Serre–Tate (multiplicative) coordinate of  $\widehat{S}_\mathfrak{p}^{\text{ord}}$ . We have  $\text{Aut}_{SCH}(\widehat{\mathbb{G}}_m) = \text{Aut}_{SCH}(\text{Spec}(W_p[t, t^{-1}])) = \mathbb{G}_m(W[t, t^{-1}]) = W[t, t^{-1}]^\times$  (the scheme automorphism group). Similarly, writing  $t = 1 + T$ , we have  $\text{Aut}(\widehat{\mathbb{G}}_m) = \text{Aut}(\text{Spf}(W_p[[T]])) = \widehat{\mathbb{G}}_m(W[[T]]) = 1 + \mathfrak{m}_W[[T]]$  for the maximal ideal  $\mathfrak{m}_W[[T]]$  of

$W[[T]]$ . Here  $\text{Aut}(\widehat{\mathbb{G}}_m)$  stands for the automorphism group of the formal scheme  $\widehat{\mathbb{G}}_m$ . Note that  $\widehat{\mathbb{G}}_m \otimes_{\mathbb{Z}_p} O_{\Sigma^{ord}} = \text{Spf}(W[[O_{\Sigma^{ord}}]]) = \text{Spf}(W[[T_\varphi]]_{\varphi \in \Sigma^{ord}})$ , where  $T_\varphi = t_\varphi - 1$  for the Serre–Tate  $\varphi$ -coordinate  $t_\varphi$ . If  $\phi \in G_p$ , then  $\phi(T) = (\phi_\varphi(T))_{\varphi \in \Sigma^{ord}}$ ; so, we have  $\phi^* dT_\varphi = \sum_\psi \frac{\partial \phi_\varphi}{\partial T_\psi} dT_\psi$ . Then having  $\phi^*(dT_\varphi) \in W_p dT_\varphi$  implies  $\frac{\partial \phi_\varphi}{\partial T_\psi} = 0$  if  $\varphi \neq \psi$ . Thus  $\phi_\varphi$  only involves the parameter  $T_\varphi$ ; so,  $G_p = \widehat{\mathbb{G}}_m(W[[T]])^{\Sigma^{ord}}$  by sending  $\phi$  to  $(\phi_\varphi(T_\varphi))_{\varphi \in \Sigma^{ord}}$ . Let  $\mathbf{1} \in \widehat{S}^{ord}$  be the  $W_p$ -point given by  $t_\varphi = 1$  for all  $\varphi \in \Sigma^{ord}$  (i.e., the origin). The point  $x \in V(\mathcal{W}_\Xi)$  gives rise to  $x \in \widehat{S}^{ord}(W_p)$  and  $x \in (\widehat{S}^{ord})^{p\text{-an}}(\mathbb{C}_p)$ . The group of  $O$ -linear automorphisms  $\mathcal{T}_\mathbf{1} = \mathcal{T}_{p,\mathbf{1}} := \text{Aut}_{O\text{-lin}}(\widehat{S}^{ord})$  is in the stabilizer of  $\mathbf{1}$ . The translation automorphism  $T_x : y \mapsto xy \in \widehat{S}^{ord}$  is an element in  $G_p$ . Thus we can translate the group structure via  $T_x$  so that  $x$  becomes the identity element. Then we have  $\mathcal{T}_{p,x} = T_x := T_x \mathcal{T}_\mathbf{1} T_x^{-1} \subset G_p$  fix  $x$ . We have again  $\mathcal{T}_{p,x} = \prod_{\mathfrak{p} \in \Sigma_p^{ord}} \mathcal{T}_{\mathfrak{p},x}$  with  $\mathcal{T}_{\mathfrak{p},x} \cong O_{\mathfrak{p}}^\times$  which preserve the  $\mathfrak{p}$ -component  $\widehat{S}_{\mathfrak{p}}^{ord} := \widehat{\mathbb{G}}_m \otimes_{\mathbb{Z}_p} O_{\mathfrak{p}} \subset \widehat{S}^{ord}$ .

**Lemma 4.2.** *Suppose that  $p \in \Xi$  is unramified in  $F$ . If  $g \in G(\mathbb{A}^{(\Xi),\infty})$  leaves stable  $\widehat{S}_{\mathfrak{p}}^{ord}$  inducing an automorphism of formal schemes and fixes the  $\mathfrak{p}$ -component  $x_{\mathfrak{p}} \in \widehat{S}_{\mathfrak{p}}^{ord}$  of  $x$ , then the action of  $g$  on  $\widehat{S}_{\mathfrak{p}}^{ord}$  belongs to  $\mathcal{T}_{\mathfrak{p},x}$ .*

*Proof.* The action of  $g$  on each test object  $(A_s, \eta_s^{(\Xi)})$  at  $s$  is given by

$$(A_s, \eta_s^{(\Xi)}) \mapsto (A_{sg}, \eta_{sg}^{(\Xi)}) \xleftarrow{\alpha} (A_s, \eta_s^{(\Xi)} \circ g)$$

with an  $O$ -linear prime to  $\Xi$ -isogeny  $\alpha$ . Thus it brings the  $O$ -linear group structure on the deformation space  $\widehat{S}_x^{ord}$  (of  $(A_x, \eta_x^{(\Xi)})$ ) with identity  $x$  to the  $O$ -linear group structure on  $\widehat{S}_{xg}^{ord}$  with identity  $xg$  just by the construction of the Serre–Tate coordinate; in other words, we have the commutative diagram of the extensions

$$\begin{array}{ccccc} A_x[\mathfrak{p}^\infty]^\circ & \xhookrightarrow{\quad} & A_x[\mathfrak{p}^\infty] & \twoheadrightarrow & A_x[\mathfrak{p}^\infty]^{et} \\ \wr \downarrow \alpha & & \wr \downarrow \alpha & & \wr \downarrow \alpha \\ A_{xg}[\mathfrak{p}^\infty]^\circ & \xhookrightarrow{\quad} & A_{xg}[\mathfrak{p}^\infty] & \twoheadrightarrow & A_{xg}[\mathfrak{p}^\infty]^{et}. \end{array}$$

The vertical arrows are  $O$ -linear isomorphisms as  $\alpha$  is prime to  $\Xi$ -isogeny. Thus if  $g$  leaves  $\widehat{S}_{\mathfrak{p}}^{ord}$  stable and fixes  $x_{\mathfrak{p}}$ , it must preserve the  $O$ -linear group structure of  $\widehat{S}_{\mathfrak{p}}^{ord}$  with identity  $x_{\mathfrak{p}}$ . This shows the action of  $g$  on  $\widehat{S}_{\mathfrak{p}}^{ord}$  belongs to  $\mathcal{T}_{\mathfrak{p},x}$ .  $\square$

Since the stabilizer of the  $\varphi$ -component  $x_\varphi$  of  $x$  in  $\mathfrak{H}^I$  is  $\mathcal{T}_{\varphi,x}$ , we have immediately

**Lemma 4.3.** *If  $g \in G(\mathbb{A}^{(\Xi)})$  leaves stable the  $\varphi$ -component  $\mathfrak{H}_\varphi = \mathfrak{H}$  of  $\mathfrak{H}^I$  and fixes the  $\varphi$ -component  $x_\varphi \in \mathfrak{H}_\varphi$ , then the action of  $g$  on  $\mathfrak{H}_\varphi$  belongs to  $\mathcal{T}_{\varphi,x}$ .*

**Lemma 4.4.** *Let  $x \in V(\mathcal{W}_\Xi)$ , and suppose that  $v \in \Xi \cup \{\infty\}$  is unramified in  $F$ . Then for each  $\varphi \in \Sigma_v^{ord}$ , there always exists a  $\varphi$ -coordinate  $\tau_\varphi = \tau_{\varphi,v}$  with  $\tau_\varphi(x) = 0$  convergent on an open rigid affinoid neighborhood  $D_x(\tau_\varphi)$  of  $x$  satisfying  $\tau_\varphi \circ a = a_v^\varphi \tau_\varphi$  for  $a = (a_v)_{v \in \Sigma_v^{ord}} \in \mathcal{T}_{v,x} = O_{\Sigma_v^{ord}}^\times$  if  $v < \infty$  and  $\tau_\varphi \circ a = a_\varphi \tau_\varphi$  for  $a \in \mathcal{T}_{\infty,x} = (\mathbb{C}^\times / \mathbb{R}^\times)^I$  if  $v = \infty$ , where  $v$  is the place of  $F$  induced by  $i_v \circ \varphi : F \hookrightarrow \mathbb{C}_p$ . The coordinate  $\tau_{\varphi,v}$  is unique up to scalar multiple.*

If  $x_{\mathfrak{p}} \neq \mathbf{1}_{\mathfrak{p}} \in \widehat{S}_{\mathfrak{p}}^{ord}$ , the automorphism  $\phi \in \mathcal{T}_{p,x} \cap \text{Aut}(\widehat{S}_{\mathfrak{p}}^{ord})$  may not extend to the global  $V$  as the translation  $T_{x_{\mathfrak{p}}} \neq 1$  is possibly a transcendental action only defined on the formal scheme  $\widehat{S}_{\mathfrak{p}}^{ord} = \widehat{\mathbb{G}}_m \otimes_{\mathbb{Z}_p} O_{\mathfrak{p}}$ . Therefore,  $\tau_{\sigma,p}$  in the lemma may not be a  $\widehat{\rho}$ -eigen coordinate for any choice of  $\widehat{\rho}$ .

*Proof.* The result for  $v = \infty$  is well known (see the argument towards the end of the proof of Theorem 4.5). If  $v < \infty$ , by Serre–Tate deformation theory, we have  $t_v$  centered at  $\mathbf{1}$  such that  $t_v \circ a = t_v^a$  for  $a \in \mathcal{T}_\mathbf{1}$ . Then  $\tau_{\varphi,v} := \log_p \circ \varphi_v \circ t_v \circ T_x^{-1}$  does the job, where  $\varphi_v : W_v \otimes_{\mathbb{Z}} O = W_v^I \rightarrow \mathbb{C}_v$  is the  $\varphi$ -projection. Uniqueness up to scalar multiple is clear from  $\tau_\varphi \circ a = a_v^\varphi \tau_\varphi$ .  $\square$

Since  $(A, \lambda, \theta, \bar{\eta}^{(\Xi)})$  gives the point  $x \in V(\mathcal{W}_\Xi)$ , we may think of its  $p$ -fiber

$$(A_0, \bar{\lambda}_0, \theta_0, \eta_0^{(\Xi)})_{/\mathbb{F}_p} = (A, \lambda, \theta, \bar{\eta}^{(\Xi)}) \times_{\mathcal{W}_\Xi} \bar{\mathbb{F}}_p.$$

Let  $M = \text{End}^0(A_0/\mathbb{F}_p)$ , which is a CM quadratic extension of  $F$  in which  $\mathfrak{p}$  splits by Lemma 2.2. As before, we define  $\hat{\rho}: T_M(\mathbb{Z}(\Xi)) \rightarrow G(\mathbb{A}^{(\Xi), \infty})$  by  $\tilde{\theta}_{\mathfrak{p}}(\alpha) \circ \eta_0^{(\Xi)} = \eta_0^{(\Xi)} \circ \hat{\rho}(\alpha)$ .

**Theorem 4.5.** *Let the notation be as above. Let  $i_p \circ \sigma$  (resp.  $i_l \circ \sigma$ ) induce  $\mathfrak{p} \in \Sigma_p^{ord}$  (resp.  $\mathfrak{l} \in \Sigma_l^{ord}$ ) for  $p, l \in \Xi$  for  $(A, \bar{\lambda}, \theta, \eta^{(\Xi)})$  sitting over  $x \in V(\mathcal{W}_\Xi)$ . Assume that  $p$  and  $l$  are unramified in  $F/\mathbb{Q}$ , and suppose that we have a  $p$ -analytic  $\hat{\rho}$ -eigen  $\sigma$ -coordinate  $\tau_p = \tau_{\sigma, p}$  of the form  $\tau_p = \log_p \circ \sigma_{\mathfrak{p}} \circ t_{\mathfrak{p}}$  with  $\tau_p(x) = 0$  around  $x \in V(\mathcal{W}_\Xi) \subset V(\mathbb{C}_p)$ , where  $\sigma_{\mathfrak{p}}: W_p \otimes_{\mathbb{Z}} O = W_p^I \rightarrow \mathbb{C}_p$  is the  $\sigma$ -projection. Then for  $v \in \{p, l, \infty\} =: \Xi_0$ , we have a  $v$ -analytic  $\hat{\rho}$ -eigen  $\sigma$ -coordinate  $\tau_v = \tau_{\sigma, v}$  with  $\tau_v(x) = 0$ . Identifying  $\mathbb{C}_p = \mathbb{C}_l = \mathbb{C}$  (as in the introduction), in the formal completion of  $\mathcal{O}_{V/\mathbb{C}_v, x}$  along the  $\mathbb{C}_v$ -point  $x \in V(\mathbb{C}_v)$ , the coordinates  $\{\tau_v\}_{v \in \Xi_0}$  are proportional (i.e., they are non-zero constant multiple each other).*

Write  $(\underline{\omega}^{2\sigma})^{v\text{-an}}$  for the  $v$ -analytic line bundle on  $(\hat{S}^{ord})^{v\text{-an}}$  coming from  $\underline{\omega}^{2\sigma}$  by a  $v$ -analytic version of GAGA theorems (e.g., [dJ] §7.1.11 and [C] Example 2.4.6). We write  $\mathfrak{v}$  for  $\mathfrak{l}$  and  $\mathfrak{p}$  according to  $v = l$  and  $p$ , and  $\Sigma_{\mathfrak{v}}^{ord}$  for the set of all embeddings  $\sigma: F \rightarrow \bar{\mathbb{Q}}$  such that  $i_{\mathfrak{v}} \circ \sigma$  induces  $\mathfrak{v}$ . To avoid confusion, if necessary, we write the  $v$ -adic deformation space  $\hat{S}$  of  $A \otimes_{W_v} \bar{\mathbb{F}}_v$  as  $\hat{S}_v$ .

*Proof.* Since  $\hat{S}_p = \hat{S}_p^{ord} \times \hat{S}_p^{ll}$  (see Proposition 1.2), the co-tangent space  $\Omega_{(\hat{S})^{p\text{-an}}/\mathbb{C}_p}(x)$  at  $x$  of  $(\hat{S})^{p\text{-an}}$  is a direct sum of  $\Omega_{(\hat{S}^{ord})^{p\text{-an}}/\mathbb{C}_p}(x^{ord})$  and  $\Omega_{(\hat{S}^{ll})^{p\text{-an}}/\mathbb{C}_p}(x^{ll})$  for the projection  $x^?$  of  $x$  to  $(\hat{S}^?)^{p\text{-an}}$ . Then we can further decompose:

$$\Omega_{(\hat{S}^{ord})^{p\text{-an}}/\mathbb{C}_p}(x^{ord}) = \bigoplus_{\mathfrak{p}_\varphi \in \Sigma^{ord}} \Omega_{(\hat{S}_{\mathfrak{p}_\varphi}^{ord})^{p\text{-an}}/\mathbb{C}_p}(x_{\mathfrak{p}_\varphi}) \quad \text{and} \quad \Omega_{(\hat{S}_{\mathfrak{p}_\varphi}^{ord})^{p\text{-an}}/\mathbb{C}_p}(x_{\mathfrak{p}_\varphi}) = \bigoplus_{\varphi \sim \mathfrak{p}_\varphi} (\underline{\omega}^{2\varphi})^{p\text{-an}}(x),$$

where  $x_{\mathfrak{p}_\varphi}$  is the projection of  $x$  to  $\hat{S}_{\mathfrak{p}_\varphi}^{ord}$  and in the second identity,  $\varphi$  runs over  $\varphi \in I$  such that  $i_{\mathfrak{p}} \circ \varphi$  induces the  $p$ -adic place  $\mathfrak{p}_\varphi$ .

Since  $\tau_p(x) = 0$ , the Serre–Tate  $\mathfrak{p}$ -coordinate  $t_{\mathfrak{p}}$  is equal to 1 at  $x$ ; so,

$$A[\mathfrak{p}^\infty] \cong (\mu_{p^\infty} \otimes_{\mathbb{Z}_p} O_{\mathfrak{p}}) \oplus (F_{\mathfrak{p}}/O_{\mathfrak{p}})$$

as Barsotti–Tate  $O$ -modules over  $W_p$ . Then the group  $T_M(\mathbb{Z}(\Xi))$  acts via  $\hat{\rho}$  on the  $\mathfrak{p}$ -component  $\hat{S}_{\mathfrak{p}}^{ord}$ , and  $T_M(\mathbb{Z}(\Xi))$  fixes the  $\mathfrak{p}$ -component  $x_{\mathfrak{p}}$  (this fact of  $T_M(\mathbb{Z}(\Xi))$  fixing  $x_{\mathfrak{p}}$  might not be true for  $\mathfrak{p}_\varphi \neq \mathfrak{p}$ ). Here  $(\underline{\omega}^{2\sigma})^{p\text{-an}}(x)$  is the  $\sigma$ -eigenspace as  $O$ -modules in  $\Omega_{(\hat{S}_{\mathfrak{p}_\varphi}^{ord})^{p\text{-an}}/\mathbb{C}_p}(x_{\mathfrak{p}_\varphi})$ . Since  $T_M(\mathbb{Z}(\Xi))$  fixes  $x_{\mathfrak{p}}$ ,  $(\underline{\omega}^{2\sigma})^{p\text{-an}}(x) \subset \Omega_{(\hat{S}_{\mathfrak{p}_\varphi}^{ord})^{p\text{-an}}/\mathbb{C}_p}(x_{\mathfrak{p}})$  for any  $\sigma$  associated with  $\mathfrak{p}$  is the  $i_{\mathfrak{p}}(\alpha^{\sigma(1-c)})$ -eigenspace of  $\hat{\rho}(\alpha)$  ( $\alpha \in T_M(\mathbb{Z}(\Xi))$ ) in  $\Omega_{(\hat{S}_{\mathfrak{p}_\varphi}^{ord})^{p\text{-an}}/\mathbb{C}_p}(x_{\mathfrak{p}})$  for an extension  $\sigma: M \hookrightarrow \bar{\mathbb{Q}}$  of  $\sigma: F \hookrightarrow \bar{\mathbb{Q}}$ . Taking the canonical lift  $A_1/W_\Xi$  of  $A_0/\mathbb{F}_p$  sitting over  $x_1 \in V(\mathcal{W}_\Xi)$ , the extension  $\sigma$  is determined by complex multiplication by  $M$  on  $\underline{\omega}^\sigma(x_1)/W_\Xi$  which is the  $\sigma$ -eigenspace (under real multiplication by  $F$ ) of  $H^0(A_1, \Omega_{A_1/W_\Xi})$  canonically isomorphic to  $\underline{\omega}^\sigma(x)$  after extending scalars to  $W_p$ . Thus for  $\sigma$  with  $\mathfrak{p} = \mathfrak{p}_\sigma$ ,  $T_M(\mathbb{Z}(\Xi))$  acts on  $(\underline{\omega}^{2\sigma})^{p\text{-an}}(x)$  via the algebraic character  $i_{\mathfrak{p}} \circ \sigma(1-c) \in X^*(T_M)$ . Since  $(\underline{\omega}^{2\sigma})^{p\text{-an}}(x) = \underline{\omega}^{2\sigma}(x) \otimes_{W_\Xi} \mathbb{C}_p$  for the algebraic co-tangent space  $\underline{\omega}^{2\sigma}(x)/W_\Xi$  and the action of  $\hat{\rho}$  on the rigid analytic line bundle  $(\underline{\omega}^{2\sigma})^{p\text{-an}}$  is induced by its action on the algebraic line bundle  $\underline{\omega}^{2\sigma}$  over  $\mathcal{W}_\Xi$ , the action via  $\hat{\rho}$  must preserve the algebraic  $\underline{\omega}^{2\sigma}(x)$ . Therefore it preserves also

$$(\underline{\omega}^{2\sigma})^{v\text{-an}}(x) = \underline{\omega}^{2\sigma}(x) \otimes_{W_\Xi} \mathbb{C}_v$$

for every  $v \in \{p, l, \infty\}$ . Since the action of  $\hat{\rho}(T_M(\mathbb{Z}(\Xi)))$  on the  $v$ -analytic  $\sigma$ -co-tangent bundle  $(\underline{\omega}^{2\sigma})^{v\text{-an}}$  over  $V(\mathbb{C}_v)$  preserves its fiber over  $x$ ,  $\hat{\rho}(T_M(\mathbb{Z}(\Xi)))$  fixes  $x_{\mathfrak{v}}$  reversing our argument.

We see this fact of  $\hat{\rho}(T_M(\mathbb{Z}(\Xi)))$  fixing  $x_{\mathfrak{v}}$  group-theoretically now. Because of the  $v$ -adic Lie group structure on  $\hat{S}_{\mathfrak{v}}^{ord}$  for  $v = p, l$ , the tangent space of  $\hat{S}_{\mathfrak{v}}^{ord}$  at the origin

$$\text{Tan}_{\mathfrak{v}} := \bigoplus_{\varphi \sim \mathfrak{v}} \text{Tan}_{\varphi, \mathfrak{v}} \quad \text{for} \quad \text{Tan}_{\varphi, \mathfrak{v}} = \text{Hom}_{W_v}(\underline{\omega}^{2\varphi}(x) \otimes_{W_\Xi} W_v, W_v)$$

is a Lie algebra over  $W_v$ , and we have the  $v$ -adic Lie algebra  $L_v := \text{Lie}(\widehat{S}_v^{\text{ord}})$  over  $\mathbb{Z}_v$  canonically inside the tangent space  $\text{Tan}_v$ ; i.e., we have

$$L_v \otimes_{\mathbb{Z}_v} W_v = \text{Tan}_v.$$

Since  $\widehat{S}_v^{\text{ord}} = \widehat{G}_m \otimes_{\mathbb{Z}_v} O_v$ , the Lie algebra  $L_v$  is isomorphic to the formal co-character group

$$X_*(\widehat{S}_v^{\text{ord}}) := \text{Hom}_{\text{formal group}}(\widehat{S}_v^{\text{ord}}, \widehat{G}_m) \cong O_v$$

and hence the action via  $\widehat{\rho}$  preserves  $L_v$ , and the action via  $\widehat{\rho}$  on  $\text{Tan}_v$  is the scalar extension of the action on  $L_v$  to  $\text{Tan}_v$ . We have the  $v$ -adic exponential map  $\exp_v : L_v \rightarrow \widehat{S}_v^{\text{ord}}$  (from the  $v$ -adic formal Lie algebra into the  $v$ -adic formal Lie group) which is equivariant under the action of  $\widehat{\rho}$ . Thus the  $\mathfrak{v}$ -component  $\widehat{S}_v^{\text{ord}}$  is stable under the action of  $T_M$  via  $\widehat{\rho}$ , and the fixed point in  $\widehat{S}_v^{\text{ord}}$  of the action is the  $\mathfrak{v}$ -component  $x_v$  of  $x$ . The action of  $\alpha \in T_M(\mathbb{Z}(\Xi))$  on  $\text{Tan}_{\sigma, \mathfrak{v}}$  is via multiplication by  $i_v(\alpha^{\sigma(1-c)})$  for a suitable extension of  $\sigma$  to  $M$  (still denoted by  $\sigma$ ). Thus its action on  $L_v$  is via multiplication by  $\alpha^{1-c} = \varprojlim_n (\alpha^{1-c} \bmod \mathfrak{v}^n) \in O_v$ ; so, we can find a linear coordinate  $T_{\sigma, \mathfrak{v}}$  of  $\text{Tan}_{\sigma, \mathfrak{v}}$  with  $T_{\sigma, \mathfrak{v}} \circ \widehat{\rho}(\alpha) = \alpha^{\sigma(1-c)} T_{\sigma, \mathfrak{v}}$ . Its image under exponential map gives rise to the coordinate  $t_\sigma$  on  $\widehat{S}_v^{\text{ord}}$  (with  $t_\sigma \circ \widehat{\rho}(\alpha) = t_\sigma^{\alpha^{\sigma(1-c)}}$  and  $t_\sigma(x) = 1$ ), whose  $v$ -adic logarithm is proportional to the  $v$ -adic logarithm of the Serre–Tate  $\sigma$ -coordinate. Then  $\tau_v = \tau_{\sigma, \mathfrak{v}} := \log_v \circ t_\sigma$  is the desired  $\widehat{\rho}$ -eigen  $\sigma$ -coordinate, and  $x_v$  is the fixed point of  $\widehat{\rho}$  on  $\widehat{S}_v^{\text{ord}}$ , which induces a homomorphism  $\rho_v : T_M(\mathbb{Z}(\Xi)) \rightarrow \text{Aut}_{O\text{-lin}}(\widehat{S}_v^{\text{ord}}) = \mathcal{T}_{v, x}$ . To see  $\tau_v(x) = 0$  in more down-to-earth terms, note that  $\mathcal{T}_{v, x} = O_v^\times$  and  $\rho_v(\alpha) = \alpha^{\sigma(1-c)} \in O_v^\times$ . Since the image of  $\rho_v$  is  $\mathfrak{v}$ -adically dense, by the continuity of the action, we have  $\tau_{\sigma, \mathfrak{v}} \circ a = a \tau_{\sigma, \mathfrak{v}}$  for all  $a \in O_v^\times = \mathcal{T}_{v, x}$ . Then by Lemma 4.4,  $\tau_{\sigma, \mathfrak{p}}$  and  $\tau_{\sigma, \mathfrak{l}}$  are proportional each other as formal functions on the formal completion of  $V_{K/\mathbb{C}_v}$  along  $x \in V_K(\mathbb{C}_v)$  identifying  $\mathbb{C}_v = \mathbb{C}_l = \mathbb{C}_p$ . In particular  $\tau_{\sigma, \mathfrak{v}}(x) = 0$ , and this re-confirms that the coordinate  $\tau_v := \tau_{\sigma, \mathfrak{v}}$  is the desired one (up to scalar multiple).

The proof for  $v = \infty$  is almost identical to the above proof for finite places. Here it is. We can bring  $\mathfrak{H}^I$  isomorphically onto  $\mathfrak{D}^I$  for the unit open disk  $\mathfrak{D}$  in  $\mathbb{C}$  by a linear fractional transformation by a matrix in  $SL_2(\mathbb{C})^I$  so that  $x$  is sent to the origin 0, and the action of the torus  $\mathcal{T}_{\infty, x/\mathbb{R}}$  in  $PSL_2(\mathbb{R})^I_{/\mathbb{R}}$  fixing  $x$  (isomorphic to  $(\mathbb{C}^\times/\mathbb{R}^\times)^I$ ) on the coordinate  $w_\sigma$  of  $\mathfrak{D}$  at  $\sigma \in I$  is given by  $w_\sigma \circ \alpha = \alpha^{1-c} w_\sigma$  for  $\alpha = (\alpha_\varphi)_{\varphi \in I} \in (\mathbb{C}^\times)^I$  (under the identification  $\mathcal{T}_x(\mathbb{C}) = (\mathbb{C}^\times/\mathbb{R}^\times)^I$ ). Here  $\mathfrak{D}$  is the symmetric domain of  $SU(1, 1)$  (the signature  $(1, 1)$  special unitary group) isomorphic to  $SL_2(\mathbb{R})$ , and  $T_1(\mathbb{C}) \subset SL_2(\mathbb{R})^I$  is identified with the diagonal torus in  $SU(1, 1)^I$ . To explain this in down-to-earth terms, write  $z$  for the standard coordinate of  $\mathfrak{H}^I \subset \mathbb{C}^I$ . If  $\xi = z_\varphi(x)$ , the holomorphic isomorphism:  $\mathfrak{H} \ni z = (z_\varphi)_\varphi \mapsto w = (w_\varphi)_\varphi = \left( \frac{z_\varphi - \xi}{z_\varphi + \xi} \right)_\varphi \in \mathfrak{D}$  does the job. Though  $\mathfrak{D}$  is not a Lie group, it is an open neighborhood of the identity of the Lie group  $\mathbb{C}^\times = \mathbb{G}_m(\mathbb{C})$ , and it is a Lie semi-group. Then the argument using the tangent space at  $x$  as above (identifying it with the Lie algebra of  $\mathcal{T}_{\infty, x}$ ) done for finite places  $v = p, l$  is valid also for  $v = \infty$  without any change. Thus we can take  $w_\sigma$  to be  $\tau_{\sigma, \infty} = \tau_\infty$ . Again it is proportional to  $\tau_v$  over the formal completion.  $\square$

**Corollary 4.6.** *Let the notation and the assumption be as in Theorem 4.5. Let  $l \in \Sigma_l^{\text{ord}}$  for a prime  $l \in \Xi$  different from  $p$ . If the prime  $l$  in  $F$  is induced by  $i_l \circ \sigma$ , we have  $\text{End}_F^0(A \otimes_{\mathcal{W}_\Xi} \overline{\mathbb{F}}_l) \cong M \cong \text{End}_F^0(A \otimes_{\mathcal{W}_\Xi} \overline{\mathbb{F}}_p)$  as  $F$ -algebras (via  $\theta$ ).*

*Proof.* Let  $A_\Xi = A \otimes_{\mathcal{W}_\Xi} \overline{\mathbb{F}}_l$ , and write  $M' = \text{End}_F^0(A_\Xi)$  with  $R' = \text{End}_O(A_\Xi)$  and  $M = \text{End}_F^0(A_0)$  with  $R = \text{End}_O(A_0)$ . Fix a level  $l$ -structure  $\eta_l : \mu_{l^\infty} \otimes_{\mathbb{Z}_l} O_l \cong A[l^\infty]^\circ$ . By Theorem 4.5, we have  $\tau_l(x) = 0$ ; so,  $t_l(x) = 1$ , which implies that we get an  $O$ -linear isomorphism of Barsotti–Tate  $O$ -modules over  $W_l$ :

$$\eta_l \oplus {}^t \eta_l^{-1} : (\mu_{l^\infty} \otimes_{\mathbb{Z}_l} O_l) \oplus (F_l/O_l) \cong A[l^\infty].$$

Again by Theorem 4.5, we have  $\tau_l \circ \widehat{\rho}(\alpha) = i_l(\alpha^{\sigma(1-c)}) \tau_l$  for  $\alpha \in R_{(\Xi)}^\times = T_M(\mathbb{Z}(\Xi))$ , and  $\widehat{\rho}(R_{(\Xi)}^\times)$  leaves stable  $\widehat{S}_l^{\text{ord}}$  and fixes  $x_l \in \widehat{S}_l^{\text{ord}}$ . Thus the image of  $\widehat{\rho} : T_M \hookrightarrow G(\mathbb{A}^{(\Xi^\infty)})$  is the unique torus (over  $\mathbb{A}^{(\Xi^\infty)}$ ) in  $G$  such that its image in  $\overline{\mathcal{E}}^{(\Xi)}$  is made of automorphisms of  $Sh^{(\Xi)}$  preserving  $\widehat{S}_l^{\text{ord}}$  and fixing  $x_l$  by Lemma 4.2 (in short,  $\widehat{\rho}(T_M(\mathbb{Z}(\Xi))) \subset \mathcal{T}_{l, x} \cap \overline{\mathcal{E}}^{(\Xi)}$  under a slight abuse of notation). This

implies by universality of  $Sh^{(\Xi)}$  that we have an endo-isogeny  $\tilde{\theta}_{\mathfrak{L}}(\alpha) \in \text{End}_F^0(A_{\mathfrak{L}})$  for each  $\alpha \in R_{(\Xi)}^{\times}$  such that  $\tilde{\theta}_{\mathfrak{L}}(\alpha) \circ \eta_{\mathfrak{L}}^{(\Xi)} = \eta_{\mathfrak{L}}^{(\Xi)} \circ \hat{\rho}(\alpha)$  and the isogeny action of  $\tilde{\theta}_{\mathfrak{L}}(\alpha)$  is given by  $t_{\sigma, \iota} \mapsto t_{\sigma, \iota}^{\alpha^{\sigma(1-c)}}$  for  $\iota$ -adic Serre–Tate  $\sigma$ -coordinate  $t_{\sigma, \iota}$ . In other words, the triple  $(E = M, \alpha_q = \tilde{\theta}_{\mathfrak{L}}(\alpha)|_{A_{\mathfrak{L}}[q^\infty]}, \hat{\rho})$  satisfies the assumption of Lemma 4.1 (3), and hence, by the  $O$ -linear embedding  $\tilde{\theta}_{\mathfrak{L}} : M \hookrightarrow \text{End}_F^0(A_{\mathfrak{L}}/\mathbb{F}_{\mathfrak{L}})$ , we have  $M \hookrightarrow M'$ , which implies  $\tilde{\theta}_{\mathfrak{L}} : M \cong M'$ , since  $M' = \text{End}_F^0(A_{\mathfrak{L}})$  is a quadratic extension of  $F$ .  $\square$

## 5. LOCAL INDECOMPOSABILITY OF AN AVR M

Let  $(A, \lambda, \theta)_{/\mathcal{W}_p}$  be an AVR M by multiplication by  $O$  with good reduction. Thus the triple  $(A, \lambda, \theta)$  is defined over a number field  $k$  in which  $p$  is unramified. We suppose that  $\Sigma^{ord} := \Sigma_p^{ord}$  is non-empty (so,  $A_0 = A \otimes_{\mathcal{W}} \overline{\mathbb{F}}_p$  is at least partially ordinary). By Serre–Tate theory, we have  $\widehat{S}^{ord} \cong \widehat{\mathbb{G}}_m \otimes_{\mathbb{Z}_p} O_{\Sigma^{ord}}$ . We fix an isomorphism  $\eta_{\mathfrak{p}} : \mu_{p^\infty} \otimes_{\mathbb{Z}_p} O_{\mathfrak{p}} \cong A[\mathfrak{p}^\infty]$  to fix the identification  $\widehat{S}_{\mathfrak{p}}^{ord} \cong \widehat{\mathbb{G}}_m \otimes_{\mathbb{Z}_p} O_{\mathfrak{p}}$  for  $\mathfrak{p} \in \Sigma_p^{ord}$ . For a finite set of primes  $\Xi$  including  $p$  (which will be specified later), we will choose suitably an isomorphism  $\eta^{(\Xi)} : (F_{\mathbb{A}}^{(\Xi), \infty})^2 \cong T^{(\Xi)}A \otimes_{\mathbb{Z}} \mathbb{Z}^{(\Xi)}$  defined over  $\mathcal{W}_{\Xi}$  in the proof of the following theorem. Since  $p$  is unramified in  $k$  and  $p \in \Xi$ , by extending scalars to a finite extension of  $k$  in the fraction field of  $\mathcal{W}_{\Xi}$  if necessary, for a neat open compact subgroup  $K \subset G(\mathbb{A}^{(\Xi), \infty})$ , we may assume that  $(A, \bar{\lambda}, \eta^{(\Xi)})$  (resp.  $(A, \bar{\lambda}, \bar{\eta}^{(\Xi)})$ ) gives rise to a point  $x \in V(\mathcal{W}_{\Xi})$  (resp.  $x \in V_K(\mathcal{W}_k)$ ) for a geometrically irreducible component  $V$  of  $Sh^{(\Xi)}$ . Let  $M = \text{End}_F^0(A_0)$  which is a CM field (see Lemma 2.2). Define  $\hat{\rho} : T_M(\mathbb{Z}^{(\Xi)}) \rightarrow G(\mathbb{A}^{(\Xi), \infty})$  by  $\tilde{\theta}_{\mathfrak{P}}(\alpha) \circ \eta_0^{(\Xi)} = \eta_0^{(\Xi)} \circ \hat{\rho}(\alpha)$ , where  $(A_0, \lambda_0, \theta_0, \eta_0^{(\Xi)}) = (A, \lambda, \theta, \eta^{(\Xi)}) \times_{\mathcal{W}_p} \overline{\mathbb{F}}_p$  and  $\tilde{\theta}_{\mathfrak{P}}$  is an extension of  $\theta_0$  to  $M$ . Then  $\hat{\rho}(\alpha)$  acts on  $\widehat{S}^{ord}$  by  $t_{\mathfrak{p}} \mapsto t_{\mathfrak{p}}^{\alpha_p^{1-c}}$  for an embedding  $M \ni \alpha \mapsto \alpha_{\mathfrak{p}} \in F_{\mathfrak{p}}$  and on  $V$  through  $\bar{\mathcal{E}}^{(\Xi)}$ , where  $c$  is the generator of  $\text{Gal}(M/F)$ . The embedding  $\alpha \mapsto \alpha_{\mathfrak{p}}$  induces  $M_{\mathcal{P}} = F_{\mathfrak{p}}$  for a prime  $\mathcal{P}$  of  $M$  over  $\mathfrak{p}$ , and  $A_0[\mathcal{P}]$  is connected. We write  $\sigma_{\mathfrak{p}} : F_{\mathfrak{p}} \rightarrow \mathbb{C}_p$  for the embedding induced by the  $\sigma$ -projection of  $W_p \otimes_{\mathbb{Z}} O = W_p^I$  to  $W_p \subset \mathbb{C}_p$ . If the place  $\mathfrak{p}$  is associated to  $i_p \circ \sigma$ , we write  $t_{\sigma} = \sigma_{\mathfrak{p}} \circ t_{\mathfrak{p}}$  and  $\tau = \tau_{\sigma, p} = \log_p \circ t_{\sigma}$ ; so,  $\tau \circ \hat{\rho}(\alpha) = i_p(\alpha^{\sigma(1-c)})\tau$ . By definition, if  $i_p \circ \sigma$  gives rise to  $p$ -adic place  $\mathfrak{p} \in \Sigma_p^{ord}$ ,  $t_{\sigma}(A) = 1$  ( $\Leftrightarrow \tau(A) = 0$ ) is equivalent to semi-simplicity of the  $\mathfrak{p}$ -adic Tate module  $T_{\mathfrak{p}}A$  as an  $I_p$ -module for the inertia group  $I_p$  in  $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$  corresponding to  $i_p \circ \sigma$ . We want to prove

**Theorem 5.1.** *Let the notation be as above. Suppose that  $O$  is unramified at  $p$  and that the place of  $k$  induced by  $i_p$  is unramified over  $\mathbb{Q}$ . If  $A/\overline{\mathbb{Q}} = A \times_{\mathcal{W}} \overline{\mathbb{Q}}$  does not have complex multiplication,  $T_{\mathfrak{p}}A$  is indecomposable as an  $I_p$ -module for each  $\mathfrak{p} \in \Sigma_p^{ord}$ .*

*Proof.* On the contrary to the conclusion of the theorem, we assume that  $T_{\mathfrak{p}}A$  for  $\mathfrak{p} \in \Sigma_p^{ord}$  is a semi-simple  $I_p$ -module (so,  $t_{\mathfrak{p}}(A) = 1$ ). As before,  $i_p \circ \sigma$  induces the place  $\mathfrak{p}$  for  $\sigma \in I$ . Note that  $A$  is defined over a number field  $k \subset \overline{\mathbb{Q}}$  unramified at  $p$  as  $A$  is defined over unramified  $\mathcal{W}$  and  $A$  is projective. If  $p$  is inert in  $F$ , then  $t_{\sigma}(A) = 1$  implies  $t_p(A) = 1$ ; so,  $A$  is the canonical lift of  $A_0$ ; so,  $A$  has complex multiplication, a contradiction. Thus we may assume that  $F$  has at least two primes over  $p$ , though we do not use this fact in the rest of the proof.

We write primes of  $F$  by lower case Gothic letters and primes of  $k$  by upper case Gothic letters. The corresponding roman character is the rational prime below. Pick a prime  $\mathfrak{q}$  of  $O$ . Identify  $T_{\mathfrak{q}}A \cong O_{\mathfrak{q}}^2$  and write  $r_{\mathfrak{q}} : \text{Gal}(\overline{\mathbb{Q}}/k) \rightarrow GL_2(O_{\mathfrak{q}})$  for the Galois representation realized on  $T_{\mathfrak{q}}A$ .

Since  $A$  is isotypic as is well known (i.e.,  $A$  is isogenous to a product of copies of an absolutely simple AVR M over a finite extension of  $k$ ; see [GME] §5.3.1), we may assume that  $A$  is absolutely simple. Then  $D := \text{End}^0(A/k)$  is a (possibly commutative) division central simple algebra over a totally real field or a CM field  $Z$  with integer ring  $\mathfrak{r}$  (as  $A \otimes_k \overline{\mathbb{Q}}$  does not have complex multiplication; see [GME] Lemma 5.3.2). Since  $F$  is totally real,  $Z \subset F$  has to be totally real. If  $D$  is commutative,  $D = F = Z$  is a totally real field, and if  $D$  is not commutative, it is a division quaternion algebra over  $Z$ . By a theorem of Faltings (solution of Tate’s conjecture; see [ARG] II), for  $\text{End}_{Z_{\mathfrak{q}}}^0(T_{\mathfrak{q}}A) := \text{End}_{\mathfrak{r}_{\mathfrak{q}}}(T_{\mathfrak{q}}A) \otimes_{\mathfrak{r}_{\mathfrak{q}}} Z_{\mathfrak{q}}$ , the algebra  $Z_{\mathfrak{q}}[r_{\mathfrak{q}}(\text{Gal}(\overline{\mathbb{Q}}/k'))] \subset \text{End}_{Z_{\mathfrak{q}}}^0(T_{\mathfrak{q}}A)$  generated by  $r_{\mathfrak{q}}(\text{Gal}(\overline{\mathbb{Q}}/k'))$  over  $Z_{\mathfrak{q}}$  is the commutant of  $\text{End}(A/k) \otimes_{\mathbb{Z}} Z_{\mathfrak{q}}$  for any finite extension  $k'/k$ , where  $T_{\mathfrak{q}}A = \varprojlim_n A[\mathfrak{q}]^n$  and  $Z_{\mathfrak{q}}$  is the  $\mathfrak{q}$ -adic completion of  $Z \subset F$  (i.e., the closure of  $Z$  in  $F_{\mathfrak{q}}$ ). Let  $C_{\mathfrak{q}} := Z_{\mathfrak{q}}[r_{\mathfrak{q}}(\text{Gal}(\overline{\mathbb{Q}}/k'))] \subset$

$\text{End}_{Z_q}^0(T_q A)$ . Then  $C_q$  is the commutant of  $D_q = D \otimes_Z Z_q$  in  $\text{End}_{Z_q}^0(T_q A)$ . Hence  $C_q$  has center  $Z_q$  and is independent of the choice of  $k'$ . In particular,  $C_q \otimes_{Z_q} D_q \cong \text{End}_{Z_q}^0(T_q A) \cong M_n(Z_q)$  ( $n \times n$  matrix algebra) for a suitable  $n$  (cf. [BAL] VIII.10.2). This implies that the Brauer classes over  $Z_q$  of  $C_q$  and  $D_q$  are inverse each other (cf. [BAL] VIII.10.4). Since  $D$  is either a quaternion algebra over  $Z$  or  $D = F = Z$ , the Brauer class of  $D_q$  has order at most 2 in the Brauer group over  $Z_q$ . Thus  $C_q$  is either isomorphic to a division quaternion algebra over  $Z_q$  or  $M_2(F_q) = M_2(Z_q)$ .

Set  $r = r_{\mathfrak{p}}$  (so  $\mathfrak{q} = \mathfrak{p}$ ). Since  $M \ni \alpha \mapsto \alpha^{\sigma(1-c)} \in F_{\mathfrak{p}}$  implies  $M_{\mathfrak{p}} = F_{\mathfrak{p}} \oplus F_{\mathfrak{p}} \hookrightarrow C_{\mathfrak{p}}$  (i.e.,  $\mathfrak{p}$  splits in  $M$ ); so,  $C_{\mathfrak{p}} \cong M_2(F_{\mathfrak{p}})$ . Thus  $\text{Im}(r)$  contains an open subgroup of  $SL_2(Z_q) \subset C_q^{\times}$ . This open image property follows from Ribet's argument in [Ri] IV. Indeed, assuming that  $\text{End}^0(A \otimes_k \overline{\mathbb{Q}}) = F$  (absolutely simple AVRМ but non CM) and that  $A$  has a place of multiplicative reduction, Ribet computed the Lie algebra of  $\text{Im}(r)$  and showed the Lie algebra contains  $\mathfrak{sl}_2(\mathbb{Z}_p)$  (which implies the open image property by exponentiating the Lie algebra). His assumptions are used to show that  $C_{\mathfrak{p}} = M_2(F_{\mathfrak{p}})$ , and, using Faltings' result, Ribet's argument still works, and we get the open image property (see [BGK] Theorem C for more details of this point).

Since there are at least two non-isomorphic ramified semi-simple quadratic extensions of  $\mathbb{Q}_p$ , we have at least four non-isomorphic maximal tori in  $SL(2)_{/\mathbb{Q}_p}$  defined over  $\mathbb{Q}_p$  having non-isomorphic projection to  $PGL(2)$ . Thus we can find a torus  $T \subset GL(2)$  defined over  $O_{\mathfrak{p}}$  such that

- (p1)  $T(O_{\mathfrak{p}}) \cap SL_2(\mathbb{Z}_p)$  is an anisotropic maximal torus (i.e.,  $T(O_{\mathfrak{p}})$  is isomorphic to the norm 1 subgroup of the multiplicative group of a field extension of  $F_{\mathfrak{p}}$ ),
- (p2)  $T(O_{\mathfrak{p}})$  remains an anisotropic maximal torus (i.e., for example, if  $T(O_{\mathfrak{p}}) \cap SL_2(\mathbb{Z}_p)$  comes from a ramified quadratic extension of  $\mathbb{Q}_p$ ,  $T(O_{\mathfrak{p}})$  remains anisotropic as  $F_{\mathfrak{p}}/\mathbb{Q}_p$  is unramified).

We pick  $\alpha \in T(O_{\mathfrak{p}}) \cap \text{Im}(r) \cap SL_2(\mathbb{Z}_p)$  so that  $T$  is the centralizer of  $\alpha$  (i.e.,  $\alpha$  has two distinct eigenvalues in  $C_{\mathfrak{p}}$ ). Since the isomorphism class of the centralizer of  $\alpha$  (as tori over  $O_{\mathfrak{p}}$ ) is determined by  $\alpha \bmod p^j$  for sufficiently large  $j$ , by Chebotarev density, changing  $T$  in its isomorphism/conjugacy class if necessary, we may choose a prime  $\mathfrak{L}$  outside  $p$  so that

- $A$  has good reduction at  $\mathfrak{L}$ ,
- $r(\text{Frob}_{\mathfrak{L}})$  commutes with  $\alpha$  (i.e.  $r(\text{Frob}_{\mathfrak{L}}) \in T(F_{\mathfrak{p}})$ ).

For the eigenvalues  $\lambda$  of  $r(\text{Frob}_{\mathfrak{L}})$ , the quadratic extensions  $M' = F[\lambda]$  is non-isomorphic to  $M = \text{End}_F^0(A_{0/\overline{\mathbb{F}}_p})$  over  $F$  (i.e.,  $\mathfrak{p}$  is non split in  $M'$  but  $\mathfrak{p}$  splits in  $M$ ). Since  $A$  has  $\Sigma_l^{\text{ord}} \neq \emptyset$  for primes  $\mathfrak{L}$  of  $k$  of Dirichlet density 1 (after replacing  $k$  by its finite extension if necessary; see Section 7 in the text and [O] 2.7 and [N] Introduction), we may assume that  $\Sigma_l^{\text{ord}}$  contains a prime induced by  $i_l \circ \sigma$  (i.e., we may assume that  $\mathfrak{l}$  splits in  $M'$ ); so, we have  $\text{End}_F^0(A \otimes_{\mathcal{W}_l} \overline{\mathbb{F}}_l) = M'$ .

Let  $\Xi = \{p, l\}$  for the residual characteristic  $l$  of  $\mathfrak{l}$ . Then  $A_0$  has complex multiplication by a CM field  $M/F$ . Thus  $A_0[q^{\infty}] \cong (F_q/O_q)^2$  over  $\overline{\mathbb{F}}_p$  for any prime  $q \notin \Xi$ . Since  $A_0$  is defined over  $\mathbb{F}_{\mathfrak{p}}$ , it has the Frobenius endomorphism  $\phi$  over  $\mathbb{F}_{\mathfrak{p}}$  and  $M = F[\phi] = \text{End}_F^0(A_{0/\mathbb{F}_{\mathfrak{p}}})$  is a CM quadratic extension of  $F$  (by Lemma 2.2). Thus we can find an  $R$ -linear isomorphism  $\eta_0^{(\Xi)} : \widehat{R}^{(\Xi)} \cong T^{(\Xi)} A_0$  for  $R = O[\phi]$ , where  $\widehat{R} = \prod_q R_q$  and  $\widehat{R}^{(\Xi)} = \prod_{q \notin \Xi} R_q$ . Thus, identifying  $\widehat{R}^{(\Xi)} \otimes_{\mathbb{Z}} \mathbb{Q} = (F_{\mathbb{A}}^{(\Xi), \infty})^2$ , we may define an embedding  $\widehat{\rho} : M^{\times} \hookrightarrow G(\mathbb{A}^{(\Xi), \infty})$  by  $\widehat{\theta}_{\mathfrak{p}}(\alpha) \circ \eta_0^{(\Xi)} = \eta_0^{(\Xi)} \circ \widehat{\rho}(\alpha)$ . Since  $A[q^{\infty}]$  for  $q \notin \Xi$  is étale over  $\mathcal{W}_{\Xi}$ , this level structure  $\eta_0^{(\Xi)}$  lifts uniquely to a level structure  $\eta^{(\Xi)}$  of  $A$ . Then the point  $x_0 \in \text{Sh}^{(\Xi)}(\overline{\mathbb{F}}_p)$  carrying  $(A_0, \lambda_0, \eta_0^{(\Xi)})$  is fixed by  $\widehat{\rho}(T_M(\mathbb{Z}_{(\Xi)}))$  by Lemma 4.1.

Suppose that  $r_{\mathfrak{p}}|_{I_p}$  is semi-simple for  $\mathfrak{p} \in \Sigma^{\text{ord}}$  for  $\mathfrak{p}$  induced by  $\sigma : F \rightarrow \overline{\mathbb{Q}}_p$ . Thus by semi-simplicity, the following connected-étale exact sequence

$$0 \rightarrow A[\mathfrak{p}^{\infty}]^{\circ}(\overline{\mathbb{Q}}_p) \rightarrow A[\mathfrak{p}^{\infty}](\overline{\mathbb{Q}}_p) \rightarrow A[\mathfrak{p}^{\infty}]^{\text{et}}(\overline{\mathbb{Q}}_p) \rightarrow 0$$

of  $I_p$ -modules splits, and hence the connected -étale sequence of Barsotti-Tate  $O$ -modules

$$0 \rightarrow A[\mathfrak{p}^{\infty}]_{/W_p}^{\circ} \rightarrow A[\mathfrak{p}^{\infty}]_{/W_p} \rightarrow A[\mathfrak{p}^{\infty}]_{/W_p}^{\text{et}} \rightarrow 0$$

also splits (cf. [T]). In particular, we have  $\mathfrak{p} = \mathcal{P}\overline{\mathcal{P}}$  for prime ideals  $\mathcal{P} \neq \overline{\mathcal{P}}$  in  $R = \text{End}_O(A_{0/\overline{\mathbb{F}}_p}) \subset M$  so that  $A_0[\mathfrak{p}^{\infty}]^{\circ} = A_0[\mathcal{P}^{\infty}]$  and  $A_0[\mathfrak{p}^{\infty}]^{\text{et}} = A_0[\overline{\mathcal{P}}^{\infty}]$ . The isomorphism  $\eta_{\mathfrak{p}} : \mu_{p^{\infty}} \otimes_{\mathbb{Z}_p} O_{\mathfrak{p}} \cong A[\mathfrak{p}^{\infty}]^{\circ}$  is defined over  $\mathcal{W}_p$ . Recall  $M^{\times} \ni \alpha \mapsto \alpha_{\mathfrak{p}} \in F_{\mathfrak{p}}$  given by  $\widehat{\theta}_{\mathfrak{p}}(\alpha) \circ \eta_{\mathfrak{p}} = \eta_{\mathfrak{p}} \circ \alpha_{\mathfrak{p}}$ . By definition, we also

have  $t_p(A) = 1$ ; so,  $\tau = \log_p \circ \sigma \circ t_p$  satisfies  $\tau \circ \hat{\rho}(\alpha) = i_p(\alpha^{\sigma(1-c)})\tau$  for an extension  $\sigma : M \hookrightarrow \overline{\mathbb{Q}}_p$  of  $\sigma$ . Thus by Theorem 4.5, the corresponding coordinate  $\tau_l = \tau_{\sigma,l}$  on the  $l$ -adic deformation space satisfies the same invariance property (i.e., they are  $\hat{\rho}$ -eigen coordinate and  $\tau_l(A) = 0$ ). By our choice,  $M' = \text{End}_F^0(A \otimes_{\mathcal{W}_l} \overline{\mathbb{F}}_l)$  with  $M \not\cong M'$ , which is a contradiction as  $M' = \text{End}_F^0(A \otimes_{\mathcal{W}_l} \overline{\mathbb{F}}_l) \cong \text{End}_F^0(A_{0/\overline{\mathbb{F}}_p}) = M$  (as  $F$ -algebras) by Corollary 4.6.  $\square$

## 6. CM PERIODS AND PROPORTIONALITY CONSTANTS

In the proof of the indecomposability in the previous section, proportionality of  $\tau_p$  and  $\tau_l$  proven in Theorem 4.5 played an important role. We study the proportionality constant of our  $\hat{\rho}$ -eigen  $\varphi$ -coordinate at each  $v \in \Xi$  in terms of CM periods of Katz and Shimura, starting with a CM abelian variety (not with the non-CM abelian variety we did so far). We show that the constant is the ratio of the square of  $v$ -adic CM periods. This section is independent of the proof of the local indecomposability (and the reader with interest only in local indecomposability may skip this section).

Let  $x_0 \in Sh^{(\Xi)}(\overline{\mathbb{F}}_p)$  be in the ordinary locus. The Serre–Tate deformation space of  $x_0$  is then isomorphic to  $\hat{S}_p = \hat{S}_p^{\text{ord}} = \widehat{\mathbb{G}}_m \otimes_{\mathbb{Z}} O$  (as  $p$  is assumed to be unramified in  $F/\mathbb{Q}$ ) embedded into  $Sh_{W_p}^{(\Xi)}$ . Let  $x_1 \in Sh^{(\Xi)}$  be the origin  $\mathbf{1} \in \hat{S}_p \hookrightarrow Sh_{W_p}^{(\Xi)}$ , and let  $(A_1, \overline{\lambda}_1, \theta_1, \eta_1^{(\Xi)})_{/\mathcal{W}_k}$  be the CM abelian variety (the canonical lift) sitting over  $x_1$  (which is originally defined over  $W_p$  but it descends to  $\mathcal{W}_k$  for a number field  $k$  by the theory of complex multiplication [ACM]), where  $\theta_1 : M \cong \text{End}_F^0(A_{1/\mathcal{W}_k})$ . This is the CM point we study. Suppose that  $A_1 \times_{\mathcal{W}_k} \overline{\mathbb{F}}_l$  is also ordinary for  $l \in \Xi$  different from  $p$ . So we may assume that  $\Xi = \{p, l\}$ , and we suppose that  $l$  is unramified in  $F/\mathbb{Q}$ .

At  $v = p, l$ , we fix  $\eta_v : \mu_{v^\infty} \otimes_{\mathbb{Z}} O \hookrightarrow A_1[v^\infty]$  defined over  $\mathcal{W}_v$ . This is possible as  $\mathcal{W}_v$  is a strict henselization of  $\mathbb{Z}_{(v)}$ . This level structure (together with  $\eta_1^{(\Xi)}$ ) gives rise to a point on the  $v$ -adic Igusa tower  $Ig_{v,\infty} := \text{Isom}_{\mathcal{W}_v}(\mu_{v^\infty} \otimes_{\mathbb{Z}} O_{/Sh_{\mathcal{W}_v}^{\text{ord}}}, \mathbf{A}_{/Sh_{\mathcal{W}_v}^{\text{ord}}}[v^\infty]^\circ)$  for the universal abelian scheme  $\mathbf{A}$  over the ordinary locus  $Sh_{\infty}^{\text{ord}}$  of  $Sh^{(v)}$ . We write simply again  $x_1$  for this point on  $Ig_{v,\infty}$ . By fixing  $\eta_v$ , we have a unique identification of  $\hat{S}_v$  with  $\widehat{\mathbb{G}}_m \otimes_{\mathbb{Z}} O$ . Define  $\hat{\rho} : T_M(\mathbb{Z}(\Xi)) \rightarrow G(\mathbb{A}^{(\Xi),\infty})$  by  $\theta_1(\alpha) \circ \eta_1^{(\Xi)} = \eta_1^{(\Xi)} \circ \hat{\rho}(\alpha)$ . Then write  $\tau_{v,\varphi}$  for the  $\hat{\rho}$ -eigen  $\varphi$ -coordinate of  $(\hat{S}_v)^{v\text{-an}}$  if  $v = p, l$ . In other words, if  $v = p$ , for the  $\varphi$ -projection  $\varphi_p : W_p \otimes_{\mathbb{Z}} O \rightarrow W_p$ , we have  $\tau_{p,\varphi} = \log_p \circ \varphi_p \circ t_{p,\varphi}$  for the Serre–Tate coordinate  $t_{p,\varphi} : \hat{S}_{p,\varphi}^{\text{ord}} = \widehat{\mathbb{G}}_m \otimes_{\mathbb{Z}_p} O_{p,\varphi}$ .

By ordinarity, any prime factor  $\mathfrak{v}|v$  (for  $v = p, l$ ) in  $O$  splits in  $M$ ; in particular,  $v$  is unramified in  $M/\mathbb{Q}$ . Write  $\Phi$  for the CM type of  $A_1$ ; so,  $\Phi$  is a collection of a half of embeddings of  $M$  into  $\overline{\mathbb{Q}}$  such that  $\sum_{\phi \in \Phi} \phi$  as a representation of  $M$  is isomorphic to  $\Omega_{A_1/k} \otimes_k \overline{\mathbb{Q}}$ . Supposing that  $k$  is large containing all conjugates of  $M$  in  $\overline{\mathbb{Q}}$ , we decompose  $H^0(A_1, \Omega_{A_1/\mathcal{W}_k}) = \bigoplus_{\phi \in \Phi} \mathcal{W}_k \omega_\phi$  with nowhere vanishing differentials  $\omega_\phi$  having an eigen-property  $\theta_1(\alpha)^* \omega_\phi = \phi(\alpha) \omega_\phi$  for  $\alpha \in M = \text{End}_F^0(A_1)$ . As is well known,  $R := \theta_1^{-1}(\text{End}_O(A_{1/\mathcal{W}_k}))$  is an  $O$ -order of  $M$ , in which any prime  $\mathfrak{v}|v$  of  $R$  splits (see Lemma 2.2).

Fixing complex uniformization  $L \hookrightarrow \mathbb{C}^\Phi \rightarrow A_1(\mathbb{C})$  for a proper  $R$ -ideal  $L \subset R$ . Writing the variable of  $\mathbb{C}^\Phi$  as  $u = (u_\phi)$ ,  $\omega_{\infty,\phi} := du_\phi \in H^0(A_1, \Omega_{A_1/\mathbb{C}})$  satisfies  $\theta_1(\alpha)^* \omega_{\infty,\phi} = \phi(\alpha) \omega_{\infty,\phi}$  for  $\alpha \in M$ . Thus  $\omega_\phi$  and  $\omega_{\infty,\phi}$  are proportional, getting Shimura’s CM period  $\Omega_{\infty,\phi} \in \mathbb{C}^\times$  by  $\omega_\phi = \Omega_{\infty,\phi} \omega_{\infty,\phi}$ . Since  $\int_\gamma \omega_{\infty,\phi} \in \phi(L) \subset \mathcal{W}_k$  for all  $\gamma \in \pi_1(A_1(\mathbb{C}), \mathbf{0})$  for the origin  $\mathbf{0} \in A_1$ , we have the classical algebraic period identity  $\int_\gamma \omega_\phi \in \mathcal{W} \cdot \Omega_{\infty,\phi}$ . We put  $\Omega_\infty = (\Omega_{\infty,\phi})_{\phi \in \Phi} \in (\mathbb{C}^\times)^\Phi$ .

For  $v = p, l$ , the level structure  $\eta_v$  induces  $\hat{\eta}_v : \widehat{\mathbb{G}}_m \otimes_{\mathbb{Z}} O = \varprojlim_n (\mu_{v^n} \otimes_{\mathbb{Z}} O) \cong \hat{A}_1$  defined over  $W_v$  for the formal completion  $\hat{A}_1$  of  $A_1$  along the origin  $\mathbf{0} \in A_1(\overline{\mathbb{F}}_v)$ . Thus, writing  $\mathbb{G}_m = \text{Spec}(W[t, t^{-1}])$ , the push forward  $\hat{\eta}_{v,*}(\frac{dt}{t} \otimes 1)$  can be written as a unique sum  $\hat{\eta}_{v,*}(\frac{dt}{t} \otimes 1) = \sum_{\phi \in \Phi} \omega_{v,\phi}$  satisfying  $\theta_1(\alpha)^* \omega_{v,\phi} = \phi(\alpha) \omega_{v,\phi}$  for  $\alpha \in M = \text{End}_F^0(A_1)$ . We then define Katz’s  $v$ -adic period  $\Omega_{v,\phi}$  by  $\omega_\phi = \Omega_{v,\phi} \omega_{v,\phi}$ . We put  $\Omega_v = (\Omega_{v,\phi})_{\phi \in \Phi} \in W_v^\Phi$  (see [K1]). Thus we get

**Proposition 6.1.** *Under our identification  $\mathbb{C}_p = \mathbb{C}$ , the proportionality constant of  $\omega_{p,\phi}$  and  $\omega_{\infty,\phi}$  both in  $H^0(A_1, \Omega_{A_1/\mathbb{C}}) = H^0(A_1, \Omega_{A_1/\mathbb{C}_p})$  is given by  $\Omega_{\infty,\phi}/\Omega_{p,\phi}$  for each  $\phi \in \Phi$ ; in other words, we have*

$$\Omega_{p,\phi} \omega_{p,\phi} = \omega_\phi = \Omega_{\infty,\phi} \omega_{\infty,\phi} = \Omega_{\infty,\phi} du_\phi$$

for  $\omega_\phi \in H^0(A_1, \Omega_{A_1/\mathcal{W}_\Xi})$ .

Assuming that  $A_1$  is also ordinary at  $l \in \Xi$ , applying the above proposition to  $\{p, \infty\}$  and then to  $\{l, \infty\}$ , we get

**Corollary 6.2.** *Under our identification  $\mathbb{C}_p = \mathbb{C}_l$ , the proportionality constant of  $\omega_{p,\phi}$  and  $\omega_{l,\phi}$  both in  $H^0(A_1, \Omega_{A_1/\mathbb{C}_p}) = H^0(A_1, \Omega_{A_1/\mathbb{C}_l})$  is given by  $\Omega_{l,\phi}/\Omega_{p,\phi}$  for each  $\phi \in \Phi$ ; in other words, we have*

$$\Omega_{p,\phi}\omega_{p,\phi} = \omega_\phi = \Omega_{l,\phi}\omega_{l,\phi}$$

for  $\omega_\phi \in H^0(A_1, \Omega_{A_1/\mathcal{W}_\Xi})$ .

We have three Kodaira–Spencer maps:

$$(6.1) \quad (\pi_*\Omega_{\mathbf{A}/Sh_{\mathcal{W}_k}^{(p)}}[\varphi])^{\otimes 2} = \underline{\omega}_{\mathcal{W}_k}^{2\varphi} \cong \Omega_{Sh^{(p)}/\mathcal{W}_k}[\varphi], \quad (\pi_*\Omega_{\mathbf{A}/\widehat{S}_p^{ord}}[\varphi])^{\otimes 2} = \underline{\omega}_{W_p}^{2\varphi} \cong \Omega_{\widehat{S}_p/W_p}[\varphi],$$

$$\text{and } (\pi_*\Omega_{\mathbf{A}/\widehat{S}_l^{ord}}[\varphi])^{\otimes 2} = \underline{\omega}_{W_l}^{2\varphi} \cong \Omega_{\widehat{S}_l/W_l}[\varphi].$$

Here  $\mathbf{A}$  denotes the universal abelian scheme corresponding to the bases  $Sh_\infty^{ord}$ ,  $\widehat{S}_p^{ord}$  and  $\widehat{S}_l^{ord}$  whose origin corresponding to  $x_1$ . As before, “[ $\varphi$ ]” indicates the  $\varphi$ -eigenspace under the natural action of  $O$ . Taking the fiber at  $x_1$  of (6.1), we get

$$(6.2) \quad (H^0(A_1, \Omega_{A_1/\mathcal{W}_k}[\varphi]))^{\otimes 2} = \underline{\omega}^{2\varphi}(x_1)_{\mathcal{W}_k} \cong \Omega_{Sh^{(p)}/\mathcal{W}_k}[\varphi](x_1),$$

$$(H^0(A_1, \Omega_{A_1/W_p}[\varphi]))^{\otimes 2} = \underline{\omega}^{2\varphi}(x_1)_{W_p} \cong \Omega_{\widehat{S}_p/W_p}[\varphi](x_1),$$

$$(H^0(A_1, \Omega_{A_1/W_l}[\varphi]))^{\otimes 2} = \underline{\omega}^{2\varphi}(x_1)_{W_l} \cong \Omega_{\widehat{S}_l/W_l}[\varphi](x_1).$$

We can identify  $\Phi$  with  $I$  by  $\phi \mapsto \phi|_F =: \varphi$ . Then  $\varphi$ -eigenspace under the action of  $O$  is identical to the  $\phi$ -eigenspace under the action of  $R$ . The data

$$\{\omega_\phi \in H^0(A_1, \Omega_{A_1/\mathcal{W}_k}[\varphi]), \omega_{p,\phi} \in H^0(A_1, \Omega_{A_1/W_p}[\varphi]), \omega_{l,\phi} \in H^0(A_1, \Omega_{A_1/W_l}[\varphi])\}$$

give rise to the corresponding differentials

$$\{\omega_\phi^{\otimes 2} = \omega_\phi \otimes \omega_\phi \in \Omega_{\mathbf{A}/Sh_{\mathcal{W}_k}^{(p)}}[\varphi](x_1), \omega_{p,\phi}^{\otimes 2} \in \Omega_{\mathbf{A}/\widehat{S}_p/W_p}[\varphi](x_1), \omega_{l,\phi}^{\otimes 2} \in \Omega_{\mathbf{A}/\widehat{S}_l/W_l}[\varphi](x_1)\}.$$

Define  $\widehat{\rho} : T_M(\mathbb{Z}(\Xi)) \rightarrow G(\mathbb{A}(\Xi), \infty)$  by  $\theta_1(\alpha) \circ \eta_1^{(\Xi)} = \eta_1^{(\Xi)} \circ \widehat{\rho}(\alpha)$ . Write  $\tau_{v,\varphi}$  for the additive Serre–Tate  $\widehat{\rho}$ -eigen  $\varphi$ -coordinate of  $(\widehat{S}_p)^{v\text{-an}}$  for  $v = l, p$ . Then we define the proportionality constant  $C_\varphi^{v,w}$  to be  $\tau_{v,\varphi} = C_\varphi^{v,w} \cdot \tau_{w,\varphi}$ . The constant  $0 \neq C_\varphi^{v,w}$  is determined by comparing  $d\tau_{v,\varphi}$  and  $d\tau_{w,\varphi}$  in  $\Omega_{Sh^{(p)}/\mathbb{C}_v}[\varphi](x_1) = \Omega_{Sh^{(p)}/\mathbb{C}_w}[\varphi](x_1)$ , in which we have algebraic  $\omega_\phi^{\otimes 2}$ . Note that, by our construction and [K] Main Theorem 3.7.1, we have  $d\tau_{v,\varphi} = \frac{dt_{v,\varphi}}{t_{v,\varphi}} = \omega_{v,\phi}^{\otimes 2}$  with  $t_{v,\varphi} = \varphi \circ t_{\mathbf{v},\varphi}$  for the unique continuous extension  $\varphi : F_{\mathbf{v},\varphi} \hookrightarrow \mathbb{C}_v$  of the original  $\varphi : F \hookrightarrow \overline{\mathbb{Q}}$ . Thus from Corollary 6.2, we get

**Theorem 6.3.** *Under our identification  $\mathbb{C}_p = \mathbb{C}_l$ , the proportionality constant  $C_\varphi^{p,l}$  of  $\tau_{p,\varphi}$  and  $\tau_{l,\varphi}$  is given by  $\Omega_{l,\phi}^2/\Omega_{p,\phi}^2$  for each  $\varphi \in I$ , where  $\phi \in \Phi$  corresponds to  $\varphi$  by  $\phi|_F = \varphi$ ; in other words, we have*

$$\Omega_{p,\phi}^2 d\tau_{p,\varphi} = \omega_\phi^{\otimes 2} = \Omega_{l,\phi}^2 d\tau_{l,\varphi} \quad \text{in } \Omega_{Sh^{(\Xi)}/\mathbb{C}_p}(x_1) = \Omega_{Sh^{(\Xi)}/\mathbb{C}_l}(x_1)$$

for  $\omega_\phi^{\otimes 2} \in \underline{\omega}^{2\varphi}$ .

**Question 6.4.** *If we start with non-CM AVRMs  $(A, \overline{\lambda}, \theta)_{/k}$ , we can associate two  $p$ -adic “periods”  $\tau_{p,\varphi}(A)$  for each  $\varphi \in \Sigma_p^{ord}$  and  $\Omega(A)_{p,\varphi}$  given by  $\Omega(A)_{p,\varphi}\omega_{p,\varphi} = \omega_\varphi$  choosing a generator  $\omega_\varphi$  of  $H^0(A, \Omega_{A/\mathcal{W}_k}[\varphi])$  over  $\mathcal{W}_k$ . It is an interesting question if these periods are related to some  $p$ -adic  $L$ -values coming out of the abelian scheme  $A_{/k}$ .*

If  $A$  is an ordinary abelian variety with CM, we have  $\tau_{p,\varphi}(A) = 0$  (as  $t_p(A) = 1$ ). Thus, we get only one nontrivial period, the Katz CM period  $\Omega(A)_{p,\varphi}$ . By a result of Shimura, the value of an algebraic modular form  $f$  at the CM abelian variety “ $A$ ” with complex invariant differentials  $\{du_\phi\}_\phi$  is equal to a monomial of  $\{\Omega(A)_{\infty,\varphi}\}_\varphi$  up to an algebraic number which is the value of  $f$  at  $(A, \omega_\phi)_\phi$ . Similarly, by a result of Katz, the value of  $f$  at the CM abelian variety “ $A$ ” with  $p$ -adic invariant differentials  $\{\omega_{p,\phi}\}$  of an algebraic modular form is equal to the same monomial of  $\{\Omega(A)_{p,\varphi}\}_\varphi$  times

$f((A, \omega_\phi)_\phi)$ . An elementary treatment of the theory of Katz and Shimura can be found in [EAI] Section 1.3. If  $A$  is non CM and has ordinary good reduction,  $\tau_{p,\varphi}(A)$  plays a role similar to the Tate period of an abelian variety with multiplicative reduction, though we do not know much about its arithmetic meaning yet.

## 7. DENSITY OF PARTIALLY ORDINARY PRIMES

For the reader's convenience, we give a sketch of a simple proof of the density result for partially ordinary primes used in the proof of Theorem 5.1. We consider the set  $\mathcal{S}^{ord}$  of prime ideals of  $k$  unramified over  $\mathbb{Q}$  at which  $A$  has partially ordinary good reduction for a given non CM absolutely simple AVRМ  $(A, \lambda, \theta)$ . Here  $(A, \lambda, \theta)$  is defined over the number field  $k \subset \overline{\mathbb{Q}}$ , and  $A$  has partially ordinary good reduction at  $\mathfrak{P}$  if  $A$  extends to an abelian scheme over  $\mathfrak{D}/\mathfrak{P}$  and  $A[\mathfrak{p}](\overline{\mathbb{F}}_p) \neq 0$  for an algebraic closure  $\overline{\mathbb{F}}_p$  of  $\mathfrak{D}/\mathfrak{P}$ . First suppose that  $k/\mathbb{Q}$  is a Galois extension. Then the set of primes of  $k$  split over  $\mathbb{Q}$  has Dirichlet density 1. Let  $\mathcal{S}_f^{ord} \subset \mathcal{S}^{ord}$  be the set of all primes in  $\mathcal{S}^{ord}$  with residual degree  $f$ . We have  $\mathcal{S}^{ord} = \bigsqcup_{f=1}^{[k:\mathbb{Q}]}$   $\mathcal{S}_f^{ord}$ . If a prime  $\mathfrak{P}$  in  $\mathfrak{D}$  has residual degree 1, the reduction  $A_{\mathfrak{P}} = A \otimes_{\mathfrak{D}} \mathfrak{D}/\mathfrak{P}$  has the  $p$ -power Frobenius endomorphism  $\phi$  whose eigenvalues  $\alpha$  satisfies  $|\alpha^\varphi| = \sqrt{p}$  for all  $\varphi \in \text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ . Since  $a = \alpha + \alpha^c$  for the complex conjugation  $c$  (i.e., the Rosati involution of  $\text{End}_O(A_{\mathfrak{P}})$  with respect to  $\lambda$ ) is in  $O$  (as  $A_{\mathfrak{P}}$  is defined over  $\mathfrak{D}/\mathfrak{P} = \mathbb{F}_p$ ). Thus we get  $|a^\varphi| \leq 2\sqrt{p}$ . Taking the norm to  $\mathbb{Q}$ , we have  $|N_{F/\mathbb{Q}}(a)| \leq 2^d p^{d/2}$ . Thus if  $p|(a)$  in  $\mathfrak{D}$ , we have  $p^d |N_{F/\mathbb{Q}}(a)$ ; so,  $p^d \leq |N_{F/\mathbb{Q}}(a)| \leq 2^d p^{d/2}$  as long as  $a \neq 0$ . Thus if  $a \neq 0$ , we have a prime  $\mathfrak{p} \nmid a$  in  $O$  above  $p$  for  $p \geq 5$ . Then as is well known, this implies that  $A[\mathfrak{p}](\overline{\mathbb{F}}_p) \neq 0$  (i.e.,  $\mathfrak{p} \in \mathcal{S}_p^{ord}$ ). For any  $p$ -adic Galois representation  $\rho : \text{Gal}(\overline{\mathbb{Q}}/k) \rightarrow GL_m(\mathbb{Q}_p)$ , fix a Haar measure on  $\text{Im}(\rho)$  with volume 1. By Chebotarev density (e.g., [ALR] I-8, Corollary 2 combined with Exercise there and [Se] §2.1), the density of primes of  $k$  whose Frobenius falling in a closed set of volume 0 is equal to 0. Applying this to the Galois representation  $\rho_l$  on the  $l$ -adic Tate module of  $A/k$  taking a prime  $l$  of  $F$ . By the result of [BGK] Theorem C, the set of  $\mathfrak{P}$  whose Frobenius has trace 0 has volume 0. Thus for non CM AVRМ  $A$ , the set of primes  $\mathfrak{P}$  with  $a = 0$  has Dirichlet density 0 by the quantitative Chebotarev density theorem (though Serre works over  $\mathbb{Q}$ , his argument generalizes to a number field  $k$  without any modification). If  $k/\mathbb{Q}$  is not a Galois extension, we just replace  $k$  by its Galois closure and apply the above argument. Since as in Theorem 5.1, we say that  $A$  does not have CM if  $A \times_k \overline{\mathbb{Q}}$  does not have CM (i.e.,  $\text{End}_F^0(A \times_k \overline{\mathbb{Q}}/\overline{\mathbb{Q}}) = F$ ), being non CM is insensitive to any base change finite over  $k$ ; in particular, the abelian variety  $A$  over the Galois closure of  $k$  does not have CM. Thus we have

**Proposition 7.1.** *Let  $(A, \lambda, \theta)$  be a non CM AVRМ defined over  $k \subset \overline{\mathbb{Q}}$ . Replacing  $k$  by its finite extension if necessary, the set  $\mathcal{S}_1^{ord}$  for a given AVRМ has Dirichlet density equal to 1.*

It is interesting to know the Dirichlet density of the set  $\{l | \mathfrak{L} \in \mathcal{S}_f^{ord}\}$  of rational primes if  $f > 1$  (this question is not about the density over  $k$  but over  $\mathbb{Q}$ ). The result [O] 2.7 tells us something about  $\mathcal{S}_2^{ord}$ .

## REFERENCES

- [ABV] D. Mumford, *Abelian Varieties*, TIFR Studies in Mathematics, Oxford University Press, New York, 1994.
- [ACM] G. Shimura, *Abelian Varieties with Complex Multiplication and Modular Functions*, Princeton University Press, Princeton, NJ, 1998.
- [ALR] J.-P. Serre, *Abelian  $l$ -Adic Representations and Elliptic Curves*, Research Notes in Mathematics **7**, A K Peters, 1998
- [ARG] G. Cornell and J. H. Silverman, editors, *Arithmetic Geometry*, Springer, New York, 1986.
- [BAL] N. Bourbaki, *Algèbre*, Hermann, Paris, 1958.
- [BDJ] K. Buzzard, F. Diamond and F. Jarvis, On Serre's conjecture for mod  $l$  Galois representations over totally real fields. *Duke Math. J.* **155** (2010), 105–161
- [BGK] G. Banaszak, W. Gajda and P. Krasoń, On Galois representations for abelian varieties with complex and real multiplications. *J. Number Theory* **100** (2003), 117–132
- [BL] S. Bosch and W. Lütkebohmert, Formal and rigid geometry. I. Rigid spaces. *Math. Ann.* **295** (1993), 291–317.
- [C] B. Conrad, Several approaches to non-Archimedean geometry. In “ $p$ -adic geometry”, Univ. Lecture Ser., **45** (2008), 9–63, Amer. Math. Soc., Providence, RI.

- [dJ] A. J. De Jong, Crystalline Dieudonné module theory via formal and rigid geometry. Inst. Hautes Études Sci. Publ. Math. No. **82** (1995), 5–96
- [dJ1] A. J. De Jong, Erratum to: “Crystalline Dieudonné module theory via formal and rigid geometry” Inst. Hautes Études Sci. Publ. Math. No. **87** (1998), 175.
- [DLS] S. Helgason *Differential geometry, Lie groups, and symmetric spaces*, Pure and Applied Mathematics, **80**, Academic Press, New York-London, 1978.
- [E] M. Emerton, A  $p$ -adic variational Hodge conjecture and modular forms with complex multiplication, preprint
- [EAI] H. Hida, *Elliptic Curves and Arithmetic Invariants*, book manuscript, 2012, to appear in Springer Monograph in Mathematics.
- [EGA] A. Grothendieck and J. Dieudonné, *Eléments de Géométrie Algébrique*, Publications IHES **4** (1960), **8** (1961), **11** (1961), **17** (1963), **20** (1964), **24** (1965), **28** (1966), **32** (1967).
- [GIT] D. Mumford, *Geometric Invariant Theory*, Ergebnisse **34**, Springer, New York, 1965.
- [G] E. Ghate, Ordinary forms and their local Galois representations, in the proceedings of a conference held in Hyderabad: Algebra and number theory, Hindustan Book Agency (2005), 226–242
- [GV] E. Ghate and V. Vatsal, On the local behaviour of ordinary  $\mathbb{L}$ -adic representations. Ann. Inst. Fourier (Grenoble) **54** (2004), 2143–2162.
- [GV1] B. Balasubramanyam, E. Ghate and V. Vatsal, On local Galois representations attached to ordinary Hilbert modular forms, preprint 2012
- [GME] H. Hida, *Geometric Modular Forms and Elliptic Curves*, second edition, World Scientific, Singapore, 2012.
- [H06] H. Hida, Automorphism groups of Shimura varieties of PEL type, Documenta Math. **11** (2006), 25–56.
- [H10] H. Hida, The Iwasawa  $\mu$ -invariant of  $p$ -adic Hecke  $L$ -functions, Annals of Math. **172** (2010), 41–137.
- [H11a] H. Hida, Constancy of adjoint  $\mathcal{L}$ -invariant, J. of Number Theory **131** (2011), 1331–1346.
- [H11b] H. Hida, Irreducibility of the Igusa tower over unitary Shimura varieties, Clay Mathematics Proceedings **13** (2011), 187–203.
- [IAT] G. Shimura, *Introduction to the Arithmetic Theory of Automorphic Functions*, Princeton University Press, Princeton, NJ, and Iwanami Shoten, Tokyo, 1971.
- [K] N. M. Katz, Serre–Tate local moduli, In “Surfaces Algébriques,” Lecture Notes in Math. **868** (1978), 138–202.
- [K1] N. M. Katz,  $p$ -adic  $L$ -functions for CM fields, Inventiones Math. **49** (1978), 199–297.
- [M] T. Miyake, On automorphism groups of the fields of automorphic functions, Annals of Math. (2) **95** (1972), 243–252.
- [N] R. Noot, Abelian varieties-Galois representation and properties of ordinary reduction, Compositio Math. **97** (1995), 161–171.
- [NAA] S. Bosch, U. Günzter, R. Remmert, Non-Archimedean analysis, Springer-Verlag, 1984.
- [O] A. Ogus, Hodge cycles and crystalline cohomology, in: *Hodge Cycles, Motives, and Shimura Varieties*, Chapter VI, Lecture Notes in Math. **900** (1982) 357–414.
- [PAF] H. Hida,  *$p$ -Adic Automorphic Forms on Shimura Varieties*, Springer Monographs in Mathematics, 2004, Springer.
- [Ra] M. Rapoport, Compactifications de l’espace de modules de Hilbert-Blumenthal, Compositio Math. **36** (1978), 255–335.
- [Ri] K. A. Ribet, Galois action on division points of abelian varieties with real multiplications, American J. Math. **98** (1976), 751–804.
- [Se] J.-P. Serre, Quelques applications du théorème de densité de Chebotarev. Inst. Hautes Études Sci. Publ. Math. **54** (1981), 323–401 (Œuvres III, No. 125).
- [Sh] G. Shimura, On canonical models of arithmetic quotients of bounded symmetric domains, Annals of Math. **91** (1970), 144–222; II, **92** (1970), 528–549 ([70a–b] in Collected Papers II).
- [T] J. Tate,  $p$ -divisible groups, Proc. Conf. on local fields, Driebergen 1966, Springer 1967, 158–183.
- [Z] B. Zhao, Local indecomposability of Hilbert modular Galois representations, preprint, 2012 (posted in web: arXiv:1204.4007v1 [math.NT])

DEPARTMENT OF MATHEMATICS, UCLA, LOS ANGELES, CA 90095-1555, U.S.A.

*E-mail address:* hida@math.ucla.edu