

Structure theorems, conjectures and questions on Iwasawa modules, past and present

Haruzo Hida

Department of Mathematics, UCLA,
Los Angeles, CA 90095-1555, U.S.A.

Colloquium talk at Renming U, Beijing, Fall, 2023, China.

We first discuss Conjectures/Problems (in cyclotomic theory) Iwasawa described in [one of his final unpublished manuscripts](#) [U3]. Then if time allows, we indicate which of his problems has generalizations in more general settings of adjoint Selmer groups via the theory of modular forms. In the general case, the Iwasawa algebra is replaced by a p -adic Hecke algebra (which is a universal deformation ring by a theorem of Taylor–Wiles). We encounter new interesting problems related to Iwasawa’s question in the general case. We fix a prime $p \geq 5$ throughout this lecture.

§1. Iwasawa's cyclotomic tower.

Iwasawa studied p -cyclotomic fields (in 1950s): the field $F_n = \mathbb{Q}(\zeta_{p^n})$ generated by the group $\zeta_{p^n}(\mathbb{C}) = \left\{ \exp\left(\frac{2\pi ia}{p^n}\right) \right\}_{a=1}^{p^n}$ of p^n -th roots of unity, and put $F_\infty = \bigcup_n F_n \subset \mathbb{C}$. Its Galois group

$$G_n := \text{Gal}(F_n/\mathbb{Q}) \xrightarrow[\nu_n]{\sim} (\mathbb{Z}/p^n\mathbb{Z})^\times, \quad G_\infty := \text{Gal}(F_\infty/\mathbb{Q}) \xrightarrow[\nu]{\sim} \mathbb{Z}_p^\times$$

with ν_n^{-1} sending $a \in (\mathbb{Z}/p^n\mathbb{Z})^\times$ to the field automorphism σ_a taking $\zeta_{p^n} = \exp\left(\frac{2\pi i}{p^n}\right)$ to $\zeta_{p^n}^a = \exp\left(\frac{2\pi ia}{p^n}\right)$.

Let \mathcal{H}_n be the maximal unramified abelian extension of F_n (Hilbert's class field). Then $\text{Gal}(\mathcal{H}_n/F_n)$ is canonically isomorphic to

$$Cl_n := \frac{\{\text{fractional ideals of } F_n\}}{\{(\alpha) \mid \alpha \in F_n^\times\}} \quad (\text{the class group})$$

such that a prime $\mathfrak{l} \nmid p$ goes to $\text{Frob}_\mathfrak{l}$ (Frobenius at \mathfrak{l}).

§2. **Minus part of p -class group.** Put $Cl_n^- = Cl_n / Cl_n^{c+1}$ for complex conjugation c and $Cl_n^+ = \text{Ker}(Cl_n \rightarrow Cl_n^-)$. Write H_n for the subfield of \mathcal{H}_n such that $\text{Gal}(H_n/F_n) \cong C_n^-$ (p -Hilbert's class field).

We know a formula of the order of Cl_n^- due to Dirichlet(1839; Gauss 1801?) and Kummer:

Theorem: $|Cl_n^-| = 2p^n \prod_{\chi: (\mathbb{Z}/p^n\mathbb{Z})^\times \rightarrow \mathbb{C}^\times; \chi(-1)=-1} \frac{1}{2} L(0, \chi^{-1})$
with $L(0, \chi^{-1}) = -\frac{1}{p^n} \sum_{a=1}^{p^n} \chi^{-1}(a)a$.

Here $L(s, \chi) = \sum_{n=1}^{\infty} \frac{\chi(n)}{n^s}$ (Dirichlet L function).

The character χ extends to a algebra homomorphism χ of the group algebra $\mathbb{Z}[G_n]$ to $\mathbb{Z}[\chi]$. For $\theta = \theta_n := \sum_{a=1}^{p^n} \frac{a}{p^n} \sigma_a^{-1} \in \mathbb{Q}[G_n]$ (Stickelberger's element), we have $\chi(\theta_n) = -L(0, \chi)$.

§3. **Stickelberger's theorem.** By $\chi(\theta_n) = -L(0, \chi)$, we imagine that θ would kill Cl_n^- . The group algebra $\mathbb{Z}[G_n]$ acts on Cl_n by $\mathfrak{a}^\xi = \prod_{\sigma} \mathfrak{a}^{\sigma a_{\sigma}}$ for $\xi := \sum_{\sigma} a_{\sigma} \sigma \in \mathbb{Z}[G_n]$. We put $\mathfrak{S}_n := (\theta_n) \cap \mathbb{Z}[G_n]$ which is called the Stickelberger ideal of F_n . By Kummer (1847) for F_1 and by Stickelberger (1890),

\mathfrak{S}_n annihilates Cl_n^- (but often $\mathbb{Z}[G_n]/\mathfrak{S}_n \not\cong Cl_n^-$).

The obstruction may be Cl_n^+ if non-trivial. We look at the p -primary part $C_n^{\pm} := Cl_n^{\pm} \otimes_{\mathbb{Z}} \mathbb{Z}_p$. Kummer (and Vandiver) conjectured triviality of C_1^+ :

KVC conjecture: $C_1^+ = \{1\}$ (no p -obstruction ?).

We do not have a convincing evidence except for numerical computation for primes up to more than **two billion** (up to 2,147,483,648, W. Hart, D. Harvey, W. Ong, 2017).

§4. **Iwasawa algebra.** For $m > n$ and $R_n = \mathbb{Z}_p[G_n]$:

- $R_m \xrightarrow{\text{Res}_{m,n}} R_n$ given by $\sigma \mapsto \sigma|_{F_n}$ with $\theta_m \mapsto \theta_n$ and

$$R_\infty := \mathbb{Z}_p[[\mathbb{Z}_p^\times]] = \varprojlim_n R_n \cong \mathbb{Z}_p[\zeta][[T]] \quad \text{for primitive } \zeta^{p-1} = 1,$$

- $C_m^- \rightarrow C_n^-$ by the norm maps $\mathfrak{a} \mapsto \prod_{\sigma \in \text{Ker}(\text{Res}_{m,n})} \mathfrak{a}^\sigma$ and the R_∞ -module $C_\infty^- := \varprojlim_n C_n^- \cong \text{Gal}(H_\infty/F_\infty)$ ($H_\infty = \bigcup_n H_n$).

Since $\mathbb{Z}_p[\zeta] \cong \prod_{j=1}^{p-1} \mathbb{Z}_p$ by the j -th projection $\omega_j(\zeta) = \zeta^j$, we have

$$R_\infty \cong \prod_{j=0}^{p-2} \Lambda \quad \text{for } \Lambda = \mathbb{Z}_p[[T]] \quad (\text{one variable power series ring}).$$

The R_∞ -module C_∞^- is called an **Iwasawa module**. and Λ is called the **Iwasawa algebra**. Then the odd j -component $L_j(T) := \theta_\infty^{(j)}(T)$ is the p -adic L-function: $L_j((1+p)^s - 1)|_{s=1-k} = L(1-k, \omega_{j-k+1})$ for $0 < k \in \mathbb{Z}$ (Iwasawa 1969).

§5. Cyclicity, 1969

Iwasawa's theorem: Let $X_j := \omega_j C_\infty^-$ for an odd j . Then as a module over Λ , under KVC, $X_j \cong \Lambda/(L_j(T)) \Rightarrow C_n^- \cong R_n/\mathfrak{S}_n R_n$.

Cyclicity Conjecture: $X_j \cong \Lambda/(L_j(T))$ up to finite error always.

The zeros of the Riemann zeta function are expected to be all simple and lie in a special line: $\frac{1}{2} + \mathbb{R}\sqrt{-1}$?

Semi-simplicity Cojecture: *The zeros of $L_j(T)$ are simple and lie on \mathbb{Z}_p .*

The following version has been verified up to two billion.

Linearity Question/conjecture: *The number of zeros of L_j is less than or equal to 1; i.e., $L_j(T) = (T - \alpha)U$ or U with $U \in \Lambda^\times$.*

Plainly

Linearity \Rightarrow Semi-simplicity \Rightarrow Cyclicity.

These Conjectures are **open** and made in [U3] in 1987.

§6. Cyclotomic theory to a 3-dimensional theory.

For a cusp form f of level N with $f|T(n) = \lambda(T(n))f$, we have a compatible system $r := \{r_{\mathfrak{p}}\}_{\mathfrak{p}|p}$ of 2-dim Galois representations with $\text{Tr}(r_{\mathfrak{p}}(\text{Frob}_l)) = \lambda(T(l))$ for primes $l \nmid Np$. For simplicity, we assume that $N = p$, hereafter. Take $r_{\mathfrak{p}}$ with absolutely irreducible $\bar{\rho} = r_{\mathfrak{p}} \bmod \mathfrak{p}$ ($\mathbb{F} := O/\mathfrak{p}$) with reducible $\bar{\rho}|_{\text{Gal}(\bar{\mathbb{Q}}_p/\mathbb{Q}_p)} \sim \begin{pmatrix} \bar{\epsilon} & * \\ 0 & \bar{\delta} \end{pmatrix}$ ($\bar{\epsilon} \neq \bar{\delta}$). Take a complete local $W(\mathbb{F})$ -algebra A and write $W = W(\mathbb{F})$. A representation ρ of $\mathcal{G} := \text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q})$ with values in $\text{GL}_2(A)$ is a **p -ordinary** deformation of $\bar{\rho}$ if $\rho \bmod \mathfrak{m}_A = \bar{\rho}$ and

(ord $_p$) $\rho|_{\text{Gal}(\bar{\mathbb{Q}}_p/\mathbb{Q}_p)} \sim \begin{pmatrix} \epsilon & * \\ 0 & \delta \end{pmatrix}$ with δ unramified ($\delta \bmod \mathfrak{m}_A = \bar{\delta}$), ρ unramified over $\bar{\mathbb{Q}}^{\text{Ker}(\bar{\rho})}$ outside p .

There exists a universal pair (R, ρ) with $R \in CL_W$ and a p -ordinary deformation $\rho : \mathcal{G} \rightarrow \text{GL}_2(R)$ **dependent** on \mathfrak{p} such that

$$\text{Hom}_{W\text{-alg}}(R, A) = \{\rho : \mathcal{G} \rightarrow \text{GL}_2(A) : p\text{-ordinary} \mid (\rho \bmod \mathfrak{m}_A) \sim \bar{\rho}\}$$

by $\rho \leftrightarrow \varphi$ if $\rho \sim \varphi \circ \rho$ for $\varphi \in \text{Hom}_{CL_W}(R, A)$.

§7. Selmer groups.

The group X_j^\vee has an interpretation of the unramified subgroup of $H^1(\mathbb{Q}, \omega_j(R_\infty)^\vee)$ (“ \vee ” denotes Pontryagin dual). Let $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ act on $\mathfrak{sl}_2(A)$ by $x \mapsto \rho(\sigma)x\rho(\sigma)^{-1}$ and write rank 3 representation as $\text{Ad}(\rho)$. So we get a new compatible system $\text{Ad}(r) = \{\text{Ad}(r_p)\}$, and a very unramified subgroup $\text{Sel}(\text{Ad}(\rho))$ of $H^1(\mathbb{Q}, \text{Ad}(\rho) \otimes A^\vee)$. We replace (Λ, X_j) by the universal algebra R and the Pontryagin dual $\text{Sel}(\text{Ad}(\rho))^\vee$.

By myself and Taylor–Wiles, R is **free and finite** over Λ , and $\text{Spec}(R)$ has densely populated points of modular Galois deformations r' . There is a p -adic L-function $L_p : \text{Spec}(R) \ni \rho \mapsto L_p(\rho) \in \mathbb{C}_p$ interpolating $L(1, \text{Ad}(r'))/\Omega_{r'}$ ($L_p \in R$) for modular r' of weight $k > 2$. Here is the **Selmer class number formula**:

If $\bar{\rho}$ is irreducible over $\mathbb{Q}[\zeta_p]$,

$$|\text{Sel}(\text{Ad}(r_p))| = |L_p(r_p)|_p^{-1} \quad \text{and} \quad |L_p(r'_p)|_p = \left| \frac{L(1, \text{Ad}(r'))}{\Omega_{r'}} \right|_p.$$

§8. Cyclicity theorem:

if $\bar{\rho}$ is absolutely irreducible over $\mathbb{Q}[\zeta_p]$, $\text{Sel}(Ad(\rho))^\vee \cong A/(L_p(\rho))$ for primes p outside an explicit finite set E of p -ordinary primes dependent on r .

Under $k > 2$, if $E := \{p \mid \frac{L(1, Ad(r))}{\Omega_r} = L_p(r_p)\}$, $\text{Sel}(Ad(\rho)) = 0$ for $p \notin E$. If $k = 2$, $E := \{p \mid \frac{L(1, Ad(r))}{\Omega_r}\}$. For infinitely many (level raising) primes p , $\frac{L(1, Ad(r))}{\Omega_r} \neq L_p(r_p)$, and $\text{Sel}(Ad(\rho))$ is infinite but cyclic if $p \notin E$. If $k = 1$, r is an Artin representation independent of p (by Deligne–Serre). If r is induced from a real quadratic field F , $\text{Sel}(Ad(\rho))$ is infinite for all ordinary p splitting in F but is cyclic for all p prime to class number h of $\overline{\mathbb{Q}}^{\text{Ker}(Ad(r))}$ ($E = \{p \mid h\}$). See my book published in 2022 from World Scientific.

If $p \nmid L_p(r_p)$, then $R \cong \Lambda$, and hence R is regular.

§9. **Questions on Selmer irregular primes.** Assume $k = 1$. Then for primes with positive density, r_p is p -ordinary.

Question. *Is $\text{Sel}(\text{Ad}(r_p) \otimes \chi)$ cyclic for almost all p with r_p satisfying (ord_p) for an even Dirichlet character χ ?*

Question. *For a fixed pair (r, χ) , when are irregular primes for $\text{Ad}(r) \otimes \chi$ infinitely many?*

A question asked by C. Khare and G. Böckle is

Question. *What is the distribution of primes p with regular R assuming r_p is p -ordinary?*

Suppose $r = \text{Ind}_F^{\mathbb{Q}} \varphi$ for a real quadratic field F . If p splits into pp' in F prime to the class number h_F , regularity of R is equivalent to $\varepsilon^{p-1} \not\equiv 1 \pmod{p^2}$ for the fundamental unit ε of F . Primes with $\varepsilon^{p-1} \equiv 1 \pmod{p^2}$ (**Wall-Sun-Sun primes**) are conjectured to be infinitely many (but perhaps zero density) by analytic number theorists (no example up to 9.7×10^{17} when $F = \mathbb{Q}[\sqrt{5}]$, but some for $F \neq \mathbb{Q}[\sqrt{5}]$).