

Viscosity solutions for a model of contact line motion

Karl Glasner and Inwon C. Kim

August 21, 2007

Abstract

In this paper we consider a free boundary problem which is used to describe the motion of contact lines of a liquid droplet on a flat surface. The elliptic nature of the equation for droplet shape and the monotonic dependence of contact line velocity on contact angle allows us to introduce a notion of “viscosity” solutions for this problem. Unlike similar free boundary problems, a comparison principle is only available for a modified short-time approximation because of the constraint that conserves volume. We use this modified problem to construct viscosity solutions to the original problem under a weak geometric restriction on the free boundary shape. We also prove uniqueness provided there is an upper bound on front velocity.

1 Introduction

This paper is concerned with solutions of the free boundary problem in $\mathbb{R}^N \times [0, \infty)$

$$(P) \quad \left\{ \begin{array}{l} -\Delta u(\cdot, t) = \lambda(t; u) \text{ in } \{u(\cdot, t) > 0\} \\ \int_{\{u(\cdot, t) > 0\}} u(x, t) dx = V_0 \\ V = F(|Du|) \text{ on } \partial\{u > 0\} \\ u(\cdot, 0) = u_0 \end{array} \right.$$

Here $V = V(x, t)$ denotes the outward normal velocity of the *free boundary* $\partial\{u > 0\}$ of u at (x, t) . In spatial dimension $N = 2$, this problem describes the motion of a liquid droplet of height $u(x, t)$ and volume V_0 on a planar

surface [Gr, Ho, G1]. In this context the *positive phase* $\{u > 0\}$ denotes the wet region and the free boundary denotes the contact line between the drop and the surface. The first equation in (P) defines the shape of a quasi-static droplet. The second equation is a volume conservation condition which is enforced by a suitable choice of the Lagrange multiplier $\lambda(t; u)$ (which is physically the hydrostatic pressure). The third equation in (P) defines the contact line motion by a relationship between the free boundary normal velocity $V = u_t/|Du|$ and the “apparent” contact angle $|Du|$.

The initial condition for the evolution is specified by an open, bounded set $\Omega_0 \subset \mathbb{R}^N$ and volume V_0 . The initial droplet shape $u_0(x) : \mathbb{R}^N \rightarrow \mathbb{R}^+$ is the smallest (weak) solution of

$$\begin{cases} -\Delta u_0 = \lambda_0 > 0 \text{ in } \Omega_0 \\ u_0 = 0 \text{ in } \mathbb{R}^N - \bar{\Omega}_0 \end{cases} \quad (1)$$

where by linearity $\lambda_0 > 0$ can be chosen to satisfy any volume constraint. Also, given $\lambda_0 > 0$, u_0 is uniquely determined by

$$u_0 := \inf\{v : -\Delta v \geq \lambda_0 \text{ in } \Omega_0, v \geq 0 \text{ in } \mathbb{R}^N\}.$$

Many formulas for the constitutive velocity relation F appear in the literature (e.g. [T]). The present paper focuses on the most widely used one [V, C])

$$F(|Du|) = |Du|^3 - 1. \quad (2)$$

The techniques which we use, however, only rely on the fact that F is continuous and strictly increasing.

The free boundary problem (P) has been used as a fundamental model for contact line motion for the last 30 years. Mathematical understanding of this problem has been slowly accruing in the form of numerical methods [G1, Hu], stability calculations [Ho] and homogenized dynamics [G2]. On the other hand, very little is known for (P) in terms of rigorous analysis. To the best of our knowledge the short-time existence of classical solutions has not been established. Furthermore, no notion of weak or generalized solutions has yet been put forth. There are, however, compelling reasons to consider non-classical solutions to this free boundary problem. Numerical (and even physical) experiments indicate that the free boundary evolution with initially convex positive phase develops corners (see Figure 1). Of course, other more standard topological singularities of the positive phase, such as splitting and reconnection, are possible as well (in fact we demonstrate this




Figure 1: Development of a non-smooth corner in the free boundary, using the numerical method in [G1].

must happen for certain initial data, see Lemma 6.10). Our results address only the former type of singularity. There is a good reason for this: during, for example, splitting of the free boundary, the model itself breaks down since separate volume constraints for each connected component would be required. While there may be a more general model that admits changes in topology, we do not address this here.

Originally invented by Crandall and Lions [CL] for Hamilton-Jacobi equations, viscosity solutions allow for singularities of their level sets, and enjoy strong stability properties under various limits. The notion of viscosity solutions has been applied to a variety of free boundary problems that satisfy a comparison principle, which states that if one solution is smaller than the other at one time, then the order is preserved for later times (see, e.g., [K2]). For example, in [K1] a notion of viscosity solutions were introduced for Hele-Shaw and Stefan problems with zero surface tension.

In this paper we define a notion of “viscosity” solutions for problem (P) , and we show that it is well-posed in the sense of existence and uniqueness of solutions. Furthermore, under a moderate geometric restriction (see condition (I) in section 3.3, (J) in section 5, and Appendix B), solutions will exist for all time. In our case, solutions of (P) do not satisfy a comparison principle directly due to the volume constraint that gives rise to a dynamic Lagrange multiplier $\lambda(t; u)$, and thus a straightforward definition of viscosity solutions is more difficult. In particular, the comparison principle we employ only holds for discrete time intervals of an approximating problem (P_n^M) (see section 3) which relaxes the constraint and fixes λ over small time intervals.

The paper is organized as follows. In section 2 we define viscosity solutions for problem (P) and a modified problem (P^M) which puts an upper bound on the free boundary velocity. We also outline the strategy for con-

structing solutions of (P) by approximating problems (P_n^M) . In section 3 the small-time-approximation problem (\tilde{P}) is defined, and a comparison principle and existence theorem for this problem is given. We also introduce a geometric restriction (I) and discuss settings for which it is satisfied to yield global-in-time existence. In section 4 we use the results in section 3 to show existence of a weak solution for (P_n^M) , and derive regularity properties for u_n^M . In section 5 we use the equicontinuity of $\{u_n^M\}$ to show that $\{u_n^M\}$ converges to viscosity solutions as $n, M \rightarrow \infty$. In section 6 we prove that u^M can be obtained as the local uniform limit of the whole sequence $\{u_n^M\}$ as $n \rightarrow \infty$, and the solution u^M of (P^M) is unique. In Appendix A we prove the comparison principle and the existence result for solutions of (\tilde{P}) stated in section 3. Finally in Appendix B we show that (I) holds for all times when the initial data is (a) symmetric with respect to two axes or (b) symmetric with respect to one axes and convex in two dimensions.

2 Definitions and preliminaries

Consider a domain $D \subset \mathbb{R}^N$ and a time interval $I \subset \mathbb{R}^+$. For a nonnegative real valued function $u(x, t)$ defined for $(x, t) \in D \times I$, we will use the notation

$$\Omega(u) = \{(x, t) \in D \times I : u(x, t) > 0\}, \quad \Omega_t(u) = \{x \in D : u(x, t) > 0\};$$

$$\Gamma(u) = \partial\Omega(u) - \partial(D \times I), \quad \Gamma_t(u) = \partial\Omega_t(u) - \partial D.$$

We call $\Omega(u)$ and $\Gamma(u)$ respectively the *positive phase* and the *free boundary* of u .

For $x \in \mathbb{R}^N$ we also denote $B_r(x)$ as the ball of radius r with center x in \mathbb{R}^N .

2.1 Viscosity solutions

We first define the notion of viscosity solutions for problem (P) , with open, bounded initial positive phase Ω_0 and initial shape given by (1).

Definition 2.1. *A nonnegative function $u(x, t)$ in $Q := \mathbb{R}^N \times [0, \infty)$ is a viscosity solution of (P) in Q with initial positive phase Ω_0 and volume V_0 if the following is true:*

1. *u is continuous with $u(\cdot, 0) = u_0(x)$.*

2. At each $t > 0$ $-\Delta u = \lambda(t; u)$ in $\Omega_t(u)$ where $\lambda(t) := \lambda(t; u)$ is chosen such that

$$\int u(x, t) dx = \int u_0(x) dx = V_0.$$

3. For every $\phi \in C^{2,1}(\Sigma)$ such that $u - \phi$ has a local maximum in $\overline{\Omega(u)} \cap \{t \leq t_0\} \cap \Sigma$ at (x_0, t_0) and with $|D\phi|(x_0, t_0) \neq 0$, then

$$\left(\phi_t - |D\phi|(|D\phi|^3 - 1) \right)(x_0, t_0) \leq 0.$$

4. For every $\phi \in C^{2,1}(\Sigma)$ such that $u - \phi$ has a local minimum in $\overline{\Omega(u)} \cap \{t \leq t_0\} \cap \Sigma$ at (x_0, t_0) and with $|D\phi|(x_0, t_0) \neq 0$, then

$$\left(\phi_t - |D\phi|(|D\phi|^3 - 1) \right)(x_0, t_0) \geq 0.$$

Note that classical solutions of (P) are also viscosity solutions.

One can similarly define viscosity solutions of a problem which has an imposed upper bound on velocity

$$(P)^M \quad \begin{cases} -\Delta u(\cdot, t) = \lambda(t; u) & \text{in } \{u > 0\} \\ \int_{\{u > 0\}} u \, dx = V_0 \\ V = \frac{u_t}{|Du|} = \min(|Du|^3 - 1, M) & \text{on } \partial\{u > 0\} \end{cases}$$

Since the upper bound is arbitrary, there is no loss of generality in the physical problem where one expects finite speeds. This modification considerably simplifies our analysis of proving uniqueness in section 6.

Note that (P) does not satisfy a comparison principle: since $\Omega_s \subset \Omega_t$ implies $\lambda(t) \leq \lambda(s)$, then one cannot use the maximum principle to conclude that $u(x, t) \leq u(x, s)$. Therefore the usual viscosity solution approach must be modified. To do this, we consider the “discrete time approximation” problem

$$(P)_n^M \quad \begin{cases} -\Delta u_n^M(\cdot, t) = \lambda_{n,M}(kt_n) & \text{in } \{u_n^M > 0\} \cap [kt_n, (k+1)t_n) \\ V = \min(F(|Du_n^M|), M) & \text{on } \partial\{u_n^M > 0\}, \\ u^M(x, 0) = u_0(x) \end{cases}$$

where $t_n = 2^{-n}$ and $\lambda_{n,M}(kt_n)$ is chosen so that

$$(2.1) \quad \int u_n^M(\cdot, kt_n) dx = V_0 \text{ for } k = 0, 1, 2, \dots,$$

This problem will satisfy the desired comparison principle in each time interval $[kt_n, (k+1)t_n)$. Note that if $\Gamma_t(u_n)$ and u_n^M change continuously in time, then by (2.1)

$$(2.2) \quad u_n^M(x, kt_n) = \frac{\lambda_{n,M}(kt_n)}{\lambda_{n,M}((k-1)t_n)} \lim_{\{t \uparrow kt_n\}} u_n^M(x, t)$$

To construct a viscosity solution of (P) , we first construct the solution u_n^M of $(P)_n^M$ by finding a viscosity solution in $[kt_n, (k+1)t_n)$ on each interval and restart at $t = (k+1)t_n$ using (2.2). We will then show that u_n^M and $\Omega_t(u_n^M)$ converge uniformly as n, M go to infinity to a viscosity solution of the original problem.

3 The small-time problem and a comparison principle

Let $F(s) : [0, \infty) \rightarrow \mathbb{R}$ be a continuous, increasing function of s .

As a small-time approximation of (P) , we consider

$$(\tilde{P}) \quad \begin{cases} -\Delta u(\cdot, t) = \lambda \text{ in } \{u > 0\} \\ V = \frac{u_t}{|Du|} = F(|Du|) \text{ on } \partial\{u > 0\} \end{cases}$$

where λ is a prescribed constant, rather than determined by an additional constraint. For purposes of this section only, we allow $F(s) : [0, \infty) \rightarrow \mathbb{R}$ to be any continuous, monotonically increasing function. In particular, if F is replaced with $\min(F, M)$, then a solution u_n^M of $(P)_n^M$ will also solve (\tilde{P}) with $\lambda = \lambda(kt_n)$ on intervals $[kt_n, (k+1)t_n)$.

Let $Q = \mathbb{R}^N \times (0, \infty)$ and let Σ be a cylindrical domain $D \times (a, b) \subset \mathbb{R}^N \times \mathbb{R}$, where D is an open subset of \mathbb{R}^N .

Definition 3.1. A nonnegative upper semi-continuous function u defined in Σ is a viscosity subsolution of (\tilde{P}) if

(a) for each $a < T < b$ the set $\overline{\Omega(u)} \cap \{t \leq T\}$ is bounded; and

(b) for every $\phi \in C^{2,1}(\Sigma)$ such that $u - \phi$ has a local maximum in $\overline{\Omega(u)} \cap \{t \leq t_0\} \cap \Sigma$ at (x_0, t_0) ,

$$(i) \quad -\Delta\phi(x_0, t_0) \leq \lambda \quad \text{when } u(x_0, t_0) > 0.$$

$$(ii) \quad \left(\phi_t - |D\phi|F(|D\phi|) \right)(x_0, t_0) \leq 0 \quad \text{if } (x_0, t_0) \in \Gamma(u) \text{ when } -\Delta\phi(x_0, t_0) > \lambda.$$

Note that because u is only upper semi-continuous there may be points of $\Gamma(u)$ at which u is positive.

Definition 3.2. A nonnegative lower semi-continuous function v defined in Σ is a viscosity supersolution of (\tilde{P}) if for every $\phi \in C^{2,1}(\Sigma)$ such that $v - \phi$ has a local minimum in $\Sigma \cap \{t \leq t_0\}$ at (x_0, t_0) ,

$$(i) \quad -\Delta\phi(x_0, t_0) \geq \lambda \quad \text{if } v(x_0, t_0) > 0,$$

$$(ii) \quad \text{If } (x_0, t_0) \in \Gamma(v), |D\phi|(x_0, t_0) \neq 0 \text{ and } -\Delta\phi(x_0, t_0) < \lambda,$$

then

$$\left(\phi_t - |D\phi|F(|D\phi|) \right)(x_0, t_0) \geq 0.$$

For a nonnegative real valued function $f(x, t)$ in a cylindrical domain $D \times (a, b)$ we define

$$f^*(x, t) := \limsup_{(\xi, s) \in D \times (a, b) \rightarrow (x, t)} f(\xi, s).$$

and

$$f_*(x, t) := \liminf_{(\xi, s) \in D \times (a, b) \rightarrow (x, t)} f(\xi, s).$$

Definition 3.3. A lower semicontinuous function u is a viscosity solution of (\tilde{P}) if u is a viscosity supersolution and u^* is a viscosity subsolution of (\tilde{P}) . u is a viscosity solution of (\tilde{P}) with initial positive phase Ω_0 if $\Omega_0(u^*) = \Omega_0(u) = \Omega_0$.

For later use we show that free boundaries of solutions of (\tilde{P}) do not “jump” at any positive time.

Lemma 3.4. Let v solve (\tilde{P}) in Q . Then for $x_0 \in \Gamma_{t_0}(v)$ with $t_0 > 0$, there exists $x_n \in \Gamma_{t_n}(v)$, $t_n < t_0$ such that $x_n \rightarrow x_0$ and $t_n \rightarrow t_0$ as $n \rightarrow \infty$.

Proof. 1. Suppose not. Then there exists $r > 0$ and a sequence of t_k converging to t_0 such that for large k , either (i) $v(\cdot, t_k) = 0$ in $B_r(x_0)$ or (ii) $B_r(x_0) \subset \Omega_{t_k}(v)$.

2. If (i) holds, we construct a barrier function $\phi(x, t)$ in $\Sigma := \mathbb{R}^N \times [t_k, t_0]$ such that

$$\begin{cases} -\Delta\phi(\cdot, t) = \lambda \text{ in } \mathbb{R}^N/B_{r(t)}(x_0) \\ \phi(\cdot, t) = 0 \text{ on } \partial B_{r(t)}(x_0), \end{cases}$$

where $r(t) = r + (t_0 - t)/(2(t_0 - t_k))$. (Note that $r(t)$ is positive for $t_k \leq t \leq t_0$.) If t_k is sufficiently close to t_0 , ϕ is a supersolution of (\tilde{P}) in Σ with $\phi \in C^{2,1}(\bar{\Omega}(\phi))$ and with smooth positive phase. Using Definition 3.1, one can check that $v \leq \phi$ in Σ and in particular x_0 lies in the interior of the zero set of $v(\cdot, t_0)$, a contradiction.

3. If (ii) holds, we construct the barrier $\varphi(x, t)$ in Σ such that

$$-\Delta\varphi(\cdot, t) = \lambda \text{ in } B_{r(t)}(x_0), \quad \varphi(\cdot, t) = 0 \text{ in } \mathbb{R}^N/B_{r(t)}(x_0)$$

where $r(t)$ is given above. If t_k is sufficiently close to t_0 , φ is a subsolution of (\tilde{P}) in Σ with $\varphi \in C^{2,1}(\bar{\Omega}(\varphi))$ and with smooth positive phase. Hence using Definition 3.2, one can check that $v \geq \varphi$ in Σ and in particular x_0 lies in the interior of $\Omega_{t_0}(v)$, a contradiction. \square

3.1 Convolutions

An important tool for the analysis in the rest of the paper is the inf- and sup-convolution over space balls. These are employed to obtain larger or smaller sub- and supersolutions, respectively, from existing sub- and supersolutions.

Lemma 3.5. (a) *If u is a viscosity subsolution of (\tilde{P}) , then the sup-convolution*

$$\tilde{u}(x, t) := \sup_{y \in B_{r-ct}(x)} u(y, t)$$

is a viscosity subsolution of (\tilde{P}) with $F(|Du|)$ replaced by $F(|Du|) - c$, as long as $r - ct > 0$.

(b) *If u is a supersolution of (\tilde{P}) then the inf-convolution*

$$\tilde{u}(x, t) = \inf_{y \in B_{r-ct}(x)} u(y, t)$$

is a viscosity supersolution of (\tilde{P}) with $F(|Du|)$ replaced by $F(|Du|) + c$, as long as $r - ct > 0$.

Proof. We only prove (a).

1. First suppose $\tilde{u}(\cdot, t) - \phi(\cdot, t)$ has a local maximum at $x_0 \in \Omega_t(\tilde{u})$. By definition of \tilde{u} , $u(\cdot, t) - \phi(\cdot + (x_0 - y_0), t)$ has a local maximum at y_0 , where $y_0 \in B_{r-ct}(x_0)$ and $\tilde{u}(x_0, t) = u(y_0, t)$. Since u is a viscosity solution of (\tilde{P}) , it follows that $-\Delta\phi(x_0, t) \leq \lambda$. Hence our claim is proved.

2. Next suppose that $\tilde{u}(\cdot, t) - \phi(\cdot, t)$ has a local maximum zero in $\bar{\Omega}(\tilde{u}) \cap \{t \leq t_0\}$ at $(x_0, t_0) \in \Gamma(\tilde{u})$ with $|D\phi|(x_0, t_0) \neq 0$. By definition of \tilde{u} , $\tilde{u} - \tilde{\phi}$, where

$$\tilde{\phi}(x, t) := \phi(x + (1 - c|x_0 - y_0|^{-1}(t - t_0))(x_0 - y_0), t),$$

has a local maximum in $\bar{\Omega}(u) \cap \{t \leq t_0\}$ at $(y_0, t_0) \in \Gamma(u)$, where $|y_0 - x_0| = r - ct_0$ and $\tilde{u}(x_0, t) = u(y_0, t)$. Note that $B_{r-ct_0}(y_0)$ lies in $\Omega_{t_0}(\tilde{u})$ and touches $\Gamma_{t_0}(\tilde{u})$ at x_0 . Since \tilde{u} touches ϕ from below, it follows that $y_0 - x_0$ is parallel to the direction of $D\phi(x_0, t_0)$. Since u is a viscosity solution of (\tilde{P}) , it follows that

$$\begin{aligned} \frac{\tilde{\phi}_t}{|D\tilde{\phi}|}(y_0, t_0) &= \frac{\phi_t}{|D\phi|}(x_0, t_0) + c \\ &\leq F(|D\tilde{\phi}|)(y_0, t_0) = F(|D\phi|)(x_0, t_0). \end{aligned}$$

□

3.2 Comparison Principle

Here we state the comparison principle for viscosity solutions of (\tilde{P}) .

Definition 3.6. We say that a pair of functions $u_0, v_0 : \bar{D} \rightarrow [0, \infty)$ are (strictly) separated (denoted by $u_0 \prec v_0$) in D if

(i) the support of u_0 , $\text{supp}(u_0) = \overline{\{u_0 > 0\}}$ restricted in \bar{D} is compact and

(ii) The functions are strictly ordered in the support of u_0 :

$$u_0(x) < v_0(x) \text{ in } \{u_0 \geq 0\}.$$

Variations of the following theorem, whose proof is deferred to appendix A, will be used later in the paper.

Theorem 3.7. (Comparison principle) Let u, v be respectively viscosity sub- and supersolutions of (\tilde{P}) in $\Sigma = D \times (a, b)$ with $u(\cdot, 0) \prec v(\cdot, 0)$ in D . If $u(\cdot, t) \prec v(\cdot, t)$ on ∂D for $a < t < b$, then $u(\cdot, t) \prec v(\cdot, t)$ in D for $t \in [0, T)$.

3.3 A geometric restriction and global existence

As discussed in the introduction, one cannot expect viscosity solutions to exist for all time in every circumstance. This fact will be encoded into a restriction on the shape of the positive phase, which is the following: We say that a domain $\Omega \in \mathbb{R}^N$ is *star-shaped* with respect to a point $p_0 \in \Omega$ if the line segments connecting p_0 to boundary points $q \in \partial\Omega$ lie in Ω .

The following theorem, whose proof is deferred to the appendix, establishes existence for small times of star-shaped solutions to problem (\tilde{P}) . We will later prove short-time existence for the full problem as well.

Theorem 3.8. *There exists a viscosity solution of (\tilde{P}) in Q with initial positive phase Ω_0 if Ω_0 is star-shaped with respect to $B_r(0)$ for some $r > 0$.*

For long time existence for the full problem (P) , we need to ensure that star-shapedness is preserved. Below we prove this is true provided the free boundary does not collapse in on the "center"; that is, there must always be some ball in Ω_t so that Ω_t is star-shaped with respect to points in that ball. In particular, this allows us to side-step issues involved with topological changes, such as when the free boundary pinches off. The precise requirement is the following: there exists $r > 0$ so that solutions $v(x, t)$ satisfy

$$(I) \quad \begin{cases} \Omega_0(v) \text{ is star-shaped with respect to } B_r(0) \text{ and} \\ B_r(0) \subset \Omega_t(v). \end{cases}$$

We will in general invoke requirement (I) when referring to approximating solutions u_n^M (see Definition 4.1), but we could just as well suppose that (I) holds for the limits as $n, M \rightarrow \infty$, that is, it holds for viscosity solutions to the full problem. We briefly detail some natural cases where (I) is expected to hold for u_n^M :

1. If the free boundary always spreads, then $B_r(0)$ will always be in $\Omega_t(u_n^M)$, and therefore the free boundary will always be star-shaped. Conversely, a contracting free boundary would still satisfy (I) with possibly different r up to the point at which $B_r(0)$ was entirely outside the positive phase.
2. A convex positive phase is star-shaped with respect to every ball, and therefore remains that way if it is contracting. In other words, convexity is preserved for strictly contracting free boundaries. We also suspect, but cannot prove, that this is the case for expanding free boundaries.

3. If the initial data has certain symmetries, (I) is guaranteed for all times. Details of this are given in appendix B.
4. Since there is a lower bound on the free boundary velocity, $B_r(0)$ will at least stay inside Ω_t for a short time. For short-time existence, we can therefore always assume (I) holds, so long as the initial data is star-shaped.

We will now prove that star-shapedness is preserved as long as (I) holds in problem (\tilde{P}) , and later observe the same is true for the full problem. Therefore if (I) is preserved by the evolution, we will be able to obtain global existence and uniqueness.

Lemma 3.9. *Suppose that v solves (\tilde{P}) with $F(|Dv|) = \min(|Dv|^3 - 1, M)$ and condition (I) is satisfied. Then $\Omega_t(v)$ is star-shaped for all $t > 0$.*

Proof. 1. Let $x_0 \in B_r(0)$. We claim that for all x ,

$$(3.1) \quad v(x, t) \leq (1 + \epsilon)^2 v\left(\frac{x - x_0}{(1 + \epsilon)} + x_0, t\right) \text{ for any } \epsilon > 0.$$

For $t \in [0, c/(2M + 2)]$ define

$$(3.2) \quad \tilde{v}(x, t) = \inf_{y \in B_{c\epsilon - (2M+2)\epsilon t}(x)} (1 + \epsilon)^2 v\left(\frac{y - x_0}{(1 + \epsilon)} + x_0, t\right)$$

where c (which only depends on r), is chosen small enough so that $v \prec \tilde{v}$ at $t = 0$.

Notice this is just a inf-convolution of a rescaled version of v , which is easily checked to be a supersolution, so Lemma 3.5 applies. Therefore,

$$\begin{aligned} \frac{\tilde{v}_t}{|D\tilde{v}|} &= (1 + \epsilon) \min(|Dv|^3 - 1, M] + (2M + 2)\epsilon \\ &\geq (1 + \epsilon) \min((1 - 3\epsilon)|D\tilde{v}|^3 - 1, M] + (2M + 2)\epsilon \\ &> \min(|D\tilde{v}|^3 - 1, M] \end{aligned}$$

Moreover $-\Delta\tilde{v}(\cdot, t) \geq \lambda$ in $\Omega_t(\tilde{v})$ due to Lemma 3.5. Hence \tilde{v} is a supersolution of (\tilde{P}) . Now Theorem 3.7 applies to v and \tilde{v} in $\mathbb{R}^N \times [0, c/(2M + 2)]$ to yield $v \leq \tilde{v}$ for $0 \leq t \leq t_1 := c/(2M + 2)$ which yields (3.1). Since $\epsilon > 0$ in (3.1) is arbitrary, it follows that $v(\cdot, t + t_1)$ satisfies (I). Hence one can repeat this process indefinitely on intervals of length $\frac{c}{2M+2}$. \square

4 Construction of u_n^M

Our next goal is to construct solutions u_n^M of the approximating problem (P_n^M) with star-shaped initial positive phase Ω_0 and initial volume V_0 , under condition (I). By definition of (P_n^M) , u_n^M is in general discontinuous in time at the endpoints of time intervals $I_k := [kt_n, (k+1)t_n)$, and as mentioned before the comparison principle only holds for u_n^M in small time intervals I_k and thus a conventional notion of viscosity solutions will not apply. It is therefore necessary to first establish a weak notion of solutions for (P_n^M) .

Definition 4.1. u_n^M is a weak solution of (P_n^M) with initial positive phase Ω_0 and volume V_0 for $0 \leq t \leq (l+1)t_n$ where $l \in \mathbb{N}$ if the following holds for $k = 0, 1, \dots, l$:

- (i) $u_n^M(\cdot, 0) = u_0$
- (ii) $u_n^M(\cdot, t + kt_n)$ is a viscosity solution of (\tilde{P}) in $(0, t_n]$ with initial positive phase $\Omega_{kt_n}(u_n^M)$ and $\lambda = \lambda_{n,M}(kt_n)$, where $\lambda_{n,M}(kt_n)$ satisfies (2.1).
- (iii) $\Omega_{kt_n}(u_n^M)$ is continuous from below, that is:

$$d(\Omega_t(u_n^M), \Omega_{kt_n}(u_n^M)) \rightarrow 0 \text{ for } t \uparrow kt_n.$$

Note that, due to (I) and Lemma 3.9, u_n^M has its positive phase star-shaped in space with respect to $B_r(0)$ as long as it exists. It follows that the family of domains $\{\Omega_t(u_n^M)\}_{n,t}$ is uniformly Lipschitz in space if they are uniformly bounded (see Remark below Lemma 4.4). Using this fact, in Proposition 4.5 we will show that for weak solutions u_n^M satisfying (I), $\Omega(u_n^M)$ is uniformly Hölder continuous in time. This establishes equi-continuity for the family of functions $\{u_n^M\}$ to obtain convergence to a solution of $(P)^M$ in section 5.

First we give an upper bound for $\lambda(kt_n)$ in terms of the circumradius of $\Omega_{kt_n}(u_n^M)$.

Lemma 4.2. Suppose u_n^M exists and satisfies (I) with $r > 0$ for $0 \leq t \leq kt_n$. Then for $x_0 \in \Gamma_{kt_n}(u_n^M)$, $\lambda(kt_n) \leq \frac{C_0}{|x_0|}$, where C_0 only depend on r, V_0 and N .

Proof. Due to Lemma 3.9, u_n^M is star-shaped with respect to $B_r(0)$. Therefore $\Omega_{kt_n}(u_n^M)$ contains a cone with vertex x_0 , axis parallel to x_0 and bottom $B_r(0) \cap \{x \cdot x_0 = 0\}$. It follows that the function $f(x)$ solving

$$-\Delta f = 1 \text{ in } \Omega_{kt_n}(u_n^M), \quad f = 0 \text{ on } \Gamma_{kt_n}(u_n^M)$$

is bigger than the superpositions of $h(x - kr\frac{x_0}{|x_0|})$, $k = 1, \dots, |x_0|/2r$, where h solves $-\Delta h = 1$ in $B_{r/2}(0)$ with $h = 0$ on $\partial B_{r/2}(0)$. Thus

$$\int f(x)dx \geq \frac{|x_0|}{2r} \int h(x)dx \geq C|x_0|r^{N+2}.$$

Multiplying by $\lambda(kt_n)$ and noting the definition of V_0 , we can obtain the desired bound. \square

Lemma 4.3. *Suppose u_n^M satisfies (I), with $r > 0$ independent of n and M , for $0 \leq t \leq T$. Define $R(t, n, M) := \sup\{|x| : x \in \Omega_t(u_n^M)\}$. Then $R(t, n, M) \leq R(T)$ for $t < T$, where $R(T)$ is independent of n, M .*

Proof. Let f solve $-\Delta f = 2C_0$ in $B_1(0)$ with $f = 0$ on $\partial B_1(0)$, where C_0 is as given in Lemma 4.2. Let A be the value of $|Df|^3$ on $\partial B_1(0)$ (Note that f is radially symmetric). Next define

$$h(x, t) := R(t)f\left(\frac{x}{R(t)}\right) \text{ with } R(t) = R(0) + At,$$

where $R(0)$ is big enough such that $B_{R(0)}(0)$ contains $\bar{\Omega}_0$. Note that

$$-\Delta h = \frac{2C_0}{R(t)} \text{ in } B_{R(t)}(0)$$

with $h = 0$ in $\partial B_{R(t)}(0)$ and

$$(4.2) \quad \frac{h_t}{|Dh|} = R'(t) \geq |Dh|^3 > |Dh|^3 - 1 \text{ on } \Gamma_t(h).$$

We claim that $\Omega_t(u_n^M)$ is always strictly contained in $B_{R(t)}(0)$. To see this, suppose not. Then due to the definition of u_n^M and Lemma 3.4, $\Gamma_t(u_n^M)$ intersects $\partial B_{R(t)}(0)$ from inside of the ball for the first time at $t = t_0 \in (kt_n, (k+1)t_n]$. Choose the smallest ball $B_R(0)$ containing $\Omega_{kt_n}(u_n^M)$. If $R(kt_n)/2 \leq R \leq R(kt_n)$, then $t_0 = kt_n$ and by Lemma 4.2, $\lambda(kt_n) \leq C_0/R(t_0)$ and thus $u_n^M(\cdot, t_0) \leq h(\cdot, t_0)$. This and (4.2) yields that h is a supersolution of (\tilde{P}) with $\lambda = \lambda(kt_n)$ on $(kt_n, (k+1)t_n)$, and Theorem 3.7 leads to a contradiction.

Hence $R < R(kt_n)/2$ and $t \in (kt_n, (k+1)t_n]$.

$$(4.3) \quad -\Delta u_n^M(\cdot, t) = \lambda(kt_n) \leq C_0/R \text{ for } t \in (kt_n, (k+1)t_n].$$

Again Theorem 3.7 yields that $u_n^M \leq \tilde{h}$ in $\mathbb{R}^N \times (kt_n, (k+1)t_n]$, where $\tilde{h}(x, t) = (R + At)f(x/(R + At))$. Since $\tilde{h}(\cdot, t) \prec h(\cdot, t)$ on $[kt_n, (k+1)t_n]$ if $At_n \leq R(kt_n)/2$, we obtain a contradiction for sufficiently small n . \square

Remark Lemma 4.3 and the star-shapedness of $\Omega_t(u_n^M)$ yield that for each $t > 0$, $\Omega_t(u_n^M)$ is a Lipschitz domain, whose Lipschitz constant is uniformly bounded for $0 \leq t \leq T$ independently of n and M . This yields the following proposition:

Proposition 4.4. *Suppose u_n^M satisfies (I) for $0 \leq t \leq T$ with $r = r(T) > 0$. Then the distance function $d_n^M(\cdot, t)$ to $\Gamma(u_n^M)$ is locally uniformly Hölder continuous in time for $0 \leq t \leq T$, independently of M and n .*

Proof. 1. Due to the previous lemma, $\Omega_t(u_n^M) \subset B_{R(t)}(0)$. Moreover for $x_0 \in \Gamma_{t_0}(u_n^M)$ with $t_0 \in (kt_n, (k+1)t_n]$, there is a cone that touches the free boundary on which u_n^M is zero:

$$u_n^M(\cdot, t_0) = 0 \text{ in } \mathcal{C} := \{y : \frac{(y - x_0)}{|y - x_0|} \cdot \frac{x_0}{|x_0|} \geq \frac{|x_0|}{\sqrt{r^2 + |x_0|^2}}\}.$$

2. Let $B_R(0)$ be the smallest ball which contains $\Omega_{kt_n}(u_n^M)$. Due to Lemma 4.2, $\lambda(kt_n) \leq C_0/R$. Moreover arguing as in the proof of Lemma 4.3 it follows that $\Omega_t(u_n^M) \subset B_{R+At}(0)$ for $t > kt_n$, where A only depends on r, V_0 and N .

Fix $0 < m \ll 1$ and let g solve

$$-\Delta g = C_0/(R-1) \text{ in } B_{R+1}(0) - (\mathcal{C} + mx_0),$$

with $g = 0$ on the boundary. Then for any $r > 0$

$$(4.4) \quad \sup_{x \in B_r((1+m)x_0)} g(x) \leq C_0 r^\alpha,$$

where $C_0 = C_0(N)$ and $0 < \alpha < 1$ only depends on $R(t), r$ and N and thus independent of n and M . Note that, since $\Omega_t(u_n^M)$ is star-shaped and the normal velocity of $\Gamma(u_n^M)$ is bigger than -1 , $\Omega_t(u_n^M)$ does not shrink more than distance $1/A$ from $\Omega_{kt_n}(u_n^M)$ by $t = kt_n + 1/A$. This and Lemma 4.2 yields that, if we choose $A > 1$,

$$-\Delta u_n^M(\cdot, t) \leq C_0/(R-1) \text{ for } kt_n \leq t \leq kt_n + 1/A.$$

Due to (4.3), it then follows that

$$(4.5) \quad u_n^M(x, t) \leq g(x) \text{ for } kt_n \leq t \leq kt_n + 1/A$$

as long as $u_n^M((1+m)x_0, t) = 0$.

3. Next we construct a barrier $\phi(x, t)$ of the form

$$\begin{cases} -\Delta\phi(\cdot, t) = C_0/R & \text{in } (B_{2m} - B_{r(t)})((1+m)x_0) \\ \phi(\cdot, t) = 0 & \text{on } \partial B_{r(t)}((1+m)x_0) \\ \phi(\cdot, t) = C_0(2m)^\alpha \text{ on } B_{2m}((1+m)x_0) \end{cases}$$

where C_0 is as given in (4.4) and

$$r(t) = m/|x_0| - C_1 m^{3\alpha-3}(t - t_0) \text{ with } C_1 = c(N)C_0^3$$

in the domain

$$\mathcal{S} := (B_{2m} - B_{r(t)})((1+m)x_0) \times [t_0, t_1], \quad t_1 := t_0 + (2|x_0|C_1)^{-1}m^{4-3\alpha}.$$

It then follows that, on $\partial B_{r(t)}((1+m)x_0) \times [t_0, t_1]$, ϕ satisfies

$$\frac{\phi_t}{|D\phi|} \geq |D\phi|^3 > |D\phi|^3 - 1.$$

if $c(N)$ is a sufficiently large dimensional constant. Due to (4.5), Theorem 3.7 applies to u_n^M and ϕ in \mathcal{S} as long as $u_n^M((m+1)x_0, t) = 0$. But $u_n^M((m+1)x_0, t) = 0$ as long as $u_n^M \leq \phi$. Thus we conclude that $u_n^M \leq \phi$ in \mathcal{S} .

4. In particular above argument yields that for any $m > 0$ if $x_0 \in \Gamma_{t_0}(u_n^M)$, then for any $m > 0$, $\Gamma(u_n^M)$ does not reach $(1+m)x_0$ until $t_1 = t_0 + C(r, T, N)m^{4-3\alpha}$. On the other hand a parallel argument, based on the fact $V = |Du_n^M|^3 - 1 \geq -1$, yields that $\Gamma(u_n^M)$ does not reach $(1-m)x_0$ until $t_1 = t_0 - m$. Since $m > 0$ can be chosen arbitrarily small, we can conclude that $\Gamma(u_n^M)$ for $t \leq T$ is Hölder continuous in time with Hölder constant $1/(4-3\alpha)$, where $\alpha = \alpha(r, T, N)$.

5. Let x be arbitrary. For times $t_1 < t_2$, choose $x_2 \in \Gamma_{t_2}(u_n^M)$ so that $|x - x_2| = d(x, t_2)$, and choose x_1 to be the unique point on $\Gamma_{t_1}(u_n^M)$ parallel to x_2 . Using step 4 we have

$$d(x, t_1) \leq |x - x_1| \leq |x_2 - x_1| + |x - x_2| \leq C(r, T, N)|t_2 - t_1|^\alpha + d(x, t_2)$$

so that $d(x, t_1) - d(x, t_2) \leq C(r, T, N)|t_2 - t_1|^\alpha$. We can analogously show $d(x, t_2) - d(x, t_1) \leq C(r, T, N)|t_2 - t_1|^\alpha$, which verifies uniform Hölder continuity. □

We now prove the main result of this section.

Theorem 4.5 (Existence of u_n^M). *Suppose any weak solution u_n^M of (P_n^M) in $\mathbb{R}^n \times [0, t_0]$, $t_0 \leq T$ satisfies (I) for $0 \leq t \leq t_0$, with $r = r(T) > 0$. Then there exists a weak solution u_n^M of (P_n^M) with initial positive phase Ω_0 and volume V_0 for $0 \leq t \leq T$. Moreover $\Omega_t(u_n^M)$ is star-shaped with respect to $B_r(0)$ and $\Gamma_t(u_n^M)$ is locally uniformly Hölder continuous in time, independently of n and M .*

Proof. We use induction on l . Suppose we have constructed u_n^M in $\mathbb{R}^N \times [0, lt_n]$. Due to Proposition 4.4, $\Omega_t(u_n^M)$ uniformly converges to $\Omega_{lt_n}(u_n^M)$ as $t \rightarrow lt_n$. Since $\Omega_{lt_n}(u_n^M)$ is star-shaped, $\lambda(lt_n)$ and $u_n^M(\cdot, lt_n)$ is well-defined and is continuous in space. Due to Theorem 3.8 there exists a viscosity solution u_n^M of (\tilde{P}) with $\lambda = \lambda(lt_n)$ in $(lt_n, (l+1)t_n]$ with initial positive phase $\Omega_{lt_n}(u_n^M)$. Now the induction can be continued to show that u_n^M can be solved for $0 \leq t \leq T$. The rest of the theorem is due to Lemma 3.9 and Proposition 4.4. \square

5 Convergence of u_n^M and existence of u^M and u

In this section we prove the existence of the viscosity solution u of our original problem (P) , by passing to limits in n and M , and verifying the result is a viscosity solution. First we fix M and send $n \rightarrow \infty$. Due to Theorem 4.5, for $0 \leq t \leq T$ the distance function $d_n^M(\cdot, t)$ to the set $\Gamma_t(u_n^M)$ is locally uniformly Lipschitz continuous in space and locally uniformly Hölder continuous in time, independently of n and M . Hence due to Arzela-Ascoli, d_n^M converges locally uniformly to d^M in $\mathbb{R}^n \times [0, T]$ along a subsequence. It then follows that

- (a) $\Omega_t^M := \{d^M(\cdot, t) > 0\}$ is star-shaped with respect to $B_r(0)$,
- (b) $\Gamma_t^M := \{d^M(\cdot, t) = 0\}$, and the limiting distance function $d^M(\cdot, t) = 0$ is locally Lipschitz in space and locally uniformly Hölder continuous in time,
- (c) $\Omega_0^M = \Omega_0$.

Let $u^M(x, t)$ solve

$$-\Delta u^M(\cdot, t) = \lambda(t; u^M) \text{ in } \Omega_t^M.$$

with zero boundary data on Γ_t^M , where $\lambda(t; u^M)$ is the volume preserving constant such that

$$\int u^M(x, t) dx = V_0.$$

Then $u^M(\cdot, 0) = u_0$.

Proposition 5.1. $u^M(x, t)$ is a viscosity solution of $(P)^M$ in $\mathbb{R}^n \times [0, T]$ with initial positive phase Ω_0 and volume V_0 .

Proof. 1. First observe that $\lambda_{n,M}(kt_n)$ converges to $\lambda(t; u^M)$, locally uniformly in time, because $\Omega_t(u_n^M)$ locally uniformly converges to $\Omega_t(u^M)$ and $\Gamma_t(u_n^M)$ is locally uniformly Hölder continuous in time independently of n (in the sense of the corresponding distance function). Therefore u_n^M locally uniformly converges to u^M .

2. Suppose $u^M - \phi$ has a local maximum zero in $\bar{\Omega}(u^M) \cap (B_r(x_0) \times [t_0 - r, t_0])$ for some $r > 0$ at $(x_0, t_0) \in \Gamma(u^M)$ with $\phi \in C^{2,1}(Q)$, $|D\phi|(x_0, t_0) \neq 0$. Assume that, for some $\epsilon > 0$,

$$(5.1) \quad \left(\frac{\phi_t}{|D\phi|} - \min(|D\phi|^3 - 1, M) \right)(x_0, t_0) > \epsilon.$$

We may assume that this maximum is strict in $B_r(x_0) \times [t_0 - r, t_0]$ otherwise one can replace ϕ by $\phi + \epsilon(x - x_0)^4 + \epsilon(t - t_0)^2$ to make it strict. Since ϕ is smooth with $|D\phi|(x_0, t_0) \neq 0$, we may assume that a space-time ball of radius r , $B_r^{N+1}(P_0)$ with $P_0 \in \mathbb{R}^{N+1}$, lies in the zero set of u^M and touches $\Gamma(u^M)$ at (x_0, t_0) (See Figure 2). Moreover due to (6.1) the outward normal vector ν of the ball $B_r^{N+1}(P_0)$ at (x_0, t_0) is given by $\nu = (\nu_1, b) \in \mathbb{R}^N \times \mathbb{R}$, where $|\nu_1| = 1$ and the *slope* b of the ball at (x_0, t_0) satisfies

$$b \geq \min(|D\phi|^3 - 1, M)(x_0, t_0) + \epsilon.$$

Let us tilt and shift the ball so that the new ball \tilde{B}^{N+1} passes through $(x_0 - a\nu_1, t_0)$ with slope $b - \epsilon/2$ for $a \ll \epsilon$. Note that if a is small compared to ϵ and τ ,

$$(5.2) \quad d(\tilde{B}^{N+1} \cap \{t = t_0 - \tau\}, \Omega(u^M)) > O(\epsilon).$$

Now let us choose a, τ, δ such that $a \ll \tau \ll \delta \ll r, \epsilon$ and define $h(x, t)$ in the domain

$$\Sigma := (1 + a)\tilde{B}^{N+1} \cap [t_0 - \tau, t_0]$$




Figure 2: Exterior ball $B_r^{N+1}(P_0)$ at the contact point P_0 .

such that

$$\begin{cases} -\Delta_x h(x, t) = \lambda(t_0) + \delta & \text{in } ((1+a)\tilde{B}^{N+1} - \tilde{B}^{N+1}) \cap [t_0 - \tau, t_0] \\ h(x, t) = \phi(x, t) & \text{on } (1+a)(\partial\tilde{B}^{N+1}) \cap [t_0 - \tau, t_0] \\ h(x, t) = 0 & \text{in } \tilde{B}^{N+1} \cap [t_0 - \tau, t_0]. \end{cases}$$

Since ϕ is smooth with $|D\phi|(x_0, t_0) \neq 0$, if r is chosen small enough (5.1) yields that

$$(5.3) \quad \frac{h_t}{|Dh|} > \min(|Dh|^3 - 1, M) \text{ on } \Gamma(h) \cap \Sigma.$$

Moreover due to (5.2) $u^M \prec h$ on the parabolic boundary of Σ . Since u_n^M and $\Omega(u_n^M)$ locally uniformly converges to u^M and $\Omega(u^M)$, it follows that u_n^M crosses h from below for the first time at (y_n, s_n) in Σ with $s_n \in (kt_n, (k+1)t_n]$ for some $k \in \mathbb{N}$, for sufficiently large n . This contradicts Theorem 3.7 if n is large enough that $\lambda_{n,M}(kt_n) \leq \lambda(t_0) + \delta$.

3. The above arguments prove that u^M is a viscosity subsolution of (P^M) . A parallel argument would similarly prove u^M is a viscosity supersolution of (P^M) . \square

So far we have proved the existence of viscosity solutions of $(P)^M$. By a similar process, we can send $M \rightarrow \infty$ to obtain the most general existence result.

Theorem 5.2. *Suppose u_n^M satisfies (I) for $0 \leq t \leq T$ with $r = r(T)$. Then along a subsequence u^M and $\Omega(u^M)$ locally uniformly converges to u and $\Omega(u)$ in $\mathbb{R}^n \times [0, T]$ as $M \rightarrow \infty$. The limit function u is a viscosity solution of (P) in $\mathbb{R}^n \times [0, T]$ with initial positive phase Ω_0 and volume V_0 .*

Corollary 5.3. *Suppose u_n^M satisfies (I) for $0 \leq t \leq T$ with $r = r(T)$. Then there exists a viscosity solution u of (P) in $\mathbb{R}^N \times [0, T]$ with initial positive phase Ω_0 and volume V_0 . Moreover $\Omega(u)$ is star-shaped in space with respect to $B_r(0)$ and $\Gamma(u)$ is Lipschitz continuous in space and Hölder continuous in time.*

Since $\Omega(u)$ is only Lipschitz, difficulties arise in the analysis due to the lack of upper bound on the free boundary velocity. For this reason we will prove uniqueness result for only the modified problem (P^M) in next section.

6 Uniqueness of u^M

In this section we show that u^M given in Corollary 5.3 is the unique viscosity solution of $(P)^M$. Recall that u_n^M is a weak solution of (P_n^M) with initial positive phase Ω_0 and volume V_0 .

Proposition 6.1. *Suppose u_n^M satisfies (I) for $0 \leq t \leq T$ with $r = r(T)$. Then for $k \geq n$ and for $0 \leq t \leq T$, there exists $A > 0$ depending only on r, δ, M, T and the spatial dimension N such that*

$$A_n(t)^{-2N-2} u_k^M(A_n(t)x, t) \leq u_n^M(x, t) \leq A(t)^{2N+2} u_k^M\left(\frac{x}{A_n(t)}, t\right),$$

where $A_n(t) := 1 + Ae^{At}t_n$.

Proof. 1. For simplicity we set $r = \delta = 1/2$ in (4.1). A parallel argument holds for the general case.

2. Let $A > M > 0$ to be chosen later. For each $t \in [0, \frac{1}{(6N+6)A}]$, we claim that

$$(6.1) \quad (1+A_n)^{-2N-2} u_k^M((1+A_n)x, t) \prec u_n^M(x, t) \prec (1+A_n)^{2N+2} u_k^M((1+A_n)^{-1}x, t)$$

where $A_n = At_n$. At $t = 0$ the inequality is true due to the star-shaped initial data. Suppose the second inequality in (6.1) is violated for the first time at $t = t_0 \in (0, \frac{1}{(6N+6)A}]$.

Due to (6.1),

$$(1 + At_n)^{-1} \Omega_t(u_k^M) \subseteq \Omega_t(u_n^M) \subseteq (1 + At_n) \Omega_t(u_k^M) \text{ for } 0 \leq t \leq t_0.$$

Thus by definition of $\lambda_{n,M}$ and $\lambda_{k,M}$,

$$(1 + At_n)^{-N} \lambda_{n,M}(lt_n) \leq \lambda_{k,M}(lt_n) \leq (1 + At_n)^N \lambda_{n,M}(lt_n)$$

for any $lt_n \leq t_0, l = 0, 1, \dots$

3. We claim that for any $x \in \Omega_t(u_k^M)$

$$(6.2) \quad d(\Omega_{lt_n}(u_k^M), x) \leq At_n \text{ for } lt_n \leq t \leq (l+1)t_n,$$

where A depends on M and the Lipschitz constant L of $\Omega_{lt_n}(u_k^M)$. Note that, due to Lemma 3.9 and Lemma 4.2, $L = L(r, T, N)$.

To verify (6.2), pick any point x_0 such that $d(\Omega_{lt_n}(u_k^M), x_0) = At_n$. Due to the fact that $\Omega_{lt_n}(u_k^M)$ is Lipschitz, if we choose $A = C_0 M$ where C_0 depends only on L , then there is a ball

$$B := B_{2M}(x_0) \subset \{u_k^M(\cdot, lt_n) = 0\}$$

which touches $\Gamma_{lt_n}(u_k^M)$. Let us define

$$\Lambda := \sup_{(l-1)t_n \leq mt_k \leq (l+1)t_n} \lambda_{k,M}(mt_k).$$

Note that $\Lambda \leq r^{-N}$ by (4.1).

Consider $\phi(x, t)$: a nonnegative function in $\Sigma := 3B \times [lt_n, (l+1)t_n]$ such that for $lt_n \leq t \leq (l+1)t_n$

$$\begin{cases} -\Delta \phi(\cdot, t) = \Lambda & \text{in } (3B - (2 - \frac{t}{t_n})B) \\ \phi(\cdot, t) = \sup_{\Sigma} u_k^M & \text{on } \partial(3B) \\ \phi(\cdot, t) = 0 & \text{in } (2 - \frac{t}{t_n})B \end{cases}$$

Then ϕ is a supersolution of (\tilde{P}) in Σ with $\lambda = \Lambda$ and $F(|Du|) = M$. Moreover $u_k^M \prec \phi$ on the parabolic boundary of Σ and $u_k^M(\cdot, t) \leq \phi(\cdot, t)$ as long as $\Omega_t(u_k^M) \cap 3B \subset \Omega_t(\phi)$. Hence Theorem 3.7 applied to u_k^M and ϕ in each time interval $(mt_k, (m+1)t_k) \cap [lt_n, (l+1)t_n]$ would then yield that $u_k^M \leq \phi$ in Σ . In particular x_0 lies outside of $\Omega(u_k^M)$ for $lt_n \leq t \leq (l+1)t_n$. This yields (6.2).

4. Due to (6.2),

$$(6.3) \quad (1 + At_n)^{-2N} \lambda_{k,M}(mt_k) \leq \lambda_{l,M}(lt_n) \leq (1 + At_n)^{2N} \lambda_{k,M}(mt_k)$$

for $mt_k \in [(l-1)t_n, (l+1)t_n] \cap [0, t_0]$, where $m, l = 0, 1, 2, \dots$

Due to (6.3) and Lemma 3.5, one can now check that for $0 \leq t \leq t_0 \leq \frac{1}{(6N+6)A}$,

$$(6.4) \quad \tilde{u}_k^M(x, t) := (1 + At_n)^{2N+2} \inf_{B_{At_n - (6N+6)A^2t_n t}} u_k^M((1 + At_n)^{-1}x, t)$$

satisfies the free boundary motion law

$$\begin{aligned} V &\geq \frac{(1 + At_n)(u_k^M)_t}{|Du_k^M|} + (6N + 6)A^2t_n \\ &= (1 + At_n) \min[|Du_k^M|^3 - 1, M] + (6N + 6)A^2t_n \\ &\geq \min[(1 + At_n)^{6N+1}|Du_k^M|^3 - 1, M] - (6N + 2)At_nM + (6N + 6)A^2t_n \\ &\geq \min[|D\tilde{u}_k^M|^3 - 1, M] \end{aligned}$$

if t_n is sufficiently small. The first inequality is due to Lemma 3.5 and the last inequality holds since $A > M$. (For rigorous argument one needs to use the definition of viscosity solutions of (\tilde{P}) . See for example the proof of Proposition 5.5 in [CJK].)

Observe that due to (6.3) and Lemma 3.5, for any $l, m = 0, 1, \dots$

$$\begin{aligned} -\Delta \tilde{u}_k^M(\cdot, t) &\geq (1 + At_n)^{2N} \lambda_{k,M}(mt_k) \\ &\geq \lambda(lt_n) \text{ for } t \in (mt_k, (m+1)t_k] \cap [lt_n, (l+1)t_n] \end{aligned}$$

Note that $u_n^M(\cdot, 0) \prec \tilde{u}_k^M(\cdot, 0)$ since Ω_0 is star-shaped with respect to zero and contains $B_1(0)$. Thus Theorem 3.7 applied to u_n^M and \tilde{u}_k^M on each time interval $[mt_k, (m+1)t_k]$ gives for $0 \leq t \leq t_0$,

$$u_n^M(\cdot, t) \prec \tilde{u}_k^M(\cdot, t) \leq (1 + At_n)^{2N+2} u_k^M((1 + At_n)^{-1}x, t).$$

This contradicts our hypothesis at $t = t_0$. Similar arguments lead to a contradiction if we assume that the first inequality breaks for the first time at $t_1 \in [0, \frac{1}{(6N+6)A}]$. Thus (6.1) holds for $0 \leq t \leq t_1 := \frac{1}{(6N+6)A}$.

5. Next we show that for $t_1 \leq t \leq t_1(1 + 1/2)$

$$(1 + 2At_n)^{-2N-2} u_k^M((1 + 2At_n)x, t) \leq u_n^M(x, t) \leq (1 + 2At_n)^{2N+2} u_k^M((1 + 2At_n)^{-1}x, t)$$

For example if the second inequality breaks, then we compare u_n^M with

$$\tilde{u}_k^M(x, t) := (1 + 2At_n)^{2N+2} \inf_{B_{At_n - 12(N+1)A^2 t_n t}} u_k^M((1 + 2At_n)^{-1}x, t)$$

using similar arguments as in step 2. Note that due to (6.1) and the fact that $\Omega_t(u_n)$ is star-shaped and contains $B_1(0)$,

$$u_n^M(x, t_1) \prec (1 + At_n)^{2N+2} u_k^M((1 + At_n)^{-1}x, t_1) \leq \tilde{u}_k^M(x, t_1)$$

6. One can repeat the argument for each interval

$$[t_1(1 + 1/2 + \dots + 1/n), t_1(1 + 1/2 + \dots + 1/n + 1)].$$

This proves the lemma since

$$t_1(1 + 1/2 + \dots + 1/n) \sim t_1(\log n).$$

□

Note that the proof presented above can be used as long as one of the functions being compared, u_k^M in above proof, satisfies (I). Thus we obtain the following corollary.

Corollary 6.2. *Suppose u_n^M satisfies (I) for $0 \leq t \leq T$ with $r = r(T)$. Then the whole sequence $\{u_n^M\}$ converges locally uniformly as $n \rightarrow \infty$ to a viscosity solution u^M of $(P)^M$ for $0 \leq t \leq T$ with initial positive phase Ω_0 and volume V_0 .*

Remarks.

1. Besides proving the uniqueness of the limit, Proposition 6.1 provides an estimate on differences between discrete-time approximation solutions u_n^M in terms of the discrete time interval size t_n .
2. Note that we need to keep track of both inequalities in the lemma in each time interval to guarantee that $\lambda_{n,M}(t)$ and $\lambda_{k,M}(t)$ stay close together.

Now let v be any other viscosity solution with initial data u_0 defined in the previous section. Parallel arguments as in the proof of Proposition 6.1 yield the following:

Lemma 6.3. *Suppose u_n^M satisfies (I) for $0 \leq t \leq T$ with $r = r(T)$. Then for the same $A_n(t)$ given in Proposition 6.1,*

$$(1+A_n(t))^{-2N-2}u_n^M((1+A_n(t))x, t) \leq v(x, t) \leq (1+A_n(t))^{2N+2}u_n^M\left(\frac{x}{1+A_n(t)}, t\right)$$

for $0 \leq t \leq T$.

Applying the same argument between two viscosity solutions u^M and v^M of (P^M) (in this case the time step size $t_n > 0$ is replaced by arbitrary small constants in the arguments) yields the following corollary.

Corollary 6.4. *Suppose u_n^M satisfies (I) for $0 \leq t \leq T$ with $r = r(T)$. Then u^M is the unique viscosity solution of $(P)^M$ in $\mathbb{R}^n \times [0, T]$ with initial data u_0 .*

Remark To prove uniqueness results for the original problem (P) , one needs some type of bound on free boundary velocity. At least for star-shaped spreading droplets, we expect solutions of (P) to have smooth positive phase for positive times and locally uniformly bounded free boundary velocity for any positive time interval. Such results have been proved for the Hele-Shaw problem with zero surface tension (see [CJK]).

A Comparison principle and existence for (\tilde{P})

Here we prove Theorem 3.7 stated in section 2 and the existence of the viscosity solutions of (\tilde{P}) with star-shaped initial positive phase Ω_0 .

Most arguments presented here are similar to the proofs of Theorem 2.2 and Theorem 4.7 of [K1]. We only sketch the outline of the proof below.

Sketch of the proof of Theorem 3.7

1. For $r, \delta > 0$ and $0 < h \ll r$, define the sup-convolution of u

$$Z(x, t) := (1 + \delta) \sup_{|(y, s) - (x, t)| < r} u(y, (1 + \delta)^3 s)$$

and the inf-convolution of v

$$W(x, t) := (1 - \delta) \inf_{|(y, s) - (x, t)| < r - ht} v(y, (1 - \delta)^3 s)$$

in the domain

$$\Sigma := \tilde{D} \times [r, r/h], \tilde{D} := \{x : x \in D, d(x, \partial D) \geq r\}.$$




Figure 3: Interior and exterior balls at the contact point P_0 .

By upper semi-continuity of $u - v$, $Z(\cdot, r) \prec W(\cdot, r)$ for sufficiently small $r, \delta > 0$. By our hypothesis and the upper semi-continuity of $u - v$,

$$Z(\cdot, t) \prec W(\cdot, t) \text{ on } \partial\tilde{D} \text{ for } r \leq t \leq r/h$$

for sufficiently small δ and r . Moreover Lemma 3.5 yields that Z and W are respectively sub- and supersolutions of (\tilde{P}) in $\tilde{D} \times [r, r/h]$.

2. If our theorem is not true for u and v , then Z crosses W from below for the first time at $P_0 := (x_0, t_0) \in \tilde{D} \times [r, r/h]$ for $h \ll r$. Due to the maximum principle of harmonic functions and Lemma 3.4, $P_0 \in \Gamma(Z) \cap \Gamma(W)$. Note that by definition $\Omega(Z)$ and $\Omega(W)$ has respectively an interior ball B_1 and exterior ball B_2 at P_0 of radius r in space-time. (see Figure 3.)

Let us call H the tangent hyperplane to the interior ball of Z at P_0 . Since $Z \leq W$ for $t \leq t_0$ and $P_0 \in \Gamma(Z) \cap \Gamma(W)$, it follows that

$$B_1 \cap \{t \leq t_0\} \subset \Omega(Z) \cap \Omega(W); \quad B_2 \cap \{t \leq t_0\} \subset \{Z = 0\} \cap \{W = 0\}$$

$$\text{with } \bar{B}_1 \cap \bar{B}_2 \cap \{t \leq t_0\} = \{P_0\}.$$

Moreover, due to Lemma 3.5

$$(A.1) \quad \frac{Z_t}{|DZ|}(x, t) \leq F(|DZ|)(x, t) \leq \quad \text{on } \Gamma(Z)$$

and

$$(A.2) \quad \frac{W_t}{|DW|}(x, t) \geq F(|DW|)(x, t) + h \text{ on } \Gamma(W)$$

In particular due to (A.1) the arguments of Lemma 2.5 in [K1] applies for Z to yield that H is not horizontal. In particular $B_1 \cap \{t = t_0\}$ and $B_2 \cap \{t = t_0\}$ share the same normal vector ν_0 , outward with respect to B_1 , at P_0 .

Formally speaking, it follows that $\frac{Z_t}{|DZ|}(x_0, t_0) < \infty$ and

$$\frac{Z_t}{|DZ|}(P_0) \leq F(|DZ|)(P_0) \leq F(|DW|)(P_0) \leq \frac{W_t}{|DW|}(P_0) - h,$$

where the second inequality follows since $F(r)$ is increasing in r and $Z(\cdot, t_0) \leq W(\cdot, t_0)$ in a neighborhood of x_0 . Above inequality says that the free boundary speed of Z is strictly less than that of W at P_0 , contradicting the fact that $\Gamma(Z)$ touches $\Gamma(W)$ from below at P_0 .

For rigorous argument one can construct radially symmetric barrier functions based on the exterior and interior ball properties of Z and W at P_0 to derive a version of (A.1) and yield a contradiction. For details see the proof of Theorem 2.2 in [K1]. □

Next we prove Theorem 3.8.

Proof of Theorem 3.8. 1. We apply Perron's method. Without loss of generality we assume that $F(r) \geq F(0) \geq -1$. Since Ω_0 is star-shaped with respect to $B_h(0)$, there exists $C > 0$ such and $0 < \alpha < 1$ such that for any $r > 0$ and $x_0 \in \Gamma_0$

$$(A.3) \quad \sup_{x \in B_r(x_0)} u_0(x) \leq Cr^\alpha.$$

Let us define

$$U_1(x, t) = \begin{cases} (1 - t/h^2)^2 u_0\left(\frac{x}{1 - t/h^2}\right) & \text{for } 0 \leq t \leq h^2 \\ 0 & \text{for } t \geq h^2 \end{cases}$$

and

$$U_{2,r}(x, t) = (1 + r)^2 \inf_{y \in B_{r(t)}(x)} u_0((1 + r)^{-1}y),$$

where $r(t) = r - C_1 r^{\alpha-1}t$ for $0 \leq t \leq (C_1)^{-1}r^\alpha$. Note that due to (A.3) $U_{2,r}$ is a supersolution of (\tilde{P}) for sufficiently large C_1 . Moreover $U_1(x, t)$ is a subsolution of (\tilde{P}) due to the fact that $F \geq -1$ and $\Omega(u_0)$ contains $B_h(0)$.

Let $z(x, t) \in \mathcal{P}$ if and only if $z(x, t)$ is a viscosity subsolution of (\tilde{P}) with $z(\cdot, 0) \leq u_0(x)$ and $U_1 \leq z$ in $\mathbb{R}^N \times [0, \infty)$. Let

$$U(x, t) := \sup\{z(x, t) : z \in \mathcal{P}\}$$

Arguing as in the proof of Theorem 4.7 in [K1] yields that U^* and U_* are respectively viscosity subsolution and supersolution of (\tilde{P}) . Moreover due to Theorem 3.7, $U^* \leq U_{2,r}$ for $0 \leq t \leq r^\alpha$ for any $r > 0$. In particular $U_* = U^* = u_0$ at $t = 0$. In other words, U_* is a viscosity solution of (\tilde{P}) . \square

B Global-time existence and uniqueness for solutions with symmetry

The purpose of this section is to illustrate some examples where u_n^M satisfies (I) for all $t \geq 0$ with $r = r(t)$. For simplicity, we set $\int_{\Omega_0} u = 1$ and $\Omega_t(u) \subset B_1(0)$.

B.1 Reflection comparison

Lemma B.1 (Strong comparison principle). *Let u, v be resp. viscosity sub- and supersolutions of (\tilde{P}) in $\Sigma = D \times (a, b)$ with $u \leq v$ at $t = a$ and on $\partial D \times (a, b)$. In addition suppose that u satisfies (I) for $a \leq t \leq b$. Then $u(\cdot, t) \leq v(\cdot, t)$ in D for $a < t < b$.*

Proof. For simplicity let $a = 0$. Let us define

$$\tilde{u}(x, t) := (1 + \epsilon)^{-2} \sup_{y \in B_{r\epsilon}(x)} u((1 + \epsilon)y, t).$$

Observe that \tilde{u} is a subsolution of (\tilde{P}) due to Lemma 3.5. Also observe that, since $\Omega_t(u)$ is star-shaped with respect to $B_r(0)$ and $-\Delta u = \lambda$ in $\Omega(u)$,

$$\tilde{u}(x, t) \prec v(x, t) \text{ on the parabolic boundary of } \Sigma.$$

Hence Theorem 3.7 yields that $\tilde{u} \prec v$ for $0 \leq t \leq b$ for any $\epsilon > 0$, and thus $u \leq v$. \square

Recall that u_n^M solves (P_n^M) with given initial positive phase $\Omega_0(u_n^M) = \Omega_0$. (see Definition 4.1.)

Lemma B.2. *Suppose u_n^M satisfies (I) for $0 \leq t \leq T$. Let H be any hyperplane in \mathbb{R}^n and let $\phi_H(x)$ be the reflection of x with respect to H . Let D_1 and D_2 be the half-planes in \mathbb{R}^n divided by H . If*

$$u_n^M(x, t_0) \leq u_n^M(\phi_H(x), t_0) \text{ in } D_1$$

then

$$u_n^M(x, t) \leq u_n^M(\phi_H(x), t) \text{ in } D_1 \text{ for } t_0 < t \leq T.$$

Proof. Set $v(x, t) := u_n^M(\phi_H(x), t)$. Then v solves the following equation in $(kt_n, (k+1)t_n]$, $k = 1, 2, \dots$

$$\begin{cases} -\Delta v(\cdot, t) = \lambda_n^M(kt_n) & \text{in } \Omega(v) \\ v_t = |Dv| \min(|Dv|^3 - 1, M) & \text{on } \Gamma(v). \end{cases}$$

Moreover $v = u_n^M$ on $H = \partial D_1 = \partial D_2$. Since u_n^M has a compact support in any finite time period (Lemma 4.3), Lemma B.1 applies to $u_n^M \leq v$ in D_1 in $(kt_n, (k+1)t_n]$ for $t_0 < t \leq T$. \square

Corollary B.3. *Suppose that $\Omega_0(u_n^M) \subset B_R(0)$, $R > 1$ and u_n^M satisfies (I) for $0 \leq t \leq T$. Then $\Omega_t(u_n^M) \subset B_{3R}(0)$ for $0 \leq t \leq T$.*

Proof. For $x_0 \in (B_{3R}(0) - B_{2R}(0))$, define

$$(B.1) \quad \mathcal{C}(x_0) := \{y : y \cdot x_0 \leq -\frac{1}{2}|x_0||y|\}.$$

If we pick a hyperplane H normal to a vector $y_0 \in \mathcal{C}(x_0)$ containing x_0 , then

$$u_n^M(\phi_H(x), 0) = 0 \leq u_n^M(x, 0) \text{ in } D_1 := \{ty + H, t > 0\}.$$

Hence Lemma B.1 yields that u_n^M is monotone increasing in the cone of directions $\mathcal{C}(x_0)$.

Suppose $\Gamma_t(u_n^M)$ touches $\partial B_{3R}(0)$ for the first time at x_0 at $t = t_0$. Then due to (B.1) we have

$$(x_0 + \mathcal{C}(x_0)) \cap (B_{3R}(0) - B_{2R}(0)) \subset \Omega_{t_0}(u).$$

Since $\int u_n^M(\cdot, kt_n) dx = 1$, we obtain $\lambda \leq R^{-n-2}$. On the other hand $u_n^M(\cdot, t_0) = 0$ outside of $B_{3R}(0)$. Now comparing $u_n^M(\cdot, t_0)$ with $f(x) := (3R)^{-n} - (3R)^{-n-2}x^2$ yields $|Du_n^M|(x_0, t_0) < 1$. Therefore the outward normal velocity of $\Gamma(u_n^M)$ at (x_0, t_0) is strictly negative, contradicting the definition of t_1 . \square

For $B_r(x) \subset \Omega_0$, let

$$(B.2) \quad t(x, r) = \sup\{t : B_r(x) \subset \Omega_t(u_n^M)\}.$$

Note that, due to Lemma 3.4, $B_r(x)$ is touched by $\Gamma_t(u)$ for the first time at $t = t(x, r)$.

Lemma B.4. Suppose u_n^M satisfies (I) for $0 \leq t \leq T$. Suppose $\Omega_0(u)$ was star-shaped with respect to $B_r(x_0)$ with $t_0 := t(x_0, r) \leq T$. Let $y \in \partial B_r(x_0) \cap \Gamma_{t_0}(u)$ and let H be the hyperplane normal to $y - x_0$ containing x_0 . Then

$$u_n^M(\phi_H(x), t) \leq u_n^M(x, t) \text{ in } D_1 \times [t_0, T],$$

where D_1 is the half-plane which is divided by H containing $B_r(x_0)$.

Proof. Note that, since $\Omega_{t_0}(u_n^M)$ is star-shaped with respect to $B_r(x_0)$ with $y_0 \in \partial B_r(x_0) \cap \Gamma_{t_0}(u_n^M)$,

$$(B.3) \quad u_n^M(\phi_H(x), t_0) = 0 \text{ in } D_1.$$

Now we can conclude due to Lemma B.2. \square

B.2 Example 1: Two symmetric axis

Let e_1, \dots, e_n be an orthonormal basis in \mathbb{R}^n .

Theorem B.5. Suppose Ω_0 is star-shaped with respect to $B_r(0)$ and is symmetric with respect to e_1 and e_2 -axis. Then for any $T > 0, n$ and M , u_n^M satisfies (I) with $r = r(T) > 0$ for $0 \leq t < T$.

Remark Due to Lemma B.2, $\Omega_t(u_n^M)$ stays symmetric with respect to e_1 and e_2 axis.

Proof. Define $t_0 = t(0, r) > 0$. If $t_0 = \infty$ then we are done, so suppose t_0 is finite. Then $B_r(0) \subset \Omega(u_0)$ and $\Gamma_{t_0}(u_n^M)$ touches $\bar{B}_r(0)$ at some point $x_0 \in \partial B_r(0)$.

Let $\nu = -x_0$ and let H be the plane which is orthogonal to ν . Let D_1 be the half-plane divided by H which does not contain $B_r(0)$. Note that up to $t = t_0$, $\Omega_t(u_n^M)$ is star-shaped with respect to $B_r(0)$. Hence $u_n^M(\cdot, t_0) = 0$ in D_1 . By symmetry, $u_n^M(\cdot, t_0) = 0$ in the reflected image of D_1 with respect to e_1 and e_2 -axis. Thus $\Omega_{t_0}(u_n^M)$ lies between two parallel hyper-planes with width at most $2r$ (see Figure 4). Recall that due to Corollary B.3 $\Omega_t(u_n^M) \subset B_R(0)$ for some $R > 0$. Thus it follows that

$$\text{Vol}(\Omega_{t_0}(u_n^M)) \leq C(n)R^{n-1}r.$$

Recall that $\Omega_{t_0}(u_n^M) = 0$ in D_1 . If we choose r sufficiently small it follows that $|Du_n^M|(x_0, t_0) > 1$ (a detailed argument is given in the proof of Theorem B.7, step 3, Case 1). This means that $\Omega(u_n^M)$ is strictly expanding at (x_0, t_0) , contradicting the definition of t_0 . \square




Figure 4: Parallel hyperplanes bounding $\Omega_t(u_n^M)$.

Above theorem in particular states that a droplet with two symmetric axis, if it satisfies (I) initially, never changes its topology at a later time, however thin and long it is. On the contrary we will show below that a dumbbell-shaped droplet changes its topology in finite time.

Lemma B.6. *Suppose u_n^M solve (P_n^M) with initial positive phase*

$$\Omega_0 = B_1(-3e_1) \cup B_1(3e_1) \cup \{x = (x_1, x') : |x'| \leq r, |x_1| \leq 3\}.$$

If r is smaller than a dimensional constant, then $\Omega(u_n^M)$ changes its topology before $t = 1/2$.

Proof. First observe that, since the free boundary velocity is greater than -1 , for $0 \leq t \leq 1/2$ we have $B_{1/2}(\pm 3e_1) \subset \Omega_t(u_n^M)$. Hence $\lambda(u_n^M; t) < C(n)$ for $0 \leq t \leq 1/2$ and for some $C(n) \geq 1$. Pick $T = T(n)$ sufficiently small $\Omega_t(u_n^M) \subset B_1(0)$ for $0 \leq t \leq T$. Now one can compare u_n^M with

$$h(x, t) = C(n) \min[(|x|^2 - 100)_+, (r(t)x_1^4 - 3^4|x'|^2 + 3^4r^2)_+]$$

where $r(t) = (1 - (10C(n))^3 t)^{-1/3}$ for $0 \leq t \leq t_0 := (10C(n))^{-4}$. One can choose $C(n)$ sufficiently large such that $t_0 \leq T$.

Observe that $\Omega(u_0) \subset \Omega_0(h)$, $-\Delta h(\cdot, t) \geq C(n)$. Also a straightforward computation yields that

$$h_t = 4C(n)r'(t)x_1^3 \geq |Dh|(-\frac{1}{2} + |Dh|^3) \text{ on } \Gamma_t(h)$$

for $0 \leq t \leq t_0$, if $0 \leq r \leq 4^{-4}$.

Hence if we set

$$\tilde{h}(x, t) = \inf_{y \in B_{t/2}(x)} h(y, t),$$




Figure 5: The lines l_0, l_1 and their reflections.

Then \tilde{h} is a supersolution of (\tilde{P}) with $\lambda = C(n)$. Now due to Theorem 3.7 we have $u_n^M \prec \tilde{h}$ in $\mathbb{R}^n \times [0, t_0]$.

If $r \leq \frac{1}{2}t_0$ then it follows that $\Omega_{t_0}(\tilde{h})$ is no longer simply connected, and therefore so is $\Omega_{t_0}(u_n^M)$ (Change of topology occurred before $t = t_0 \leq 1/2$.) \square

B.3 Example 2: One axis symmetry with convexity

Here we set the dimension $N = 2$.

Theorem B.7. *Suppose $\Omega_0 \subset \mathbb{R}^2$ is convex and symmetric to e_1 axis. Then for any $T > 0, n$ and M , u_n^M satisfies (I) with $r = r(T) > 0$ for $0 \leq t \leq T$.*

Proof. 1. Let

$$\mathcal{S} = \{B_r(y) \subset \bar{\Omega}_0 : y \in \Omega_0 \cap \{x = (x_1, 0, \dots, 0)\}\}.$$

Then for each ball in \mathcal{S} there is the first time $\Gamma_t(u_n^M)$ touches the ball. Let t_0 be the supremum of these times. Then $\Gamma_{t_0}(u_n^M)$ touches $y_0 \in \partial B_r(x_0)$ for some $B_r(x_0) \subset \mathcal{S}$. We may assume that $(y_0 - x_0) \cdot e_1 \leq 0$. Let l_0 be the line normal to $y_0 - x_0$ with $y_0 \in l_0$.

2. First assume that $B_r(x_1) \subset \Omega_0$, where $x_1 := x_0 + r^{1/2}e_1$. Then when the free boundary hits the boundary of $B_r(x_1)$ for the first time at $t = t_1 \leq t_0$ it should not cross $B_r(x_0)$. Therefore the first touching point $y_1 \in \partial B_r(x_1)$ satisfies $(y_1 - x_1) \cdot e_1 \geq 0$. Let l_1 be the line normal to $y_1 - x_1$ with $y_1 \in l_1$ and let e_1 point to the right, horizontally.

3. Due to (1), $u_n^M(\cdot, t_0) = 0$ on the left side of l_0 . Moreover $u_n^M(\cdot, t_1) = 0$ on the right side of l_1 . By symmetry $u_n^M(\cdot, t_1) = 0$ on the right side of \tilde{l}_1 : the reflection of l_1 with respect to e_1 -axis (see Figure 5). Let θ_0 be the angle between l_0 and e_1 , and θ_1 be the angle between l_1 and e_1 .

Case 1: $\theta_1 < r^{1/2}$. By above argument $u_n^M(\cdot, t_1) = 0$ outside of the cone of angle $r^{1/2}$ with e_1 -axis. Since $\Omega_t(u_n^M) \subset B_R(0)$ for some R , $\Omega_t(u_n^M)$ is

confined in a cone of angle $r^{1/2}$ and height $6R$. Let $\lambda_0 := \lambda(t_1; u_n^M) = \lambda_0$ and

$$h(x) := \frac{\lambda_0}{2}([3Rr^{1/2}]^2 - (x_2)^2).$$

Since $-\Delta h = \lambda_0$ and

$$\Omega_{t_1}(u_n^M) \subset \Omega(h) = \{x : |x_2| \leq 3Rr^{1/2}\},$$

we have $u_n^M \leq h$. Thus

$$\int u_n^M(\cdot, t_1) dx \leq \int_{\Omega_{t_1}(u_n^M)} h(x) dx \leq (\lambda_0 R^2 r) \text{Vol}(\Omega_{t_1}(u_n^M)) \leq \lambda_0 (3Rr^{1/2})^3.$$

Since $\int u_n^M(\cdot, kt_n) dx = 1$, we obtain $\lambda_0 \geq (3Rr^{1/2})^{-3}$ if $r < (3R)^{-6}$. Note that $B_r(x_1) \subset \Omega_{t_1}(u_n^M)$, and thus

$$u_n^M(\cdot, t_1) \geq f(x) = (3R)^{-3} r^{1/2} - \frac{(3Rr^{1/2})^{-3}}{2} (x - x_1)^2.$$

In particular $|Du_n^M|(y_1, t_1) \geq |Df|(y_1) = (3R)^{-3} r^{-1/2} > 1$ if $r < (3R)^{-6}$. This contradicts the fact that the outward normal velocity of $\Gamma(u_n^M)$ at (x_1, t_1) is nonnegative.

Case 2: $\theta_1 > r^{1/2}$. Note that, up to $t = t_1$, $\Omega_t(u_n^M)$ is star-shaped with respect to both $B_r(x_0)$ and $B_r(x_1)$. Hence it follows that $\Omega_t(u_n^M)$ contains the strip

$$\Sigma = \{x : x \in B_r(z), z = x_0 + te_1, t \in [0, r^{1/2}]\}.$$

Let ϕ be the reflection with respect to the line parallel to e_2 and goes through x_1 . Then

$$u_n^M(\phi(x), t_1) \leq u_n^M(x, t_1) \text{ in } D_3 = \{x : (x - x_1) \cdot e_1 \leq 0\}.$$

Hence, by Lemma B.1 we have

$$(B.4) \quad u_n^M(\phi(x), t_0) \leq u_n^M(x, t_0).$$

If we combine (B.4) with the fact that $u_n^M(x, t_0) = 0$ on the right hand side of l_0 , then it follows that $\Omega_{t_0}(u_n^M)$ is contained in the channel of width at most $2r^{1/2}$ and height $6R$. Now the same argument as in Case 1 yields a contradiction if $r < (3R)^{-6}$.

4. Lastly suppose

$$y_1 \in B_r(x_0 + \tau e_1) \cap \partial\Omega_0 \text{ for some } \tau \in [0, r^{1/2}).$$

Let l_3 be the line parallel to e_2 containing y_1 . Since Ω_0 is convex and symmetric with respect to e_1 -axis, we have $u_0(x) = 0$ on the right side of l_3 . Therefore Lemma B.1 yields that for $t > 0$

$$u_n^M(\cdot, t) \leq u_n^M(\phi(x), t), \text{ on the right side of } l_3,$$

where $\phi(x)$ is the reflection of x with respect to l_3 . Now one can proceed as in Case 2 to derive a contradiction. □

Remark One class of initial configurations covered in above theorem are circular sectors

$$\Omega(u_0) = \{re^{i\theta} : 0 \leq r \leq R, 0 \leq \theta \leq \theta_0\}.$$

Acknowledgments

K.G. acknowledges support under NSF award DMS-0405596. I. K acknowledges support under NSF award DMS-0627896.

References

- [C] R. G. COX, *The dynamics of the spreading of liquids on a solid surface. part 1. viscous flow*, J. Fluid Mech., 168 (1986), pp. 169–194.
- [CL] M. G. CRANDALL AND P. L. LIONS, *Viscosity solutions of Hamilton-Jacobi equations*, Trans. Amer. Math. Soc., 277 (1983), pp. 1–42.
- [CJK] S. CHOI, D. S. JERISON AND I. C. KIM, *Regularity properties of initially Lipschitz free boundaries in One-Phase Hele-Shaw problem*, Amer. J. Math., 129 (2007), pp. 527–582.
- [G1] K. B. GLASNER, *A boundary integral formulation of quasi-steady fluid wetting*, J. Comput. Phys., 207 (2005), pp. 529–541.
- [G2] K. B. GLASNER, *Homogenization of contact line dynamics*, Int. Free Bound., to appear (2006).
- [Gr] H. P. GREENSPAN, *On the motion of a small viscous droplet that wets a surface*, J. Fluid Mech., 84 (1978), pp. 125–143.

- [Ho] L. M. HOCKING AND M. J. MIKSIS, *Stability of a ridge of fluid*, J. Fluid Mech., 247 (1993), pp. 157–177.
- [Hu] J. K. HUNTER, Z. LI, AND H. ZHAO, *Reactive autophobic spreading of drops*, J. Comput. Phys., 183 (2002), pp. 335–366.
- [K1] I. C. KIM, *Uniqueness and Existence result of Hele-Shaw and Stefan problem*, Arch. Rat. Mech. Anal., 168 (2003), pp. 299–328.
- [K2] I. C. KIM, *A free boundary problem with mean curvature*, Comm. PDE., 30 (2005). pp. 121–138.
- [T] L. TANNER, *The spreading of silicone oil drops on horizontal surfaces*, J. Phys. D., 12 (1979), pp. 1473–1484.
- [Va] J. L. VAZQUEZ, *The free boundary problem for the heat equation with fixed gradient condition*, *Free boundary problems: theory and applications*. Proceedings of the International Conference held in Toledo, June 21–26, 1993. Pitman Research Notes in Mathematics Series, 323. Longman, Harlow, 1996, pp. 277–302.
- [V] O. V. VOINOV, *Hydrodynamics of wetting*, [English translation] Fluid Dynamics, 11 (1976), pp. 714–721.