

A variational approach to a quasi-static droplet model

Natalie Grunewald* and Inwon Kim†

Abstract

We consider a quasi-static droplet motion based on contact angle dynamics on a planar surface. We derive a natural time-discretization and prove the existence of a weak global-in-time solution in the continuum limit. The time discrete interface motion is described in comparison with *barrier* functions, which are classical sub- and super-solutions in a local neighborhood. This barrier property is different from standard viscosity solutions since there is no comparison principle for our problem. In the continuum limit the barrier properties still hold in a modified sense.

Contents

1	Introduction	1
2	Construction of a time discrete solution	4
3	The barrier properties for time discrete solutions	8
4	The continuum limit and existence of weak solutions	15

1 Introduction

The motion of liquid drops on a planar surface is a widely studied topic. We consider a *quasi-stationary* free boundary model, derived in [8], [10] and [12]. The model is contact angle driven, i.e. the motion of the boundary of the wetted region is due to a deviation of the contact angle from the ideal contact angle. It is also quasi-stationary in the sense that the actual profile of the drop adjusts itself to the wetted region by minimizing a “surface energy” under a volume constraint.

We derive a natural time discretization by exploiting a formal gradient flow structure of the model. The time-discrete solutions satisfy barrier properties similar to standard viscosity solutions. These barrier properties stay valid in a modified sense as the time step size goes to zero.

Let us begin by a formal introduction of the model. The profile of the droplet is given by the height function $u : \mathbb{R}^N \times (0, T) \rightarrow \mathbb{R}$ with $N = 2$, the *positive phase* $\{u > 0\}$ denotes the wetted region and the

*Inst. für Angewandte Mathematik, Universität Bonn, Germany, grunewald@iam.uni-bonn.de

†Dept. of Mathematics, UCLA, Los Angeles CA 90095, USA, ikim@math.ucla.edu

free boundary $\partial\{u > 0\}$ denotes the contact line between drop, air and surface. It should be pointed out that our analysis is performed in general space dimension N . Throughout the paper we denote the spatial derivative of u by Du .

The motion of the droplet is described by contact angle dynamics - the free boundary $\partial\{u > 0\}$ evolves by a relationship between the outward normal velocity V and the *contact angle* $|Du|$ of the droplet with the surface. In this paper the normal velocity is given by

$$V = |Du|^2 - 1 \quad \text{on } \partial\{u > 0\}.$$

The square of the contact angle in the velocity law seems natural, as it is the only power for which we directly have a gradient flow structure like the one considered in this paper. For discussion of the contact angle dynamics in form of more general free boundary velocities we refer to [18].

On the other hand the shape of the drop adjusts to the wetted region by obeying two constraints: First the volume in each component ω_i of the drop is kept constant over time. Secondly the liquid/vapor interface is minimal in the sense that it minimizes the Dirichlet integral, leading to the Euler–Lagrange equation

$$-\Delta u(\cdot, t) \equiv \lambda$$

in each connected components of $\{u > 0\}$. This equation, a simplification of minimal surface equation, defines the shape of a quasi-static droplet. By choosing a suitable Lagrange multiplier $\lambda = \lambda_I(x, t)$, the volume of droplets in each component can be preserved.

Summarizing above discussion we arrive at the following free boundary problem:

$$(P) \quad \begin{cases} -\Delta u(\cdot, t) = \lambda_i(t) & \text{in } \omega_i(t); \\ V = |Du|^2 - 1 & \text{on } \partial\omega_i(t); \\ \int_{\omega_i(t)} u(\cdot, t) dx \equiv c_i, \end{cases}$$

where, as mentioned above, V is the outward normal velocity of the connected component of the support of the drop $\omega_i(t)$, so for $|Du| \neq 0$ one has $V = \frac{\partial_t u}{|Du|}$. As the overall volume is conserved we have $\sum_i c_i \equiv 1$.

Several serious challenges arise in developing a global notion of solutions for the model described above:

Most importantly, (P) *does not satisfy the comparison principle* between solutions, even in the case of single components. For example consider two sets $D_1 \subset D_2 \subset \mathbb{R}^n$ with the droplet profile $u_i(x, 0)$ supported in D_i for $i = 1, 2$. Suppose we have the same volume constraint, i.e., $\int u_i = 1$. Since we assume a quasi-stationary profile for u_i , they satisfy the first equation in (P) :

$$-\Delta u_i(x, 0) = \lambda_i \text{ in } D_i,$$

Due to the volume constraint and the fact that $D_1 \subset D_2$, it is clear that $\lambda_1 > \lambda_2$. Therefore it may be the case that $u_1 > u_2$ in some parts of D_1 . Also the fact that $\lambda_1 > \lambda_2$ and the second equation in (P)

suggests the possibility that the free boundary velocity of $\{u_1 > 0\}$ is bigger than in $\{u_2 > 0\}$ in some parts, and therefore the evolution of D_1 and D_2 by (P) may reverse the inclusion order between the sets.

Due to the failure of comparison principle, the viscosity solutions approach applied to mean curvature flow (see [7] and [4] for example) does not apply here, even if we assume that there is no topology change. Observe that if λ is independent of time then standard viscosity solution theory as in [14] applies. Based on this observation a discrete-time approximation with fixed λ in each time step was carried out in [9]. This way a unique weak solution is obtained for star-shaped initial data, for short times (as long as the wet region stays star-shaped). However approximating (P) with fixed λ in small time intervals (apparently) does not work well with topology changes.

On the other hand *topology changes* seem unavoidable. *Splitting* of droplets into multiple components is generic for non-convex droplets, even if we start the evolution with a simply connected droplet. *Merging* of different parts of the droplet also naturally occurs. (Recall that our model is quasi-stationary. This means that the dynamics inside the liquid phase is not modeled. In some sense when a topology change occurs we "fast forward" the time so that the droplet becomes quasi-stationary again.) In addition to the topological changes, we expect corner or cusp formation on the interface, due to merging, splitting, and also shrinking of droplets (see [9]).

Lastly, there is a bifurcation (*non-uniqueness*) of solutions in the event of merging. More precisely, two stationary drops touching each other at exactly one point can either decide to stay as they are, or see each other and develop into one big drop. A similar bifurcation was also observed, for solutions of a flame propagation model ([17]).

Our goal is to introduce a global-time notion of weak solution which describes (P) past topological changes and singularities. We take a variational approach, based on the following observation. Formally speaking the droplet evolution (P) is a gradient flow for the energy

$$E(\omega) := \int_{\omega} |Du_{\omega}|^2 dx + |\omega|, \tag{1.1}$$

where $|\omega|$ denotes the (Lebesgue) measure of ω . The gradient flow takes place on the manifold of possible supports of the droplet. The droplet height u itself is then part of a tangent bundle above the manifold. We refer to Section 2 for detailed discussion of this structure.

In Section 2 we approximate the solution (P) by a time-discrete gradient flow (JKO) scheme, originated by [13]. This scheme defines the solution in the next time step as a minimizer of a composited functional. This functional consists partly of the energy and partly of the distance to the previous time step. See Section 2 for details. Such approach was taken before by Almgren, Taylor and Wang [1] and Luckhaus and Sturzenhecker [15] for mean curvature motion. In [4] it was shown that a particular selection of the discrete scheme in [1] converges to viscosity solution of the mean curvature flow in the sense of [7].

As mentioned above our problem lacks the comparison property even in simple settings, which prevents us to develop any connection to standard viscosity solutions approach. However it is still possible to describe the evolution of solutions by *barrier properties* (Proposition 3.1 and 3.3) of the time-discrete weak solutions. Roughly speaking this means that the time-discrete solutions evolve with the free boundary velocity given by (P) , at "regular" points of the interface.

In the continuum limit we show that a global-in-time weak solution (see Definition 4.4) exists. At the moment, we are only able to describe the limiting free boundary behavior in terms of the liminf and limsup of the time-discrete solutions. We refer to Section 4 for definition of weak solutions (Definition 4.4) and precise statements (Theorem 4.6).

2 Construction of a time discrete solution

We consider a generalized version of (P) with curvature:

$$(P_\epsilon) \quad \begin{cases} -\Delta u(\cdot, t) = \lambda_i(t) & \text{in } \omega_i(t); \\ V = |Du|^2 - 1 - \epsilon\kappa & \text{on } \partial\omega_i(t); \\ \int_{\omega_i(t)} u(\cdot, t) dx \equiv \int u(x, 0) dx. \end{cases}$$

Here $\epsilon \geq 0$ and $\kappa = -\nabla \cdot \left(\frac{Du}{|Du|} \right)$ denotes the mean curvature of the interface, positive if the positive phase $\{u(\cdot, t) > 0\}$ is convex. The curvature term in $(P)_\epsilon$ is introduced to use the structure of Caccioppoli sets in the variational arguments in Section 3. However the regularized problem $(P)_\epsilon$ and their properties are also of independent interest.

Let us start with the definitions:

Definition 2.1 Let $B := \{x \in \mathbb{R}^n : |x| < R\}$ with R a sufficiently large constant.

(a) Let us define the set of Caccioppoli sets

$$Cacc := \{\omega \subset B; \omega \text{ is a Borel set with finite perimeter}\}.$$

(b) For any $\omega \in Cacc$ and any volume c

$$u_{\omega, c} := \operatorname{argmin} \left\{ \int_{\omega} |Du|^2 dx : u \in H^1(\omega), \operatorname{supp} u \subseteq \omega, \int_{\omega} u dx = c \right\}.$$

Remark 2.2 Note that the minimizer $u_{\omega, c}$ exists for any $c > 0$ and any set $\omega \in Cacc$ that admits one H^1 -function u with $\operatorname{supp} u \subseteq \omega$.

Definition 2.3 For a nonnegative function $u \in H^1(\mathbb{R}^n)$ and for $x \in \Omega(u)$ we define

$$\lambda(u)(x) := \frac{\int_{\omega} |Du|^2 dx}{\int_{\omega} u dx} \tag{2.1}$$

where ω is the connected component of $\Omega(u)$ which contains x . If $\int_{\omega} u dx = 0$ we set $\lambda(u)(x) = 0$.

Note that $-\Delta u_{\omega, c} = \lambda(u_{\omega, c})(x) \equiv \lambda(u_{\omega, c})$ in its positive set, if ω has a single component with smooth boundary.

Problem (P_ϵ) is a formal gradient flow on $Cacc$ for the energy

$$E_\epsilon(\omega) := \int_\omega |Du_{\omega,1}|^2 dx + |\omega| + \epsilon per(\omega), \quad (2.2)$$

where $|\omega|$ and $per(\omega)$ respectively denote the Lebesgue measure and the perimeter of ω . To see this we calculate the differential of E_ϵ for some normal velocity field \tilde{v} applied to $\partial\omega$ and $\tilde{\delta}u$ the change of u introduced by \tilde{v} :

$$\begin{aligned} diff E_\epsilon(\omega) \cdot \tilde{v} &= \int_\omega 2Du_\omega \cdot D\tilde{\delta}u dx + \int_{\partial\omega} (1 + |Du_\omega|^2) \tilde{v} dS + \int_{\partial\omega} \epsilon\kappa \tilde{v} dS \\ &= - \int_\omega 2\Delta u_\omega \tilde{\delta}u dx + \int_{\partial\omega} -2|Du_\omega| \tilde{\delta}u + (1 + |Du_\omega|^2 + \epsilon\kappa) \tilde{v} dS \\ &= \lambda \int_\omega \tilde{\delta}u dx + \int_{\partial\omega} -2|Du_\omega|^2 \tilde{v} + (1 + |Du_\omega|^2 + \epsilon\kappa) \tilde{v} dS \\ &= \int_{\partial\omega} (1 - |Du_\omega|^2 + \epsilon\kappa) \tilde{v} dS. \end{aligned}$$

This gives (P_ϵ) for the Riemannian structure

$$g_\omega(v, \tilde{v}) := \int_{\partial\omega} v \tilde{v} dS \quad \forall v, \tilde{v} \in T_\omega Cacc, \quad (2.3)$$

on $Cacc$, by the volume conservation and Definition 2.1. As the distance connected to (2.3) are difficult to model, we introduce a modified “distance”, which was originally introduced in [1] and [15] (also see e.g. [5] and [4].)

$$\widetilde{dist}^2(\omega_0, \omega_1) := \int_{\omega_0 \Delta \omega_1} dist(x, \partial\omega_0) dx.$$

Here $dist$ is the distance function, and $\omega_0 \Delta \omega_1$ denotes the symmetric difference between the two sets. Note that \widetilde{dist}^2 is not a (squared) distance function (it lacks e. g. symmetry), but an approximation of the distance connected to (2.3).

Following [15] and the JKO-scheme [13], ω_h^{i+1} is determined from the previous set ω_h^i by

$$\omega_h^{i+1} = \underset{\omega \in Cacc}{argmin} \left\{ \frac{1}{h} \widetilde{dist}^2(\omega_h^i, \omega) + E_\epsilon(\omega) \right\}.$$

Lemma 2.4 *For fixed $h > 0$, fixed volume c and any $\omega^0 \in Cacc$ there exists at least one minimizer $\omega_c^{min} \in Cacc$ of*

$$\mathcal{F}(\omega) := \frac{1}{h} \widetilde{dist}^2(\omega^0, \omega) + E_\epsilon(\omega).$$

Note that we do not show uniqueness. We also do not expect uniqueness for (P) or (P_ϵ) , see Section 1. The dependence on c is suppressed in the notation of E_ϵ .

Proof. There exist sets $\omega \subset B$ such that $\mathcal{F}(\omega) < \infty$ (e.g. spheres around ω^0) and $\mathcal{F}(\omega) \geq 0$. Therefore there exists a minimizing sequence $\{\omega_k\} \subset Cacc$ such that

$$\mathcal{F}(\omega_k) \xrightarrow{k \rightarrow \infty} \inf\{\mathcal{F}(\omega) : \omega \subset B\}.$$

By the definition of $E_\epsilon(\omega)$ we have $|\omega_k| + \epsilon per(\omega_k) < C$ and therefore the indicator functions χ_{ω_k} are uniformly bounded in BV–norm. Thus (see e.g. [6], p.176) there exists a subsequence and a function $\chi \in BV(B)$ such that

$$\chi_{\omega_k} \rightarrow \chi \quad \text{in } L^1(B)$$

Since χ_{ω_k} take values in $\{0,1\}$ so does χ and there exists a set $\omega_c^{min} \subset B$ such that $\chi = \chi_{\omega_c^{min}}$. It remains to show that $\mathcal{F}(\omega_c^{min}) \leq \inf \mathcal{F}(\omega_k)$. This is direct for the part of the energy $|\omega_k| + \epsilon per(\omega_k)$, by the lower semi continuity of the perimeter and the L^1 convergence of χ_{ω_k} . For the remaining part of the energy we have to take into account the convergence of the corresponding droplet with volume c , $u_{\omega_k,c}$. By the boundedness of the H^1 –norm of $u_{\omega_k,c}$

$$u_{\omega_k,c} \rightarrow \tilde{u} \quad \text{in } L^2(B).$$

Where $\int \tilde{u} = c$ and

$$u_{\omega_k,c} = u_{\omega_k,c} \chi_{\omega_k} \rightarrow \tilde{u} \chi_{\omega_c^{min}} \text{ a.e. in } B.$$

Therefore by the lower semi-continuity of H^1 –norm and Definition 2.1

$$\inf \int_{\omega_k,c} |Du_{\omega_k,c}|^2 \geq \int_{\omega_c^{min}} |D\tilde{u}|^2 \geq \int_{\omega_c^{min}} |Du_{\omega_c^{min},c}|^2.$$

On the other hand \widetilde{dist}^2 is continuous with respect to the L^1 –topology of the indicator functions:

$$\begin{aligned} \left| \widetilde{dist}^2(\omega^0, \omega) - \widetilde{dist}^2(\omega^0, \bar{\omega}) \right| &= \left| \int_{\omega^0 \Delta \omega} dist(x, \partial\omega^0) dx - \int_{\omega^0 \Delta \bar{\omega}} dist(x, \partial\omega^0) dx \right| \\ &= \left| \int_{(\omega \Delta \bar{\omega}) \Delta \omega^0} dist(x, \partial\omega^0) dx \right|. \end{aligned}$$

This vanishes as $\|\chi_\omega - \chi_{\bar{\omega}}\|_{L^1(\mathbb{R}^N)} = |\omega \Delta \bar{\omega}| \rightarrow 0$, by the boundedness of the distance function in B . \square

Definition of the time–discrete evolution. We define a time–discrete evolution of (P) . Roughly speaking we do the minimization in Lemma 2.4 for each component of the drop separately. If two components merge at the next time step, we go back and do the same minimization step but for the two components together. Splitting of a component is already taken care of in the minimization in Lemma 2.4, as ω^{min} might have several components. To be more precise: for fixed $h > 0$ and $i \in \mathbb{N}$ take the previous state $\omega_h^i \in Cacc$ with possibly infinitely many connected components $\omega_h^{i,k} \in Cacc, k \in \mathbb{N}$. For each connected component (in the classical sense) we have some droplet $u_{\omega_h^{i,k}, c_k}$ by Remark 2.2 and Lemma 2.4. Then ω_h^{i+1} is given by

$$\omega_h^{i+1} := \bigcup_k \omega_{c_k}^{min} \quad \text{if for any } l \neq m : \omega_{c_l}^{min} \cap \omega_{c_m}^{min} = \emptyset, \quad (2.4)$$

where $\omega_{c_k}^{min}$ is a minimizer in Lemma 2.4 for the connected component $\omega_h^{i,k}$.

If $\omega_{c_l}^{min} \cap \omega_{c_m}^{min} \neq \emptyset$ for only one pair (l, m) and if it does not intersect with other components $(\omega_{c_k}^{min}, k \neq l, m)$, then we define

$$\omega_h^{i+1} := \left(\bigcup_{k \neq l, m} \omega_{c_k}^{min} \right) \cup \omega_{c_l+c_m}^{min}.$$

where $\omega_{c_l+c_m}^{min}$ is a minimizer in Lemma 2.4 for initial set $\omega_h^{i,l} \cup \omega_h^{i,m}$.

In general the process of sorting out merging components is non-unique: we will prescribe the following process to proceed without ambiguity. Let us first consider the maximal index set I_1 such that each element $\omega_{c_k}^{min}$ with $k \in I_1$ intersects with $\omega_{c_1}^{min}$.

Next take the first element $\omega_{c_k}^{min}$ with $k \notin I_1$ and repeat the process to create the second index set I_2 . If I_2 intersects with I_1 , then we replace I_1 with $I_1 \cup I_2$. If not, check whether

$$\omega_{I_1}^{min} := \omega_{\sum_{c_k}^{min}}, \quad k \in I_1$$

intersects with $\omega_{I_2}^{min}$. If yes then still replace I_1 with $I_1 \cup I_2$. If no, then proceed to create the third index set I_3 , and check against $\omega_{I_1}^{min}$ and $\omega_{I_2}^{min}$. This way we end up with a sequence of (disjoint) index sets I_1, I_2, \dots such that $\omega_{I_k}^{min}$ are all disjoint. Then

$$\omega_h^{i+1} := \bigcup_k \omega_{I_k}^{min}.$$

Now define

$$u_{I_k} := u_{\omega_{I_k}^{min}, \sum_{j \in I_k} c_j}$$

and

$$u_h(\cdot, t) := \sum_k u_{I_k} \quad \text{for } t \in [ih, (i+1)h]. \quad (2.5)$$

This way u_h is a H^1 -function in B at any time $t \in [0, T]$. Thus

$$u_h \in L_{loc}^2(H^1(B)).$$

The total volume of u_h at time t is $\int u_h(\cdot, t) dx = \sum_k c_k = 1$.

As the JKO-scheme is constructed to describe a time-discrete gradient flow, we have the energy decrease for free: Suppressing in the notation the dependence of the energy on the volumes in each component, we have:

Lemma 2.5 *The time evolution defined in (2.4) and (2.5) satisfies*

$$E_\epsilon(\omega_h^i) \geq \frac{1}{h} \widetilde{dist}^2(\omega_h^i, \omega_h^{i+1}) + E_\epsilon(\omega_h^{i+1}). \quad (2.6)$$

Proof. Equation (2.6) is obvious for any components $\bigcup_{k \in I_j} \omega_h^{i,k}$ and the corresponding minimizer $\omega_{I_j}^{min} = \omega_{I_j}^{i+1}$, as Lemma 2.4 can be tested with the set $\bigcup_{k \in I_j} \omega_h^{i,k}$. Furthermore

$$\begin{aligned} E_\epsilon(\omega_h^i) &= \sum_j E_\epsilon\left(\bigcup_{k \in I_j} \omega_h^{i,k}\right) && \geq \frac{1}{h} \sum_j \widetilde{dist}^2\left(\bigcup_{k \in I_j} \omega_h^{i,k}, \omega_{I_j}^{min}\right) + \sum_j E_\epsilon(\omega_{I_j}^{min}) \\ &&& \geq \frac{1}{h} \widetilde{dist}^2(\omega_h^i, \omega_h^{i+1}) + E_\epsilon(\omega_h^{i+1}). \end{aligned}$$

□

3 The barrier properties for time discrete solutions

In this section we show, that for fixed time step $h > 0$ the discrete-time solution constructed above satisfies the free boundary motion law in time scale h , in the sense that it is comparable to smooth sub- and super-solutions of (P_ϵ) in local neighborhoods. A more precise statement will follow in Propositions 3.1 and 3.3 for which we need the following notation:

Let us denote the *positive phase* of a function $u(x, t) : \mathbb{R}^N \times [0, \infty) \rightarrow \mathbb{R}^+$ and its boundary by:

$$\Omega_t(u) := \{u(\cdot, t) > 0\} \quad \text{and} \quad \Gamma_t(u) := \partial\{u(\cdot, t) > 0\},$$

and the positive phase in space-time by:

$$\Omega(u) := \{u > 0\} \subset \{B \times [0, \infty)\} \quad \text{and} \quad \Gamma(u) := \partial\Omega(u).$$

Next we show the barrier properties for the time discrete solutions. We begin with the barrier property for u_h being a super-solution. That is, u_h can be compared to a barrier function ϕ that is below. If ϕ is not fast enough at the boundary and not curved enough in the interior, then the ordering will persist:

Proposition 3.1 (Super-solution barrier property) *Let u_h be defined by (2.5). Given a ball $B_r(x_0)$ in B let*

$$\lambda := \inf_{x \in B_r(x_0)} \{\lambda(u_h(0, \cdot))(x), \lambda(u_h(h, \cdot))(x)\},$$

where $\lambda(u)(x)$ is as defined by (2.1).

Suppose there exists a smooth function ϕ with $|D\phi| \neq 0$ in $B_r(x_0) \times [0, h]$. Further suppose that for some small $\delta > 0$

$$-\Delta\phi(\cdot, t) < \lambda - \delta \quad \text{in} \quad B_r(x_0) \times [0, h], \tag{3.1}$$

$$\frac{\phi_t}{|D\phi|} - (|D\phi|^2 - 1 - \epsilon\kappa_\phi) < -\delta \quad \text{on} \quad \Gamma(\phi) \cap (B_r(x_0) \times [0, h]),$$

where $\kappa_\phi := -\nabla \cdot (\frac{D\phi}{|D\phi|})$ is the mean curvature of the corresponding level set of ϕ . Then for sufficiently small $h > 0$ - depending on δ, r , the minimum of $|D\phi|$ and the C^2 -norm of ϕ in $B_r(x_0) \times [0, h]$ - the following holds:

If $\phi \leq u_h$ on the parabolic boundary of $B_r(x_0) \times [0, h]$, then $\phi(\cdot, h) \leq u_h(\cdot, h)$ in $B_r(x_0)$.

Note that $\frac{\phi_t}{|D\phi|} = V$, where V is the outward normal velocity of $\partial\{\phi > 0\}$ with respect to the positive set of ϕ . Therefore, Proposition 3.1 shows that a function ϕ which is a *sub-solution* of (P_ϵ) can not cross the discrete time solution u_h . Thus, u_h is a super-solution. We also mention that a local barrier function like the ones in Proposition 3.1 can always be extended to a global barrier function satisfying (3.1), which is not restricted to a ball B_r .

We begin by a lemma which states that the support of ϕ cannot cross ω_h too much.

Lemma 3.2 *Under the assumptions of Proposition 3.1 and for sufficiently small $h > 0$, there exists a constant $C > 0$ independent of $h > 0$ such that*

$$(\phi(x, t) - Ch^{1/2})_+ \leq u_h(x, t) \text{ in } B_r(x_0) \times [0, h]. \quad (3.2)$$

In particular, if ϕ crosses u_h on $B_r(x_0) \times [0, h]$, we can choose $-h^2 < \tau < Ch^{1/2}$ such that $\varphi(x, t) := (\phi(x, t) - \tau)_+$ crosses ω_h from below by $o(h^2)$, i.e.,

$$0 < |(\Omega_h(\varphi) - \omega_h) \cap B_r(x_0)| = o(h^2). \quad (3.3)$$

Proof. Once (3.2) is proved, our second claim follows from the fact that $\Omega_h(\phi)$ does cross ω_h and ϕ is smooth with $|D\phi| > 0$ near $\Omega(\phi)$.

To prove (3.2), first observe that

$$\Omega_h(\phi) \subset \{\Omega_0(\phi) \cup \{x \in B : d(x, \Gamma_0(\phi)) \leq C_1 h\}\},$$

where

$$C_1 = \sup_{B_r(x_0) \times [0, h]} |\phi_t| / |D\phi|.$$

Therefore, we have for φ

$$\Omega_h(\varphi) \subset \{x \in \Omega_0(\phi) : d(x, \Gamma_0(\phi)) \geq C_2 \tau - C_1 h\}$$

with

$$C_2 = \inf_{B_r(x_0) \times [0, h]} |D\phi|^{-1}(x, t).$$

Thus, (3.2) follows by the comparison principle if we show

$$S := \{x \in (\Omega_0(\phi) \cap B_r(x_0)) : d(x, \Gamma_0(\phi)) \geq C_3 h^{1/2}\} \subset \omega_h. \quad (3.4)$$

To prove (3.4), let us define

$$\hat{\omega}_h := \omega_h \cup S \text{ and } \Sigma := \hat{\omega}_h - \omega_h = S - \omega_h.$$

Suppose that $|\Sigma| \neq 0$, otherwise we are done. Since $\Omega_0(\phi) \subset \omega_0$, we have by the smoothness of ϕ

$$\widetilde{dist}^2(\omega_0, \hat{\omega}_h) - \widetilde{dist}^2(\omega_0, \omega_h) \leq -C_4 h |\Sigma|.$$

where C_4 is proportional to the size of C_3 . On the other hand, since the Dirichlet energy decreases when the domain increases,

$$\begin{aligned} E_\epsilon(\hat{\omega}_h) - E_\epsilon(\omega_h) &\leq |\Sigma| + \epsilon \text{per}(\hat{\omega}_h) - \epsilon \text{per}(\omega_h) \\ &\leq |\Sigma| + \epsilon |\partial S - \omega_h| - \epsilon |\partial \omega_h \cap S| \\ &\leq C_5 |\Sigma|, \end{aligned}$$

where C_5 depends on ϕ . The last inequality follows from

$$\begin{aligned} |\partial S - \omega_h| - |\partial \omega_h \cap S| &\leq \int_{\partial \Sigma} -\frac{D\phi}{|D\phi|}(x, h) \cdot \eta \, dS \\ &= -\int_{\Sigma} \nabla \cdot \left(\frac{D\phi}{|D\phi|}\right)(x, h) \, dx \\ &= \int_{\Sigma} \kappa_\phi \, dx. \end{aligned}$$

Here η is the outward normal vector at $x \in \partial\Sigma$ and κ_ϕ is the mean curvature of the level set of ϕ . We conclude that if C_3 is chosen sufficiently large, then

$$\widetilde{dist}^2(\omega_0, \hat{\omega}_h) + h E_\epsilon(\hat{\omega}_h) < \widetilde{dist}^2(\omega_0, \omega_h) + h E_\epsilon(\omega_h).$$

This contradicts the minimizing property of ω_h . \square

Proof of Proposition 3.1:

1) Suppose the proposition is not true. Then $\phi(x_0, h) > u_h(x_0, h)$ at some point $x_0 \in \Omega_h(u_h)$. Due to the maximum principle for harmonic functions, this implies that $\Omega_h(\phi) \cap (B \setminus \omega_h) \neq \emptyset$ for one of the components ω_h in $\Omega_h(u_h)$. Let ω_0 be a corresponding component in $\Omega_0(u_h)$ which gives rise to ω_h .

For notational simplicity, we prove the proposition assuming that ω_h is indeed the only component generated by ω_0 , i.e. ω_0 has not splitted into multiple components and ω_h is generated by only one component: the proof for the general case is parallel.

2) Let us define

$$\varphi(x, t) := (\phi(x, t) - \tau)_+$$

where $-h^2 < \tau < Ch^{1/2}$ is as given in Lemma 3.2. In the proof we use φ instead of ϕ , which is possible without violating the assumptions according to Lemma 3.2. Next set

$$\tilde{\omega}_h := \omega_h \cup (\Omega_h(\varphi) \cap B_r(x_0)).$$

We claim that

$$\widetilde{dist}^2(\omega_0, \tilde{\omega}_h) + h E_\epsilon(\tilde{\omega}_h) < \widetilde{dist}^2(\omega_0, \omega_h) + h E_\epsilon(\omega_h), \quad (3.5)$$

which yields a contradiction to the minimizing property of ω_h .

3) To prove (3.5) first observe that

$$\begin{aligned} \widetilde{dist}^2(\omega_0, \tilde{\omega}_h) - \widetilde{dist}^2(\omega_0, \omega_h) &= \int_{\tilde{\omega}_h \Delta \omega_h} \text{signdist}(x, \partial\omega_0) dx \\ &\leq \int_{\tilde{\omega}_h \Delta \omega_h} \text{signdist}(x, \Gamma_0(\varphi)) dx, \end{aligned}$$

where *signdist* is the signed distance function, that is negative inside the set. Here the first equality is due to straightforward computation, and the inequality is due to the fact that $\Omega_0(\varphi)$ is a subset of ω_0 . By construction of φ , for each point $x \in \tilde{\omega}_h \Delta \omega_h$ there exists a time t^* with $0 \leq t^* \leq h + o(h^2)$ such that $x \in \Gamma_{t^*}(\varphi)$. Therefore, as $\frac{\varphi_t}{|D\varphi|}(0, \cdot)$ denotes the outward normal velocity of $\Gamma(\varphi)$,

$$\text{signdist}(x, \Gamma_0(\varphi)) \leq h \frac{\varphi_t}{|D\varphi|} + o(h). \quad (3.6)$$

Next we consider the energy difference

$$E_\epsilon(\omega_h) - E_\epsilon(\tilde{\omega}_h) = I + II + III$$

where

$$I = \int |Du_h|^2(\cdot, h) - \int |D\tilde{u}_h|^2, \quad II = - \int_{\tilde{\omega}_h \Delta \omega_h} 1 \, dx, \quad III = \epsilon \operatorname{per}(\omega_h) - \epsilon \operatorname{per}(\tilde{\omega}_h).$$

Here $\tilde{u}_h(x) := u_{\tilde{\omega}_h, f} u_h$. In the next step we will show that

$$I \geq \int_{\tilde{\omega}_h \Delta \omega_h} |D\varphi|^2(\cdot, h) \, dx \quad \text{and} \quad III \geq \int_{\tilde{\omega}_h \Delta \omega_h} -\epsilon \kappa_\varphi \, dx$$

This proves our claim by (3.1) and (3.6). Note that (3.1) is strict and therefore extends to a small region inside.

4) Let us estimate III . Note that, as before,

$$\begin{aligned} \operatorname{per}(\omega_h) - \operatorname{per}(\tilde{\omega}_h) &\geq \int_{\partial\omega_h \setminus \partial\tilde{\omega}_h} -\frac{D\varphi}{|D\varphi|}(\cdot, h) \cdot \eta \, dS - \int_{\partial\tilde{\omega}_h \setminus \partial\omega_h} -\frac{D\varphi}{|D\varphi|}(\cdot, h) \cdot \tilde{\eta} \, dS \\ &= \int_{\tilde{\omega}_h \Delta \omega_h} \nabla \cdot \left(\frac{D\varphi}{|D\varphi|} \right)(\cdot, h) \, dx \\ &= \int_{\tilde{\omega}_h \Delta \omega_h} -\kappa_\varphi \, dx \end{aligned}$$

where $\tilde{\eta} = -D\varphi/|D\varphi|(x, h)$ is the outward normal vector at $x \in \partial\tilde{\omega}_h$, η is the outward normal vector at $x \in \partial\omega_h$, and κ_φ is the mean curvature of the level sets of φ .

It remains to estimate I . To this end let us define two auxiliary functions, \bar{u} and v :

$$\begin{aligned} -\Delta \bar{u} &= \lambda \quad \text{in } \tilde{\omega}_h \quad \text{with} \quad \operatorname{supp}(\bar{u}) = \overline{\tilde{\omega}_h}, \\ -\Delta v &= 0 \quad \text{in } \omega_h \quad \text{with} \quad v = \varphi(\cdot, h) \quad \text{on } \partial\omega_h. \end{aligned} \tag{3.7}$$

We remark that \bar{u} is defined by approximation from outside and v is defined by approximation from inside, i.e.

$$\bar{u}(x) := \inf\{f(x) : -\Delta f = \lambda \text{ in } \{f > 0\} \text{ with } \tilde{\omega}_h \subset \overline{\{f > 0\}}\},$$

and

$$v(x) := \sup\{f(x) : -\Delta f = 0 \text{ in } \omega_h \text{ with } f < \phi \text{ on } B \setminus \omega_h\}.$$

Let us define $c := \int_{\omega_h} u_h(\cdot, h)$, $\bar{c} := \int_{\tilde{\omega}_h} \bar{u}$ and $\tilde{u} := \frac{\epsilon}{\bar{c}} \bar{u}$. Then the following holds:

$$\begin{aligned} \int_{\omega_h} |Du_h(\cdot, h)|^2 - \int_{\tilde{\omega}_h} |D\tilde{u}_h|^2 &= \lambda \int_{\omega_h} u_h(\cdot, h) - \lambda(\tilde{u}) \int_{\tilde{\omega}_h} \tilde{u} \\ &= \frac{\epsilon}{\bar{c}} \lambda (\int_{\tilde{\omega}_h} \bar{u} - \int_{\omega_h} u). \end{aligned} \tag{3.8}$$

Furthermore, $\bar{u} \geq \max((u_h + v)|_{\omega_h}, \varphi)(\cdot, h)$ since

$$\tilde{\omega}_h = \Omega_h(\max((u_h + v)|_{\omega_h}, \varphi)) \quad \text{and} \quad \max(-\Delta(u_h + v)|_{\omega_h}, -\Delta\varphi)(\cdot, h) \leq \lambda.$$

For the same reason, on the reduced boundary of ω_h we have, for the inward normal η ,

$$\partial_\eta(u_h(\cdot, h) + v) \geq \partial_\eta \varphi(\cdot, h). \tag{3.9}$$

Thus,

$$\begin{aligned}
\lambda \left(\int_{\tilde{\omega}_h} \bar{u} - \int_{\omega_h} u_h(\cdot, h) \right) &\geq \lambda \int_{\omega_h} v + \lambda \int_{\omega_h \Delta \tilde{\omega}_h} \varphi(\cdot, h) \\
&\geq - \int_{\omega_h} \Delta(u_h(\cdot, h) + v) v - \int_{\omega_h \Delta \tilde{\omega}_h} (\Delta \varphi \varphi)(\cdot, h) \\
&\geq \int_{\omega_h} (D(u_h(\cdot, h) + v)) Dv + \int_{\partial \omega_h} \partial_\eta(u_h(\cdot, h) + v) v \\
&\quad + \int_{\omega_h \Delta \tilde{\omega}_h} |D\varphi|^2(\cdot, h) - \int_{\partial \omega_h} (\partial_\eta \varphi \varphi)(\cdot, h) \\
&\geq \int_{\omega_h} |Dv|^2 + \int_{\omega_h \Delta \tilde{\omega}_h} |D\varphi|^2(\cdot, h),
\end{aligned}$$

by (3.9) and (3.7). Thus together with (3.8) we have

$$I \geq \frac{c}{\bar{c}} \int_{\omega_h \Delta \tilde{\omega}_h} |D\varphi|^2(\cdot, h).$$

Lastly, note that as $\tau \rightarrow \tau_0$, $\lambda(\tilde{u})$ converges to $\lambda(u_h)$, due to the minimizing property of ω_h . Hence $\bar{c} \rightarrow c$ and we can choose $\tau - \tau_0$ given in (3.3) small enough that $\bar{c} \leq c(1 + o(h))$ to conclude. \square

By a parallel argument in the proof of Proposition 3.1, u_h can also be compared with barriers which are *super-solutions* of (P_ϵ) :

Proposition 3.3 (Sub-solution – barrier property) *Let u_h be defined by (2.5). Given a ball $B_r(x_0)$ in B , let*

$$\lambda := \inf_{x \in B_r(x_0)} \{ \lambda(u_h(0, \cdot))(x), \lambda(u_h(h, \cdot))(x) \},$$

where $\lambda(u)(x)$ is as defined in (2.1).

Suppose there exists a smooth function ϕ with $|D\phi| \neq 0$ in $B_r(x_0) \times [0, h]$. Further suppose that for some small $\delta > 0$

$$-\Delta\phi(\cdot, t) > \lambda + \delta \quad \text{and} \quad \frac{\phi_t}{|D\phi|} - (|D\phi|^2 - 1 - \epsilon\kappa_\phi) > \delta \quad \text{in } B_r(x_0) \times [0, h]. \quad (3.10)$$

Then for sufficiently small $h > 0$ – depending on δ, r , the minimum of $|D\phi|$ and the C^2 -norm of ϕ in $B_r(x_0) \times [0, h]$ – the following holds:

If $u_h \leq \phi_+ := \max(\phi, 0)$ on the parabolic boundary of $B_r(x_0) \times [0, h]$, then $u_h(\cdot, h) \leq \phi(\cdot, h)_+$ in $B_r(x_0)$.

Proof. The proof is analogous to the proof of Proposition 3.1. We still present it, as the estimation of the Dirichlet integral has a non-trivial difference from the previous proof.

Suppose the above proposition is not true. Then $\phi(\cdot, h)$ crosses $u_h(\cdot, h)$ from above at some point in $B_r(x_0)$. As before, the maximum principle for harmonic functions states that then $\Omega_h(\phi) \cap \omega_h$ is

nonempty for a component ω_h of $\Omega_h(u_h)$. Set ω_0 be the component of $\Omega_0(u_h)$ which generates ω_h . Again we construct a contradiction to the minimizing property of ω_h and u_h . With a parallel argument to Lemma 3.2 one can change ϕ to $\varphi := (\phi + \tau)_+$, $h^2 \leq \tau \leq Ch^{1/2}$, such that $u_h(x, h) \leq (\phi(x, h) + \tau)_+$ and

$$0 < |(\omega_h - \Omega_h(\varphi)) \cap B_r(x_0)| = o(h^2).$$

This time we denote:

$$\tilde{\omega}_h = ((\omega_h \cap \Omega_h(\varphi)) \cap B_r(x_0)) \cup (\omega_h \cap (B \setminus B_r(x_0))).$$

We claim that

$$\widetilde{dist}^2(\omega_0, \tilde{\omega}_h) + h E_\epsilon(\tilde{\omega}_h) < \widetilde{dist}^2(\omega_0, \omega_h) + h E_\epsilon(\omega_h).$$

First observe that this time

$$\begin{aligned} \widetilde{dist}^2(\omega_0, \tilde{\omega}_h) - \widetilde{dist}^2(\omega_0, \omega_h) &= - \int_{\tilde{\omega}_h \Delta \omega_h} \text{signdist}(x, \partial\omega_0) dx \\ &\leq - \int_{\tilde{\omega}_h \Delta \omega_h} \text{signdist}(x, \Gamma_0(\varphi)) dx. \end{aligned}$$

By integration of the velocity of $\Gamma_t(\varphi)$ we have

$$-\text{signdist}(x, \Gamma_0(\varphi)) \leq -h \frac{\varphi_t}{|D\varphi|} + o(h). \quad (3.11)$$

Next we consider the energy difference

$$E_\epsilon(\omega_h) - E_\epsilon(\tilde{\omega}_h) = I + II + III \quad (3.12)$$

where

$$I = \int |Du_h|^2(\cdot, h) - \int |D\tilde{u}_h|^2, \quad II = \int_{\tilde{\omega}_h \Delta \omega_h} 1 dx, \quad III = \epsilon \text{per}(\omega_h) - \epsilon \text{per}(\tilde{\omega}_h).$$

Here $\tilde{u}(x)$ solves $-\Delta\tilde{u} = \tilde{\lambda}$ with support $\tilde{\omega}_h$, where $\tilde{\lambda}$ is chosen such that $\int \tilde{u} = \int u_h(\cdot, h)$. We will show that

$$I \geq - \int_{\tilde{\omega}_h \Delta \omega_h} |D\varphi|^2(\cdot, h) dx \quad \text{and} \quad III \geq \int_{\tilde{\omega}_h \Delta \omega_h} \epsilon \kappa_\varphi dx$$

This proves our claim by (3.10), (3.12) and (3.11).

First let us estimate III :

$$\begin{aligned} \text{per}(\omega_h) - \text{per}(\tilde{\omega}_h) &\geq \int_{\partial\omega_h \setminus \partial\tilde{\omega}_h} \frac{D\varphi}{|D\varphi|}(\cdot, h) \cdot \eta dS - \int_{\partial\tilde{\omega}_h \setminus \partial\omega_h} \frac{D\varphi}{|D\varphi|}(\cdot, h) \cdot \tilde{\eta} dS \\ &= - \int_{\tilde{\omega}_h \Delta \omega_h} \nabla \cdot \left(\frac{D\varphi}{|D\varphi|} \right)(\cdot, h) dx \\ &= \int_{\tilde{\omega}_h \Delta \omega_h} \kappa_\varphi dx \end{aligned}$$

where $\tilde{\eta} = -D\varphi/|D\varphi|(x, h)$ is the outward normal vector at $x \in \partial\tilde{\omega}_h$, η is the outward normal vector at $x \in \partial\omega_h$, and κ_φ is the mean curvature of the level sets of φ .

It remains to estimate I . We again consider the two auxiliary functions, \bar{u} and v defined by (3.7). As before we have for $c := \int_{\omega_h} u_h(\cdot, h)$ and $\bar{c} := \int_{\tilde{\omega}_h} \bar{u}$:

$$\int_{\omega_h} |Du_h|^2(\cdot, h) - \int_{\tilde{\omega}_h} |D\bar{u}|^2 = \frac{\varepsilon}{\bar{c}} \lambda (\int_{\tilde{\omega}_h} \bar{u} - \int_{\omega_h} u_h(\cdot, h)). \quad (3.13)$$

But this time the inequality $(\min[\bar{u}, \varphi(\cdot, h)] - v)_+ \geq u_h(\cdot, h)$ holds on ω_h , as

$$\omega_h = \text{supp} (\min[\bar{u}, \varphi(\cdot, h)] - v)_+$$

and

$$-\Delta(\min[\bar{u}, \varphi(\cdot, h)] - v) \geq \lambda.$$

For the same reason we have for the outward normal η of $\tilde{\omega}_h$

$$\partial_\eta(u_h(\cdot, h) + v) \geq \partial_\eta \varphi(\cdot, h). \quad (3.14)$$

Thus, as $\min(\bar{u}, \varphi) = \bar{u}$ in $\tilde{\omega}_h$ and $\min(\bar{u}, \varphi) = \varphi$ in $\tilde{\omega}_h \Delta \omega_h$, using the smoothness of φ it follows that

$$\begin{aligned} \lambda (\int_{\tilde{\omega}_h} \bar{u} - \int_{\omega_h} u_h(\cdot, h)) &\geq \lambda \int_{\tilde{\omega}_h} v - \lambda \int_{\omega_h \Delta \tilde{\omega}_h} \varphi(\cdot, h) \\ &\geq - \int_{\tilde{\omega}_h} \Delta(u_h(\cdot, h) + v) v + \int_{\omega_h \Delta \tilde{\omega}_h} (\Delta \varphi \varphi)(\cdot, h) \\ &\quad + \int_{\omega_h \Delta \tilde{\omega}_h} (-\Delta \varphi - \lambda) \varphi(\cdot, h) \\ &\geq \int_{\tilde{\omega}_h} (D(u_h(\cdot, h) + v)) Dv - \int_{\partial \tilde{\omega}_h} \partial_\eta(u_h(\cdot, h) + v) v \\ &\quad - \int_{\omega_h \Delta \tilde{\omega}_h} |D\varphi|^2(\cdot, h) + \int_{\partial \tilde{\omega}_h} (\partial_\eta \varphi \varphi)(\cdot, h) + \int_{\omega_h \Delta \tilde{\omega}_h} o(h) \\ &\geq \int_{\omega_h} |Dv|^2 - \int_{\omega_h \Delta \tilde{\omega}_h} |D\varphi|^2(\cdot, h) + \int_{\omega_h \Delta \tilde{\omega}_h} o(h) \end{aligned}$$

by (3.14) and (3.7). Thus together with (3.13) we have

$$I \geq \frac{c}{\bar{c}} \int_{\omega_h \Delta \tilde{\omega}_h} |D\varphi|^2.$$

Lastly we need to show that

$$\bar{c} \rightarrow c \text{ as } |\tau - \tau_0| \rightarrow 0.$$

To see this, first note that $u_h(x, h) \leq (\phi(x, h) + \tau_0)_+$. In particular

$$u_h(\cdot, h) \leq C|\tau - \tau_0| \quad \text{on } \partial \tilde{\omega}_h - \partial \omega_h \subset \partial \{x : \phi(x, h) + \tau \geq 0\}$$

where C depends on the C^2 -norm of ϕ . It follows that $u_h(\cdot, h)|_{\omega_h} \leq \bar{u}_h + C|\tau - \tau_0|$, and therefore $c \leq \bar{c} + O(\tau - \tau_0)$. Hence we conclude. \square

4 The continuum limit and existence of weak solutions

In this section we show that in the limit $h \rightarrow 0$ and $\epsilon = h$ the time discrete solution u_h converges to a weak solution $u(\cdot, t) \in H^1(B)$ of (P) in the sense that the liminf and limsup -envelopes satisfy the barrier property at infinitesimal time scale (see Definition 4.4). We begin by defining viscosity sub- and supersolutions for a given multiplier function $\lambda(x, t) : B \times [0, \infty) \rightarrow [0, \infty)$.

Definition 4.1 A lower semi-continuous function $u : B \times [0, \infty) \rightarrow \mathbb{R}$ is a viscosity super-solution on $[t_1, t_2]$ with respect to $\lambda(x, t)$ if following holds:

For a given function $\phi \in C^{2,1}(\overline{\{\phi > 0\}})$ with $|D\phi| \neq 0$ in $B_r(x_0) \times [t_1, t_2]$, suppose that $\phi \leq u$ on the parabolic boundary of $B_r(x_0) \times [t_1, t_2]$ with

$$\begin{aligned} -\Delta\phi(\cdot, t) &< \lambda(\cdot, t) \quad \text{in } B_r(x_0) \times [t_1, t_2], \\ \frac{\phi_t}{|D\phi|} - (|D\phi|^2 - 1) &< 0 \quad \text{in } \Gamma(\phi) \cap \{B_r(x_0) \times [t_1, t_2]\}. \end{aligned}$$

Then $\phi \leq u$ in $B_r(x_0) \times [t_1, t_2]$.

For the subsolution part, in the context of our limit as $h \rightarrow 0$, we have to take into account the possibility that $\{u_h > 0\}$ leave thin segments or isolated points in the limit, which are not traceable from the limit of u_h . We get around this difficulty by including a set Σ in the definition:

Definition 4.2 Let $u : B \times [0, \infty) \rightarrow \mathbb{R}^+$ be upper semi-continuous, and let Σ be a closed subset of $B \times [0, \infty)$ containing $\Omega(u)$. Then the pair (u, Σ) is a viscosity sub-solution on $[t_1, t_2]$ with respect to $\lambda(x, t)$ if the following holds:

For a given function $\phi \in C^{2,1}(\overline{\{\phi > 0\}})$ with $|D\phi| \neq 0$ in $B_r(x_0) \times [t_1, t_2]$, suppose that

$$\begin{aligned} -\Delta\phi(\cdot, t) &> \lambda(\cdot, t) \quad \text{in } B_r(x_0) \times [t_1, t_2], \\ \frac{\phi_t}{|D\phi|} - (|D\phi|^2 - 1) &> 0 \quad \text{in } \Gamma(\phi) \cap \{B_r(x_0) \times [t_1, t_2]\}. \end{aligned}$$

If $u \leq \phi$ and $\Sigma \subset \Omega(\phi)$ on the parabolic boundary of $B_r(x_0) \times [t_1, t_2]$, then $u \leq \phi$ and $\Sigma \subset \Omega(\phi)$ in $B_r(x_0) \times [t_1, t_2]$.

Let us go back to the time discrete solutions u_h . Define

$$\mathcal{G} := \{k2^{-n} : k, n \in \mathbb{N}\} \text{ and } h = h(n) = 2^{-n}, n \in \mathbb{N}.$$

Then u_h is defined on grid times $t \in \mathcal{G}$ by (2.5), with the choice of $\epsilon = h$. Due to the Dirichlet energy bound, along a subsequence

$$u_h(\cdot, t) \rightarrow u(\cdot, t) \quad \text{weakly in } H^1(\mathbb{R}^N) \text{ for each } t \in \mathcal{G}. \quad (4.1)$$

We then choose a common subsequence of $h(n)$ such that (4.1) holds along the same sequence for each time. We obtain a weak form of convergence in the continuum limit $h \rightarrow 0$ along a subsequence.

Unfortunately a stronger, point-wise convergence of u_h cannot be obtained without extra regularity of u_h such as equicontinuity in time. Instead we consider the limit infimum and supremum:

$$u_*(x, t) := \lim_{r \rightarrow 0} \inf_{\{|x-y| \leq r, |s-t| \leq r, h \leq r\}} u_h(y, s) \quad (4.2)$$

and

$$u^*(x, t) := \lim_{r \rightarrow 0} \sup_{\{|x-y| \leq r, |s-t| \leq r, h \leq r\}} u_h(y, s). \quad (4.3)$$

Let us also define

$$\Sigma := \{(x, t) \mid \exists \text{ a sequence } (x_h, t_h) \rightarrow (x, t) \text{ such that } x_h \in \Omega(u_h(\cdot, t_h))\}. \quad (4.4)$$

Note that Σ contains $\Omega(u^*)$. Σ is a closed set, including ‘‘traces’’ of supports of $u_h(\cdot, t)$ which may degenerate into zero set of u^* in the limit $h \rightarrow 0$. Let us denote $\Sigma(s) := \Sigma \cap \{t = s\}$.

Next we define appropriate limits for the multipliers to be used for u^* and u_* .

Definition 4.3 For $\omega \subset B$, let us define

$$\lambda^{in}(\omega, c) := \lim_{\delta \rightarrow 0} \lambda(u_{\omega_\delta, c})$$

and

$$\lambda^{out}(\omega, c) := \lim_{\delta \rightarrow 0} \lambda(u_{\omega^\delta, c}),$$

with $\omega^\delta := \{x : d(x, \omega) \leq \delta\}$ and $\omega_\delta := \{x : B_\delta(x) \subset \omega\}$.

Clearly $\lambda^{in} \geq \lambda^{out}$, as $\omega_\delta \subset \omega^\delta$. Now we are ready to define our weak solution:

Definition 4.4 For functions $u_1, u_2 : B \times [0, \infty)$ and a closed set $\Sigma \subset B \times [0, \infty)$, the triple (u_1, u_2, Σ) is a weak solution of (P) if the following holds:

- (a) $u_1 \leq u_2$ and $\{u_2 > 0\} \subset \Sigma$;
- (b) u_1 is a viscosity supersolution with respect to $\lambda_1(x, t) := \lambda^{out}(\omega, c_1)$, where ω is the connected component of $\Omega_t(u_2)$ containing x and

$$c_1 := \int_\omega u_1(\cdot, t) dx.$$

- (c) (u_2, Σ) is a viscosity subsolution with respect to $\lambda_2(x, t) := \lambda^{in}(\omega, c_2)$, where ω is the connected component of $\Omega_t(u_1)$ containing x and

$$c_2 := \int_D u_2(\cdot, t) dx, \text{ where } D \text{ is the connected component of } \Omega_t(u_2) \text{ containing } \omega.$$

Roughly speaking, λ_1 and λ_2 are respectively the smallest and the largest possible value of the multiplier one can obtain by the lim sup and lim inf operation at a given point (x, t) . These definitions are tailored for $u_1 = u_*$ and $u_2 = u^*$.

Remark 4.5 *Classical solutions (i.e. $u \in C^{2,1}(\bar{\Omega}(u))$ with smooth $\Gamma(u)$ satisfying (P) in the classical sense), if they exist, would be weak solutions of (P) in our definition with $u_1 = u_2 = u$ and $\Sigma = \Omega(u)$.*

In the rest of the section we will show the following:

Theorem 4.6 *The triple (u_*, u^*, Σ) defined in (4.2) - (4.4) is a weak solution of (P).*

Remark 4.7 1. In [9] it was proven that starting from a star-shaped initial data, there is a unique star-shaped weak solution $(u, u, \Omega(u))$ of (P) for a short time, and for global time with additional symmetries in the initial data. Short-time existence of any nature for general smooth initial data is an open problem.

2. For free boundary problems which satisfy a comparison principle (such as the mean-curvature flow or (P) with fixed λ), the sub-solution would stay below the super-solution, which would then yield that $u^* \leq u_*$. This in turn yields $u^* = u_*$ and in particular the uniform convergence of u_h to a weak solution readily follows. Unfortunately for us this line of argument cannot be applied since (P) does not satisfy a comparison principle.

Proposition 4.8 *Let us define $\lambda_1(x, t)$ and $\lambda_2(x, t)$ as in Definition 4.4 with $u_1 = u_*$ and $u_2 = u^*$.*

(a) *Suppose $(x, t) \in \Sigma$. There is $x_h \in \Omega_{t_h}(u_h)$ such that $(x_h, t_h) \rightarrow (x, t)$. Let w^* be the connected component of $\Sigma(t)$ containing x and let w_h be the corresponding connected component containing x_h . Then*

$$\lambda_1(x, t) \leq \liminf_{h \rightarrow 0} \lambda(u_h(\cdot, t))(x)$$

(b) *Suppose $(x, t) \in \bar{\Omega}(u_*)$. Then there is $x_h \in \Omega_{t_h}(u_h)$ such that $(x_h, t_h) \rightarrow (x, t)$. Let w_* be the connected component of $\Omega(u_*)$ containing x , and let w_h be the corresponding connected component containing x_h . Then*

$$\limsup_{h \rightarrow 0} \lambda(u_h(\cdot, t))(x) \leq \lambda_2(x, t).$$

Proof. To prove (a), first note that for fixed δ we have that $u_h(\cdot, t_h)$ converges uniformly to zero outside of $\omega^{*,\delta} := \{x : d(x, \omega^*) \leq \delta\}$. Therefore, we can lower u_h to its essential part: there exists $\varepsilon_h \rightarrow 0$ such that $\tilde{u}_h := (u_h - \varepsilon_h)_+$ satisfies $\Omega_t(\tilde{u}_h) \subset \omega^{*,\delta}$. Moreover we have, by definition of c_1 ,

$$\liminf_{h \rightarrow 0} \int_{\omega^{*,\delta}} u_h dx \geq c_1.$$

Therefore,

$$\begin{aligned}
\lambda_1(x, t) &\leq \lambda(u_{\omega^*, \delta, c_1}(\cdot, t))(x) \\
&\leq \liminf_{h \rightarrow 0} \frac{\int_{\omega^*, \delta} |D\tilde{u}_h|^2(\cdot, t) dx}{c_1} \\
&= \liminf_{h \rightarrow 0} \lambda(u_h(\cdot, t))(x).
\end{aligned}$$

To prove (b), note that for any $\delta > 0$, there exists $h_0 < \delta$, such that $\omega_\delta := \{x : B_\delta(x) \subset \omega_*\}$ is contained in ω_{h_0} : Suppose $\omega_\delta \not\subset \omega_h$ for some $\delta > 0$ for any small $h > 0$. Then there exists a sequence of points x_h converging to a point \bar{x} in ω_δ such that $u_h(t_h, x_h) = 0$. This is a contradiction to the fact $\omega_\delta \subset \Omega_t(u_*)$. Therefore $\omega_\delta \subset \omega_h$ at least for a sequence of h converging to zero. Furthermore, by definition of c_2 ,

$$\limsup_{h \rightarrow 0} \int_{\omega_\delta} u_h(\cdot, t) dx \leq c_2.$$

And

$$\limsup_{h \rightarrow 0} \lambda(u_h(\cdot, t))(x) \leq \lim_{\delta \rightarrow 0} \lambda(u_{\omega_\delta, c_2})(x) \leq \lambda_2(x, t).$$

□

Proof of Theorem 4.6 The proof carries over the barrier properties of the time discrete solutions. We will only show that u_* is a viscosity supersolution of (P) with respect to $\lambda_1(x, t)$. The subsolution part can be shown via parallel arguments.

Suppose there exists a smooth function ϕ as in Definition 4.1 in $S := B_r(x_0) \times [t_1, t_0]$ such that ϕ crosses u_* from below at (x_0, t_0) : i.e. $u_* - \phi$ has a minimum zero at (x_0, t_0) . By using $\tilde{\phi}(x, t) := (\phi(x, t) - \sigma(x - x_0)^2 + \sigma(t - t_0))_+$ with small $\sigma > 0$ if necessary, one may assume that the minimum is strict in S . Then for small $h > 0$ the function $u_h - \phi$ also has a minimum at (x_h, t_h) in S with $(x_h, t_h) \rightarrow (x_0, t_0)$ as $h \rightarrow 0$. Since by Definition 4.1

$$-\Delta\phi(x, t) < \lambda_1(x, t),$$

we have that by Proposition 4.8(a) there exists $\delta > 0$ such that

$$-\Delta\phi < \lambda(u_h)(\cdot, t_0) \text{ in } B_\delta(x_0) \text{ for } 0 < h < \delta.$$

The above inequality as well as the second inequality in Definition 4.1 yield that ϕ satisfies (3.1) for h and r sufficiently small. Hence Proposition 3.1, applied to ϕ and u_h at (x_h, t_h) in S , yields a contradiction.

□

Acknowledgements

Natalie Grunewald was supported by the German Science Foundation, DFG, through fellowship GR 3391/1-1. Inwon Kim is supported by NSF DMS 0700732. The authors thank Karl Glasner for helpful discussions. Natalie Grunewald thanks UCLA for its hospitality during her stay in the last year, during which most of this paper was written.

References

- [1] F. Almgren, J.E. Taylor, L. Wang *Curvature driven flows, a variational approach*, SIAM J. Control and Optimization, **31** (1993) pp. 387–437.
- [2] I. Athanasopolous, L.A. Caffarelli, C. Keng, S. Salsa, *An Area–Dirichlet Integral Minimization Problem*, Comm. Pure and Applied Math., **LIV** (2001) pp. 479–499.
- [3] G. Belettini, V. Caselles, A. Chambolle and M. Novaga, *Crystalline Mean Curvature Flow of Convex Sets*, Arch. Rational Mech. Anal., **179** (2005) pp. 109–152.
- [4] A. Chambolle, *An algorithm for Mean Curvature Motion*, Interfaces and Free Boundaries, **6** (2004) pp.195–218.
- [5] S. Esedoglu and P. Smereka, *A variational formulation for a level set representation of multiphase flow and area preserving curvature flow*, Commun. Math. Sci., **6** (2008) pp.125–148.
- [6] L.C. Evans, R.F. Gariepy, *Measure Theory and Fine Properties of Functions*. CRC-Press, (1992).
- [7] L. C. Evans, J. Spruck, *Motion of level sets by mean curvature*, J. Differential Geometry, **33** (1991) pp. 635–681.
- [8] K. B. Glasner, *A boundary integral formulation of quasi-steady fluid wetting*, J. Comput. Phys., **207** (2005) pp. 529–541.
- [9] K. B. Glasner and I. C. Kim, *Global-time solutions for a model of contact line motion*, to appear in Interfaces and Free boundaries., **11** (2009) pp. 37–60.
- [10] H. P. Greenspan, *On the motion of a small viscous droplet that wets a surface*, J. Fluid Mechanics., **84** (1978) pp. 125–143.
- [11] M. Günther, G. Prokert, *On a Hele-Shaw-type Domain Evolution with convected Surface Energy Density*, SIAM J. Math. Anal., **37** (2005) pp. 372–410.
- [12] L. M. Hocking and M. J. Miksis, *Stability of a ridge of fluid*, J. Fluid Mech., **247** (1993) pp. 157–177.
- [13] R. Jordan, D. Kinderlehrer, and F. Otto. *The variational formulation of the Fokker-Planck equation*, SIAM J. Math. Anal., **29** (1998) pp.1–17.
- [14] I. C. Kim, *Uniqueness and Existence result of Hele-Shaw and Stefan problem*, Arch. Rat. Mech. Anal., **168** (2003) pp. 299–328.
- [15] S. Luckhaus, T. Sturzenhecker, *Implicit time discretization for the mean curvature flow equation*, Calc. Var. and PDE., **3** (1995) pp. 253–271.
- [16] F. Otto. *The geometry of dissipative evolution equations: the porous medium equation*, Comm. PDE., **26** (2001) pp.101–174.

- [17] A. Petrosyan and A.Yip, *Nonuniqueness in a free boundary problem from combustion*, J. Geom. Anal., **18** (2008) pp.10981126.
- [18] L. Tanner, *The spreading of silicone oil drops on horizontal surfaces*, J. Phys. D., **12** (1979) pp. 1473–1484.