

**Exercise #6, page 446:**

In the proof of necessary conditions for a local minimizer for non-linear minimization problems with linear inequality constraints (page 440) we needed the following linear algebra result:

*Let  $A$  be an  $m \times n$  matrix, with  $m \leq n$  and of rank  $m$  (all rows of  $A$  are linearly independent). Let  $e_1 = [1 \ 0 \ 0 \ \dots \ 0]^T \in R^m$  be the first vector of the canonical basis. Then there is a vector  $p \in R^n$  such that  $Ap = e_1$ .*

**Proof.** Let  $v_1, \dots, v_m$  be the  $m$  rows of the matrix  $A$ . Let

$$S = \text{Span}\{v_2, \dots, v_m\}$$

be the linear subspace of  $R^n$  spanned by the linearly independent vectors  $\{v_2, \dots, v_m\}$ . We know that  $R^n$  is an inner product space, and any vector  $v \in R^n$  can be written as  $v = p + q$ , with  $q \in S$  and  $p \in S^\perp$  (the orthogonal subspace to  $S$ ).

Therefore  $v_1 = p + q$ , with  $q \in S$ ,  $p \in S^\perp$ . We have that  $p \neq \vec{0}$  (indeed, if by contradiction  $p = \vec{0}$ , then  $v_1 = q \in S$ , impossible, because  $v_1$  is not a linear combination of  $v_2, \dots, v_m$ ).

We also have  $v_1 \cdot p = (p + q) \cdot p = p \cdot p + q \cdot p = p \cdot p$ . Let  $\hat{p} = \frac{p}{\|p\|^2}$ , with  $\|p\|^2 = p \cdot p$ . Then  $v_1 \cdot \hat{p} = v_1 \cdot \frac{p}{\|p\|^2} = 1$  and  $v_2 \cdot \hat{p} = 0, \dots, v_m \cdot \hat{p} = 0$ .

This implies that  $A\hat{p} = e_1$ .