

269C: HW #4

Due on Friday, June 3rd.

[1] Let K be a tetrahedron with vertices a^i , $i = 1, \dots, 4$, and let a^{ij} denote the midpoint on the straight line $a^i a^j$, $i < j$. Consider $v \in P_2(K)$ with the degrees of freedom $v(a^i)$, $v(a^{ij})$, $i, j = 1, \dots, 4$, $i < j$. Show that the corresponding finite element space V_h satisfies $V_h \subset C^0(\Omega)$, assuming continuity at all degrees of freedom.

[2] Let K be a triangle with vertices a^i , $i = 1, 2, 3$, and let a^{ij} , $i < j$, denote the midpoints of the sides of K . Let a^{123} denote the center of gravity of K . Consider $v \in P_4(K)$ with the degrees of freedom:

$$\begin{aligned} &v(a^i), \\ &\frac{\partial v}{\partial x_j}(a^i), \quad i = 1, 2, 3, \quad j = 1, 2, \\ &v(a^{ij}), \quad i, j = 1, 2, 3, \quad i < j, \\ &v(a^{123}), \quad \frac{\partial v}{\partial x_j}(a^{123}), \quad j = 1, 2, \end{aligned}$$

(typo in Exercise 3.8 in the textbook).

Show that the functions in the corresponding finite element space V_h are continuous, assuming continuity at all degrees of freedom.

[3] Consider the PDE (in the distributional sense)

$$-\Delta u + k^2 u = f \quad \text{in } R^n,$$

where k is a constant. Let $s \in R$. Show that, for all $f \in H^s(R^n)$, there exists a unique $u \in H^{s+2}(R^n)$, solution of the PDE, with $k \in R$, $k \neq 0$.

Hint: use the Fourier transform (see handout for notations of Sobolev spaces with arbitrary exponent s).

[4] Let $I = [0, h]$ and let $\pi v \in P_1(I)$ be the linear interpolant that agrees with $v \in C^2(I)$ at the end points of I . Using the technique of the proof of Thm. 4.1, prove estimates for $\|v - \pi v\|_{L^\infty(I)}$ and $\|v' - (\pi v)'\|_{L^\infty(I)}$, cf. (1.12) and (1.13).