

Math 270C: Assignment 6

Due Friday, March 3rd, 2006
(late homework accepted)

Instructor: Luminita Vese

[1] Consider the Krylov subspace $K(A, \vec{b}, p) = \text{span}\{\vec{b}, A\vec{b}, \dots, A^{p-2}\vec{b}, A^{p-1}\vec{b}\}$, where A is an n by n matrix. If A is non-singular and $A^p\vec{b} \in K(A, \vec{b}, p)$ then show

- (a) $A(K(A, \vec{b}, p)) \subseteq K(A, \vec{b}, p)$.
- (b) there exists a solution $\vec{x} \in K(A, \vec{b}, p)$ such that $A\vec{x} = \vec{b}$.

[2] Consider the second equivalent version of the conjugate gradient method, where A is symmetric and positive definite, as given in the handout Algorithm 10.2.1 [Conjugate Gradients], taken from Golub-Van Loan, page 527.

Assume now that A is only non-singular. Apply Algorithm 10.2.1 first to solve the system $AA^T y = b$, using \hat{p}_k for the directions in y_k . Then by change of variables $x_k = A^T y_k$ and $p_k = A^T \hat{p}_k$, recover the CGNE algorithm (see also Golub-Van Loan, pages 527 and 546).

[3] Implement the ADI method for the model problem from [4], assignment 3. Use again $h = 1/32$, and $\omega_{opt} = 2\frac{\sin(\pi h)}{h^2} \approx 200.7391$. Plot the error versus iterations. Comment your results.

(on the web page you will find a pseudo-code for “Symmetric, Tridiagonal, Positive Definite System Solver”)