

Connections with the finite dimensional case and with the KKT conditions

Example: Nonlinear inequality constrained optimization

Let $V = V^* = \mathbb{R}^n$, $Y = Y^* = \mathbb{R}^m$. Assume $f : \mathbb{R}^n \rightarrow \mathbb{R}$ convex and l.s.c. Let functions $c_1, \dots, c_m : \mathbb{R}^n \rightarrow \mathbb{R}$ be convex and l.s.c. Assume that there is a $u_0 \in \mathbb{R}^n$ such that $c_i(u_0) \leq 0$ for all $i = 1, \dots, m$. Denote by $c(u) = (c_1(u), \dots, c_m(u))$.

The primal problem is

$$\inf_{u \in \mathbb{R}^n, c_i(u) \leq 0, i=1, \dots, m} f(u).$$

The perturbation is defined by

$$\Phi(u, p) = \begin{cases} f(u) & \text{if } c(u) \leq p \\ +\infty & \text{otherwise} \end{cases},$$

or

$$\Phi(u, p) = \begin{cases} f(u) & \text{if } c_i(u) \leq p_i, i = 1, \dots, m \\ +\infty & \text{otherwise} \end{cases}.$$

Then $\Phi(u, p)$ can also be expressed as $\Phi(u, p) = f(u) + \chi_{E_p}(u)$, where χ_{E_p} is the indicator function of the set

$$E_p = \{u \in \mathbb{R}^n : c(u) \leq p\}.$$

It is possible to show that the dual problem is

$$\sup_{p^* \leq 0} \inf_{u \in \mathbb{R}^n} \left\{ -\langle p^*, c(u) \rangle + f(u) \right\}.$$

Also, it is possible to show that the Lagrangian defined by

$$-L(u, p^*) := \sup_{p \in \mathbb{R}^m} \{ \langle p^*, p \rangle - \Phi(u, p) \}$$

becomes

$$L(u, p^*) = f(u) - \sum_{i=1}^m p_i^* c_i(u) \text{ if } p_i^* \leq 0.$$

Then we have the KKT theorem:

Theorem: \bar{u} is a solution of the primal problem if and only if there is a $\bar{p}^* \in \mathbb{R}^m$, $\bar{p}^* \leq 0$ such that

$$L(\bar{u}, p) \leq L(\bar{u}, \bar{p}^*) \leq L(u, \bar{p}^*) \text{ for all } u \in \mathbb{R}^n, p \in \mathbb{R}^m, p \leq 0,$$

and the extremality relation holds $\langle \bar{p}^*, c(\bar{u}) \rangle = 0$, which implies that for all i , $1 \leq i \leq m$,

$$\text{either } \left\{ c_i(\bar{u}) < 0 \text{ and } p_i^* = 0 \right\} \text{ or } \left\{ c_i(\bar{u}) = 0 \text{ and } \bar{p}_i^* < 0 \right\}.$$

Remark: The usual notations in the finite dimensional case where $f(u) = f(x)$, for $u = x \in \mathbb{R}^n$, $p = \lambda \in \mathbb{R}^m$, and we substituted $C_i(x) \geq 0$ by $-C_i(x) = c_i(x) \leq 0$. Thus the Lagrange multiplier $\bar{p}^* = \lambda^*$ here is negative, while with $C_i(x) \geq 0$ we have $\bar{p}^* = \lambda^* \geq 0$.