

MOTIVIC DECOMPOSITION OF CERTAIN SPECIAL LINEAR GROUPS

ALEXANDER S. MERKURJEV

ABSTRACT. We compute the motive of the algebraic group $G = \mathbf{SL}_1(D)$ for a central simple algebra D of prime degree over a perfect field. As an application we determine certain motivic cohomology groups and differentials in the motivic spectral sequence of G .

1. INTRODUCTION

In this paper we study the motive in the triangulated category of geometric mixed effective motives $DM_{gm}^{eff}(F)$ over a perfect field F and the motivic cohomology of the algebraic group $\mathbf{SL}_1(D)$ of reduced norm 1 elements in a central simple algebra D of prime degree l .

In [18], A. Suslin computed the K -cohomology groups of the (split) special linear group \mathbf{SL}_n and the symplectic groups \mathbf{Sp}_{2n} using higher Chern classes in K -cohomology. O. Pushin in [15] constructed higher Chern classes in motivic cohomology and found decompositions of the motives of the groups \mathbf{SL}_n and \mathbf{GL}_n into direct sums of Tate motives. S. Biglari computed in [1] the motives of certain split reductive groups over \mathbb{Q} . In particular, he showed that

$$(1.1) \quad M(\mathbf{SL}_n)_{\mathbb{Q}} \simeq \coprod_{i=0}^{n-1} \mathrm{Sym}^i(\mathbb{Q}(2)[3] \oplus \mathbb{Q}(3)[5] \oplus \cdots \oplus \mathbb{Q}(n)[2n-1]).$$

A. Huber and B. Kahn determined the motives over \mathbb{Z} of split reductive groups in [9].

The motives of non-split algebraic groups are more complicated. The slices of the slice filtration of the motive $M(\mathbf{GL}_l(D))$ for a division algebra D of prime degree were computed by E. Shinder in [17].

In this paper we study the motive of the group $G = \mathbf{SL}_1(D)$, where D is a central simple algebra of a prime degree l . As a warm-up, consider the simplest case $l = 2$. The variety of G is then an open subscheme of a 3-dimensional projective isotropic quadric X given by the homogeneous quadratic equation $\mathrm{Nrd} = t^2$, where Nrd is the reduced norm form of D . The surface $Y = X \setminus G$, given by $\mathrm{Nrd} = 0$, is isomorphic to $S \times S$, where S is the Severi-Brauer variety

Key words and phrases. Central simple algebras, Severi-Brauer varieties, special linear groups, motives, motivic cohomology. *Mathematical Subject Classification (2010):* 20G15; 19E15, 14F42.

The work of the author has been supported by the NSF grant DMS #1160206 and the Guggenheim Fellowship.

of D (a conic curve in the case $l = 2$). Computing the motives of X and Y as in [17, §4], we get an exact triangle

$$M(G) \rightarrow \mathbb{Z} \oplus M(S)(1)[2] \oplus \mathbb{Z}(3)[6] \rightarrow M(S)(1)[2] \oplus M(S)(2)[4] \rightarrow M(G)[1].$$

Canceling out the summands $M(S)(1)[2]$, we obtain an isomorphism

$$(1.2) \quad M(G) \simeq \mathbb{Z} \oplus N(2)[3],$$

where the motive N is defined by the exact triangle

$$\mathbb{Z}(1)[2] \rightarrow M(S) \rightarrow N \rightarrow \mathbb{Z}(1)[3]$$

with the first morphism dual to the canonical one $M(S) \rightarrow \mathbb{Z}$.

In the general case, when l is an arbitrary prime, since the group G and its motive are split over a field extension of degree l , the torsion part of motivic cohomology of G is l -torsion. We work over the coefficient ring $\mathbb{Z}[\frac{1}{(l-1)!}]$, just inverting insignificant integers.

As in the case $l = 2$, the motive of G can be computed out of motive of the Severi-Brauer variety S of the algebra D . Let N be the motive defined by the exact triangle

$$\mathbb{Z}(l-1)[2l-2] \rightarrow M(S) \rightarrow N \rightarrow \mathbb{Z}(l-1)[2l-1].$$

As $(l-1)!$ is invertible in the coefficient ring, one can define symmetric $\mathbf{Sym}^i(M)$ and alternating powers $\mathbf{Alt}^i(M)$ of any motive M for $i = 0, 1, \dots, l-1$. The main result of the paper is the following theorem generalizing (1.1) and (1.2) (see Theorem 11.1).

Theorem. *Let D be a central simple algebra of prime degree l over a perfect field F . Then there is an isomorphism*

$$M(\mathbf{SL}_1(D)) \xrightarrow{\sim} \prod_{i=0}^{l-1} \mathbf{Sym}^i(N(2)[3]) = \prod_{i=0}^{l-1} (\mathbf{Alt}^i N)(2i)[3i]$$

in the category $DM_{gm}^{eff}(F)$ of motives over F with coefficients in $\mathbb{Z}[\frac{1}{(l-1)!}]$.

The most difficult part of the proof is the construction of a morphism $M(G) \rightarrow N(2)[3]$ in $DM_{gm}^{eff}(F)$. The main players of the proof are the groups $H^{3,2}(G) \simeq \mathbb{Z}$ and the Chow group $\mathbf{CH}^{l+1}(G) = H^{2l+2, l+1}(G) \simeq \mathbb{Z}/l\mathbb{Z}$ (when D is not split). These groups are related by a pair of homomorphisms

$$(1.3) \quad H^{3,2}(G) \leftarrow \mathbf{Hom}(M(G), N(2)[3]) \rightarrow H^{2l+2, l+1}(G).$$

We prove that there is a morphism $M(G) \rightarrow N(2)[3]$ with the images in (1.3) generating the two side cyclic groups. This is done in Section 10.

Using Theorem 1 and the exact triangle (Corollary 6.5)

$$(\mathbf{Alt}^{i-1} N)(l-1)[2l-2] \rightarrow \mathbf{Alt}^i M(S) \rightarrow (\mathbf{Alt}^i N) \rightarrow (\mathbf{Alt}^{i-1} N)(l-1)[2l-1],$$

we can compute inductively the motivic cohomology of G . As an application, in Section 12 we compute the motivic cohomology $H^{p,q}(G)$ with $2q - p \leq 1$. We also compute certain differentials in the motivic spectral sequence of G .

ACKNOWLEDGEMENTS. I thank Shahram Biglari and Bruno Kahn for helpful discussions.

2. MOTIVIC COHOMOLOGY

The base field F is assumed to be perfect. We fix a prime integer l and work over the *coefficient ring* \mathbf{Z} that is either the ring of integers \mathbb{Z} or $\mathbb{Z}[\frac{1}{(l-1)!}]$, or the localization $\mathbb{Z}_{(l)}$ of \mathbb{Z} by the prime ideal generated by l . Note that $\mathbf{Z}/l\mathbf{Z} = \mathbb{Z}/l\mathbb{Z}$.

We write $DM(F) := DM_{gm}^{eff}(F)$ for the *triangulated category of (geometric mixed effective) motives* with coefficients in \mathbf{Z} (see [19]). If p and $q \geq 0$ are integers, $\mathbf{Z}(q)[p]$ denotes the *Tate motive* and $M(X)$ the motive of a smooth variety X over F . We have $M(\text{Spec } F) = \mathbf{Z} := \mathbf{Z}(0)[0]$. For a motive M and an integer $q \geq 0$, we write $M(q)$ for $M \otimes \mathbf{Z}(q)$.

For a motive M in $DM(F)$ define the *motivic cohomology* by

$$H^{p,q}(M) := \text{Hom}(M, \mathbf{Z}(q)[p]),$$

where Hom is taken in the category $DM(F)$. If X is a smooth variety, simply write $H^{p,q}(X)$ for $H^{p,q}(M(X))$. We have

$$(2.1) \quad H^{p,q}(X) = 0 \quad \text{if } p > 2q \quad \text{or} \quad p > q + \dim(X).$$

In particular, $H^{p,q}(F) = 0$ if $p > q$. Moreover, $H^{p,p}(F) = K_p^M(F)$, the Milnor K -groups of F (see [12, Lecture 5]).

The bi-graded group $\coprod_{p,q} H^{p,q}(X)$ has a natural structure of a graded commutative ring (with respect to p , [12, Theorem 15.9]).

There is a canonical isomorphism between $H^{2p,p}(X)$ and the *Chow groups* $\text{CH}^p(X)$ of (rational equivalence) classes of algebraic cycles on a smooth variety X of codimension p ([12, Lecture 18]). We also write $\text{CH}^p(M) := H^{2p,p}(M)$ for every motive M .

The *cancelation* theorem (see [21]) states that the canonical morphism

$$\text{Hom}(M, N) \xrightarrow{\sim} \text{Hom}(M(1), N(1))$$

is an isomorphism for every two motives M and N .

The natural functor from the category of smooth projective varieties over F to $DM(F)$ extends uniquely to a canonical functor from the category $\text{Chow}(F)$ of Chow motives over F to $DM(F)$ (see [19, Proposition 2.1.4]). The motives in $DM(F)$ coming from $\text{Chow}(F)$ are called *pure* motives.

Let M be any motive and X a smooth projective variety of pure dimension d over F . The two canonical morphisms (given by the diagonal of X in the category of Chow motives)

$$\mathbf{Z}(d)[2d] \rightarrow M(X \times X) \rightarrow \mathbf{Z}(d)[2d]$$

together with the cancelation theorem define the two mutually inverse isomorphisms (see [9, Appendix B])

$$(2.2) \quad \text{Hom}(M, M(X)) \rightleftharpoons \text{Hom}(M \otimes M(X), \mathbf{Z}(d)[2d]) = \text{CH}^d(M \otimes M(X)).$$

In particular, if Y is another smooth projective variety, then

$$\mathrm{Hom}_{DM(F)}(M(Y), M(X)) \simeq \mathrm{CH}^d(Y \times X) = \mathrm{Hom}_{\mathrm{Chow}(F)}(M(Y), M(X)).$$

We say that a motive M is of *degree* d if M is a direct summand of a motive of the form $M(X)(q)[p]$ with $2q-p = d$, where X is a smooth projective variety. The pure motives are of degree 0. The following statement is an immediate consequence of (2.1).

Lemma 2.3. *Let M and N be motives of degree d and e respectively. If $d > e$, then $\mathrm{Hom}(M, N) = 0$. \square*

The *coniveau* spectral sequence for a smooth variety X over F ,

$$E_1^{p,q} = \coprod_{x \in X^{(p)}} H^{q-p, n-p} F(x) \Rightarrow H^{p+q, n}(X),$$

where $X^{(p)}$ is the set of points in X of codimension p , yields isomorphisms

$$H^{i+n, n}(X) \simeq A^i(X, K_n) \quad \text{when } n - i \leq 2$$

with the K -cohomology groups $A^i(X, K_n)$ defined in [16].

If X is a variety over F , we write X_{sep} for the variety $X \otimes_F F_{\mathrm{sep}}$ over a separable closure F_{sep} of F .

3. SEVERI-BRAUER VARIETIES

Let D be a central simple algebra of degree n over F and S the *Severi-Brauer* variety $\mathrm{SB}(D)$ of right ideals in D of rank n . This is a smooth projective variety of dimension $n - 1$ (see [6]). If D is *split*, i.e., $D = \mathrm{End}(V)$ for an n -dimensional vector space V over F , then S is isomorphic to the projective space $\mathbb{P}(V)$. Therefore, in the split case,

$$M(S) \simeq \mathbf{Z} \oplus \mathbf{Z}(1)[2] \oplus \cdots \oplus \mathbf{Z}(l-1)[2l-2].$$

Let $I \rightarrow S$ be the *tautological* vector bundle of rank n (with the fiber over a right ideal the ideal itself). We have $D = \mathrm{End}(I)^{op} = \mathrm{End}(I^\vee)$, where I^\vee is the vector bundle dual to I . In the split case, when $S = \mathbb{P}(V)$,

$$I = V^\vee \otimes L_t = \mathrm{Hom}(V, L_t),$$

where $L_t \rightarrow \mathbb{P}(V)$ is the tautological line bundle. The sheaf of sections of L_t is $\mathcal{O}(-1)$.

In the split case, when $S = \mathbb{P}(V)$, let $s \in \mathrm{CH}^1(S)$ be the class of a hyperplane section. We have

$$\mathrm{CH}^i(\mathbb{P}(V)) = \begin{cases} \mathbb{Z}s^i, & i = 0, 1, \dots, n-1; \\ 0, & \text{otherwise.} \end{cases}$$

The ring $\mathrm{End}(M(S)) = \mathrm{CH}^{n-1}(S \times S)$ is canonically isomorphic to the product \mathbf{Z}^n of n copies of \mathbf{Z} with the idempotents $s^i \times s^{n-1-i}$.

In the non-split case we have the following statement (see [13, Corollary 8.7.2]):

Lemma 3.1. *When D is a division algebra of prime degree l , the natural map $\mathrm{CH}^*(S) \rightarrow \mathrm{CH}^*(S_{\mathrm{sep}})$ is injective and it identifies the Chow group of S as follows*

$$\mathrm{CH}^i(S) = \begin{cases} \mathbf{Z} 1, & i = 0; \\ l \mathbf{Z} s^i, & i = 1, \dots, l-1; \\ 0, & \text{otherwise.} \end{cases} \quad \square$$

Any of the two projections $p : S \times S \rightarrow S$ is the projective bundle of I^\vee , i.e., $S \times S = \mathbb{P}_S(I^\vee)$. Let L be the tautological line bundle of this projective bundle. By the projective bundle theorem [19, Proposition 3.5.1], we have:

$$(3.2) \quad \mathrm{CH}^{l-1}(S \times S) = \mathrm{CH}^{l-1}(S) \cdot 1 \oplus \mathrm{CH}^{l-2}(S) \cdot \xi \oplus \cdots \oplus \mathrm{CH}^0(S) \cdot \xi^{l-1},$$

where ξ is the first Chern class of L in $\mathrm{CH}^1(S \times S)$. Consider the composition

$$\mathbf{type} : \mathrm{CH}^{l-1}(S \times S) = \mathrm{End} M(S) \rightarrow \mathrm{End} M(S_{\mathrm{sep}}) \xrightarrow{\sim} \mathbf{Z}^l.$$

Proposition 3.3. *Let D be a division algebra of degree l and $S = \mathrm{SB}(D)$. Then the ring homomorphism*

$$\mathbf{type} : \mathrm{End} M(S) \rightarrow \mathbf{Z}^l$$

is injective. Its image consists of all tuples (a_1, a_2, \dots, a_l) such that

$$a_1 \equiv a_2 \equiv \cdots \equiv a_l \pmod{l}.$$

Proof. It follows from Lemma 3.1 and (3.2) that \mathbf{type} is injective and $[\mathrm{Im}(\mathbf{type}) : l \mathbf{Z}^l] = l$. Therefore, the identity in \mathbf{Z}^l and $l \mathbf{Z}^l$ generate $\mathrm{Im}(\mathbf{type})$. \square

We will also need the following lemma.

Lemma 3.4. *Let M_1 and M_2 be direct sum of shifts of $M(S)$ (with arbitrary coefficients) and $f : M_1 \rightarrow M_2$ a morphism in $\mathrm{DM}(F)$. If f is an isomorphism over a field extension, then f is also an isomorphism.*

Proof. Write M_1 and M_2 as direct sums of the homogeneous (degree k) components $M_1^{(k)}$ and $M_2^{(k)}$ respectively. By Lemma 2.3, the morphism f is given by a triangular matrix with the diagonal terms $f_k : M_1^{(k)} \rightarrow M_2^{(k)}$. By assumption, the matrix is invertible over a splitting field L , hence all f_k are isomorphisms over L . Note that f_k is a shift of a morphism of pure motives that are direct sums of shifts of $M(S)$. By [4, Corollary 92.7], all f_k are isomorphisms. Therefore, the triangular matrix is invertible and hence f is an isomorphism. \square

4. THE MOTIVE N

Let D be a central simple algebra of prime degree l over F and $S = \mathrm{SB}(D)$. The motive N is defined by the triangle

$$(4.1) \quad \mathbf{Z}(l-1)[2l-2] \rightarrow M(S) \xrightarrow{\kappa} N \xrightarrow{\varepsilon} \mathbf{Z}(l-1)[2l-1]$$

in $DM(F)$ with the first morphism of pure motives given by the identity in $CH^0(S)$. We have

$$(4.2) \quad N_{\text{sep}} \simeq \mathbf{Z} \oplus \mathbf{Z}(1)[2] \oplus \cdots \oplus \mathbf{Z}(l-2)[2l-4],$$

therefore, $\text{Hom}(M(S_{\text{sep}}), N_{\text{sep}}) \simeq \mathbf{Z}^{l-1}$.

Consider the map

$$\mathbf{type} : \text{Hom}(M(S), N) \rightarrow \text{Hom}(M(S_{\text{sep}}), N_{\text{sep}}) \simeq \mathbf{Z}^{l-1}.$$

For example, $\mathbf{type}(\kappa) = (1, 1, \dots, 1)$.

Proposition 4.3. *Let D be a division algebra of degree l and $S = \text{SB}(D)$. Then the homomorphism*

$$\mathbf{type} : \text{Hom}(M(S), N) \rightarrow \mathbf{Z}^{l-1}$$

is injective. Its image consists of all tuples $(a_1, a_2, \dots, a_{l-1})$ such that

$$a_1 \equiv a_2 \equiv \cdots \equiv a_{l-1} \pmod{l}.$$

Proof. Let $\varphi \in \text{Ker}(\mathbf{type})$. The triangle (4.1) yields an exact sequence

$$\text{CH}^{l-1}(S) \rightarrow \text{End } M(S) \rightarrow \text{Hom}(M(S), N) \rightarrow H^{2l-1, l-1}(S).$$

The last term is zero as $2l-1 > 2(l-1)$. Therefore, $\varphi = \kappa \circ \sigma$ for some $\sigma \in \text{End } M(S)$. By assumption, $\mathbf{type}(\sigma) = (0, \dots, 0, a)$, where $a \equiv 0$ modulo l in view of Proposition 3.3. Then σ comes from $\text{CH}^{l-1}(S) = l\mathbf{Z}$ by Lemma 3.1 and hence $\varphi = 0$. This proves injectivity. The second statement follows from Proposition 3.3. \square

Lemma 4.4. *There is an isomorphism*

$$N \otimes M(S) \simeq M(S) \oplus M(S)(1)[2] \oplus \cdots \oplus M(S)(l-2)[2l-4].$$

In particular, $N \otimes M(S)$ is a pure motive.

Proof. The triangle (4.1) is split after tensoring with $M(S)$. Indeed, the morphism $M(S)(l-1)[2l-2] \rightarrow M(S) \otimes M(S)$ has a left inverse given by the class of the diagonal in $\text{CH}^{2l-2}(S \times S \times S)$. \square

Lemma 4.5. *We have $\text{CH}^i(N) = 0$ if $i > l$.*

Proof. In the exact sequence induced by (4.1)

$$H^{2i-2l+1, i-l+1}(F) \rightarrow \text{CH}^i(N) \rightarrow \text{CH}^i(S)$$

the first and the last terms are trivial as $2i-2l+1 > i-l+1$ and $\dim(S) < l$. \square

Since $\text{Hom}(\mathbf{Z}(q)[p], \mathbf{Z}) = 0$ if $q > 0$, the natural morphism $M(S) \rightarrow \mathbf{Z}$ factors uniquely through a morphism $\nu : N \rightarrow \mathbf{Z}$.

5. HIGHER CHERN CLASSES

Let X be a smooth variety. The higher Chern classes with values in motivic cohomology were constructed in [15]:

$$c_{j,i} : K_j(X) \rightarrow H^{2i-j,i}(X).$$

We will be using the classes

$$c_i := c_{1,i+1} : K_1(X) \rightarrow H^{2i+1,i+1}(X).$$

Proposition 5.1 ([17, §4.1]). *Let L be a vector bundle over a smooth variety X and $\alpha \in K_1(X)$. Then*

$$c_i(\alpha \cdot [L]) = \sum_{j=0}^i (-1)^j \binom{i}{j} c_{i-j}(\alpha) h^j,$$

where $h \in \text{CH}^1(X) = H^{2,1}(X)$ is the first (classical) Chern class of L .

Let $E \rightarrow X$ be a vector bundle of rank n . We write $\mathbf{SL}(E)$ for the group scheme over X of determinant 1 automorphisms of E .

Let a be the generic element of $\mathbf{SL}(E)$ (see [18, §4]). We also write a for the corresponding element in $K_1(\mathbf{SL}(E))$. We have $c_0(a) = 0$ since $\det(a) = 1$. For a sequence $\mathbf{i} = (i_1, i_2, \dots, i_k)$ with $1 \leq i_1 < i_2 < \dots < i_k \leq n-1$, set $d_{\mathbf{i}} = i_1 + i_2 + \dots + i_k$ and $e_{\mathbf{i}} = k$. Let

$$c_{\mathbf{i}}(\alpha) := c_{i_1}(\alpha) c_{i_2}(\alpha) \cdots c_{i_k}(\alpha) \in H^{2d_{\mathbf{i}}+e_{\mathbf{i}}, d_{\mathbf{i}}+e_{\mathbf{i}}}(\mathbf{SL}(E)).$$

Proposition 5.2. *Let $E \rightarrow X$ be a vector bundle of rank n . Then the $H^{*,*}(X)$ -module $H^{*,*}(\mathbf{SL}(E))$ is free with basis $\{c_{\mathbf{i}}(\alpha)\}$ over all sequences \mathbf{i} .*

Proof. This follows from [15, Proposition 3]. □

Write $\tilde{c}_{\mathbf{i}}(\alpha)$ for the composition

$$M(\mathbf{SL}(E)) \xrightarrow{\text{diag}} M(\mathbf{SL}(E)) \otimes M(\mathbf{SL}(E)) \xrightarrow{j \otimes c_{\mathbf{i}}(\alpha)} M(X)(d_{\mathbf{i}} + e_{\mathbf{i}})[2d_{\mathbf{i}} + e_{\mathbf{i}}],$$

where $j : M(\mathbf{SL}(E)) \rightarrow M(X)$ is the canonical morphism. The following corollary is deduced from Proposition 5.2 the same way as in [17, Proposition 4.2].

Corollary 5.3. (cf. [17, Proposition 4.4]) The morphisms $\tilde{c}_{\mathbf{i}}(\alpha)$ yield an isomorphism

$$M(\mathbf{SL}(E)) \xrightarrow{\sim} \coprod_{\mathbf{i}} M(X)(d_{\mathbf{i}} + e_{\mathbf{i}})[2d_{\mathbf{i}} + e_{\mathbf{i}}]. \quad \square$$

Remark 5.4. The natural composition

$$M(\mathbf{SL}(E)) \xrightarrow{\tilde{c}_{\mathbf{i}}(\alpha)} M(X)(d_{\mathbf{i}} + e_{\mathbf{i}})[2d_{\mathbf{i}} + e_{\mathbf{i}}] \rightarrow \mathbf{Z}(d_{\mathbf{i}} + e_{\mathbf{i}})[2d_{\mathbf{i}} + e_{\mathbf{i}}]$$

coincides with $c_{\mathbf{i}}(\alpha)$.

Corollary 5.5. *There is a canonical isomorphism*

$$M(\mathbf{SL}_n) \simeq \coprod_{\mathbf{i}} \mathbf{Z}(d_{\mathbf{i}} + e_{\mathbf{i}}) [2d_{\mathbf{i}} + e_{\mathbf{i}}]. \quad \square$$

Let $G = \mathbf{SL}(E)$ where $E \rightarrow X$ is a vector bundle of rank n over a smooth variety X . Consider the grading on $M(G)$ with respect to the value $e(\mathbf{i})$:

$$M(G)^{(k)} := \coprod_{e_{\mathbf{i}}=k} M(X)(d_{\mathbf{i}} + k) [2d_{\mathbf{i}} + k]$$

for $k = 0, 1, \dots, n-1$. Thus,

$$M(G) = \coprod_{k=0}^{n-1} M(G)^{(k)}$$

and each motive $M(G)^{(k)}$ has degree k .

Example 5.6. In the split case, we have a natural isomorphism $M(\mathbf{SL}_l)^{(1)} \simeq N(2)[3]$.

Let D be a central simple algebra of prime degree l over F and $G = \mathbf{SL}_1(D)$. Let S be the Severi-Brauer variety of D .

Corollary 5.3 yields

Corollary 5.7. *There is a canonical isomorphism*

$$M(G \times S) \simeq \coprod_{\mathbf{i}} M(S)(d_{\mathbf{i}} + e_{\mathbf{i}}) [2d_{\mathbf{i}} + e_{\mathbf{i}}].$$

In particular, $\mathrm{CH}^*(G \times S) \simeq \mathrm{CH}^*(S)$. \square

It follows from Corollary 5.5 that $M(G_{\mathrm{sep}})^{(1)} \simeq N_{\mathrm{sep}}(2)[3]$ and therefore $\mathrm{Hom}(M(G_{\mathrm{sep}}), M(S_{\mathrm{sep}})(2)[3])$ is naturally isomorphic to \mathbf{Z}^{l-1} . Consider the map

$$\mathrm{type} : \mathrm{Hom}(M(G), M(S)(2)[3]) \rightarrow \mathrm{Hom}(M(G_{\mathrm{sep}}), M(S_{\mathrm{sep}})(2)[3]) \simeq \mathbf{Z}^{l-1}.$$

By (2.2) and Corollary 5.7, we have

$$(5.8) \quad \mathrm{Hom}(M(G), M(S)(2)[3]) = H^{2l+1, l+1}(G \times S) = \prod_{i=1}^{l-1} \mathrm{CH}^i(S) c_{l-i}(\alpha).$$

Lemma 3.1 and (5.8) yield the following proposition.

Proposition 5.9. *Let D be a division algebra of degree l and $S = \mathrm{SB}(D)$. Then the homomorphism*

$$\mathrm{type} : \mathrm{Hom}(M(G), M(S)(2)[3]) \rightarrow \mathbf{Z}^{l-1}$$

is injective and $\mathrm{Im}(\mathrm{type}) = l \mathbf{Z}^{l-1}$. \square

We will need the Chow groups of G that were computed in [10].

Proposition 5.10. *Let D be a central division algebra of prime degree l and $G = \mathbf{SL}_1(D)$. There is an element $h \in \mathrm{CH}^{l+1}(G)$ such that*

$$\mathrm{CH}^*(G) = \mathbf{Z} \cdot 1 \oplus (\mathbf{Z}/l\mathbf{Z})h \oplus (\mathbf{Z}/l\mathbf{Z})h^2 \oplus \cdots \oplus (\mathbf{Z}/l\mathbf{Z})h^{l-1}. \quad \square$$

Recall that $D \simeq \mathrm{End}(I^\vee)$ and $G \times S \simeq \mathbf{SL}(I^\vee)$, where I is the tautological vector bundle over S of rank n (see Section 3).

Suppose that the algebra D is split. We can compare the generic matrices $\bar{\alpha}$ in $G = \mathbf{SL}_l$ and α in $G \times S = \mathbf{SL}(I^\vee)$. The bundle $I^\vee \otimes L_t$ over $G \times S$ is trivial, hence

$$\bar{\alpha} \times S = \alpha \otimes L_t \quad \text{in} \quad K_1(G \times S).$$

We have the Chern classes $c_i(\alpha) \in H^{2i+1, i+1}(G \times S)$ and $c_i := c_i(\bar{\alpha}) \in H^{2i+1, i+1}(G)$. We also write c_i for its image in $H^{2i+1, i+1}(G \times S)$ under the pull-back map given by the projection $G \times S \rightarrow G$.

By Proposition 5.1, we have

$$(5.11) \quad c_i = \sum_{j=0}^{i-1} \binom{i}{j} c_{i-j}(\alpha) s^j \quad \text{in} \quad H^{2i+1, i+1}(G \times S)$$

for all $i = 1, 2, \dots, l-1$, since the first Chern class of L_t is equal to $-s$, where $s \in \mathrm{CH}^1(S)$ is the class of a hyperplane section, and $c_0(\alpha) = 0$ as $\det(\alpha) = 1$. In particular, $c_1 = c_1(\alpha)$.

The group $H^{3,2}(G) = A^1(G, K_2)$ is infinite cyclic with a canonical generator, and this group does not change under field extensions. (This is true for every absolutely simple simply connected group, see [5, Part II, §9].) Therefore, we can write $H^{3,2}(G) = \mathbb{Z}c_1$ viewing c_1 as a generator of $H^{3,2}(G)$.

6. SYMMETRIC AND ALTERNATING POWERS

We consider motives with coefficients in $\mathbf{Z} = \mathbb{Z}[\frac{1}{(l-1)!}]$ in this section. Let $i = 0, 1, \dots, l-1$. The symmetric group Σ_i acts naturally on the i -th tensor power $M^{\otimes i}$ of a motive M . The elements

$$\tau_i = \frac{1}{i!} \sum_{\sigma \in \Sigma_i} \sigma \quad \text{and} \quad \rho_i = \frac{1}{i!} \sum_{\sigma \in \Sigma_i} \mathrm{sgn}(\sigma) \sigma$$

are idempotents in the group ring of Σ_i . The motives $\mathbf{Sym}^i(M) := (M, \tau_i)$ and $\mathbf{Alt}^i(M) := (M, \rho_i)$, that are split off M by the projectors τ_i and ρ_i , are called the i -th symmetric power and i -th alternating power of M respectively. We have $\mathbf{Sym}^0(M) = \mathbb{Z} = \mathbf{Alt}^0(M)$ and $\mathbf{Sym}^1(M) = M = \mathbf{Alt}^1(M)$.

We will need the following properties of symmetric and alternating powers.

Proposition 6.1 ([1, Proposition 2.3]). *Let M and N be two motives. Then*

- (1) $\mathbf{Sym}^i(M[1]) \simeq (\mathbf{Alt}^i M)[i]$ and $\mathbf{Alt}^i(M[1]) \simeq (\mathbf{Sym}^i M)[i]$,
- (2) $\mathbf{Sym}^i(M(q)) \simeq (\mathbf{Sym}^i M)(iq)$,
- (3) $\mathbf{Sym}^i(M \oplus N) = \coprod_{k+m=i} \mathbf{Sym}^k(M) \otimes \mathbf{Sym}^m(N)$ and similarly for \mathbf{Alt} .

Corollary 6.2. *We have*

$$\mathrm{Sym}^i(\mathbf{Z}(q)[p]) \simeq \begin{cases} \mathbf{Z}(iq)[ip], & \text{if } p \text{ is even;} \\ 0, & \text{if } p > 1 \text{ is odd.} \end{cases} \quad \square$$

Example 6.3. Let N be the motive $\mathbf{Z} \oplus \mathbf{Z}(1)[2] \oplus \cdots \oplus \mathbf{Z}(l-2)[2l-4]$ (see (4.2)). Then

$$\mathrm{Sym}^k(N(2)[3]) = (\mathrm{Alt}^k N)(2k)[3k] = \coprod_{e_i=k} \mathbf{Z}(d_i + k)[2d_i + k],$$

with the notation from Section 5.

Proposition 6.4 ([8, Proposition 15]). *Let $X \rightarrow Y \rightarrow Z \rightarrow X[1]$ be an exact triangle. Then there are sequences of morphisms*

$$\mathrm{Alt}^i X = T_0 \rightarrow T_1 \rightarrow \cdots \rightarrow T_i = \mathrm{Alt}^i Y,$$

$$\mathrm{Sym}^i X = V_0 \rightarrow V_1 \rightarrow \cdots \rightarrow V_i = \mathrm{Sym}^i Y$$

and exact triangles

$$T_{j-1} \rightarrow T_j \rightarrow \mathrm{Alt}^{i-j} X \otimes \mathrm{Alt}^j Z \rightarrow T_{j-1}[1],$$

$$V_{j-1} \rightarrow V_j \rightarrow \mathrm{Sym}^{i-j} X \otimes \mathrm{Sym}^j Z \rightarrow V_{j-1}[1]$$

for every $j = 1, 2, \dots, i$.

Assuming that $\mathrm{Alt}^k X = 0$ for $k > 1$, we get an exact triangle

$$X \otimes \mathrm{Alt}^{i-1} Z \rightarrow \mathrm{Alt}^i Y \rightarrow \mathrm{Alt}^i Z \rightarrow (X \otimes \mathrm{Alt}^{i-1} Z)[1].$$

Applying this to the exact triangle (4.1), we have the following proposition.

Corollary 6.5. *There is an exact triangle*

$$(\mathrm{Alt}^{i-1} N)(l-1)[2l-2] \rightarrow \mathrm{Alt}^i M(S) \rightarrow \mathrm{Alt}^i N \rightarrow (\mathrm{Alt}^{i-1} N)(l-1)[2l-1]. \quad \square$$

This proposition will be used in Section 12 to compute inductively the motivic cohomology of $\mathrm{Alt}^i N$.

The pure motive $\mathrm{Alt}^i M(S)$ is a direct summand of $M(S^i)$ and the latter is a direct sum of shifts of the motive $M(S)$. If D is a division algebra, the motive $M(S)$ is indecomposable [11, Corollary 2.22]. When the coefficient ring \mathbf{Z} is the local ring $\mathbb{Z}_{(l)}$, by uniqueness of the decomposition [3, Corollary 35], $\mathrm{Alt}^i M(S)$ is a pure motive that is a direct sum of pure shifts of $M(S)$. Moreover, since in the split case

$$\mathrm{Alt}^i M(S_{\mathrm{sep}}) = \mathbf{Z}((i(i-1)/2)[i(i-1)] \oplus \text{terms with higher shifts},$$

we must have

$$(6.6) \quad \mathrm{Alt}^i M(S) = M(S)((i(i-1)/2)[i(i-1)] \oplus \text{terms with higher shifts}.$$

7. SPLIT CASE

We are going to prove the main theorem in the split case. Let $G = \mathbf{SL}_l$ with prime l . We have

$$N = \mathbf{Z} \oplus \mathbf{Z}(1)[2] \oplus \mathbf{Z}(2)[4] \oplus \cdots \oplus \mathbf{Z}(l-2)[2l-4].$$

In fact, N is a direct summand of the motive of $S = \mathbb{P}^{l-1}$:

$$M(S) = N \oplus \mathbf{Z}(l-1)[2l-2].$$

The Chern classes $c_1(\bar{a}), c_2(\bar{a}), \dots, c_{l-1}(\bar{a})$ with values in the motivic cohomology of G , where \bar{a} is the generic matrix in \mathbf{SL}_l , define a morphism

$$\varphi_1 : M(G) \rightarrow N(2)[3].$$

For every $i = 0, 1, \dots, l-1$, consider the composition

$$\varphi_i : M(G) \xrightarrow{\text{diag}} M(G^i) \xrightarrow{\varphi^i} N(2)[3]^{\otimes i} \rightarrow \mathbf{Sym}^i(N(2)[3]),$$

where the first morphism is given by the diagonal embedding.

Proposition 7.1. *In the split case $G = \mathbf{SL}_l$, the morphism*

$$\varphi = (\varphi_i) : M(G) \rightarrow \prod_{i=0}^{l-1} \mathbf{Sym}^i(N(2)[3])$$

is an isomorphism.

Proof. By Corollary 5.5, Example 6.3 and Lemma 2.3, for every $i = 0, 1, \dots, l-1$, the morphism φ_i is zero on $M(G)^{(j)}$ for $j > i$ and yields an isomorphism

$$M(G)^{(i)} \xrightarrow{\sim} \mathbf{Sym}^i(N(2)[3]).$$

The result follows, as φ is given by an invertible triangular matrix. \square

8. COMPACTIFICATIONS OF G

Let D be a central division algebra over F and $G = \mathbf{SL}_1(D)$. By [2, §6.2] and [10, §6], G admits a smooth projective $G \times G$ -equivariant compactification X . In other words, X is a projective variety equipped with an action of $G \times G$ and containing G as an open orbit on which the group $G \times G$ acts by the left-right translations. The motive of X is split (i.e., $M(X)$ is a direct sum of Tate motives) over any field extension that splits D by [10, Theorem 6.5].

By Proposition 5.10, the group $\text{CH}^{l+1}(G)$ is cyclic of order l . Choose a generator $h \in \text{CH}^{l+1}(G)$ and let $\bar{h} \in \text{CH}^{l+1}(X)$ be any element such that $h|_G = h$. Set

$$(8.1) \quad R = \mathbf{Z} \oplus \mathbf{Z}(l+1)[2l+2] \oplus \mathbf{Z}(2l+2)[4l+4] \oplus \cdots \oplus \mathbf{Z}(l^2-1)[2l^2-2].$$

Proposition 8.2 ([10, §8]). *When $\mathbf{Z} = \mathbf{Z}_{(l)}$, the morphism $M(X) \rightarrow R$ defined by the powers of \bar{h} has a right inverse. Moreover, we have $M(X) \simeq R \oplus T$ for a pure motive T that is a direct sum of pure shifts of $M(S)$, where $S = \text{SB}(D)$.*

Note that the composition $\pi : M(G) \rightarrow M(X) \rightarrow R$ is given by the powers

$$h^i \in \mathrm{CH}^{i(p+1)}(G) = \mathrm{Hom}(M(G), \mathbb{Z}(i(p+1)[2i(p+1)]).$$

9. THE MORPHISM θ

In this section we construct a morphism $\theta : M(S)(2)[3] \rightarrow M(G)$.

As $M(G_{\mathrm{sep}})^{(1)} \simeq N_{\mathrm{sep}}(2)[3]$ by Example 5.6, there is a natural isomorphism

$$\mathrm{Hom}(M(S_{\mathrm{sep}})(2)[3], M(G_{\mathrm{sep}})^{(1)}) \simeq \mathbf{Z}^{l-1}.$$

Projecting $M(G_{\mathrm{sep}})$ onto $M(G_{\mathrm{sep}})^{(1)}$, we get a composition

$$\mathit{type} : \mathrm{Hom}(M(S)(2)[3], M(G)) \rightarrow \mathrm{Hom}(M(S_{\mathrm{sep}})(2)[3], M(G_{\mathrm{sep}})) \rightarrow \mathbf{Z}^{l-1}.$$

By Corollary 5.7, there is a canonical morphism

$$\theta : M(S)(2)[3] \rightarrow M(G \times S) \rightarrow M(G),$$

where the first morphism is the projection to the \mathbf{i} -component for $\mathbf{i} = (1)$ and the second morphism is given by the projection $G \times S \rightarrow G$.

Note that by definition of θ , the composition of $M(S)(2)[3] \rightarrow M(G \times S)$ with $\tilde{c}_k(\alpha) : M(G \times S) \rightarrow M(S)(k+1)[2k+1]$ is zero for $k > 1$. In view of (5.11) and Remark 5.4, the composition

$$M(S_{\mathrm{sep}})(2)[3] \xrightarrow{\theta_{\mathrm{sep}}} M(G_{\mathrm{sep}}) \xrightarrow{c_{\mathbf{i}}} \mathbf{Z}(i+1)[2i+1]$$

for $i = 1, 2, \dots, l-1$ coincides with $i = \binom{i}{i-1}$ times the composition

$$M(S_{\mathrm{sep}})(2)[3] \rightarrow M(G_{\mathrm{sep}} \times S_{\mathrm{sep}}) \xrightarrow{c_1 s^{i-1}} \mathbf{Z}(i+1)[2i+1].$$

The latter is equal to the morphism $s^{i-1} : M(S_{\mathrm{sep}})(2)[3] \rightarrow \mathbf{Z}(i+1)[2i+1]$ that is the identity on the summand $\mathbf{Z}(i+1)[2i+1]$. It follows that

$$\mathit{type}(\theta) = (1, 2, \dots, l-1).$$

10. A KEY LEMMA

Let D be a central simple algebra of degree l and $S = \mathrm{SB}(D)$.

Lemma 10.1. *Let Y be a variety over F such that D is split over the residue field $F(y)$ for every $y \in Y$. Then the push-forward homomorphism $\mathrm{CH}_j(Y \times S) \rightarrow \mathrm{CH}_j(Y)$ is surjective for every j .*

Proof. Let $y \in Y$ be a point of dimension j . As S is split over $F(y)$, there is a $F(y)$ -rational point $y' \in Y \times S$ in the fiber of the projection $q : Y \times S \rightarrow Y$ over y . We have $[y] = q_*([y'])$. \square

Lemma 10.2. *Let U be a smooth variety such that D is split over $F(u)$ for every $u \in U$. Then $H^{2j+1, j}(M(U) \otimes N) = 0$ for every j .*

Proof. The last term in the exact sequence induced by the triangle (4.1)

$$H^{2j,j}(U \times S) \rightarrow H^{2k,k}(U) \rightarrow H^{2j+1,j}(M(U) \otimes N) \rightarrow H^{2j+1,j}(U \times S),$$

where $k = j - l + 1$, is zero as $2j + 1 > 2j$. The first map is surjective by Lemma 10.1. \square

Let Y be a closed subvariety of a smooth variety X . We define the motive $M_Y(X)$ by the triangle

$$(10.3) \quad M(X \setminus Y) \rightarrow M(X) \rightarrow M_Y(X) \rightarrow M(X \setminus Y)[1].$$

Lemma 10.4. *Let X be a smooth irreducible variety and $Y \subset X$ an equidimensional closed subvariety such that S is split over $F(y)$ for every $y \in Y$. Then $H^{2i+1,i}(M_Y(X) \otimes N) = 0$ for every i .*

Proof. We proceed by induction on $\dim(Y)$. Choose a closed subset $Z \subset Y$ of pure codimension 1 such that $Y \setminus Z$ is smooth. In the exact triangle

$$M_{Y \setminus Z}(X \setminus Z) \rightarrow M_Y(X) \rightarrow M_Z(X) \rightarrow M_{Y \setminus Z}(X \setminus Z)[1]$$

the first term is isomorphic to $M(Y \setminus Z)(c)[2c]$ for $c = \text{codim}_X(Y)$ since $Y \setminus Z$ is smooth by [19, Proposition 3.5.4]. In the exact sequence

$$H^{2i+1,i}(M_Z(X) \otimes N) \rightarrow H^{2i+1,i}(M_Y(X) \otimes N) \rightarrow H^{2i+1,i}(M_{Y \setminus Z}(X \setminus Z) \otimes N)$$

the last term is isomorphic to $H^{2i-2c+1,i-c}(M(Y \setminus Z) \otimes N)$ which is zero by Lemma 10.2, and the first term is zero by induction. \square

Let X be a smooth $G \times G$ -equivariant compactification of $G = \mathbf{SL}_1(D)$ (see Section 8). Set $Y := X \setminus G$. By [10, Lemma 7.1], D is split by $F(y)$ for every $y \in Y$. Applying Lemma 10.4 to the exact cohomology sequence for the exact triangle (10.3) we get the following corollary.

Corollary 10.5. *The natural homomorphism*

$$\text{CH}^i(M(X) \otimes N) \rightarrow \text{CH}^i(M(G) \otimes N)$$

is surjective for every i . \square

Lemma 10.6. *The natural homomorphism*

$$\text{CH}^{l+1}(R \otimes N) \rightarrow \text{CH}^{l+1}(M(G) \otimes N)$$

induced by $\pi : M(G) \rightarrow R$ is surjective.

Proof. The group $\text{CH}^{l+1}(M(G) \otimes N)$ is l -torsion as it is split over a splitting field. Therefore, we may assume that $\mathbf{Z} = \mathbf{Z}_{(l)}$. Recall (see Section 8) that $M(X) = R \oplus T$, where R is defined in (8.1) and the pure motive T is a direct sum of shifts of $M(S)$. Moreover, π is the composition $M(G) \rightarrow M(X) \rightarrow R$.

Consider the commutative diagram

$$\begin{array}{ccccc}
\mathrm{CH}^{l+1}(R \otimes N) & \xrightarrow{\alpha} & \mathrm{CH}^{l+1}(M(G) \otimes N) & \longleftarrow & \mathrm{CH}^{2l}(M(G) \otimes N \otimes S) \\
& \searrow & \uparrow \beta & & \uparrow \\
& & \mathrm{CH}^{l+1}(M(X) \otimes N) & \longleftarrow & \mathrm{CH}^{2l}(M(X) \otimes N \otimes S) \\
& & \uparrow & & \uparrow \\
& & \mathrm{CH}^{l+1}(T \otimes N) & \xleftarrow{\gamma} & \mathrm{CH}^{2l}(T \otimes N \otimes S),
\end{array}$$

where γ (and the two homomorphisms above γ) are induced by the morphism $\mathbf{Z}(l-1)[2l-2] \rightarrow M(S)$. By Corollary 10.5, β is surjective. The pure motive T is a direct sum of shifts of $M(S)$, so is $T \otimes N$ by Lemma 4.4. It follows from Lemma 10.1 that γ is surjective.

Again by Lemma 4.4,

$$\mathrm{CH}^{2l}(M(G) \otimes N \otimes S) = \prod_{i=0}^{l-2} \mathrm{CH}^{2l-i}(G \times S) = 0.$$

as $\mathrm{CH}^j(G \times S) = 0$ for $j \geq l$ in view of Corollary 5.7. Recall that $M(X) = R \oplus T$. By diagram chase, α is surjective. \square

Consider the following key diagram:

$$\begin{array}{ccccccc}
\mathrm{Hom}(M(G), M(S)(2)[3]) & \xrightarrow{\xi} & \mathrm{Hom}(M(G), N(2)[3]) & \longrightarrow & \mathrm{CH}^{l+1}(G) & \longrightarrow & 0 \\
\downarrow \wr & & \downarrow \rho & & \downarrow \sigma & & \\
H^{2l+1, l+1}(G \times S) & \xrightarrow{\tau} & H^{3,2}(G) & \longrightarrow & \mathrm{CH}^{l+1}(M(G) \otimes N) & \longrightarrow & 0.
\end{array}$$

The rows of the diagram are induced by the exact triangle (4.1). The left vertical homomorphism is an isomorphism by (2.2). The rows are exact since $\mathrm{CH}^{l+1}(G \times S) \simeq \mathrm{CH}^{l+1}(S) = 0$ by Corollary 5.7. The morphisms ρ and σ are induced by the morphism $\nu : N \rightarrow \mathbf{Z}$ (see Section 4).

The diagram is commutative. Indeed, both compositions in the right square take a morphism $\varphi : M(G) \rightarrow N(2)[3]$ to $(\varepsilon(2)[3] \circ \varphi) \otimes \nu$. (The morphisms ε and ν are defined in Section 4.) Both compositions in the left square take a morphism $\psi : M(G) \rightarrow M(S)(2)[3]$ to $\nu(2)[3] \circ \psi$.

Now we can prove the following key lemma.

Lemma 10.7. *The homomorphism induced by the morphism $\nu : N \rightarrow \mathbf{Z}$*

$$\sigma : \mathrm{CH}^{l+1}(G) \rightarrow \mathrm{CH}^{l+1}(M(G) \otimes N)$$

is an isomorphism.

Proof. In the commutative diagram

$$\begin{array}{ccc} \mathrm{CH}^{l+1}(R) & \longrightarrow & \mathrm{CH}^{l+1}(R \otimes N) \\ \downarrow & & \downarrow \\ \mathrm{CH}^{l+1}(G) & \longrightarrow & \mathrm{CH}^{l+1}(M(G) \otimes N) \end{array}$$

the right vertical map (induced by π) is surjective by Lemma 10.6. We have $\mathrm{CH}^{l+1}(R) = \mathbf{Z}$ and by Lemma 4.5, $\mathrm{CH}^{l+1}(R \otimes N) = \mathrm{CH}^0(N) = \mathbf{Z}$, hence the top map is an isomorphism. It follows that the bottom map in the diagram is surjective. If D is split, the group $\mathrm{CH}^{l+1}(G)$ is trivial and we are done.

Suppose D is a division algebra. Since $H^{3,2}(G) = \mathbf{Z}c_1$ (see Section 5), by Proposition 5.9, the image of τ in the key diagram is equal to $l\mathbf{Z}c_1$. It follows that $\mathrm{CH}^{l+1}(M(G) \otimes N)$ is a cyclic group of order l . The group $\mathrm{CH}^{l+1}(G)$ is also cyclic of order l by Proposition 5.10. The statement follows from the surjectivity of σ . \square

It follows from Corollary 5.5 and Example 5.6 that $\mathrm{Hom}(M(G_{\mathrm{sep}}), N_{\mathrm{sep}}(2)[3])$ is naturally isomorphic to \mathbf{Z}^{l-1} . Consider the map

$$\mathbf{type} : \mathrm{Hom}(M(G), N(2)[3]) \rightarrow \mathrm{Hom}(M(G_{\mathrm{sep}}), N_{\mathrm{sep}}(2)[3]) \simeq \mathbf{Z}^{l-1}.$$

Proposition 10.8. *Let D be a division algebra of degree l and $S = \mathrm{SB}(D)$. Then the homomorphism*

$$\mathbf{type} : \mathrm{Hom}(M(G), N(2)[3]) \rightarrow \mathbf{Z}^{l-1}$$

is injective. Its image consists of all tuples $(a_1, a_2, \dots, a_{l-1})$ such that

$$a_1 \equiv 2a_2 \equiv \dots \equiv (l-1)a_{l-1} \pmod{l}.$$

Proof. Let $\beta \in \mathrm{Hom}(M(G), N(2)[3])$ have zero type. We have $\rho(\beta) = kc_1$ with k the first component of the type of β . Hence $k = 0$. It follows from Lemma 10.7 that the image of β in $\mathrm{CH}^{l+1}(G)$ is trivial, t.e., $\beta = \xi(\gamma)$ for some $\gamma \in \mathrm{Hom}(M(G), M(S)(2)[3])$ with $\mathbf{type}(\gamma) = 0$. By Proposition 5.9, $\gamma = 0$. This proves the injectivity of \mathbf{type} .

Take any $\beta \in \mathrm{Hom}(M(G), N(2)[3])$. We have

$$\mathbf{type}(\beta) = (a_1, a_2, a_3, \dots, a_{l-1})$$

for some $a_i \in \mathbf{Z}$. Composing β with $\theta : M(S)(2)[3] \rightarrow M(G)$ (see Section 9) we get a morphism $M(S)(2)[3] \rightarrow N(2)[3]$ of type $(a_1, 2a_2, 3a_3, \dots, (l-1)a_{l-1})$. By Proposition 4.3, we have $a_1 \equiv 2a_2 \equiv \dots \equiv (l-1)a_{l-1}$ modulo l .

By Proposition 5.9, the image of the map \mathbf{type} contains $l\mathbf{Z}^{l-1}$. To finish the proof it suffices to find a β such that $\mathbf{type}(\beta)$ is not divisible by l . By Lemma 10.7 and diagram chase, the map ρ is surjective. Hence there is a morphism $\beta : M(G) \rightarrow N(2)[3]$ such that the composition of β with $N(2)[3] \rightarrow \mathbf{Z}(2)[3]$ coincides with c_1 , i.e., $\mathbf{type}(\beta) = (1, \dots)$. \square

Remark 10.9. If $\alpha \in \text{Hom}(M(G), N(2)[3])$ is such that $\text{type}(\alpha)$ is not divisible by l , then $\rho(\alpha)$ is not divisible by l in $H^{3,2}(G) = \mathbf{Z}c_1$, and hence by Lemma 10.7, the image of α in $\text{CH}^{l+1}(G)$ is not zero if D is not split.

11. MAIN THEOREM

Now the coefficient ring is $\mathbf{Z} = \mathbb{Z}[\frac{1}{(l-1)!}]$. By Proposition 10.8, there is a unique morphism $\beta_1 : M(G) \rightarrow N(2)[3]$ with

$$\text{type}(\beta_1) = (1^{-1}, 2^{-1}, \dots, (l-1)^{-1}).$$

For every $i = 0, 1, \dots, l-1$ we have a composition

$$\beta_i : M(G) \xrightarrow{\text{diag}} M(G^i) \xrightarrow{\beta^i} N(2)[3]^{\otimes i} \rightarrow \text{Sym}^i(N(2)[3]).$$

Theorem 11.1. *Let D be a central simple algebra of prime degree l over a perfect field F . Then the morphism*

$$\beta = (\beta_i) : M(\mathbf{SL}_1(D)) \rightarrow \prod_{i=0}^{l-1} \text{Sym}^i(N(2)[3]) = \prod_{i=0}^{l-1} (\text{Alt}^i N)(2i)[3i]$$

in the category $DM(F)$ of motives over F with coefficients in $\mathbb{Z}[\frac{1}{(l-1)!}]$ is an isomorphism.

Proof. We first prove the theorem in the split case. The morphisms $\beta : M(G) \rightarrow N(2)[3]$ and $\varphi : M(G) \rightarrow N(2)[3]$ of type $(1, 1, \dots, 1)$ defined in Section 7 differ by an automorphism of $N(2)[3]$ of type $(1, 2, \dots, l-1)$. Therefore, the statement follows from Proposition 7.1.

Assume that D is a division algebra. We show next that $1_{M(S)} \otimes \beta$ is an isomorphism. By Corollary 5.7, the motive $M(S) \otimes M(G) = M(S \times G)$ is a direct sum of shifts of $M(S)$. The motive $M(S) \otimes N$ is a direct sum of shifts of $M(S)$ by Lemma 4.4, hence so is $M(S) \otimes (\text{Alt}^i N)$. By the first part of the proof, β is an isomorphism over a splitting field, hence so is $1_{M(S)} \otimes \beta$. By Lemma 3.4, $1_{M(S)} \otimes \beta$ is an isomorphism. It follows that

$$(11.2) \quad 1_{M(S^i)} \otimes \beta \quad \text{is an isomorphism for every } i > 0.$$

We embed the category $DM(F)$ into a larger triangulated category $DM_-^{eff}(F)$ of *motivic complexes* with coefficients in \mathbf{Z} as a full subcategory (see [19]).

Let $\check{C}(S)$ be the motive in $DM_-^{eff}(F)$ associated with the simplicial scheme given by the powers of S (see [20, Appendix B]). Using the exact triangle in the proof of [20, Proposition 8.1] we see from (11.2) that $1_{\check{C}(S)} \otimes \beta$ is an isomorphism.

It follows from Remark 10.9 that the composition

$$M(G) \xrightarrow{\beta_1} N(2)[3] \xrightarrow{\varepsilon(2)[3]} \mathbf{Z}(l+1)[2l+2]$$

represents a nontrivial element $h \in \text{CH}^{l+1}(G)$. Therefore, for every $i = 0, 1, \dots, l-1$, the composition

$$M(G) \xrightarrow{\beta_i} \text{Sym}^i(N(2)[3]) \xrightarrow{\delta_i} \text{Sym}^i(\mathbf{Z}(l+1)[2l+2]) = \mathbf{Z}(i(l+1))[2i(l+1)],$$

where $\delta_i = \mathbf{Sym}^i(\varepsilon(2)[3])$, is equal to h^i . By Section 8, we have a commutative diagram

$$\begin{array}{ccc} M(G) & \xrightarrow{\beta} & \coprod_{i=0}^{l-1} \mathbf{Sym}^i(N(2)[3]) \\ \downarrow \alpha & & \downarrow \delta \\ M(X) & \xrightarrow{\gamma} & R, \end{array}$$

where X is a smooth compactification of G and $\delta = \coprod \delta_i$.

Consider the motive $\tilde{C}(S)$ in $DM_-^{eff}(F)$ defined by the exact triangle

$$(11.3) \quad \tilde{C}(S) \rightarrow \check{C}(S) \rightarrow \mathbf{Z} \rightarrow \tilde{C}(S)[1].$$

We also have an exact triangle

$$M(G) \rightarrow M(X) \rightarrow M_Y(X) \rightarrow M(G)[1],$$

where $Y = X \setminus G$. The algebra D is split by the residue field $F(y)$ for every $y \in Y$ by [10, Lemma 7.1]. Hence, by [17, Lemma 3.4], $\tilde{C}(S) \otimes M_Y(X) = 0$. Therefore, $1_{\tilde{C}(S)} \otimes \alpha$ is an isomorphism.

By Proposition 8.2, $M(X) \simeq R \oplus T$, where T is a direct sum of shifts of $M(S)$ if $\mathbf{Z} = \mathbb{Z}_{(l)}$. Since $\tilde{C}(S) \otimes T = 0$, we have $1_{\tilde{C}(S)} \otimes \gamma$ is an isomorphism when $\mathbf{Z} = \mathbb{Z}_{(l)}$. As $\tilde{C}(S)$ vanishes over a splitting field of D of degree l , $1_{\tilde{C}(S)} \otimes \gamma$ is an isomorphism when $\mathbf{Z} = \mathbb{Z}[\frac{1}{(l-1)!}]$.

By Proposition 6.4 applied to the exact triangle (4.1), there is a Postnikov tower connecting $\mathbf{Sym}^i(N(2)[3])$ and

$$\mathbf{Sym}^i(\mathbf{Z}(l+1)[2l+2]) = \mathbf{Z}(i(l+1))[2i(l+1)]$$

with “factors” divisible by $M(S)$. Since $\tilde{C}(S) \otimes M(S) = 0$, the morphism

$$1_{\tilde{C}(S)} \otimes \mathbf{Sym}^i \varepsilon : \tilde{C}(S) \otimes \mathbf{Sym}^i(N(2)[3]) \xrightarrow{\sim} \tilde{C}(S)(i(l+1))[2i(l+1)]$$

is an isomorphism. Therefore, $1_{\tilde{C}(S)} \otimes \delta$ is an isomorphism.

It follows from the commutativity of the diagram that $1_{\tilde{C}(S)} \otimes \beta$ is an isomorphism. Finally, by 5-lemma applied to the exact triangle (11.3), the morphism β is an isomorphism. \square

12. APPLICATIONS

As an application of Theorem 11.1, we compute certain motivic cohomology of G . The Chow groups $\mathrm{CH}^i(G) = H^{2i,i}(G)$ are given in Proposition 5.10. In Theorem 12.4 below we compute the groups $H^{2i+1,i+1}(G)$.

The following Lemma is an immediate application of the exact triangle in Corollary 6.5.

Lemma 12.1. *If $p > 2q$, then*

$$H^{p,q}(\mathrm{Alt}^{i-1}N) \simeq H^{p+2l-1,q+l-1}(\mathrm{Alt}^i N). \quad \square$$

We compute the Chow groups of N .

Lemma 12.2. *We have*

$$\mathrm{CH}^i(N) = \begin{cases} \mathbf{Z}, & \text{if } i = 0; \\ l\mathbf{Z}, & \text{if } i = 1, 2, \dots, l-2; \\ F^\times / \mathrm{Nrd}(D^\times), & \text{if } i = l; \\ 0, & \text{otherwise.} \end{cases}$$

Proof. We may assume that $\mathbf{Z} = \mathbb{Z}$. Using (4.1) we get $\mathrm{CH}^i(N) \simeq \mathrm{CH}^i(S)$ for $i \leq l-2$ and apply Lemma 3.1. In the exact sequence

$$0 \rightarrow \mathrm{CH}^{l-1}(N) \rightarrow \mathrm{CH}^{l-1}(S) \rightarrow \mathrm{CH}^0(\mathbf{Z})$$

the last map is injective again by Lemma 3.1, hence $\mathrm{CH}^{l-1}(N) = 0$. In the exact sequence

$$H^{2l-1,l}(S) \rightarrow H^{1,1}(F) \rightarrow \mathrm{CH}^l(N) \rightarrow 0$$

the first map is isomorphic to $A^{l-1}(S, K_l) \rightarrow K_1^M(F) = F^\times$ and its image is equal to $\mathrm{Nrd}(D^\times)$ since the image is generated by the norms from finite field extensions that split D . By Lemma 4.5, $\mathrm{CH}^i(N) = 0$ if $i > l$. \square

Lemma 12.3. *We have*

$$H^{2i+1,i}(\mathrm{Alt}^2 N) = \begin{cases} \mathbf{Z}/l\mathbf{Z}, & \text{if } i = l-1; \\ F^\times / \mathrm{Nrd}(D^\times), & \text{if } i = 2l-1; \\ 0, & \text{otherwise.} \end{cases}$$

Proof. Using the triangle in Corollary 6.5, we get an exact sequence

$$\mathrm{CH}^i(\mathrm{Alt}^2 M(S)) \rightarrow \mathrm{CH}^{i-l+1}(N) \rightarrow H^{2i+1,i}(\mathrm{Alt}^2 N) \rightarrow 0.$$

The middle group is trivial if $i < l-1$, $l = 2l-2$ and $l > 2l-1$ by Lemma 12.2. The first map in the sequence is surjective in the split case since $\mathrm{Alt}^2 N$ is pure and $2i+1 > 2i$. As $\mathrm{CH}^{i-l+1}(N) = l\mathbf{Z}$ for $i = l, l+1, \dots, 2l-3$ by Lemma 12.2, the first map is also surjective in general for these values of i . If $i = 2l-1$, the first group is trivial as $\mathrm{Alt}^2 M(S)$ is a direct summand of $M(S \times S)$ and $\dim(S \times S) = 2l-2$.

Finally consider the case $i = l-1$. We may assume that $\mathbf{Z} = \mathbb{Z}_{(l)}$. As

$$\mathrm{Alt}^2 M(S) = M(S)(1)[2] \oplus M(S)(3)[6] \oplus \dots \oplus M(S)(l-2)[2l-4],$$

we have

$$\mathrm{CH}^{l-1}(\mathrm{Alt}^2 M(S)) = \mathrm{CH}^1(S) \oplus \mathrm{CH}^3(S) \oplus \dots \oplus \mathrm{CH}^{l-2}(S).$$

This is divisible by l when going to the split case by Lemma 3.1. Whence the case $i = l-1$. \square

Lemmas 12.1, 12.2 and 12.3 then yield

Theorem 12.4. *Let D be a central division algebra of degree l over F . Then*

$$H^{2i+1, i+1}(\mathbf{SL}_1(D)) = \begin{cases} F^\times, & \text{if } i = 0; \\ \mathbb{Z}c_1, & \text{if } i = 1; \\ l\mathbb{Z}c_i, & \text{if } i = 2, 3, \dots, l-1; \\ \mathbb{Z}/l\mathbb{Z}, & \text{if } i = k(l+1) + 1 \text{ for } k = 1, \dots, l-2; \\ F^\times / \text{Nrd}(D^\times), & \text{if } i = k(l+1) \text{ for } k = 1, \dots, l-1; \\ 0, & \text{otherwise.} \end{cases} \quad \square$$

Let $G = \mathbf{SL}_1(D)$. Note that the cup-product maps

$$F^\times \otimes \mathbb{Z}/l\mathbb{Z} = H^{1,1}(F) \otimes \text{CH}^{k(l+1)}(G) \rightarrow H^{2k(l+1)+1, k(l+1)+1}(G) = F^\times / \text{Nrd}(D^\times)$$

are natural surjections for $k = 1, \dots, l-1$.

Consider the motivic spectral sequence for G when D is not split (see [7]):

$$E_2^{p,q} = H^{p-q, -q}(G) \Rightarrow K_{-p-q}(G).$$

The K -groups of G were computed in [18, Theorem 6.1]. In particular, $K_0(G) = \mathbb{Z}$ and $K_1(G) = K_1(F) \oplus K_0(D) \oplus K_0(D^{op}) \simeq F^\times \oplus 3\mathbb{Z} \oplus 3\mathbb{Z}$. It follows that the zero-diagonal limit terms $E_\infty^{p,-p}$ are trivial if $p \neq 0$. On the other hand, by Proposition 5.10, we have

$$E_2^{p,-p} = \text{CH}^p(G) = \begin{cases} \mathbb{Z}, & \text{if } p = 0; \\ \mathbb{Z}/l\mathbb{Z}, & \text{if } p = i(l+1) \text{ for } i = 1, \dots, l-1; \\ 0, & \text{otherwise.} \end{cases}$$

Note that in the split case all the differentials coming to the zero diagonal are trivial. It follows that in the general case such differentials are l -torsion. By [14, Theorem 3.4], nontrivial differentials coming to the zero diagonal can appear only on pages $E_s^{*,*}$ with $l-1$ dividing $s-1$. It follows that the nonzero differentials appear only on page $E_l^{*,*}$ and they are

$$(12.5) \quad d : E_l^{1+k(l+1), -2-k(l+1)} \rightarrow E_l^{(k+1)(l+1), -(k+1)(l+1)} = \mathbb{Z}/l\mathbb{Z}$$

for $k = 0, 1, \dots, l-2$. These maps are all surjective and are isomorphisms for $k > 0$, thus “clearing” the zero diagonal and partially the first diagonal. The other differentials on the l -th page coming to the first diagonal are the cup-products with F^\times of the differentials (12.5). Nontrivial $E_\infty^{*,*}$ -terms on the first diagonal are F^\times and $l\mathbb{Z}$ ($l-1$ times).

Below is a fragment of the third page of the spectral sequence when $l = 3$.

$$\begin{array}{ccccccc}
 & & & & & & \mathbb{Z} \\
 & & & & & & \\
 F^\times & & 0 & & & & \\
 & & & & & & \\
 * & & \mathbb{Z} & & 0 & & \\
 & & \searrow & & \searrow & & \\
 & * & 3\mathbb{Z} & & 0 & & \\
 & & \searrow & & \searrow & & \\
 & & * & & 0 & & \mathbb{Z}/3\mathbb{Z} \\
 & & & & \searrow & & \\
 & & & & * & & F^\times/N & & 0 \\
 & & & & & & \\
 & & & & & & \\
 & & & & & * & \mathbb{Z}/3\mathbb{Z} & & 0 \\
 & & & & & \searrow & \searrow & & \\
 & & & & * & & 0 & & 0 \\
 & & & & \searrow & & \searrow & & \\
 & & & & * & & 0 & & \mathbb{Z}/3\mathbb{Z} \\
 & & & & & & \searrow & & \\
 & & & & & & * & & F^\times/N
 \end{array}$$

It follows from Theorem 12.4 that the Chern classes c_2, c_3, \dots, c_{l-1} (which are defined in the split case) are not defined over F if D is not split. (Recall that c_1 is always defined over F .) We will show that the product $c_1 c_2 \cdots c_k$ is defined over F for all $k = 1, 2, \dots, l-1$.

Lemma 12.6. *For every $i = 1, 2, \dots, l-1$, if $q < i(i-1)/2$, the group $H^{p,q}(\text{Alt}^i N)$ is trivial for every p .*

Proof. Induction on k . We may assume that $\mathbf{Z} = \mathbb{Z}_{(l)}$. The basic triangle (6.5) yields an exact sequence

$$H^{p-2l+1, q-l+1}(\text{Alt}^{i-1} N) \rightarrow H^{p,q}(\text{Alt}^i N) \rightarrow H^{p,q}(\text{Alt}^i M(S)).$$

The first term is trivial by induction as $q-l+1 < (i-1)(i-2)/2$. The last group is zero by (6.6). \square

Theorem 12.7. *Let $G = \mathbf{SL}_1(D)$ for a central simple algebra D of prime degree l . Then the product of Chern classes $c_1 c_2 \cdots c_k$ is defined over F for all $k = 1, 2, \dots, l-1$.*

Proof. We may assume that $\mathbf{Z} = \mathbb{Z}_{(l)}$. The product $c_1 c_2 \cdots c_k$ belongs to $H^{(k+1)^2-1, (k+1)(k+2)/2-1}(G)$. Consider the following direct summand of this group (see Theorem 11.1):

$$H^{(k+1)^2-1, (k+1)(k+2)/2-1}((\mathbf{Alt}^k N)(2k)[3k]) = \mathrm{CH}^{(k^2-k)/2}(\mathbf{Alt}^k N).$$

The basic triangle (6.5) yields an exact sequence

$$\begin{aligned} H^{k^2-k-2l+1, (k^2-k)/2-l+1}(\mathbf{Alt}^{k-1} N) &\rightarrow \mathrm{CH}^{(k^2-k)/2}(\mathbf{Alt}^k N) \rightarrow \\ &\mathrm{CH}^{(k^2-k)/2}(\mathbf{Alt}^k M(S)) \rightarrow \mathrm{CH}^{(k^2-k)/2-l+1}(\mathbf{Alt}^{k-1} N). \end{aligned}$$

The side terms are trivial by Lemma 12.6. The third term is isomorphic to $H^{0,0}(S) = \mathbf{Z}$ by (6.6). Therefore, the group $\mathrm{CH}^{(k^2-k)/2}(\mathbf{Alt}^k N)$ contains an element representing $c_1 c_2 \cdots c_k$ over a splitting field. \square

REFERENCES

- [1] S. Biglari, *Motives of reductive groups*, Amer. J. Math. **134** (2012), no. 1, 235–257.
- [2] M. Brion and S. Kumar, *Frobenius splitting methods in geometry and representation theory*, Progress in Mathematics, vol. 231, Birkhäuser Boston Inc., Boston, MA, 2005.
- [3] V. Chernousov and A. Merkurjev, *Motivic decomposition of projective homogeneous varieties and the Krull-Schmidt theorem*, Transform. Groups **11** (2006), no. 3, 371–386.
- [4] R. Elman, N. Karpenko, and A. Merkurjev, *The algebraic and geometric theory of quadratic forms*, American Mathematical Society, Providence, RI, 2008.
- [5] R. Garibaldi, A. Merkurjev, and Serre J.-P., *Cohomological invariants in galois cohomology*, American Mathematical Society, Providence, RI, 2003.
- [6] Ph. Gille and T. Szamuely, *Central simple algebras and Galois cohomology*, Cambridge Studies in Advanced Mathematics, vol. 101, Cambridge University Press, Cambridge, 2006.
- [7] D. Grayson, *The motivic spectral sequence*, Handbook of K-theory. Vol. 1, Springer, Berlin, 2005, pp. 39–69.
- [8] V. Guletskiĭ, *Finite-dimensional objects in distinguished triangles*, J. Number Theory **119** (2006), no. 1, 99–127.
- [9] A. Huber and B. Kahn, *The slice filtration and mixed Tate motives*, Compos. Math. **142** (2006), no. 4, 907–936.
- [10] N. Karpenko and A. Merkurjev, *Motivic decomposition of compactifications of certain group varieties*, arXiv:1402.5520 (2014).
- [11] N. Karpenko, *Upper motives of algebraic groups and incompressibility of Severi-Brauer varieties*, J. Reine Angew. Math. **677** (2013), 179–198.
- [12] C. Mazza, V. Voevodsky, and C. Weibel, *Lecture notes on motivic cohomology*, Clay Mathematics Monographs, vol. 2, American Mathematical Society, Providence, RI, 2006.
- [13] A. Merkurjev and A. Suslin, *K-cohomology of Severi-Brauer varieties and the norm residue homomorphism*, Izv. Akad. Nauk SSSR Ser. Mat. **46** (1982), no. 5, 1011–1046, 1135–1136.
- [14] A. Merkurjev, *Adams operations and the Brown-Gersten-Quillen spectral sequence*, Quadratic forms, linear algebraic groups, and cohomology, Dev. Math., vol. 18, Springer, New York, 2010, pp. 305–313.
- [15] O. Pushin, *Higher Chern classes and Steenrod operations in motivic cohomology*, K-Theory **31** (2004), no. 4, 307–321.
- [16] M. Rost, *Chow groups with coefficients*, Doc. Math. **1** (1996), No. 16, 319–393 (electronic).

- [17] E. Shinder, *On the motive of the group of units of a division algebra*, arXiv:1202.3090 (2014).
- [18] A. Suslin, *K-theory and K-cohomology of certain group varieties*, Algebraic K-theory, Adv. Soviet Math., vol. 4, Amer. Math. Soc., Providence, RI, 1991, pp. 53–74.
- [19] V. Voevodsky, *Triangulated categories of motives over a field*, Cycles, transfers, and motivic homology theories, Ann. of Math. Stud., vol. 143, Princeton Univ. Press, Princeton, NJ, 2000, pp. 188–238.
- [20] V. Voevodsky, *Motivic cohomology with $\mathbf{Z}/2$ -coefficients*, Publ. Math. Inst. Hautes Études Sci. (2003), no. 98, 59–104.
- [21] V. Voevodsky, *Cancellation theorem*, Doc. Math. (2010), no. Extra volume: Andrei A. Suslin sixtieth birthday, 671–685.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, LOS ANGELES, CA,
USA

E-mail address: merkurev at math.ucla.edu