

MIDTERM 2 (MATH 61, SPRING 2017)

Your Name: _____

UCLA id: _____

Math 61 Section: _____

Date: _____

The rules:

You MUST simplify completely and BOX all answers with an **INK PEN**.

You are allowed to use only this paper and pen/pencil. No calculators.

No books, no notebooks, no phones, no web access. You MUST write your name and UCLA id. Except for the last problem, you MUST write out your logical reasoning and/or proof in full. You have exactly 50 minutes.

Warning: those caught violating the rules get automatic 10% score deduction.

Points:

1 |

2 |

3 |

4 |

5 |

Total:

(out of 100)

Problem 1. (15 points)

Solve the following LHR: $a_{n+1} = 5a_n - 6a_{n-1}$, $a_1 = 9$, $a_2 = 20$.

Problem 2. (20 points)

Find the number of subgraphs of G isomorphic to H , where

- a) $G = C_{12}$, $H = P_5$.
- b) $G = K_{9,9}$, $H = C_6$.
- c) $G = K_{9,9}$, $H = C_7$.
- d) $G = H_3$ (the 3-cube), $H = C_4$.
- e) $G = H_3$ (the 3-cube), $H = P_4$.

Problem 3. (20 points)

For each of these sequences, either draw a simple graph with this score (degree sequence), or explain why there is no such graph.

a) $(3,3,3,1,1,1)$

b) $(4,4,4,1,1,1)$

c) $(5,5,5,3,2,1)$


d) $(4,4,\dots,4) \leftarrow 16 \text{ numbers.}$

e) $(3,3,3,3,2,2,2,2)$


Problem 4. (15 points)

Decide whether the following pairs of graphs on 8 vertices are isomorphic or non-isomorphic.

a)



b)



Important: In case of isomorphism, you must present a bijection. In case non-isomorphism, you must present an argument.

Problem 5. (30 points, 2 points each) **TRUE or FALSE?**

Circle correct answers with ink. No explanation is required or will be considered.

- T F** (1) The hypercube graph H_4 contains an Eulerian circuit.
- T F** (2) The hypercube graph H_4 contains a Hamiltonian cycle.
- T F** (3) A subgraph of a connected graph is always connected.
- T F** (4) A subgraph of a disconnected graph is always disconnected.
- T F** (5) The sum of degrees of $K_{\ell,\ell}$ is ℓ^2 .
- T F** (6) The sum of degrees of a simple graph on $n \geq 3$ vertices is $\leq n^2 - 1$.
- T F** (7) Computing the number of walks of given length in a graph can be done efficiently.
- T F** (8) Deciding whether a graph has Hamiltonian cycle can be done efficiently.
- T F** (9) The number of (shortest) grid walks $(0,0) \rightarrow (5,5)$ which do not go through $(1,3)$, $(3,4)$, $(4,1)$, $(1,5)$ is > 100 .
- T F** (10) Graph H is a subgraph of G . Graph H contains a Hamiltonian cycle. Then G contains a Hamiltonian cycle.
- T F** (11) Graph H is a subgraph of G . Graph G contains a Hamiltonian cycle. Then H contains a Hamiltonian cycle.
- T F** (12) Graph $K_{50,50}$ contains a subgraph isomorphic to K_{10} .
- T F** (13) Graph $K_{50,50}$ contains a subgraph isomorphic to $K_{10,10}$.
- T F** (14) The number of walks $1 \rightarrow 1$ of length k in a graph G on n vertices can be computed via matrix $(A_G)^n$.
- T F** (15) Isomorphic graphs have the same number of Eulerian circuits.