

HOMEWORK 3 (18.315, FALL 2005)

- 1) Let $G = K_{n_1 \dots n_r}$ be r -partite graph with parts n_1, \dots, n_r . Use the matrix-tree theorem to compute the number of spanning trees in G .
- 2) Problem 2.11 in Stanley, *EC1*. Use part b) to rederive the result of problem 1).
- 3) Problems 45, 46 from Bollobas, *MGT*, p. 376.
- 4) Let $G_{k,n}$ be the grid graph as before, and let $c(k, n, q)$ be the number of proper c -colorings of $G_{k,n}$ with q colors.
 - a) If k, n are fixed, prove that $c(k, n, q)$ can be computed in time polynomial in $(\log q)$.
 - b) If k, q are fixed, prove that $c(k, n, q)$ can be computed in time polynomial in n .
- 5) Let e_n be the expected number of cycles in a random permutation $\sigma \in S_n$. Compute e_n exactly. Conclude from here that $e_n = \theta(\log n)$.
- 6) Let $A(m)$ be the largest number of spanning trees a graph with m edges can have. Find non-trivial bounds on $A(m)$. (*Hint:* $A(m) \leq 2^m$ is a trivial bound. Similarly, a complete graph K_n gives a lower bound $A(m) \geq n^{n-2}$, where $m = \binom{n}{2}$. Can you do better?)

This Homework is due on Wednesday October 19 at 4 pm. in my office (2-390) or by e-mail.
P.S. Problem 2.11 in Stanley in the first edition is on trees and forests in graphs. Let me know if this is different in the new edition.