

OPTIMAL SINE AND SAWTOOTH INEQUALITIES

LOUIS ESSER, TERENCE TAO, BURT TOTARO, AND CHENGXI WANG

ABSTRACT. We determine the optimal inequality of the form $\sum_{k=1}^m a_k \sin kx \leq 1$, in the sense that $\sum_{k=1}^m a_k$ is maximal. We also solve exactly the analogous problem for the sawtooth (or signed fractional part) function. Equivalently, we solve exactly an optimization problem about equidistribution on the unit circle.

In this paper, we determine the optimal inequality of the form

$$\sum_{k=1}^m a_k \sin kx \leq 1$$

for each positive integer m , in the sense that $\sum_{k=1}^m a_k$ is maximal (Theorem 2.1). Namely, $\sum_{k=1}^m a_k$ is on the order of $\log m$, and we compute it exactly. This is a natural extremal problem in Fourier analysis. (We are considering inequalities that hold for all real numbers x .)

We also solve the analogous optimization problem for the *sawtooth* (or *signed fractional part*) function $g(x) = x + \lfloor \frac{1}{2} - x \rfloor$, which takes values in $(-1/2, 1/2]$. Namely, we find an optimal inequality of the form

$$\sum_{k=1}^m b_k g(kx) \leq 1$$

for each positive integer m , in the sense that $\sum_{k=1}^m b_k$ is maximal (Theorem 1.2). See the figures in sections 1 and 2 for examples of these inequalities, which show striking cancellation among dilated sine or sawtooth functions.

By linear programming duality, these inequalities are equivalent to statements about equidistribution on the unit circle. In particular: for each positive integer m and every probability measure on the real line, at least one of the dilated sawtooth functions $g(kx)$ for $k \in \{1, \dots, m\}$ must have small expected value, and we determine the optimal bound in terms of m (Theorem 1.2). It is on the order of $1/\log m$, and we compute it exactly.

These results were motivated by an application to algebraic geometry. For smooth complex projective varieties of general type, the *volume* is a positive rational number that measures the asymptotic growth of the plurigena $h^0(X, mK_X)$. Before the authors' series of papers in 2021, the varieties of general type with smallest known volume in high

2010 *Mathematics Subject Classification*. 42A05 (Primary) 11K06; 14E30; 14J40; 26D05 (Secondary).

dimensions n were those found by Ballico, Pignatelli, and Tasin, with volume roughly $1/n^n$ [1].

Using our equidistribution result for the sawtooth function, for any constant $c < 1$, version 1 of this paper on the arXiv constructed varieties of general type in all sufficiently high dimensions n with volume less than $1/e^{cn^{3/2}(\log n)^{1/2}}$. The equidistribution result was used to optimize the constant c .

Three of the authors then went further by different methods, finding n -dimensional varieties of general type with volume less than $1/2^{2^{n/2}}$ [4]. In view of that improvement, we have omitted the algebro-geometric application from this paper. There should be other ways to apply our optimization results for the sine and sawtooth functions.

0.1. Acknowledgments. LE and BT were supported by NSF grant DMS-2054553. TT was supported by a Simons Investigator grant, the James and Carol Collins Chair, the Mathematical Analysis & Application Research Fund Endowment, and by NSF grant DMS-1764034. Thanks to John Ottem, Sam Payne, and Miles Reid for their suggestions.

1. DILATED FRACTIONAL PARTS OF A RANDOM REAL NUMBER

In this section, we prove an optimal inequality for the sawtooth function (Theorem 1.2). By linear programming duality, this is equivalent to an optimal bound in a problem about equidistribution on the unit circle.

For a real number x , let $\lfloor x \rfloor$ denote the greatest integer less than or equal to x and $\lceil x \rceil$ the smallest integer greater than or equal to x . Note that

$$\lfloor x + n \rfloor = \lfloor x \rfloor + n \quad \text{and} \quad \lceil x + n \rceil = \lceil x \rceil + n \quad (1.1)$$

for any integer n , and that

$$\lceil x \rceil = -\lfloor -x \rfloor. \quad (1.2)$$

We also define the lower fractional part

$$\{x\} := x - \lfloor x \rfloor \quad (1.3)$$

which takes values in $[0, 1)$, and the upper fractional part

$$\{x\}^* := x - \lceil x \rceil + 1 \quad (1.4)$$

which takes values in $(0, 1]$. Finally, define the signed fractional part

$$g(x) := x - \left\lceil x - \frac{1}{2} \right\rceil = x + \left\lfloor \frac{1}{2} - x \right\rfloor \quad (1.5)$$

which takes values in $(-1/2, 1/2]$. We call $g(x)$ the *sawtooth function*.

For a (Borel) probability measure μ on the reals and a positive integer k , define the expectation

$$\mathbb{E}_\mu g(kx) := \int_{\mathbb{R}} g(kx) \, d\mu(x).$$

Consider the quantity

$$\min_{1 \leq k \leq m} \mathbb{E}_\mu g(kx), \quad (1.6)$$

where m is a natural number and μ is a probability measure on the reals. Since each function $g(kx)$ is pointwise bounded by $1/2$, we trivially have the bound

$$\min_{1 \leq k \leq m} \mathbb{E}_\mu g(kx) \leq \frac{1}{2}, \quad (1.7)$$

but one expects to do better as m gets large. For instance, from the Dirichlet approximation theorem one sees that

$$\min_{1 \leq k \leq m} |g(kx)| \leq \frac{1}{m+1},$$

but this does not directly allow one to improve the bound (1.7) since one cannot interchange the minimum and the expectation. As it turns out, there is an improvement in m , and the optimal value of (1.6) can be computed exactly, but it only decays like $\frac{1}{\log m}$ rather than $\frac{1}{m}$ as $m \rightarrow \infty$.

As a first attempt to control the quantity (1.6), one could try to estimate it by its unweighted mean

$$\frac{1}{m} \sum_{k=1}^m \mathbb{E}_\mu g(kx).$$

However, this quantity can be quite large: in particular, if μ is the Dirac mass at $1/2$, then this mean is equal to $\frac{\lceil m/2 \rceil}{2m}$, which is asymptotic to $\frac{1}{4}$ as $m \rightarrow \infty$. Closely related to this is the observation that the unweighted sum

$$g(x) + g(2x) + \cdots + g(mx)$$

of the $g(jx)$ can be much larger than 1, and in particular equal to $\lceil m/2 \rceil/2$ when $x = 1/2$.

However, one can obtain much better results by working with *weighted* means of the $\mathbb{E}_\mu g(kx)$, or equivalently by *weighted* linear combinations of the $g(jx)$; indeed by linear programming duality, we see that a bound of the form

$$\min_{1 \leq k \leq m} \mathbb{E}_\mu g(kx) \leq \lambda$$

for all μ holds if and only if there exist non-negative coefficients a_1, \dots, a_m with $\sum_{j=1}^m a_j \geq \frac{1}{\lambda}$ such that we have the dual inequality

$$\sum_{k=1}^m a_k g(kx) \leq 1$$

for all x . Thus to compute the minimal value of (1.6), we just need to find an optimal dual inequality.

We begin with the model case where m is a power of two, in which the dual inequality is particularly easy to establish.

Proposition 1.1. *Let r be a natural number, and set $m = 2^r$.*

(i) For every real number x , we have

$$2g(x) + \sum_{j=1}^r g(2^j x) \leq 1. \quad (1.8)$$

(ii) We have

$$\min_{1 \leq k \leq m} \mathbb{E}_\mu g(kx) \leq \frac{1}{r+2} \quad (1.9)$$

for every (Borel) probability measure μ on the real line. Moreover, this is the optimal bound: equality is attained for the measure μ_m with mass $\frac{1}{r+2}$ at each of the numbers $\frac{1}{2}, \frac{1}{4}, \frac{1}{8}, \dots, \frac{1}{2^r}$, and mass $\frac{2}{r+2}$ at $\frac{1}{2^{r+1}}$.

Proof. We begin with (i). We observe the identity

$$g(x) = \{2x\}^* - \{x\}^* \quad (1.10)$$

for all real numbers x , since both sides of this equation are 1-periodic, equal to x on $(0, 1/2]$, and equal to $x - 1$ on $(1/2, 1]$. Similarly,

$$2g(x) - \{2x\}^* = -1_{\{x\}^* > 1/2},$$

where the indicator function $1_{\{x\}^* > 1/2}$ is defined to equal 1 when $\{x\}^* > 1/2$ and vanish otherwise. Thus we have the telescoping formula

$$2g(x) + \sum_{j=1}^r g(2^j x) = \{2^{r+1}x\}^* - 1_{\{x\}^* > 1/2} \quad (1.11)$$

(see Figure 1.) This establishes (i).

Now we prove (ii). Integrating (i) against an arbitrary probability measure μ on \mathbb{R} , we conclude that

$$2\mathbb{E}_\mu g(x) + \sum_{j=1}^r \mathbb{E}_\mu g(2^j x) \leq 1.$$

Thus we have

$$\left(2 + \sum_{j=1}^r 1\right) \min_{1 \leq k \leq m} \mathbb{E}_\mu g(kx) \leq 1,$$

which gives the upper bound in (1.9).

To establish the matching lower bound using the measure μ_m , it suffices to show that

$$\sum_{j=1}^r g\left(\frac{k}{2^j}\right) + 2g\left(\frac{k}{2^{r+1}}\right) = 1$$

for all $k = 1, \dots, m$. But from (1.11) with $x = k/2^{r+1}$, the left-hand side is equal to

$$\{k\}^* - 1_{\{k/2^{r+1}\}^* > 1/2} = 1 - 0,$$

giving the claim. \square

Now we handle the general case.




FIGURE 1. The left-hand side of (1.8) when $m = 8$. We are showing that it is at most 1. Note the large number of locations where the bound of 1 is attained, including the support $\{1/2, 1/4, 1/8, 1/16\}$ of the optimal measure μ_8 .

Theorem 1.2. *Let r, m be natural numbers such that $2^r < m \leq 2^{r+1}$.*

(i) *For every real number x , we have*

$$2g(x) + \sum_{j=1}^r g(2^j x) + \sum_{\ell=2^r+1}^m \frac{m}{\ell(\ell-1)} \left(g(\ell x) + g((2^{r+1} + 1 - \ell)x) - g(x) \right) \leq 1. \quad (1.12)$$

(ii) *We have*

$$\min_{1 \leq k \leq m} \mathbb{E}_\mu g(kx) \leq \frac{2^r}{(r+1)2^r + m} \quad (1.13)$$

for every (Borel) probability measure μ on the real line. Moreover, this is the optimal bound: equality is attained for the measure μ_m with mass $\frac{2^r}{(r+1)2^r + m}$ at each of the numbers $\frac{1}{2}, \frac{1}{4}, \frac{1}{8}, \dots, \frac{1}{2^{r+1}}$ and mass $\frac{m}{(r+1)2^r + m}$ at $\frac{1}{2m}$.

In particular, the right side of (ii) is less than $\frac{\log 2}{\log m}$. So (ii) says in particular: for every probability measure μ on the real line and every positive integer m , there is a positive integer k at most m such that the expected value $\mathbb{E}_\mu g(kx)$ is at most $\frac{\log 2}{\log m}$. This is an equidistribution statement, sharpening the rough idea that the image measure of μ on \mathbb{R}/\mathbb{Z} under multiplication by some not-too-large positive integer is not concentrated too much in the first half of $[0, 1]$.




FIGURE 2. The left-hand side of (1.12) when $m = 7$. Note the large number of locations where the bound of 1 is attained, including the support $\{1/2, 1/4, 1/8, 1/14\}$ of the optimal measure μ_7 .

It follows from our argument (in particular the properties of the measure μ_m in (ii)) that statement (i) is an optimal inequality of the form $\sum_{k=1}^m a_k g(kx) \leq 1$, in the sense that it has the maximal value of $\sum a_k$ (namely, $r + 1 + \frac{m}{2^r}$) among all inequalities of this form. See Remark 1.3 on the non-uniqueness of this inequality.

Proof. By (1.11), we can rearrange (1.12) as

$$\sum_{\ell=2^{r+1}}^m \frac{m}{\ell(\ell-1)} \left(g(\ell x) + g((2^{r+1} + 1 - \ell)x) - g(x) \right) \leq 1 - \{2^{r+1}x\}^* + 1_{\{x\}^* > 1/2}.$$

Both sides are 1-periodic, so we may assume that $x \in (0, 1]$, and thus we may write

$$x = \frac{j - \theta}{2^{r+1}} \tag{1.14}$$

for some integer $1 \leq j \leq 2^{r+1}$ and some real number $0 \leq \theta < 1$. In particular, we have $\{2^{r+1}x\}^* = 1 - \theta$. From (1.5) we have

$$g(\ell x) + g((2^{r+1} + 1 - \ell)x) - g(x) = 2^{r+1}x + \left\lfloor \frac{1}{2} - \ell x \right\rfloor + \left\lfloor \frac{1}{2} + \ell x - 2^{r+1}x - x \right\rfloor - \left\lfloor \frac{1}{2} - x \right\rfloor.$$

But $-\left\lfloor \frac{1}{2} - x \right\rfloor = 1_{\{x\}^* > 1/2}$ since $x \in (0, 1]$, so we have

$$\sum_{\ell=2^{r+1}}^m \frac{m}{\ell(\ell-1)} \left(-\left\lfloor \frac{1}{2} - x \right\rfloor \right) \leq 1_{\{x\}^* > 1/2},$$

since

$$\sum_{\ell=2^{r+1}}^m \frac{m}{\ell(\ell-1)} = m \left(\frac{1}{2^r} - \frac{1}{m} \right) \leq 1. \quad (1.15)$$

So it will suffice to establish the inequality

$$\sum_{\ell=2^{r+1}}^m \frac{m}{\ell(\ell-1)} \left(2^{r+1}x + \lfloor \tfrac{1}{2} - \ell x \rfloor + \lfloor \tfrac{1}{2} + \ell x - 2^{r+1}x - x \rfloor \right) \leq \theta.$$

Writing $2^{r+1}x = j - \theta$, we can rearrange this using (1.1) as

$$\sum_{\ell=2^{r+1}}^m \frac{m}{\ell(\ell-1)} \left(-\theta + \lfloor \tfrac{1}{2} - \ell x \rfloor + \lfloor \tfrac{1}{2} + \ell x + \theta - x \rfloor \right) \leq \theta.$$

By (1.15), it is equivalent to show that

$$\sum_{\ell=2^{r+1}}^m \frac{m}{\ell(\ell-1)} \left(\lfloor \tfrac{1}{2} - \ell x \rfloor + \lfloor \tfrac{1}{2} + \ell x + \theta - x \rfloor \right) \leq \frac{m\theta}{2^r}.$$

We may cancel the factor of m on both sides. The quantity $\lfloor \tfrac{1}{2} - \ell x \rfloor + \lfloor \tfrac{1}{2} + \ell x + \theta - x \rfloor$ is clearly an integer that is bounded above by

$$\left(\tfrac{1}{2} - \ell x \right) + \left(\tfrac{1}{2} + \ell x + \theta - x \right) = 1 + \theta - x < 2,$$

and by (1.3) it is equal to 1 if and only if

$$\left\{ \tfrac{1}{2} - \ell x \right\} + \left\{ \tfrac{1}{2} + \ell x + \theta - x \right\} = \theta - x,$$

so in particular

$$0 \leq \left\{ \tfrac{1}{2} - \ell x \right\} \leq \theta - x.$$

Thus it will suffice to show that

$$\sum_{2^r < \ell \leq m: 0 \leq \{ \frac{1}{2} - \ell x \} \leq \theta - x} \frac{1}{\ell(\ell-1)} \leq \frac{\theta}{2^r}. \quad (1.16)$$

We can write the left-hand side as

$$\sum_{k=0}^{\infty} \sum_{2^r < \ell \leq m: -k \leq \frac{1}{2} - \ell x \leq -k + \theta - x} \frac{1}{\ell(\ell-1)}$$

or equivalently

$$\sum_{k=0}^{\infty} \sum_{\max(2^r+1, \frac{k-\theta+1/2}{x}+1) \leq \ell \leq \min(\frac{k+1/2}{x}, m)} \frac{1}{\ell(\ell-1)}. \quad (1.17)$$

Using the fundamental theorem of calculus to write $\frac{1}{\ell(\ell-1)} = \int_{\ell-1}^{\ell} \frac{dt}{t^2}$, we may upper bound this expression by

$$\sum_{k=0}^{\infty} \int_{[2^r, m] \cap [\frac{k-\theta+1/2}{x}, \frac{k+1/2}{x}]} \frac{dt}{t^2}$$

and so (using $m \leq 2^{r+1}$) it will suffice to establish the bound

$$\sum_{k=0}^{\infty} \int_{[2^r, 2^{r+1}] \cap [\frac{k-\theta+1/2}{x}, \frac{k+1/2}{x}]} \frac{dt}{t^2} \leq \frac{\theta}{2^r}. \quad (1.18)$$

The interval $[2^r, 2^{r+1}] \cap [\frac{k-\theta+1/2}{x}, \frac{k+1/2}{x}]$ is only non-empty when

$$\frac{k-\theta+1/2}{x} \leq 2^{r+1} \quad \text{and} \quad \frac{k+1/2}{x} \geq 2^r.$$

Hence we may restrict the k summation in (1.18) to the region

$$2^r x - \frac{1}{2} \leq k \leq 2^{r+1} x - \frac{1}{2} + \theta.$$

By (1.14) and the fact that $0 \leq \theta < 1$, we conclude that

$$\frac{j}{2} - 1 < k < j. \tag{1.19}$$

We now split into cases.

Case 1: $j = 1$. By (1.19), the only value of k that contributes to (1.18) is $k = 0$, and we can upper bound the left-hand side of (1.18) by

$$\int_{[2^r, \frac{1/2}{x}]} \frac{dt}{t^2} = \frac{1}{2^r} - 2x$$

which is precisely $\frac{\theta}{2^r}$ as desired thanks to (1.14).

Case 2: $j = 2$. Now (1.19) restricts us to $k = 1$, and we can upper bound the left-hand side of (1.18) by

$$\int_{[\frac{3/2-\theta}{x}, \frac{3/2}{x}]} \frac{dt}{t^2} = \frac{x}{3/2-\theta} - \frac{x}{3/2}$$

which by (1.14) simplifies to

$$\frac{2}{3} \cdot \frac{2-\theta}{3-2\theta} \cdot \frac{\theta}{2^r}.$$

Since $\frac{2}{3}$ and $\frac{2-\theta}{3-2\theta}$ are at most 1, we obtain the desired upper bound of $\frac{\theta}{2^r}$ (with a little room to spare).

Case 3: $j = 3$. Now (1.19) restricts us to $k = 1, 2$. The left-hand side of (1.18) is now upper bounded by

$$\int_{[2^r, \frac{3/2}{x}]} \frac{dt}{t^2} + \int_{[\frac{5/2-\theta}{x}, \frac{5/2}{x}]} \frac{dt}{t^2} = \frac{1}{2^r} - \frac{x}{3/2} + \frac{x}{5/2-\theta} - \frac{x}{5/2}$$

which by (1.14) simplifies to

$$\left(\frac{1}{3} + \frac{2}{5} \cdot \frac{3-\theta}{5-2\theta} \right) \frac{\theta}{2^r}.$$

Since $0 \leq \theta < 1$, we have $\frac{1}{3} + \frac{2}{5} \cdot \frac{3-\theta}{5-2\theta} < \frac{1}{3} + \frac{2}{5} \cdot \frac{2}{3} < 1$, and we obtain the desired upper bound of $\frac{\theta}{2^r}$ (with a little more room to spare).

Case 4: $j = 4$. Now (1.19) restricts us to $k = 2, 3$. The left-hand side of (1.18) is now upper bounded by

$$\int_{[\frac{5/2-\theta}{x}, \frac{5/2}{x}]} \frac{dt}{t^2} + \int_{[\frac{7/2-\theta}{x}, \frac{7/2}{x}]} \frac{dt}{t^2} = \frac{x}{5/2-\theta} - \frac{x}{5/2} + \frac{x}{7/2-\theta} - \frac{x}{7/2}$$

which by (1.14) simplifies to

$$\left(\frac{2}{5} \cdot \frac{4-\theta}{5-2\theta} + \frac{2}{7} \cdot \frac{4-\theta}{7-2\theta} \right) \frac{\theta}{2^r}.$$

Since $0 \leq \theta < 1$, we have

$$\frac{2}{5} \cdot \frac{4-\theta}{5-2\theta} + \frac{2}{7} \cdot \frac{4-\theta}{7-2\theta} < \frac{2}{5} \cdot \frac{3}{3} + \frac{2}{7} \cdot \frac{3}{5} < 1$$

and we again obtain the desired upper bound of $\frac{\theta}{2^r}$ (with a fair amount¹ of room to spare).

Case 5: $j > 4$. There is a finite interval $[k_1, k_2]$ of integers k for which $[2^r, m] \cap [\frac{k-\theta+1/2}{x}, \frac{k+1/2}{x}]$ is non-empty. From the decreasing nature of $\frac{1}{t^2}$, we have

$$\int_{[2^r, 2^{r+1}] \cap [\frac{k-\theta+1/2}{x}, \frac{k+1/2}{x}]} \frac{dt}{t^2} \leq \theta \int_{[2^r, 2^{r+1}] \cap [\frac{k-1+1/2}{x}, \frac{k+1/2}{x}]} \frac{dt}{t^2}$$

when $k_1 < k \leq k_2$. Thus we may bound the left-hand side of (1.18) by

$$\int_{[2^r, 2^{r+1}] \cap [\frac{k_1-\theta+1/2}{x}, \frac{k_1+1/2}{x}]} \frac{dt}{t^2} + \theta \int_{[2^r, 2^{r+1}]} \frac{dt}{t^2}.$$

For the first integral, we observe that the domain is an interval of length at most θ/x and the integrand is at most $1/2^{2r}$; meanwhile, the second integral can be evaluated as $\frac{1}{2^r} - \frac{1}{2^{r+1}}$. Putting all this together, we have upper bounded the left-hand side of (1.18) by

$$\frac{\theta}{2^{2r}x} + \frac{\theta}{2^r} - \frac{\theta}{2^{r+1}}.$$

But since $j > 4$, we have from (1.14) that

$$\frac{\theta}{2^{2r}x} \leq \frac{\theta}{2^{2r} \cdot 4/2^{r+1}} = \frac{\theta}{2^{r+1}}$$

and the claim (1.18) follows. This concludes the proof of (i).

Now we prove (ii). Integrating (i) against an arbitrary probability measure μ on \mathbb{R} , we conclude that

$$\begin{aligned} 2\mathbb{E}_\mu g(x) + \sum_{j=1}^r \mathbb{E}_\mu g(2^j x) \\ + \sum_{\ell=2^{r+1}}^m \frac{m}{l(l-1)} \left(\mathbb{E}_\mu g(\ell x) + \mathbb{E}_\mu g((2^{r+1} + 1 - \ell)x) - \mathbb{E}_\mu g(x) \right) \leq 1. \end{aligned}$$

Since

$$\sum_{\ell=2^{r+1}}^m \frac{m}{l(l-1)} = \frac{m}{2^r} - 1 < 2, \tag{1.20}$$

¹Note that the increasing ease of proof of (1.12) as j increases is consistent with the behavior exhibited in Figure 2.

the net coefficient of $\mathbb{E}_\mu g$ here is positive. Thus we have

$$\left(2 + \sum_{j=1}^r 1 + \sum_{\ell=2^{r+1}}^m \frac{m}{\ell(\ell-1)}(1+1-1)\right) \min_{1 \leq k \leq m} \mathbb{E}_\mu g(kx) \leq 1$$

which gives the upper bound in (1.13) after a brief computation using (1.20).

To establish the matching lower bound using the measure μ_m , it suffices to show that

$$\sum_{j=1}^{r+1} g\left(\frac{k}{2^j}\right) + \frac{m}{2^r} g\left(\frac{k}{2m}\right) = 1$$

for all $k = 1, \dots, m$. But from (1.10) and telescoping series we have

$$\sum_{j=1}^{r+1} g\left(\frac{k}{2^j}\right) = \{k\}^* - \left\{\frac{k}{2^{r+1}}\right\}^* = 1 - \frac{k}{2^{r+1}}$$

while since $\frac{k}{2m} \leq \frac{1}{2}$ we have

$$g\left(\frac{k}{2m}\right) = \frac{k}{2m}$$

and the claim follows. \square

Remark 1.3. The particular linear combination of the $g(kx)$ used in (1.12) was discovered after some numerical experimentation, guided by the fact that this combination should attain the bound of 1 at every point in the support of the optimal measure μ_m . However, this constraint does not fully determine the coefficients of the combination, and it would be possible to establish the bound (1.13) using other linear combinations of $g(kx)$ instead. For instance, when m is a power of two, the inequalities (1.12) and (1.8) differ, even though they both imply (1.13): see Figures 1, 3.

2. OPTIMAL EQUIDISTRIBUTION FOR THE SINE FUNCTION

The function $\sin(2\pi x)$ is somewhat analogous to the sawtooth function $g(x)$ studied in Theorem 1.2. We now solve exactly the corresponding optimization problem for the sine function in Theorem 2.1. The problem solved here is equivalent to finding the optimal inequality of the form $\sum_{k=1}^m a_k \sin kx \leq 1$ for each positive integer m .

Theorem 2.1. *Let m be a positive integer.*

(i) *We have*

$$\sum_{1 \leq k \leq m; k \text{ odd}} \frac{2}{(m+1)^2} \cot\left(\frac{\pi k}{2m+2}\right) [(m+1-k) \sin kx + k \sin(m+1-k)x] \leq 1 \quad (2.1)$$

for all real numbers x . Write this inequality as $\sum_{k=1}^m a_k \sin kx \leq 1$; then all the coefficients a_k are nonnegative.




FIGURE 3. The left-hand side of (1.12) when $m = 8$; compare with Figure 1.

(ii) *We have*

$$\min_{1 \leq k \leq m} \mathbb{E}_\mu \sin kx \leq \frac{1}{c_m} \quad (2.2)$$

for every (Borel) probability measure μ on the real line, where

$$c_m := \frac{2}{m+1} \sum_{1 \leq j \leq m; j \text{ odd}} \cot\left(\frac{\pi j}{2m+2}\right). \quad (2.3)$$

This is the optimal bound: equality is attained for the measure μ_m with mass $\frac{2}{(m+1)c_m} \cot\left(\frac{\pi j}{2m+2}\right)$ at $\frac{\pi j}{m+1}$ for every odd $1 \leq j \leq m$.

It follows from our argument (in particular the properties of the measure μ_m in (ii)) that statement (i) is the optimal inequality of the form $\sum_{k=1}^m a_k \sin kx \leq 1$, in the sense that it has the maximal value of $\sum a_k$ (namely, c_m) among all inequalities of this form. The closest relative of this inequality in the literature seems to be Vaaler's inequality, of the form $\sum_{k=1}^m b_k \sin 2\pi kx \geq x - \frac{1}{2}$ for $x \in [0, \frac{1}{2}]$ [6, Theorem 18].

By comparing the cotangent sum c_m to an integral, with the first few terms of the sum separated off for greater accuracy, one checks that c_m is close to $(2/\pi) \log(m+1)$. Precisely, by an argument due to Pinielis [5],

$$c_m = \frac{2}{\pi} \log(m+1) + \frac{2}{\pi} \left(\log\left(\frac{4}{\pi}\right) + \gamma \right) + o(1),$$




FIGURE 4. The left-hand side $f(x)$ of (2.1) when $m = 8$. We are showing that it is at most 1. The bound of 1 is attained exactly at the support $\{\pi/9, 3\pi/9, 5\pi/9, 7\pi/9\}$ of the optimal measure μ_8 .

where γ is Euler's constant. In particular, the bounds for the sine problem are of the same order of magnitude as the bounds for the sawtooth problem. If one replaces sine by cosine then the problem becomes trivial, as in this case the Dirac mass at the origin is clearly the extremizing measure and there is no decay in m .

Another formula for the constant a_k in inequality (2.1) is

$$a_k = \frac{4(m+1-k)}{(m+1)^2} \frac{\sin^2 \frac{[m/2]k\pi}{m+1}}{\sin \frac{\pi k}{m+1}}$$

for $1 \leq k \leq m$. We will not use this, however.

Proof. We first show that part (ii) follows from (i). All the coefficients in the inequality $\sum_{k=1}^m a_k \sin kx \leq 1$ from (2.1) are nonnegative, since $\cot x$ decreases from ∞ to 0 on the interval $(0, \pi/2]$. Therefore, on integrating this inequality against any Borel probability measure μ on the real line, we have

$$\sum_{k=1}^m a_k \mathbb{E}_\mu \sin kx \leq 1.$$

From the definition of a_k in terms of cotangents, it is immediate that $\sum_{k=1}^m a_k = c_m$. This proves (2.2), namely that $\min_{1 \leq k \leq m} \mathbb{E}_\mu \sin kx \leq 1/c_m$.




FIGURE 5. The left-hand side $f(x)$ of (2.1) when $m = 9$. We are showing that it is at most 1. The bound of 1 is attained exactly at the support $\{\pi/10, 3\pi/10, 5\pi/10, 7\pi/10, 9\pi/10\}$ of the optimal measure μ_9 .

Next, we show that for every $1 \leq k \leq m$, the measure μ_m defined in (ii) satisfies

$$\mathbb{E}_{\mu_m} \sin kx = \frac{1}{c_m}.$$

The proof seems easiest in terms of the Fourier transform on the cyclic group $G = \mathbb{Z}/(2m+2)$. Namely, for a complex-valued function f on G , define the Fourier transform on the dual group $\hat{G} = \mathbb{Z}/(2m+2)$ by $\hat{f}(\xi) = (1/(2m+2)) \sum_{x \in G} f(x) e^{-2\pi i \xi x / (2m+2)}$; then the inverse Fourier transform gives that $f(x) = \sum_{\xi \in \hat{G}} \hat{f}(\xi) e^{2\pi i \xi x / (2m+2)}$.

Let σ_m be the function on $G = \mathbb{Z}/(2m+2)$ defined by $\sigma_m(k) = 1$ if $1 \leq k \leq m$, $\sigma_m(k) = -1$ if $-m \leq k \leq -1$, and zero otherwise (a discrete version of a square wave). Let $\zeta = e^{2\pi i / (2m+2)}$. Then the Fourier transform of σ_m is, for $j \in \mathbb{Z}/(2m+2)$,

$$\begin{aligned} \widehat{\sigma_m}(j) &= \frac{1}{2m+2} \sum_{k=0}^{2m+1} \sigma_m(k) \zeta^{-jk} \\ &= \frac{-1}{2m+2} \sum_{k=1}^m (\zeta^{jk} - \zeta^{-jk}). \end{aligned}$$

Clearly $\widehat{\sigma_m}(0) = 0$. For $j \neq 0$ in $\mathbb{Z}/(2m+2)$, we have $\sum_{k=1}^m \zeta^{jk} = (\zeta^j - (\zeta^j)^{m+1}) / (1 - \zeta^j)$. Since $\zeta^{m+1} = -1$, that sum is $(1 + \zeta^j) / (1 - \zeta^j)$ if j is odd and -1 if j is even. Likewise, $\sum_{k=1}^m \zeta^{-jk}$ is $-(1 + \zeta^j) / (1 - \zeta^j)$ if j is odd and -1 if j is even. So the Fourier transform

above is given by $\widehat{\sigma}_m(j) = (-1/(m+1))(1+\zeta^j)/(1-\zeta^j)$ if j is odd and 0 if j is even. Equivalently, $\widehat{\sigma}_m(j) = (-i/(m+1)) \cot(\pi j/(2m+2))$ for j odd.

Therefore, applying the inverse Fourier transform tells us, in particular for $1 \leq k \leq m$, that

$$\begin{aligned} 1 &= \sigma_m(k) \\ &= \sum_{j=0}^{2m+1} \widehat{\sigma}_m(j) e^{2\pi i j k / (2m+2)} \\ &= \sum_{1 \leq j \leq m; j \text{ odd}} \frac{2}{m+1} \cot\left(\frac{\pi j}{2m+2}\right) \sin\left(\frac{\pi j k}{m+1}\right). \end{aligned}$$

(This can also be deduced from an identity due to Eisenstein and Stern, discussed in the introduction to [2].) After dividing by c_m , this says that the measure μ_m defined in (ii) has $\mathbb{E}_{\mu_m} \sin kx = 1/c_m$ for all $1 \leq k \leq m$, as we want.

It remains to prove part (i). We can relate the linear combination of sines $f(x)$ on the left side of (2.1) to the function σ_m above. First, let ν_m be the function on $\mathbb{Z}/(2m+2)$ defined by $\nu_m(j) = 2\sigma_m(j)$ if j is odd and 0 otherwise; so $\nu_m(j)$ is 2 for $1 \leq j \leq m$ and j odd, -2 for $m+2 \leq j \leq 2m-1$ and j odd, and 0 otherwise. One checks that multiplying a function $f(j)$ on $\mathbb{Z}/(2m+2)$ by $(-1)^j$ corresponds to shifting its Fourier transform by $m+1$. Therefore, the Fourier transform of ν_m is

$$\widehat{\nu}_m(j) = \widehat{\sigma}_m(j) - \widehat{\sigma}_m(m+1+j).$$

Since σ_m is an odd function, so is $\widehat{\sigma}_m$, and hence we can rewrite this formula as

$$\widehat{\nu}_m(j) = \widehat{\sigma}_m(j) + \widehat{\sigma}_m(m+1-j).$$

If m is odd, then this is

$$\widehat{\nu}_m(j) = \frac{-i}{m+1} \left(\cot \frac{\pi j}{2m+2} + \cot \frac{\pi(m+1-j)}{2m+2} \right)$$

if j is odd and 0 if j is even. If m is even, then

$$\widehat{\nu}_m(j) = \begin{cases} \frac{-i}{m+1} \cot\left(\frac{\pi j}{2m+2}\right) & \text{if } j \text{ is odd,} \\ \frac{-i}{m+1} \cot\left(\frac{\pi(m+1-j)}{2m+2}\right) & \text{if } j \text{ is even.} \end{cases}$$

This is clearly related to the function $f(x)$. To make the connection precise, consider another interpretation of the Fourier transform on $\mathbb{Z}/(2m+2)$: namely, as Fourier series on the circle $\mathbb{R}/2\pi\mathbb{Z}$ applied to linear combinations of Dirac delta functions with support in $(1/(2m+2))2\pi\mathbb{Z}$. Let S denote the square wave function

$$S(x) := 1_{0 < \{x/2\pi\} < 1/2} - 1_{1/2 < \{x/2\pi\} < 1}.$$

We sample this function at odd multiples of $\pi/(m+1)$ to create a discrete approximation ν_m to $S(x)$, basically a Dirac comb modulated by a square wave:

$$\nu_m := \frac{2\pi}{m+1} \sum_{j \in \mathbb{Z}; j \text{ odd}} S\left(\frac{\pi j}{m+1}\right) \delta_{\pi j/(m+1)}. \quad (2.4)$$

This measure is 2π -periodic. Here we multiplied the function ν_m defined earlier on $\mathbb{Z}/(2m+2)$ by $2\pi/(2m+2)$ in order to make the Fourier coefficients

$$\widehat{\nu}_m(k) := \frac{1}{2\pi} \int_0^{2\pi} e^{-ikx} d\nu(x)$$

the same as those we computed on $\mathbb{Z}/(2m+2)$. (In particular, these Fourier coefficients are periodic with period $2m+2$.)

By inspection, then, the function $f(x)$ from part (i) of the theorem is

$$f(x) = \frac{2i}{m+1} \sum_{1 \leq k \leq m} (m+1-k) \widehat{\nu}_m(k) \sin kx,$$

or equivalently (due to the odd nature of ν_m and hence $\widehat{\nu}_m$)

$$f(x) = \sum_{-m \leq k \leq m} \left(1 - \frac{|k|}{m+1}\right) \widehat{\nu}_m(k) e^{ikx}.$$

This is clearly related to the *Fejér kernel*, $F_{m+1}(x) := \frac{1}{m+1} \left(\frac{\sin(m+1)(x/2)}{\sin x/2} \right)^2 = \sum_{|k| \leq m} \left(1 - \frac{|k|}{m+1}\right) e^{ikx}$. (The original motivation for the Fejér kernel was to show that every continuous function on the circle is the uniform limit of the averaged partial sums of its Fourier series [3, Theorem 1.3.3].) Namely, since the Fourier series takes convolution to multiplication, f is the convolution of ν_m with the Fejér kernel:

$$f(x) = \frac{1}{2\pi} \int_0^{2\pi} F_{m+1}(x-y) d\nu_m(y).$$

Here $F_{m+1}(x)$ equals $m+1$ at multiples of 2π , and it vanishes at other even multiples of $\pi/(m+1)$. So from (2.4) we have

$$f\left(\frac{\pi j}{m+1}\right) = S\left(\frac{\pi j}{m+1}\right)$$

whenever j is odd. This proves:


$$f\left(\frac{\pi j}{m+1}\right) = 1 \tag{2.5}$$

whenever $1 \leq j \leq m$ is odd. More broadly, we now have an interpretation of f as an interpolation of S given by the Fejér kernel.

Next, let us show that

$$f'\left(\frac{\pi j}{m+1}\right) = 0 \tag{2.6}$$

for any odd integer j . Namely, since f is the convolution of ν with the Fejér kernel F_{m+1} , f' is the convolution of ν with the derivative F'_{m+1} . But $F'_{m+1}(x)$ vanishes at all even multiples of $\pi/(m+1)$, since F_{m+1} reaches its minimum 0 or maximum $m+1$ at those points. Since ν is supported on odd multiples of $\pi/(m+1)$, this proves (2.6).

FIGURE 6. The Fejér kernel $F_{m+1}(x)$ with $m = 8$.

Clearly f is odd and 2π -periodic; in particular

$$f(0) = f(\pi) = 0. \quad (2.7)$$

Next, since $f'(x)$ is a linear combination of the functions $\cos kx$ for $1 \leq k \leq m$, de Moivre's formula gives that $f'(x) = P(\cos x)$ for some polynomial P of degree m . In particular, f' has at most m zeros in the interval $[0, \pi]$ (counting multiplicity for the zeros in $(0, \pi)$). On the other hand, from (2.6) we see that f' has $\lfloor m/2 \rfloor + 1$ zeros in this region, at the points $\frac{\pi j}{m+1}$ with $1 \leq j \leq m+1$ odd. On the other hand, from (2.5) and Rolle's theorem (and the fact that f is not locally constant) we also see that we have $\lceil m/2 \rceil - 1$ additional zeros distinct from the preceding ones, with one additional zero strictly between $\frac{\pi j}{m+1}$ and $\frac{\pi(j+2)}{m+1}$ whenever $1 \leq j \leq j+2 \leq m$ is odd. Thus all the m zeros of f' are accounted for, and there are no further zeros; in particular all the zeros of f' in $(0, \pi)$ are simple, and f' changes sign as it crosses each zero in $(0, \pi)$. We then conclude from (2.7), (2.5), and the mean value theorem that

- $f(x)$ is strictly increasing from 0 to 1 as x goes from 0 to $\frac{\pi}{m+1}$;
- Whenever $1 \leq j \leq j+2 \leq m$ is odd, f starts at a local maximum of 1 at $x = \frac{\pi j}{m+1}$, strictly decreases to a local minimum somewhere between $\frac{\pi j}{m+1}$ and $\frac{\pi(j+2)}{m+1}$, then strictly increases back to a local maximum of 1 at $x = \frac{\pi(j+2)}{m+1}$.
- If $j = 2\lfloor \frac{m}{2} \rfloor + 1$ is the largest odd number less than $m+1$, f is strictly decreasing from 1 to 0 as x goes from $\frac{\pi j}{m+1}$ to π .

(See Figures 4, 5.) This already establishes that $f(x) \leq 1$ when $0 \leq x \leq \pi$. If we can also show that $f(x) \geq -1$ for $0 \leq x \leq \pi$, then as f is odd and 2π -periodic we will have $f(x) \leq 1$ for all x , proving the theorem. In reality, this bound will be true with substantial room to spare, as $f(x)$ is only moderately smaller than 1 on most of the interval $[0, \pi]$ (cf. the Gibbs phenomenon).

From the observations above and the oddness of f , we know that $f(x) \geq -1$ if $x \in [-\pi/(m+1), \pi/(m+1)]$ and also if $x \in [m\pi/(m+1), (m+2)\pi/(m+1)]$. So it suffices to show that $f(x + \frac{\pi}{m+1}) - f(x - \frac{\pi}{m+1})$ is nonnegative for $x \in [0, (m-1)\pi/(2m+2)]$, and nonpositive for $x \in [(m+3)\pi/(2m+2), \pi]$.

Assume that m is odd. Since f is the convolution of ν_m with the Fejér kernel F_{m+1} , we have

$$\begin{aligned} f\left(x + \frac{\pi}{m+1}\right) - f\left(x - \frac{\pi}{m+1}\right) &= \left[\nu_m\left(x + \frac{\pi}{m+1}\right) - \nu_m\left(x - \frac{\pi}{m+1}\right) \right] * F_{m+1} \\ &= \frac{4\pi}{m+1} [\delta_0 - \delta_\pi] * F_{m+1} \\ &= \frac{4\pi}{m+1} [F_{m+1}(x) - F_{m+1}(x - \pi)]. \end{aligned}$$

(This description of the change of ν_m when the input changes by $2\pi/(m+1)$ uses that m is odd.) So it suffices to show that $F_{m+1}(x) - F_{m+1}(x - \pi)$ is nonnegative for $x \in [0, (m-1)\pi/(2m+2)]$, and that it is nonpositive for $x \in [(m+3)\pi/(2m+2), \pi]$. We prove this (in fact for slightly bigger intervals) in Lemma 2.2 below. That completes the proof for m odd.

For m even, we have the somewhat messier situation that

$$\begin{aligned} &f\left(x + \frac{\pi}{m+1}\right) - f\left(x - \frac{\pi}{m+1}\right) \\ &= \left[\nu_m\left(x + \frac{\pi}{m+1}\right) - \nu_m\left(x - \frac{\pi}{m+1}\right) \right] * F_{m+1} \\ &= \frac{2\pi}{m+1} [2\delta_0 - \delta_{m\pi/(m+1)} - \delta_{(m+2)\pi/(m+1)}] * F_{m+1} \\ &= \frac{2\pi}{m+1} \left[2F_{m+1}(x) - F_{m+1}\left(x - \frac{m\pi}{m+1}\right) - F_{m+1}\left(x - \frac{(m+2)\pi}{m+1}\right) \right]. \end{aligned}$$

This is nonnegative for $x \in [0, \frac{(m-1)\pi}{2m+2}]$ and nonpositive for $x \in [\frac{(m+3)\pi}{2m+2}, \pi]$, by Lemma 2.2 below (which in fact works for slightly bigger intervals). This completes the proof of Theorem 2.1. \square

Lemma 2.2. *Let m be a positive integer, and let $F(x) = F_{m+1}(x) = \frac{1}{m+1} \left(\frac{\sin((m+1)(x/2))}{\sin x/2} \right)^2$ be the Fejér kernel.*

- (i) *If m is odd, then $F(x) - F(x - \pi)$ is nonnegative if $x \in [-\pi/2, \pi/2]$ and nonpositive if $x \in [\pi/2, 3\pi/2]$.*

- (ii) If m is even, then $2F(x) - F(x - \frac{m\pi}{m+1}) - F(x - \frac{(m+2)\pi}{m+1})$ is nonnegative if $x \in [-\frac{\pi}{2} + \frac{\pi}{m+1}, \frac{\pi}{2} - \frac{\pi}{m+1}]$ and nonpositive if $x \in [\frac{\pi}{2} + \frac{\pi}{m+1}, \frac{3\pi}{2} - \frac{\pi}{m+1}]$.

Proof. (i) Let m be an odd positive integer. By definition of the Fejér kernel $F(x)$, we have

$$F(x) - F(x - \pi) = \frac{1}{m+1} \left[\frac{\sin^2((m+1)x/2)}{\sin^2 x/2} - \frac{\sin^2(\frac{(m+1)x}{2} - \frac{(m+1)\pi}{2})}{\sin^2(x - \pi)/2} \right].$$

Here $\sin((x - \pi)/2) = -\cos(x/2)$. Since m is odd, $(m+1)\pi/2$ is an integer multiple of π , and so the numerators of the two terms are equal. We deduce that

$$F(x) - F(x - \pi) = \frac{\sin^2((m+1)x/2)}{m+1} \left[\frac{1}{\sin^2(x/2)} - \frac{1}{\cos^2(x/2)} \right].$$

For $x \in [-\pi/2, \pi/2]$, we have $\sin^2(x/2) \leq 1/2$, while $\cos^2(x/2) \geq 1/2$. It follows that $F(x) - F(x - \pi)$ is nonnegative if $x \in [-\pi/2, \pi/2]$. By applying that result to $x - \pi$ in place of x , we also find that $F(x) - F(x - \pi)$ is nonpositive if $x \in [\pi/2, 3\pi/2]$, as we want.

- (ii) Let m be an even positive integer. By definition of the Fejér kernel $F(x)$,

$$\begin{aligned} & 2F(x) - F(x - m\pi/(m+1)) - F(x - (m+2)\pi/(m+1)) \\ &= \frac{1}{m+1} \left[\frac{2\sin^2((m+1)x/2)}{\sin^2(x/2)} - \frac{\sin^2(\frac{(m+1)x}{2} - \frac{m\pi}{2})}{\sin^2(\frac{x}{2} - \frac{m\pi}{2m+2})} - \frac{\sin^2(\frac{(m+1)x}{2} - \frac{(m+2)\pi}{2})}{\sin^2(\frac{x}{2} - \frac{(m+2)\pi}{2m+2})} \right]. \end{aligned}$$

Since m is even, $m\pi/2$ and $(m+2)\pi/2$ are both integer multiples of π , and so the three \sin^2 terms in the numerators are equal. So we can rewrite the expression above as

$$\frac{\sin^2((m+1)x/2)}{m+1} \left[\frac{2}{\sin^2(x/2)} - \frac{1}{\cos^2(\frac{x}{2} + \frac{\pi}{2m+2})} - \frac{1}{\cos^2(\frac{x}{2} - \frac{\pi}{2m+2})} \right].$$

For $x \in [-\frac{\pi}{2} + \frac{\pi}{m+1}, \frac{\pi}{2} - \frac{\pi}{m+1}]$, we have $\sin^2(x/2) \leq 1/2$, while $\cos^2(\frac{x}{2} - \frac{\pi}{2m+2})$ and $\cos^2(\frac{x}{2} + \frac{\pi}{2m+2})$ are both $\geq 1/2$. It follows that the previous paragraph's expression is nonnegative for this range of x . Likewise, if $x \in [\frac{\pi}{2} + \frac{\pi}{m+1}, \frac{3\pi}{2} - \frac{\pi}{m+1}]$, then $\sin^2(x/2) \geq 1/2$ while $\cos^2(\frac{x}{2} + \frac{\pi}{2m+2})$ and $\cos^2(\frac{x}{2} - \frac{\pi}{2m+2})$ are both $\leq 1/2$. It follows that the previous paragraph's expression is nonpositive for this range of x . Lemma 2.2 is proved. \square

REFERENCES

- [1] E. Ballico, R. Pignatelli, and L. Tasin. Weighted hypersurfaces with either assigned volume or many vanishing plurigeners. *Comm. Alg.* **41** (2013), 3745–3752. 2
- [2] B. C. Berndt and B. P. Yeap. Explicit evaluations and reciprocity theorems for finite trigonometric sums. *Adv. Appl. Math.* **29** (2002), 358–385. 14
- [3] H. Dym and H. P. McKean. *Fourier series and integrals*. Academic Press (1972). 15
- [4] L. Esser, B. Totaro, and C. Wang. Varieties of general type with doubly exponential asymptotics. [arXiv:2109.13383](https://arxiv.org/abs/2109.13383) 2
- [5] I. Pinelis. The cotangent sum $\sum_{k=0}^{n-1} (-1)^k \cot\left(\frac{\pi}{4n}(2k+1)\right) = n$. <https://mathoverflow.net/q/368805> 11

- [6] J. Vaaler. Some extremal functions in Fourier analysis. *Bull. Amer. Math. Soc.* **12** (1985), 183–216.
11

UCLA DEPARTMENT OF MATHEMATICS, LOS ANGELES, CA 90095-1555.

E-mail address: `esserl@math.ucla.edu`

UCLA DEPARTMENT OF MATHEMATICS, LOS ANGELES, CA 90095-1555.

E-mail address: `tao@math.ucla.edu`

UCLA DEPARTMENT OF MATHEMATICS, LOS ANGELES, CA 90095-1555.

E-mail address: `totaro@math.ucla.edu`

UCLA DEPARTMENT OF MATHEMATICS, LOS ANGELES, CA 90095-1555.

E-mail address: `chwang@math.ucla.edu`