

# Hodge theory of classifying stacks

Burt Totaro

This paper creates a correspondence between the representation theory of algebraic groups and the topology of Lie groups. In more detail, we compute the Hodge and de Rham cohomology of the classifying space  $BG$  (defined as étale cohomology on the algebraic stack  $BG$ ) for reductive groups  $G$  over many fields, including fields of small characteristic. These calculations have a direct relation with representation theory, yielding new results there. Eventually,  $p$ -adic Hodge theory should provide a more subtle relation between these calculations in positive characteristic and torsion in the cohomology of the classifying space  $BG_{\mathbf{C}}$ .

For the representation theorist, this paper's interpretation of certain Ext groups (notably for reductive groups in positive characteristic) as Hodge cohomology groups suggests spectral sequences that were not obvious in terms of Ext groups (Proposition 9.3). We apply these spectral sequences to compute Ext groups in new cases. The spectral sequences form a machine that can lead to further calculations.

One main result is an isomorphism between the Hodge cohomology of the classifying stack  $BG$  and the cohomology of  $G$  as an algebraic group with coefficients in the ring  $O(\mathfrak{g}) = S(\mathfrak{g}^*)$  of polynomial functions on the Lie algebra  $\mathfrak{g}$  (Theorem 3.1):

$$H^i(BG, \Omega^j) \cong H^{i-j}(G, S^j(\mathfrak{g}^*)).$$

This was shown by Bott over a field of characteristic 0 [8], but in fact the isomorphism holds in any characteristic, and even for group schemes over the integers. More generally, we give an analogous description of the equivariant Hodge cohomology of an affine scheme (Theorem 2.1). This was shown by Simpson and Teleman in characteristic 0 [29, Example 6.8(c)].

Using that isomorphism, we improve the known results on the cohomology of the representations  $S^j(\mathfrak{g}^*)$ . Namely, by Andersen, Jantzen, and Donkin, we have  $H^{>0}(G, O(\mathfrak{g})) = 0$  for a reductive group  $G$  over a field of characteristic  $p$  if  $p$  is a “good prime” for  $G$  [13, Proposition and proof of Theorem 2.2], [21, II.4.22]. We strengthen that to an “if and only if” statement (Theorem 9.1):

**Theorem 0.1.** *Let  $G$  be a reductive group over a field  $k$  of characteristic  $p \geq 0$ . Then  $H^{>0}(G, O(\mathfrak{g})) = 0$  if and only if  $p$  is not a torsion prime for  $G$ .*

For example, this cohomology vanishing holds for every symplectic group  $Sp(2n)$  in characteristic 2 and for the exceptional group  $G_2$  in characteristic 3; these are “bad primes” but not torsion primes.

Finally, we address the problem of computing the Hodge cohomology and de Rham cohomology of  $BG$ , especially at torsion primes. At non-torsion primes, we have a satisfying result, proved using ideas from topology (Theorem 9.2):

**Theorem 0.2.** *Let  $G$  be a split reductive group over  $\mathbf{Z}$ , and let  $p$  be a non-torsion prime for  $G$ . Then Hodge cohomology  $H_{\mathbf{H}}^*(BG/\mathbf{Z})$  and de Rham cohomology  $H_{\text{dR}}^*(BG/\mathbf{Z})$ , localized at  $p$ , are polynomial rings on generators of degrees equal to 2 times the fundamental degrees of  $G$ . These graded rings are isomorphic to the cohomology of the topological space  $BG_{\mathbf{C}}$  with  $\mathbf{Z}_{(p)}$  coefficients.*

At torsion primes  $p$ , it is an intriguing question how the de Rham cohomology of  $BG_{\mathbf{F}_p}$  is related to the mod  $p$  cohomology of the topological space  $BG_{\mathbf{C}}$ . We show that these graded rings are isomorphic for  $G = SO(n)$  with  $p = 2$  (Theorem 11.1). On the other hand, we find that

$$\dim_{\mathbf{F}_2} H_{\text{dR}}^{32}(B \text{Spin}(11)/\mathbf{F}_2) > \dim_{\mathbf{F}_2} H^{32}(B \text{Spin}(11)_{\mathbf{C}}, \mathbf{F}_2)$$

(Theorem 12.1). It seems that no existing results on integral  $p$ -adic Hodge theory address the relation between these two rings (because the stack  $BG$  is not proper over  $\mathbf{Z}$ ), but the theory may soon reach that point. In particular, the results of Bhatt-Morrow-Scholze suggest that the de Rham cohomology  $H_{\text{dR}}^i(BG/\mathbf{F}_p)$  may always be an upper bound for the mod  $p$  cohomology of the topological space  $BG_{\mathbf{C}}$  [6].

This work was supported by NSF grant DMS-1701237. Bhargav Bhatt convinced me to change some definitions in an earlier version of this paper: Hodge and de Rham cohomology of a smooth stack are now defined as étale cohomology. Thanks to Johan de Jong, Eric Primožic, Raphaël Rouquier, and the referees for their comments. Finally, I am grateful to Jungkai Chen for arranging my visit to National Taiwan University, where this work was completed.

## 1 Notation

The *fundamental degrees* of a reductive group  $G$  over a field  $k$  are the degrees of the generators of the polynomial ring  $S(X^*(T) \otimes_{\mathbf{Z}} \mathbf{Q})^W$  of invariants under the Weyl group  $W$ , where  $X^*(T)$  is the character group of a maximal torus  $T$ . For  $k$  of characteristic zero, the fundamental degrees of  $G$  can also be viewed as the degrees of the generators of the polynomial ring  $O(\mathfrak{g})^G$  of invariant functions on the Lie algebra. Here are the fundamental degrees of the simple groups [17, section 3.7, Table 1]:

$A_l$	$2, 3, \dots, l + 1$
$B_l$	$2, 4, 6, \dots, 2l$
$C_l$	$2, 4, 6, \dots, 2l$
$D_l$	$2, 4, 6, \dots, 2l - 2; l$
$G_2$	$2, 6$
$F_4$	$2, 6, 8, 12$
$E_6$	$2, 5, 6, 8, 9, 12$
$E_7$	$2, 6, 8, 10, 12, 14, 18$
$E_8$	$2, 8, 12, 14, 18, 20, 24, 30$

For a commutative ring  $R$  and  $j \geq 0$ , write  $\Omega^j$  for the sheaf of differential forms over  $R$  on any scheme over  $R$ . For an algebraic stack  $X$  over  $R$ ,  $\Omega^j$  is a sheaf of abelian groups on the big étale site of  $X$ . (In particular, for every scheme  $Y$  over  $X$



for principal  $G$ -bundles (in the fppf topology). Concretely, for any scheme  $X$  over  $R$ , a principal  $G$ -bundle over  $X$  determines a morphism  $X \rightarrow BG$  of stacks over  $R$  and hence a pullback homomorphism

$$H^i(BG, \Omega^j) \rightarrow H^i(X, \Omega^j).$$

Note that for a scheme  $X$  over  $R$ ,  $H^i(X, \Omega^j)$  can be computed either in the Zariski or in the étale topology, because the sheaf  $\Omega^j$  (on the small étale site of  $X$ ) is quasi-coherent [32, Tags 03OY, 0DGB].

For any scheme  $X$  over a commutative ring  $R$ , there is a simplicial scheme  $EX$  whose space  $(EX)_n$  of  $n$ -simplices is  $X^{\{0, \dots, n\}} = X^{n+1}$  [11, 6.1.3]. For a group scheme  $G$  over  $R$ , the simplicial scheme  $B_{\text{simp}}G$  over  $R$  is defined as the quotient of the simplicial scheme  $EG$  by the free left action of  $G$ :

$$\text{Spec}(R) \xleftarrow{\quad} G \xleftarrow{\quad} G^2 \cdots$$

If  $G$  is smooth over  $R$ , then Hodge cohomology  $H^i(BG, \Omega^j)$  as defined above can be identified with the cohomology of  $B_{\text{simp}}G$ , because this simplicial scheme is the Čech nerve of the smooth surjective morphism  $\text{Spec}(R) \rightarrow BG$ . For  $G$  not smooth, one has instead to use the Čech nerve of a smooth presentation of  $BG$ . See for example the calculation of the Hodge cohomology of  $B\mu_p$  in characteristic  $p$ , Proposition 10.1.

It is useful that we can compute Hodge cohomology via any smooth presentation of a stack. For example, let  $H$  be a closed subgroup scheme of a smooth group scheme  $G$  over a commutative ring  $R$ , and assume that  $H$  is flat and locally of finite presentation over  $R$ . Then  $G/H$  is an algebraic space with a smooth surjective morphism  $G/H \rightarrow BH$  over  $R$ , and so we can compute the Hodge cohomology of the stack  $BH$  using the associated Čech nerve. Explicitly, that is the simplicial algebraic space  $EG/H$ , and so we have:

**Lemma 1.2.**

$$H^i(BH, \Omega^j) \cong H^i(EG/H, \Omega^j).$$

Note that the cohomology theories we are considering are not  $A^1$ -homotopy invariant. Indeed, Hodge cohomology is usually not the same for a scheme  $X$  as for  $X \times A^1$ , even over a field of characteristic zero. For example,  $H^0(\text{Spec}(k), \mathcal{O}) = k$ , whereas  $H^0(A_k^1, \mathcal{O})$  is the polynomial ring  $k[x]$ . In de Rham cohomology,  $H_{\text{dR}}^0(A^1/k)$  is just  $k$  if  $k$  has characteristic zero, but it is  $k[x^p]$  if  $k$  has characteristic  $p > 0$ .

## 2 Equivariant Hodge cohomology and functions on the Lie algebra

In this section, we identify the Hodge cohomology of a quotient stack with the cohomology of an explicit complex of vector bundles (Theorem 2.1). As a special case, we relate the Hodge cohomology of a classifying stack  $BG$  to the cohomology of  $G$  as a group scheme (Corollary 2.2). In this section, we assume  $G$  is smooth. Undoubtedly, various generalizations of the statements here are possible. In particular, we will give an analogous description of the Hodge cohomology of  $BG$  for a non-smooth group  $G$  in Theorem 3.1.

The main novelty is that these results hold in any characteristic. In particular, Theorem 2.1 was proved in characteristic zero by Simpson and Teleman [29, Example 6.8(c)]. As discussed in section 1, equivariant Hodge cohomology  $H_G^i(X, \Omega^j)$  is defined as cohomology of the quotient stack  $[X/G]$ .

**Theorem 2.1.** *Let  $G$  be a smooth affine group scheme over a commutative ring  $R$ . Let  $G$  act on a smooth affine scheme  $X$  over  $R$ . Then there is a canonical isomorphism*

$$H_G^i(X, \Omega^j) \cong H_G^i(X, \Lambda^j L_{[X/G]}),$$

where  $\Lambda^j L_{[X/G]}$  is the complex of  $G$ -equivariant vector bundles on  $X$ , in degrees 0 to  $j$ :

$$0 \rightarrow \Omega_X^j \rightarrow \Omega_X^{j-1} \otimes \mathfrak{g}_X^* \rightarrow \cdots \rightarrow S^j(\mathfrak{g}_X^*) \rightarrow 0,$$

associated to the map  $\mathfrak{g}_X \rightarrow TX$ .

Here the action of  $G$  on  $X$  gives an action of the Lie algebra  $\mathfrak{g}$  by vector fields on  $X$  by differentiating the action  $G \times_R X \rightarrow X$  at  $1 \in G(R)$ . This can be viewed as a map from the trivial vector bundle  $\mathfrak{g}_X$  over  $X$  to the tangent bundle  $TX$ . (The action of  $G$  on  $\mathfrak{g}_X$  is nontrivial, coming from the adjoint representation of  $G$  on  $\mathfrak{g}$ .) Dualizing gives the map  $\Omega_X^1 \rightarrow \mathfrak{g}_X^*$  used in Theorem 2.1.

The isomorphism of Theorem 2.1 expresses the cohomology over  $[X/G]$  of the “big sheaf”  $\Omega^j$ , which is not a quasi-coherent sheaf on  $[X/G]$ , in terms of the cohomology of a complex of quasi-coherent sheaves on  $[X/G]$ . (Here differentials are over  $R$  unless otherwise stated. The sheaf  $\Omega^j$  on the big étale site of  $[X/G]$  is not quasi-coherent for  $j > 0$  because, for a morphism  $f: Y \rightarrow Z$  of schemes over  $[X/G]$ , the pullback map  $f^* \Omega_{Z/R}^j \rightarrow \Omega_{Y/R}^j$  need not be an isomorphism.)

One might prefer to take the right side of Theorem 2.1 as a definition of Hodge cohomology for algebraic stacks. This could be done without any smoothness assumption. Namely, Olsson defined the cotangent complex  $L_{X/Y}$  as an inverse system for any quasi-compact and quasi-separated morphism  $f: X \rightarrow Y$  of algebraic stacks, correcting the approach of Laumon and Moret-Bailly [27, section 8]. One could then define Hodge cohomology of  $X$  over  $Y$  as  $R^i f_*(L\Lambda^j L_{X/Y})$  (perhaps “Hodge-completed” in the sense of [5]). For  $X$  and  $G$  smooth over  $Y = \text{Spec}(R)$ , this definition agrees with the right side of Theorem 2.1. using that

$$L_{[X/G]/R} \cong [\Omega_X^1 \rightarrow \mathfrak{g}_X^*]$$

by the transitivity triangle [27, 8.1.5]. We have preferred to take the left side of Theorem 2.1 as the definition, using “big sheaves”, because that definition is directly related to the cohomology of simplicial spaces as discussed in section 1. As a result, Theorem 2.1 makes a nontrivial connection between the two approaches.

Theorem 2.1 is useful already for  $X = \text{Spec}(R)$ , where it gives the following result, proved over a field of characteristic zero by Bott [8].

**Corollary 2.2.** *Let  $G$  be a smooth affine group scheme over a commutative ring  $R$ . Then there is a canonical isomorphism*

$$H^i(BG, \Omega^j) \cong H^{i-j}(G, S^j(\mathfrak{g}^*)).$$

The group on the left is an étale cohomology group of the algebraic stack  $BG$  over  $R$ , as discussed in section 1. On the right is the cohomology of  $G$  as a group scheme, defined by  $H^i(G, M) = \text{Ext}_G^i(R, M)$  for a  $G$ -module  $M$  [21, section 4.2].

*Proof.* (Corollary 2.2) This follows from Theorem 2.1 applied to the stack  $BG = [\text{Spec}(R)/G]$ . The deduction uses two facts. First, a quasi-coherent sheaf on  $BG$  is equivalent to a  $G$ -module [32, Tag 06WS]. Second, for a  $G$ -module  $M$ , the cohomology of the corresponding quasi-coherent sheaf on the big étale site of  $BG$  coincides with its cohomology as a  $G$ -module,  $H^*(G, M)$ , since both are computed by the same Čech complex (section 1 for the sheaf, [21, Proposition 4.16] for the module).  $\square$

*Proof.* (Theorem 2.1) The adjoint representation of  $G$  on  $\mathfrak{g}$  determines a  $G$ -equivariant vector bundle  $\mathfrak{g}_X$  on  $X$ . The action of  $G$  on  $X$  gives a morphism  $\Omega_X^1 \rightarrow \mathfrak{g}_X^*$  of  $G$ -equivariant quasi-coherent sheaves (in fact, vector bundles) on  $X$ . Consider these equivariant sheaves as quasi-coherent sheaves on  $[X/G]$ , according to [32, Tag 06WS].

We will define a map from the complex  $\Omega_X^1 \rightarrow \mathfrak{g}_X^*$  of quasi-coherent sheaves on  $[X/G]$  (in degrees 0 and 1) to the sheaf  $\Omega^1$ , in the derived category  $D([X/G]_{\text{ét}}, O_{[X/G]})$  of  $O_{[X/G]}$ -modules on the big étale site  $[X/G]_{\text{ét}}$ . To do this, define another sheaf  $S$  on the big étale site of  $[X/G]$  by: for a scheme  $U$  over  $[X/G]$ , let  $E = U \times_{[X/G]} X$  (so that  $\pi: E \rightarrow U$  is a principal  $G$ -bundle), and define  $S(U) = H^0(E, \Omega^1)^G$ . Since  $G$  is smooth over  $k$ , there is a short exact sequence of quasi-coherent sheaves on  $E$ ,  $0 \rightarrow \pi^* \Omega_U^1 \rightarrow \Omega_E^1 \rightarrow \Omega_{E/U}^1 \rightarrow 0$ . These are  $G$ -equivariant sheaves on  $E$ , and so this can be viewed as the pullback of a short exact sequence of sheaves on  $U$ , known as the *Atiyah sequence* [2, Theorem 1], [20, VII.2.4.2.13–14]:

$$0 \rightarrow \Omega_U^1 \rightarrow \pi_*(\Omega_E^1)^G \rightarrow {}^E \mathfrak{g}_U^* \rightarrow 0.$$

Here  ${}^E \mathfrak{g}_U^*$  is the vector bundle on  $U$  associated to the  $G$ -bundle  $E \rightarrow U$  and the action of  $G$  on  $\mathfrak{g}^*$ . Since the  $G$ -bundle  $E \rightarrow U$  is arbitrary, we have produced an exact sequence

$$0 \rightarrow \Omega^1 \rightarrow S \rightarrow \mathfrak{g}_X^* \rightarrow 0$$

of sheaves on the big étale site of  $[X/G]$ . By definition of the Atiyah sequence, the map from  $S(U) = H^0(E, \Omega_E^1)^G$  to  $H^0(U, {}^E \mathfrak{g}_U^*) = H^0(E, \mathfrak{g}_E^*)^G$  arises from the map  $\Omega_E^1 \rightarrow \mathfrak{g}_E^*$  given by differentiating the action of  $G$  on  $E$ .

Thus the sheaf  $\Omega^1$  on  $[X/G]$  is isomorphic in the derived category to the complex  $S \rightarrow \mathfrak{g}_X^*$  (in cohomological degrees 0 and 1) on  $[X/G]$ . Therefore, to produce the map in  $D([X/G]_{\text{ét}}, O_{[X/G]})$  promised above, it suffices to define a map  $\alpha$  of complexes of sheaves on  $[X/G]_{\text{ét}}$ :

$$\begin{array}{ccccccc} 0 & \longrightarrow & \Omega_X^1 & \longrightarrow & \mathfrak{g}_X^* & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & S & \longrightarrow & \mathfrak{g}_X^* & \longrightarrow & 0. \end{array}$$

(As above,  $\mathfrak{g}_X^*$  denotes the vector bundle on  $[X/G]$  associated to the representation of  $G$  on  $\mathfrak{g}^*$ , and  $\Omega_X^1$  denotes the vector bundle on  $[X/G]$  corresponding to the  $G$ -equivariant vector bundle of the same name on  $X$ .) It is now easy to produce the map  $\alpha$  of complexes: for any scheme  $U$  over  $[X/G]$ , with associated principal  $G$ -bundle  $E \rightarrow U$  and  $G$ -equivariant morphism  $h: E \rightarrow X$ , the map from  $\Omega_X^1(U) =$

$H^0(E, h^*\Omega_X^1)^G$  to  $S(U) = H^0(E, \Omega_E^1)^G$  is the pullback, and the map from  ${}^E\mathfrak{g}_U^*$  to itself is the identity. In words, the difference between  $\Omega_X^1(U)$  and  $S(U)$  comes from the difference between (1) the pullback to  $E$  of the sheaf of differentials of  $X$  over  $R$  and (2) the sheaf of differentials on  $E$  over  $R$ .

The commutativity of the diagram above follows from the  $G$ -equivariance of the morphism  $h: E \rightarrow X$ , since the two horizontal maps arise by differentiating the actions of  $G$  on  $X$  and on  $E$ .

For any  $j \geq 0$ , taking the  $j$ th derived exterior power over  $O_{[X/G]}$  of this map of complexes gives a map from the Koszul complex

$$0 \rightarrow \Omega_X^j \rightarrow \Omega_X^{j-1} \otimes \mathfrak{g}_X^* \rightarrow \cdots \rightarrow S^j(\mathfrak{g}_X^*) \rightarrow 0$$

(in degrees 0 to  $j$ ) of vector bundles on  $[X/G]$  to the big sheaf  $\Omega^j$ , in  $D([X/G]_{\text{et}}, O_{[X/G]})$ . (The description of the derived exterior power of a 2-term complex of flat modules as a Koszul complex follows from Illusie [18, Proposition II.4.3.1.6], by the same argument used for derived divided powers in [20, Lemme VIII.2.1.2.1].) We want to show that this map of complexes induces an isomorphism on cohomology over  $[X/G]$ .

By the exact sequence above for the big sheaf  $\Omega^1$  on  $[X/G]$ , we can identify the big sheaf  $\Omega^j$  in the derived category with a similar-looking Koszul complex:

$$0 \rightarrow \Lambda^j(S) \rightarrow \Lambda^{j-1}(S) \otimes \mathfrak{g}_X^* \rightarrow \cdots \rightarrow S^j(\mathfrak{g}_X^*) \rightarrow 0.$$

We want to show that the map  $\Lambda^j(\alpha)$  from the Koszul complex of vector bundles (in the previous paragraph) to this complex of big sheaves induces an isomorphism on cohomology over  $[X/G]$ . For each of these complexes, we have a spectral sequence from the cohomology over  $[X/G]$  of the individual sheaves to the ‘‘hypercohomology’’ over  $[X/G]$  of the whole complex [32, Tag 015J]. We have a map of spectral sequences. Therefore, to show that the map on hypercohomology is an isomorphism, it suffices to show that the map on cohomology of the individual sheaves is an isomorphism. That is, it suffices to show that for each  $0 \leq i \leq j$ , the map

$$H_G^*(X, \Omega_X^i \otimes S^{j-i}(\mathfrak{g}_X^*)) \rightarrow H_G^*(X, \Lambda^i(S) \otimes S^{j-i}(\mathfrak{g}_X^*))$$

is an isomorphism. (Equivariant cohomology is defined as cohomology of the stack  $[X/G]$ , as discussed in section 1.)

By section 1, we can compute both of these cohomology groups on the Čech nerve of the smooth surjective morphism  $X \rightarrow [X/G]$ . This simplicial space can be written as  $(X \times EG)/G$ , where all products are over  $R$ :

$$X \begin{array}{c} \longleftarrow \\ \xrightarrow{\quad} \\ \longleftarrow \\ \xrightarrow{\quad} \\ \longleftarrow \\ \xrightarrow{\quad} \end{array} X \times G \begin{array}{c} \longleftarrow \\ \xrightarrow{\quad} \\ \longleftarrow \\ \xrightarrow{\quad} \\ \longleftarrow \\ \xrightarrow{\quad} \end{array} X \times G^2 \cdots,$$

Since  $X$  is affine, all the spaces in this simplicial space are affine schemes. Therefore, for any  $0 \leq i \leq j$ ,  $H_G^*(X, \Omega_X^i \otimes S^{j-i}(\mathfrak{g}_X^*))$  is the cohomology of the complex of  $H^0$  of the sheaves  $\Omega_X^i \otimes O_{EG} \otimes S^{j-i}(\mathfrak{g}_X^*)$  over the spaces making up  $(X \times EG)/G$ . Likewise,  $H_G^*(X, \Lambda^i(S) \otimes S^{j-i}(\mathfrak{g}_X^*))$  is the cohomology of the complex of  $H^0$  of the sheaves  $\Lambda^i(S) \otimes S^{j-i}(\mathfrak{g}_X^*)$  over the spaces making up  $(X \times EG)/G$ .

Let

$$\varphi: H^0(X \times EG, \Omega_X^i \otimes O_{EG} \otimes S^{j-i}(\mathfrak{g}_X^*)) \rightarrow H^0(X \times EG, \Lambda^i(S) \otimes S^{j-i}(\mathfrak{g}_X^*))$$

be the map of complexes of  $G$ -modules arising as  $H^0$  of sheaves over the spaces making up  $X \times EG$ . The boundary maps in these complexes are alternating sums of pullbacks via the face maps in this simplicial space. By the previous paragraph, we want to show that the induced map  $\varphi^G$  on  $G$ -invariants is a quasi-isomorphism. Moreover, all of these  $G$ -modules arising as  $H^0$  are induced from representations of the trivial group, because  $X \times G^{r+1} \rightarrow (X \times G^{r+1})/G$  is a  $G$ -torsor with a section for each  $r \geq 0$ . Indeed, a choice of section of this  $G$ -torsor trivializes the torsor, and so the group of sections of a  $G$ -equivariant sheaf of  $X \times G^{r+1}$  is the subspace of invariants tensored with  $O(G)$ , as a  $G$ -module. (Note that trivializations of these  $G$ -torsors cannot be made compatible with the face maps of the simplicial space, in general.) And every tensor product  $O(G) \otimes_R M$  for a  $G$ -module  $M$  is injective as a  $G$ -module [21, Proposition 3.10]. It follows that  $H^i(G, O(G) \otimes_R M) = 0$  for  $i > 0$  [21, Lemma I.4.7].

Therefore, to show that the map  $\varphi^G$  of  $G$ -invariants is a quasi-isomorphism (as we want), it suffices to show that the map  $\varphi$  is a quasi-isomorphism. And for that, we can forget about the  $G$ -action. That is, we want to show that the map of complexes with  $r$ th term (for  $r \geq 0$ )

$$H^0(X \times G^{r+1}, \Omega_{X \times G^{r+1}}^i \otimes O_{G^{r+1}} \otimes S^{j-i}(\mathfrak{g}^*)) \rightarrow H^0(X \times G^{r+1}, \Omega_{X \times G^{r+1}}^i \otimes S^{j-i}(\mathfrak{g}^*))$$

is a quasi-isomorphism.

We can write  $\Omega_{X \times G^{r+1}}^i$  as the direct sum  $\bigoplus_{l=0}^i \Omega_X^{i-l} \otimes \Omega_{G^{r+1}}^l$ . Moreover, this splitting is compatible with pullback along the face maps of the simplicial scheme  $X \times EG$ . So the map of complexes above is the inclusion of a summand (corresponding to  $l = 0$ ). It remains to show that for every  $0 < l \leq i$ , the  $l$ th summand is a complex  $A_l$  with cohomology zero. Its  $r$ th term is

$$\Omega^{i-l}(X) \otimes_R \Omega^l(G^{r+1}) \otimes_R S^{j-i}(\mathfrak{g}^*).$$

To analyze the cohomology of the complex  $A_l$ , we use the well-known “contractibility” of  $EG$ , in the following form:

**Lemma 2.3.** *Let  $Y$  be a scheme over a ring  $R$  with  $Y(R)$  not empty. For any sheaf  $M$  of abelian groups on the big étale site of  $R$ , the cohomology of the simplicial scheme  $EY$  over  $R$  with coefficients in  $M$  coincides with the cohomology of  $\text{Spec}(R)$ :*

$$H^i(EY, M) \cong H^i(R, M).$$

*Proof.* This is the standard result that a morphism  $Y \rightarrow \text{Spec}(R)$  with a section satisfies cohomological descent, via an explicit chain homotopy. More generally, it would suffice to have sections locally on  $\text{Spec}(R)$  [1, Proposition Vbis.3.3.1].  $\square$

Returning to the proof of Theorem 2.1: we want to show that for  $l > 0$ , the complex  $A_l$  has zero cohomology. Recall that the  $r$ th term of  $A_l$  is

$$\Omega^{i-l}(X) \otimes_R \Omega^l(G^{r+1}) \otimes_R S^{j-i}(\mathfrak{g}^*),$$

with boundary maps coming from the face maps of the simplicial scheme  $X \times EG$ . By Lemma 1.2 (applied to the sheaf  $\Omega^l$  on the big étale site of  $R$  and the simplicial scheme  $EG$ ), the complex  $A_l$  has cohomology equal to  $\Omega^{i-l}(X) \otimes \Omega^l(\text{Spec } R) \otimes S^{j-i}(\mathfrak{g}^*)$  in degree 0 and zero in other degrees. Since  $l > 0$ , the cohomology in degree 0 also vanishes. The proof is complete.  $\square$



The argument works verbatim to prove a twisted version of Corollary 2.2, where the sheaf  $\Omega^j$  on  $BG$  is tensored with the vector bundle associated to any  $G$ -module. The generalization will not be needed in this paper, but we state it for possible later use.

**Theorem 2.4.** *Let  $G$  be a smooth affine group scheme over a commutative ring  $R$ . Let  $M$  be a  $G$ -module that is flat over  $R$ . Then there is a canonical isomorphism*

$$H^i(BG, \Omega^j \otimes M) \cong H^{i-j}(G, S^j(\mathfrak{g}^*) \otimes M).$$

### 3 Flat group schemes

We now describe the Hodge cohomology of the classifying stack of a group scheme  $G$  which need not be smooth, generalizing Corollary 2.2. The analog of the co-Lie algebra  $\mathfrak{g}^*$  in this generality is the co-Lie complex  $l_G$  in the derived category of  $G$ -modules, defined by Illusie [20, section VII.3.1.2]. Namely,  $l_G$  is the pullback of the cotangent complex of  $G \rightarrow \text{Spec}(R)$  to  $\text{Spec}(R)$ , via the section  $1 \in G(R)$ . (The cotangent complex  $L_{X/Y}$  of a morphism  $X \rightarrow Y$  of schemes is an object of the quasi-coherent derived category of  $X$ ; if  $X$  is smooth over  $Y$ , then  $L_{X/Y}$  is the sheaf  $\Omega_{X/Y}^1$ .)

The cohomology of  $l_G$  in degree 0 is the  $R$ -module  $\omega_G^1$ , the restriction of  $\Omega_G^1$  to the identity  $1 \in G(R)$ ; thus  $\omega_G^1$  is the co-Lie algebra  $\mathfrak{g}^*$  if  $G$  is smooth over  $R$ . The complex  $l_G$  has zero cohomology except in cohomological degrees  $-1$  and  $0$ . If  $G$  is smooth, then  $l_G$  has cohomology concentrated in degree 0. More generally, a closed immersion of  $G$  into a smooth  $R$ -group scheme  $H$  yields an explicit formula for  $l_G$  in the derived category of  $G$ -modules:  $l_G$  is the complex

$$0 \rightarrow I/\mathfrak{m}I \rightarrow \mathfrak{m}/\mathfrak{m}^2 \rightarrow 0,$$

where  $I$  is the ideal defining  $G$  in  $H$  and  $\mathfrak{m}$  is the ideal defining the point  $1$  in  $H$ , so that  $\mathfrak{m}/\mathfrak{m}^2 = \mathfrak{h}^*$  [19, section 4.2].

**Theorem 3.1.** *Let  $G$  be a flat affine group scheme of finite presentation over a commutative ring  $R$ . Then there is a canonical isomorphism*

$$H^i(BG, \Omega^j) \cong H^{i-j}(G, S^j(l_G)).$$

*This is an isomorphism of rings from  $H_{\mathbb{H}}^*(BG/R)$  to  $H^*(G, S^*(l_G))$ .*

*Proof.* As discussed in section 1, we can compute  $H^*(BG, \Omega^j)$  as the étale cohomology with coefficients in  $\Omega^j$  of the Čech nerve associated to any algebraic space  $U$  over  $R$  with a smooth surjective morphism from  $U$  to the stack  $BG$ . The assumption on  $G$  implies that  $BG$  is a quasi-compact algebraic stack over  $R$ , and so there is an affine scheme  $U$  with a smooth surjective morphism  $U \rightarrow BG$  [32, Tags 06FI and 04YA]. By Lemma 1.1,  $BG$  is smooth over  $R$ , and so  $U$  is smooth over  $R$ . Let  $E = U \times_{BG} \text{Spec}(R)$ ; then  $E$  is a smooth  $R$ -space with a free  $G$ -action such that  $U = E/G$ . Also,  $E$  is affine because  $U$  and  $G$  are affine.

By section 1,  $H^*(BG, \Omega^j)$  is the étale cohomology with coefficients in  $\Omega^j$  of the simplicial algebraic space  $EE/G$ :

$$E/G \longleftarrow E^2/G \longleftarrow E^3/G \cdots$$

By the properties of  $E$  and  $U$  above,  $E^{n+1}/G$  is an affine scheme for all  $n \geq 0$ . Since  $H^*(BG, \Omega^j)$  is the cohomology with coefficients in  $\Omega^j$  of the simplicial scheme  $EE/G$ , this is the cohomology of the cochain complex

$$0 \rightarrow \Omega^j(E/G) \rightarrow \Omega^j(E^2/G) \rightarrow \cdots .$$

As in the proof of Theorem 2.1, this complex is the  $G$ -invariants of the complex

$$0 \rightarrow H^0(E, \pi^*(\Omega_{E/G}^j)) \rightarrow H^0(E^2, \pi^*(\Omega_{E^2/G}^j)) \rightarrow \cdots ,$$

where we write  $\pi$  for the morphism  $E^{n+1} \rightarrow E^{n+1}/G$  for any  $n \geq 0$ .

For any smooth  $R$ -scheme  $X$  with a free action of  $G$ , I claim that there is a canonical exact triangle in the quasi-coherent derived category of  $G$ -equivariant sheaves on  $X$ :

$$\pi^*(\Omega_{X/G}^1) \rightarrow \Omega_X^1 \rightarrow l_G,$$

where we write  $l_G$  for the pullback of the co-Lie complex  $l_G$  from the stack  $BG$  over  $R$  to  $X$ . To deduce this from Illusie's results on the cotangent complex  $L_{X/Y}$ , let  $Y = X/G$  and  $S = \text{Spec}(R)$ , and use the transitivity exact triangle for  $X \rightarrow Y \rightarrow S$  in the derived category of  $X$  [18, II.2.1.5.2]:

$$\pi^*L_{Y/S} \rightarrow L_{X/S} \rightarrow L_{X/Y}.$$

Since  $X$  is smooth over  $S$ , so is  $Y$  (even though  $G$  need not be); so  $L_{Y/S} \cong \Omega_{Y/S}^1$  and  $L_{X/S} \cong \Omega_{X/S}^1$ . Also, since  $X \rightarrow Y$  is a  $G$ -torsor in the fppf topology,  $L_{X/Y}$  is the pullback of an object  $l_{X/Y}$  on  $Y$  [20, VII.2.4.2.8]. Furthermore,  $l_{X/Y}$  in the fppf topology is the pullback of  $l_G$  via the morphism from  $Y$  to the stack  $BG$  corresponding to the  $G$ -torsor  $X \rightarrow Y$  [20, VII.3.1.2.6].

Applying this to  $E^{n+1}/G$  for any  $n \geq 0$ , we get an exact triangle

$$\pi^*(\Omega_{EE/G}^1) \rightarrow \Omega_{EE}^1 \rightarrow l_G$$

in  $D_G(EE)$ , or equivalently

$$l_G[-1] \rightarrow \pi^*(\Omega_{EE/G}^1) \rightarrow \Omega_{EE}^1.$$

It follows that for any  $j \geq 0$ ,  $\pi^*(\Omega_{EE/G}^j)$  has a filtration in the derived category with quotients  $\pi^*(\Omega_{EE}^{j-m}) \otimes \Lambda^m(l_G[-1])$  for  $m = 0, \dots, j$ .

If  $E(R)$  is nonempty, then  $H^i(EE, \Omega^j) \cong H^i(\text{Spec}(R), \Omega^j)$ , by Lemma 2.3. That group is zero unless  $i = j = 0$ , in which case it is  $R$ . By faithfully flat descent, the same conclusion holds under our weaker assumption that  $E \rightarrow \text{Spec}(R)$  is smooth and surjective. Therefore, in the filtration above, all objects but one have zero cohomology in all degrees over  $EE$ . We deduce that the homomorphism

$$H^i(EE, \Lambda^j(l_G[-1])) \rightarrow H^i(EE, \pi^*(\Omega_{EE/G}^j))$$

is an isomorphism of  $G$ -modules for all  $i$ . By Illusie's "d ecalage" isomorphism [18, Proposition I.4.3.2.1(i)], we can write  $S^j(l_G)[-j]$  instead of  $\Lambda^j(l_G[-1])$ .

The cochain complex  $O(EE)$  has cohomology  $R$  in degree 0 and 0 otherwise, by Lemma 2.3 again. So the complex of global sections of the trivial vector bundle

$S^j(l_G)$  over  $EE$  is isomorphic, in the derived category of  $G$ -modules, to the complex of  $G$ -modules  $S^j(l_G)$ . We conclude that the complex of sections of  $\pi^*(\Omega_{EE/G}^j)$  over  $EE$  is isomorphic to  $S^j(l_G)[-j]$  in the derived category of  $G$ -modules.

Finally, we observe that each  $G$ -module in this complex,

$$M := H^0(E^{n+1}, \pi^*(\Omega_{E^{n+1}/G}^j))$$

for  $n \geq 0$ , is acyclic (meaning that  $H^{>0}(G, M) = 0$ ). More generally, for any affine  $R$ -scheme  $Y$  with a free  $G$ -action such that  $Y/G$  is affine, and any quasi-coherent sheaf  $F$  on  $Y/G$ ,  $M := H^0(Y, \pi^*F)$  is acyclic. Indeed, this holds if  $Y \rightarrow Y/G$  is a trivial  $G$ -bundle, since then  $M = O(G) \otimes F$  and so  $M$  is acyclic [21, Lemma 4.7]. We can prove acyclicity in general by pulling the  $G$ -bundle over  $Y/G$  back to a  $G$ -bundle over  $Y$ , which is trivial; then  $H^{>0}(G, M) \otimes_{O(Y/G)} O(Y)$  is 0 by [21, Proposition 4.13], and so  $H^{>0}(G, M) = 0$  by faithfully flat descent.

We conclude that the complex computing  $H^*(BG, \Omega^j)$  is the same one that computes  $H^*(G, S^j(l_G)[-j])$ .  $\square$

## 4 Good filtrations

In this section, we explain how known results in representation theory imply calculations of the Hodge cohomology of classifying spaces in many cases, via Corollary 2.2. This is not logically necessary for the rest of the paper: Theorem 9.1 is a stronger calculation of Hodge cohomology, based on ideas from homotopy theory.

Let  $G$  be a split reductive group over a field  $k$ . (A textbook reference on split reductive groups is [24, Chapter 21].) A *Schur module* for  $G$  is a module of the form  $H^0(\lambda)$  for a dominant weight  $\lambda$ . By definition,  $H^0(\lambda)$  means  $H^0(G/B, L(\lambda))$ , where  $B$  is a Borel subgroup and  $L(\lambda)$  is the line bundle associated to  $\lambda$ . For  $k$  of characteristic zero, the Schur modules are exactly the irreducible representations of  $G$ . Kempf showed that the dimension of the Schur modules is independent of the characteristic of  $k$  [21, Chapter II.4]. They need not be irreducible in characteristic  $p$ , however.

A  $G$ -module  $M$  has a *good filtration* if there is a sequence of submodules  $0 \subset M_0 \subset M_1 \subset \dots$  such that  $M = \cup M_j$  and each quotient  $M_i/M_{i-1}$  is a Schur module. One good feature of Schur modules is that their cohomology groups are known, by Cline-Parshall-Scott-van der Kallen [21, Proposition 4.13]. Namely,

$$H^i(G, H^0(\lambda)) \cong \begin{cases} k & \text{if } i = 0 \text{ and } \lambda = 0 \\ 0 & \text{otherwise.} \end{cases}$$

As a result,  $H^i(G, M) = 0$  for all  $i > 0$  when  $M$  has a good filtration.

The following result was proved by Andersen-Jantzen and Donkin [13, Proposition and proof of Theorem 2.2], [21, II.4.22]. The statement on the ring of invariants incorporates earlier work by Kac and Weisfeiler. Say that a prime number  $p$  is *bad* for a reductive group  $G$  if  $p = 2$  and  $G$  has a simple factor not of type  $A_n$ ,  $p = 3$  and  $G$  has a simple factor of exceptional type, or  $p = 5$  and  $G$  has an  $E_8$  factor. Otherwise,  $p$  is *good* for  $G$ .

**Theorem 4.1.** *Let  $G$  be a split reductive group over a field  $k$ . Assume either that  $G$  is a simply connected semisimple group and  $\text{char}(k)$  is good for  $G$ , or that  $G = GL(n)$ . Then the polynomial ring  $O(\mathfrak{g}) = S(\mathfrak{g}^*)$  has a good filtration as a  $G$ -module, and the ring of invariants  $O(\mathfrak{g})^G$  is a polynomial ring over  $k$ , with generators in the fundamental degrees of  $G$ .*

It follows that, under these assumptions,  $H^{>0}(G, S^j(\mathfrak{g}^*))$  is zero for all  $j \geq 0$ . Equivalently,  $H^i(BG, \Omega^j) = 0$  for  $i \neq j$ , by Corollary 2.2. We prove this under the weaker assumption that  $p$  is not a torsion prime in Theorem 9.1.

## 5 Künneth formula

The Künneth formula holds for Hodge cohomology, in the following form. The hypotheses apply to the main case studied in this paper: classifying stacks  $BG$  with  $G$  an affine group scheme of finite type over a field.

**Proposition 5.1.** *Let  $X$  and  $Y$  be quasi-compact algebraic stacks with affine diagonal over a field  $k$ . Then*

$$H_{\mathbb{H}}^*((X \times_k Y)/k) \cong H_{\mathbb{H}}^*(X/k) \otimes_k H_{\mathbb{H}}^*(Y/k).$$

*Proof.* Since  $X$  and  $Y$  are quasi-compact, there are affine schemes  $A$  and  $B$  with smooth surjective morphisms  $A \rightarrow X$  and  $B \rightarrow Y$  [32, Tag 04YA]. Since  $X$  and  $Y$  have affine diagonal, the fiber products  $A_X^{n+1}$  and  $B_Y^{n+1}$  are affine over the products  $A^{n+1}$  and  $B^{n+1}$  over  $k$ , and so they are affine schemes, for all  $n \geq 0$ .

The morphism  $A \times B \rightarrow X \times Y$  is smooth and surjective. Therefore, the Hodge cohomology of  $X \times Y$  is the cohomology of the Čech nerve  $C(A \times B/X \times Y)$  over  $k$ , with coefficients in  $\Omega^*$  (with zero differential). This space is the product  $C(A/X) \times C(B/Y)$  over  $k$ . By the previous paragraph, these are in fact simplicial affine schemes over  $k$ .

The quasi-coherent sheaf  $\Omega^1$  on the product of two affine schemes over  $k$  is the direct sum of the pullbacks of  $\Omega^1$  from the two factors. (No smoothness is needed for this calculation.) Therefore, the quasi-coherent sheaf  $\Omega^*$  on the product affine scheme  $A_X^{n+1} \times B_Y^{n+1}$  over  $k$  is the tensor product of the pullbacks on  $\Omega^*$  on those two schemes. So  $H^0(A_X^{n+1} \times B_Y^{n+1}, \Omega^*)$  is the tensor product of  $H^0(A_X^{n+1}, \Omega^*)$  and  $H^0(B_Y^{n+1}, \Omega^*)$  over  $k$ .

The spectral sequence of the simplicial scheme  $C(A/X) \times C(B/Y)$  with coefficients in  $\Omega^*$  reduces to one row, since all the schemes here are affine. Explicitly, by the previous paragraph, the cohomology of the product simplicial scheme is the cohomology of the tensor product over  $k$  of the two cosimplicial vector spaces  $H^0(A_X^{n+1}, \Omega^*)$  and  $H^0(B_Y^{n+1}, \Omega^*)$ . By the Eilenberg-Zilber theorem, it follows that the cohomology of the product simplicial scheme is the tensor product over  $k$  of the cohomology of the two factors. [23, Theorem 29.3]. Equivalently,

$$H_{\mathbb{H}}^*((X \times_k Y)/k) \cong H_{\mathbb{H}}^*(X/k) \otimes_k H_{\mathbb{H}}^*(Y/k).$$

□

## 6 Parabolic subgroups

**Theorem 6.1.** *Let  $P$  be a parabolic subgroup of a reductive group  $G$  over a field  $k$ , and let  $L$  be the Levi quotient of  $P$  (the quotient of  $P$  by its unipotent radical). Then the restriction*

$$H^i(BP, \Omega^j) \rightarrow H^i(BL, \Omega^j)$$

*is an isomorphism for all  $i$  and  $j$ . Equivalently,*

$$H^a(P, S^j(\mathfrak{p}^*)) \rightarrow H^a(L, S^j(\mathfrak{l}^*))$$

*is an isomorphism for all  $a$  and  $j$ .*

Theorem 6.1 can be viewed as a type of homotopy invariance for Hodge cohomology of classifying spaces. This is not automatic, since Hodge cohomology is not  $A^1$ -homotopy invariant for smooth varieties. Homotopy invariance of Hodge cohomology also fails in general for classifying spaces. For example, let  $G_a$  be the additive group over a field  $k$ . Then the Hodge cohomology group  $H^1(BG_a, \mathcal{O})$  is not zero for any  $k$ , and it is a  $k$ -vector space of infinite dimension for  $k$  of positive characteristic; this follows from Theorem 6.3, due to Cline, Parshall, Scott, and van der Kallen, together with Corollary 2.2.

*Proof.* (Theorem 6.1) Let  $U$  be the unipotent radical of  $P$ , so that  $L = P/U$ . It suffices to show that

$$H^a(P, S^j(\mathfrak{p}^*)) \rightarrow H^a(L, S^j(\mathfrak{l}^*))$$

is an isomorphism after extending the field  $k$ . So we can assume that  $G$  has a Borel subgroup  $B$  and that  $B$  is contained in  $P$ . Let  $R$  be the set of roots for  $G$ . We follow the convention that the weights of  $B$  acting on the Lie algebra of its unipotent radical are the *negative* roots  $R^-$ . There is a subset  $I$  of the set  $S$  of simple roots so that  $P$  is the associated subgroup  $P_I$ , in the notation of [21, II.1.8]. More explicitly, let  $R_I = R \cap \mathbf{Z}I$ ; then  $P = P_I$  is the semidirect product  $U_I \rtimes L_I$ , where  $L_I$  is the reductive group  $G(R_I)$  and  $U := U_I$  is the unipotent group  $U(R^- \setminus R_I)$ .

As a result, the weights of  $P$  on  $\mathfrak{p}$  are all the roots  $\sum_{\alpha \in S} n_\alpha \alpha$  such that  $n_\alpha \leq 0$  for  $\alpha$  not in  $I$ . The coefficients  $n_\alpha$  for  $\alpha$  not in  $I$  are all zero exactly for the weights of  $P$  on  $\mathfrak{p}/\mathfrak{u}$ . As a result, for any  $j \geq 0$ , the weights of  $P$  on  $S^j(\mathfrak{p}^*)$  are all in the root lattice, with nonnegative coefficients for the simple roots not in  $I$ , and with those coefficients all zero only for the weights of  $P$  on the subspace  $S^j((\mathfrak{p}/\mathfrak{u})^*) \subset S^j(\mathfrak{p}^*)$ .

We now use the following information about the cohomology of  $P$ -modules [21, Proposition II.4.10]. For any element  $\lambda$  of the root lattice  $\mathbf{Z}S$ ,  $\lambda = \sum_{\alpha \in S} n_\alpha \alpha$ , the *height*  $\text{ht}(\lambda)$  means the integer  $\sum_{\alpha \in S} n_\alpha$ .

**Proposition 6.2.** *Let  $P$  be a parabolic subgroup of a reductive group  $G$  over a field, and let  $M$  be a  $P$ -module. If  $H^j(P, M) \neq 0$  for some  $j \geq 0$ , then there is a weight  $\lambda$  of  $M$  with  $-\lambda \in \mathbf{N}R^+$  and  $\text{ht}(-\lambda) \geq j$ .*

As mentioned above, for any  $j \geq 0$ , every weight of  $P$  on  $M := \text{coker}(S^j((\mathfrak{p}/\mathfrak{u})^*) \rightarrow S^j(\mathfrak{p}^*))$  has at least one positive coefficient in terms of the simple roots. By Proposition 6.2, it follows that  $H^a(P, M) = 0$  for all  $a$ . Therefore, the homomorphism

$$H^a(P, S^j((\mathfrak{p}/\mathfrak{u})^*)) \rightarrow H^a(P, S^j(\mathfrak{p}^*))$$

is an isomorphism for all  $a$  and  $j$ . Here  $\mathfrak{p}/\mathfrak{u} \cong \mathfrak{l}$  is a representation of the quotient group  $L = P/U$ . It remains to show that the pullback

$$H^a(L, S^j((\mathfrak{p}/\mathfrak{u})^*)) \rightarrow H^a(P, S^j((\mathfrak{p}/\mathfrak{u})^*))$$

is an isomorphism. This would not be true for an arbitrary representation of  $L$ ; we will have to use what we know about the weights of  $L$  on  $S^j((\mathfrak{p}/\mathfrak{u})^*)$ .

We also use the following description of the cohomology of an additive group  $V = (G_a)^n$  over a perfect field  $k$  [21, Proposition I.4.27]. (To prove Theorem 6.1, we can enlarge the field  $k$ , and so we can assume that  $k$  is perfect.) The following description is canonical, with respect to the action of  $GL(V)$  on  $H^*(V, k)$ . Write  $V^{(j)}$  for the  $j$ th Frobenius twist of a vector space  $V$ , as a representation of  $GL(V)$ .

**Theorem 6.3.** (1) *If  $k$  has characteristic zero, then  $H^*(V, k) \cong \Lambda(V^*)$ , with  $V^*$  in degree 1.*

(2) *If  $k$  has characteristic 2, then*

$$H^*(V, k) \cong S(\oplus_{j \geq 0} (V^*)^{(j)}),$$

*with all the spaces  $(V^*)^{(j)}$  in degree 1.*

(3) *If  $k$  has characteristic  $p > 2$ , then*

$$H^*(V, k) \cong \Lambda(\oplus_{j \geq 0} (V^*)^{(j)}) \otimes S(\oplus_{j \geq 1} (V^*)^{(j)}),$$

*with all the spaces  $(V^*)^{(j)}$  in the first factor in degree 1, and all the spaces  $(V^*)^{(j)}$  in the second factor in degree 2.*

We also use the Hochschild-Serre spectral sequence for the cohomology of algebraic groups [21, I.6.5, Proposition I.6.6]:

**Theorem 6.4.** *Let  $G$  be an affine group scheme of finite type over a field  $k$ , and let  $N$  be a normal  $k$ -subgroup scheme of  $G$ . For every  $G$ -module (or complex of  $G$ -modules)  $V$ , there is a spectral sequence*

$$E_2^{ij} = H^i(G/N, H^j(N, V)) \Rightarrow H^{i+j}(G, V).$$

Theorems 6.3 and 6.4 give information about the weights of  $L$  on  $H^*(U, k)$ , that is, about the action of a maximal torus  $T \subset L$  on  $H^*(U, k)$ . The method is to write  $U$  (canonically) as an extension of additive groups  $V = (G_a)^n$  and use the Hochschild-Serre spectral sequence. We deduce that as a representation of  $L$ , all weights of  $H^{>0}(U, k)$  are in the root lattice of  $G$ , with nonnegative coefficients for the simple roots not in  $I$ , and with at least one of those coefficients positive. (This is the same sign as we have for the action of  $L$  on  $\mathfrak{u}^*$ .)

Now apply the Hochschild-Serre spectral sequence to the normal subgroup  $U$  in  $P$ :

$$E_2^{ij}(L, H^j(U, k) \otimes S^l((\mathfrak{p}/\mathfrak{u})^*)) \Rightarrow H^{i+j}(P, S^l((\mathfrak{p}/\mathfrak{u})^*)).$$

By the analysis of  $S^l(\mathfrak{p}^*)$  above, all the weights of  $L$  on the subspace  $S^l((\mathfrak{p}/\mathfrak{u})^*)$  are in the root lattice of  $G$ , and the coefficients of all simple roots not in  $I$  are equal to zero. Combining this with the previous paragraph, we find: for  $l \geq 0$  and  $j > 0$ , all

weights of  $L$  on  $H^j(U, k) \otimes S^l((\mathfrak{p}/\mathfrak{u})^*)$  have all coefficients of the simple roots not in  $I$  nonnegative, with at least one positive. By Proposition 6.2, it follows that

$$H^i(L, H^j(U, k) \otimes S^l((\mathfrak{p}/\mathfrak{u})^*)) = 0$$

for all  $i$  and  $l$  and all  $j > 0$ . So the spectral sequence above reduces to an isomorphism

$$H^i(P, S^l((\mathfrak{p}/\mathfrak{u})^*)) \cong H^i(L, S^l((\mathfrak{p}/\mathfrak{u})^*)),$$

as we wanted. Theorem 6.1 is proved.  $\square$

## 7 Hodge cohomology of flag manifolds

We use the following result, proved by Srinivas [31, section 3]:

**Proposition 7.1.** *Let  $P$  be a parabolic subgroup of a split reductive group  $G$  over a field  $k$ . Then the cycle map*

$$CH^*(G/P) \otimes_{\mathbf{Z}} k \rightarrow H_{\mathbf{H}}^*((G/P)/k)$$

*is an isomorphism of  $k$ -algebras. In particular,  $H^i(G/P, \Omega^j) = 0$  for  $i \neq j$ .*

There are many related results. In particular, Proposition 7.1 can also be deduced from the work of El Zein (who constructed the cycle map in Hodge cohomology over any field) and Gros (who constructed the pushforward homomorphism in Hodge cohomology over any perfect field) [15, Proposition 3.3.5], [16, sections II.2 and II.4]. That approach implies Proposition 7.1 more generally for any smooth proper variety with a cell decomposition. Also, Andersen gave the additive calculation of  $H^i(G/P, \Omega^j)$  over any field [21, Proposition II.6.18].

Note that Chevalley and Demazure gave combinatorial descriptions of the Chow ring of  $G/P$ , which in particular show that this ring is independent of  $k$ , and isomorphic to the ordinary cohomology ring  $H^*(G_{\mathbf{C}}/P_{\mathbf{C}}, \mathbf{Z})$  [10, Proposition 11], [12]. (That makes sense because the classification of split reductive groups and their parabolic subgroups is the same over all fields.)

## 8 Invariant functions on the Lie algebra

**Theorem 8.1.** *Let  $G$  be a simple group over a field  $k$ ,  $T$  a maximal torus in  $G$ ,  $\mathfrak{g}$  and  $\mathfrak{t}$  the Lie algebras. Assume that we are not in the case where  $\text{char}(k) = 2$  and  $G_{\bar{k}}$  is a product of copies of  $Sp(2n)$  for some positive integer  $n$ . Then the restriction  $O(\mathfrak{g})^G \rightarrow O(\mathfrak{t})^W$  is an isomorphism.*

Theorem 8.1 was proved by Springer and Steinberg for any adjoint group  $G$  [30, II.3.17'], and generalized to any simple group by Chaput and Romagny [9, Theorem 1.1]. They assumed that  $G$  is split, but that implies Theorem 8.1 by passage to the algebraic closure  $\bar{k}$ .

The statement is optimal, in that the conclusion fails for the symplectic group  $Sp(2n)$  in characteristic 2 for any positive integer  $n$  (for example, for  $SL(2)$ ), as Chaput and Romagny pointed out. In their argument, the distinctive feature of

the symplectic group is that it is the only simple group for which some roots are divisible by 2 in the weight lattice.

In particular, Theorem 8.1 applies to cases such as the spin group  $\text{Spin}(n)$  in characteristic 2 with  $n \geq 6$ , which we study further in Theorem 12.1.

Here is a related observation.

**Lemma 8.2.** *Let  $G$  be a smooth affine group over a field  $k$  whose identity component is reductive. Then there are canonical maps  $H^a(BG, \Omega^a) \rightarrow H_{\text{dR}}^{2a}(BG/k)$  and  $H^{a+1}(BG, \Omega^a) \rightarrow H_{\text{dR}}^{2a+1}(BG/k)$ . These maps are compatible with products and with pullback under group homomorphisms.*

*Proof.* By Corollary 2.2, we have  $H^a(BG, \Omega^b) = 0$  for all  $a < b$ . Therefore, the Hodge spectral sequence gives a canonical “edge map”  $H^a(BG, \Omega^a) \rightarrow H_{\text{dR}}^{2a}(BG/k)$ .

In odd degrees, there is one possible differential on the group  $H^{a+1}(BG, \Omega^a)$ :

$$d_1: H^{a+1}(BG, \Omega^a) \rightarrow H^{a+1}(BG, \Omega^{a+1}).$$

In fact, this differential is zero. To see this, let  $T$  be a maximal torus in  $G$ . Then the restriction  $O(\mathfrak{g})^G \rightarrow O(\mathfrak{t})$  is injective, because the  $G$ -conjugates of elements of  $\mathfrak{t}$  (over the algebraic closure of  $k$ ) are the semisimple elements of  $\mathfrak{g}$ , which are Zariski dense in  $\mathfrak{g}$ . Equivalently,  $H^b(BG, \Omega^b) \rightarrow H^b(BT, \Omega^b)$  is injective for all  $b$ . But  $H^a(BT, \Omega^b) = 0$  for all  $a \neq b$ , and so  $H^b(BT, \Omega^b)$  injects into  $H_{\text{dR}}^{2b}(BT/k)$ . Therefore,  $H^b(BG, \Omega^b)$  injects into  $H_{\text{dR}}^{2b}(BG/k)$ . In particular, the  $d_1$  differential into  $H^{a+1}(BG, \Omega^{a+1})$  is zero, as we want. Therefore, we have a canonical “edge map”  $H^{a+1}(BG, \Omega^a) \rightarrow H_{\text{dR}}^{2a+1}(BG/k)$ .  $\square$

## 9 Hodge cohomology of $BG$ at non-torsion primes

**Theorem 9.1.** *Let  $G$  be a reductive group over a field  $k$  of characteristic  $p \geq 0$ . Then  $H^{>0}(G, O(\mathfrak{g})) = 0$  if and only if  $p$  is not a torsion prime for  $G$ .*

**Theorem 9.2.** *Let  $G$  be a split reductive group over  $\mathbf{Z}$ , and let  $p$  be a non-torsion prime for  $G$ . Then  $H^j(BG_{\mathbf{Z}}, \Omega^i)$  localized at  $p$  is zero for  $i \neq j$ . Moreover, the Hodge cohomology ring  $H^*(BG_{\mathbf{Z}}, \Omega^*)$  and the de Rham cohomology  $H_{\text{dR}}^*(BG/\mathbf{Z})$ , localized at  $p$ , are polynomial rings on generators of degrees equal to 2 times the fundamental degrees of  $G$ . These rings are isomorphic to the cohomology of the topological space  $BG_{\mathbf{C}}$  with  $\mathbf{Z}_{(p)}$  coefficients.*

We recall the definition of torsion primes for a reductive group  $G$  over a field  $k$ . Let  $B$  be a Borel subgroup of  $G_{\bar{k}}$ , and  $T$  a maximal torus in  $B$ . Then there is a natural homomorphism from the character group  $X^*(T) = \text{Hom}(T, G_m)$  (the weight lattice of  $G$ ) to the Chow group  $CH^1(G_{\bar{k}}/B)$ . Therefore, for  $N = \dim(G_{\bar{k}}/B)$ , there is a homomorphism from the symmetric power  $S^N(X^*(T))$  to  $CH^N(G_{\bar{k}}/B)$ ; taking the degree of a zero-cycle on  $G_{\bar{k}}/B$  gives a homomorphism (in fact, an isomorphism)  $CH^N(G_{\bar{k}}/B) \rightarrow \mathbf{Z}$ . A prime number  $p$  is said to be a torsion prime for  $G$  if the image of  $S^N(X^*(T)) \rightarrow \mathbf{Z}$  is zero modulo  $p$ . Borel showed that  $p$  is a torsion prime for  $G$  if and only if the cohomology  $H^*(BG_{\mathbf{C}}, \mathbf{Z})$  has  $p$ -torsion, where  $G_{\mathbf{C}}$  is the corresponding complex reductive group [7, Proposition 4.2]. It is also equivalent to say that  $G(\mathbf{C})$  contains an elementary abelian  $p$ -subgroup that is not contained in a torus [7, Théorème 4.5].



In most cases, Theorem 9.1 follows from Theorem 4.1. Explicitly, a prime number  $p$  is torsion for a simply connected simple group  $G$  if  $p = 2$  and  $G$  has a simple factor not of type  $A_n$  or  $C_n$ ,  $p = 3$  and  $G$  has a simple factor of type  $F_4$ ,  $E_6$ ,  $E_7$ , or  $E_8$ , or  $p = 5$  and  $G$  has an  $E_8$  factor. So the main new cases in Theorem 9.1 are the symplectic groups  $Sp(2n)$  in characteristic 2 and  $G_2$  in characteristic 3. (These are non-torsion primes, but not good primes in the sense of Theorem 4.1.) In these cases, the representation-theoretic result that  $H^{>0}(G, O(\mathfrak{g})) = 0$  seems to be new. Does  $O(\mathfrak{g})$  have a good filtration in these cases?

The following spectral sequence, modeled on the Leray-Serre spectral sequence in topology, will be important for the rest of the paper.

**Proposition 9.3.** *Let  $P$  be a parabolic subgroup of a split reductive group  $G$  over a field  $k$ . Let  $L$  be the quotient of  $P$  by its unipotent radical. Then there is a spectral sequence of algebras*

$$E_2^{ij} = H_{\mathbb{H}}^i(BG/k) \otimes H_{\mathbb{H}}^j((G/P)/k) \Rightarrow H_{\mathbb{H}}^{i+j}(BL/k).$$

*Proof.* Consider  $\Omega^* = \bigoplus \Omega^i$  as a presheaf of commutative dgas on smooth  $k$ -schemes, with zero differential.

For a smooth morphism  $f: X \rightarrow Y$  of smooth  $k$ -schemes, consider the object  $Rf_*(\Omega_X^*)$  in the derived category  $D(Y)$  of étale sheaves on  $Y$ . Here the sheaf  $\Omega_X^*$  on  $X$  has an increasing filtration, compatible with its ring structure, with 0th step the subsheaf  $f^*(\Omega_Y^*)$  and  $j$ th graded piece  $f^*(\Omega_Y^*) \otimes \Omega_{X/Y}^j$ . So  $Rf_*(\Omega_X^*)$  has a corresponding filtration in  $D(Y)$ , with  $j$ th graded piece  $Rf_*(f^*(\Omega_Y^*) \otimes \Omega_{X/Y}^j) \cong \Omega_Y^* \otimes Rf_*\Omega_{X/Y}^j$ . This gives a spectral sequence

$$E_2^{ij} = H^{i+j}(Y, \Omega_Y^* \otimes Rf_*\Omega_{X/Y}^j) \Rightarrow H^{i+j}(X, \Omega_X^*).$$

Now specialize to the case where  $f: X \rightarrow Y$  is the  $G/P$ -bundle associated to a principal  $G$ -bundle over  $Y$ . The Hodge cohomology of  $G/P$  is essentially independent of the base field, by the isomorphism  $H_{\mathbb{H}}^*((G/P)/k) \cong CH^*(G/P) \otimes_{\mathbb{Z}} k$  (Proposition 7.1). Here  $CH^*(G/P)$  is a free abelian group with a fixed basis (independent of  $k$ ), as discussed in section 7, and  $G$  acts trivially on  $CH^*(G/P)$ . Therefore, each object  $Rf_*(\Omega_{X/Y}^j)$  is a trivial vector bundle on  $Y$ , with fiber  $H^j(G/P, \Omega^j)$ , viewed as a complex in degree  $j$ . So we can rewrite the spectral sequence as

$$E_2^{ij} = H^i(Y, \Omega^*) \otimes H^j(G/P, \Omega^j) \Rightarrow H^{i+j}(X, \Omega^*).$$

All differentials in the spectral sequence above preserve the degree in the grading of  $\Omega^*$ . Therefore, we can renumber the spectral sequence so that it is graded by total degree:

$$E_2^{ij} = H_{\mathbb{H}}^i(Y/k) \otimes H_{\mathbb{H}}^j((G/P)/k) \Rightarrow H_{\mathbb{H}}^{i+j}(X/k).$$

Finally, we consider the analogous spectral sequence for the morphism  $f: EG/P \rightarrow B_{\text{simp}}G$  of simplicial schemes:

$$E_2^{ij} = H_{\mathbb{H}}^i(BG/k) \otimes H_{\mathbb{H}}^j((G/P)/k) \Rightarrow H_{\mathbb{H}}^{i+j}((EG/P)/k).$$

By Lemma 1.2, the output of the spectral sequence is isomorphic to  $H_{\mathbb{H}}^*(BP/k)$ , or equivalently (by Theorem 6.1) to  $H_{\mathbb{H}}^*(BL/k)$ . This is a spectral sequence of algebras. All differentials preserve the degree in the grading of  $\Omega^*$ .  $\square$

*Proof.* (Theorem 9.1) First, suppose that  $H^{>0}(G, O(\mathfrak{g})) = 0$ ; then we want to show that  $\text{char}(k)$  is not a torsion prime for  $G$ . By Corollary 2.2, the assumption implies that  $H^j(BG, \Omega^i) = 0$  for all  $i \neq j$ . Apply Proposition 9.3 when  $P$  is a Borel subgroup  $B$  in  $G$ ; this gives a spectral sequence

$$E_2^{ij} = H_{\mathbb{H}}^i(BG/k) \otimes H_{\mathbb{H}}^j((G/B)/k) \Rightarrow H_{\mathbb{H}}^{i+j}(BT/k),$$

where  $T$  is a maximal torus in  $B$ . Under our assumption, this spectral sequence degenerates at  $E_2$ , because the differential  $d_r$  (for  $r \geq 2$ ) takes  $H^i(BG, \Omega^i) \otimes H^j(G/B, \Omega^j)$  into  $H^{i+r}(BG, \Omega^{i+r-1}) \otimes H^{j-r+1}(G/B, \Omega^{j-r+1})$ , which is zero. It follows that  $H_{\mathbb{H}}^*(BT/k) \rightarrow H_{\mathbb{H}}^*((G/B)/k)$  is surjective. Here  $H_{\mathbb{H}}^*(BT/k)$  is the polynomial ring  $S(X^*(T) \otimes k)$  by Theorem 4.1, and  $H_{\mathbb{H}}^*((G/B)/k) = CH^*(G/B) \otimes k$  by Proposition 7.1. It follows that the ring  $CH^*(G/B) \otimes k$  is generated as a  $k$ -algebra by the image of  $X^*(T) \rightarrow CH^1(G/B)$ . Equivalently,  $p$  is not a torsion prime for  $G$ .

Conversely, suppose that  $p$  is not a torsion prime for  $G$ . That is, the homomorphism  $S(X^*(T) \otimes k) \rightarrow CH^*(G/B) \otimes k$  is surjective. Equivalently,  $H_{\mathbb{H}}^*(BT/k) \rightarrow H_{\mathbb{H}}^*((G/B)/k)$  is surjective. By the product structure on the spectral sequence above, it follows that the spectral sequence degenerates at  $E_2$ . Since  $H^j(BT, \Omega^i) = 0$  for  $i \neq j$ , it follows that  $H^j(BG, \Omega^i) = 0$  for  $i \neq j$ . Equivalently,  $H^{>0}(G, O(\mathfrak{g})) = 0$ .  $\square$

*Proof.* (Theorem 9.2) Let  $G$  be a split reductive group over  $\mathbf{Z}$ , and let  $p$  be a non-torsion prime for  $G$ . We have a short exact sequence

$$0 \rightarrow H^j(BG_{\mathbf{Z}}, \Omega^i)/p \rightarrow H^j(BG_{\mathbf{F}_p}, \Omega^i) \rightarrow H^{j+1}(BG_{\mathbf{Z}}, \Omega^i)[p] \rightarrow 0.$$

By Theorem 9.1 and Corollary 2.2, the Hodge cohomology ring  $H^*(BG_{\mathbf{Z}}, \Omega^*)$  localized at  $p$  is concentrated in bidegrees  $H^{i,i}$  and is torsion-free. This ring tensored with  $\mathbf{Q}$  is the ring of invariants  $O(\mathfrak{g}_{\mathbf{Q}})^G$ , which is a polynomial ring on generators of degrees equal to the fundamental degrees of  $G$ .

To show that the Hodge cohomology ring over  $\mathbf{Z}_{(p)}$  is a polynomial ring on generators in  $H^{i,i}$  for  $i$  running through the fundamental degrees of  $G$ , it suffices to show that the Hodge cohomology ring  $H_{\mathbb{H}}^*(BG/\mathbf{F}_p)$  is a polynomial ring in the same degrees. Given that, the other statements of the theorem will follow. Indeed, the statement on Hodge cohomology implies that the de Rham cohomology ring  $H_{\text{dR}}^*(BG/\mathbf{Z})$  localized at  $p$  is also a polynomial ring, on generators in 2 times the fundamental degrees of  $G$ . The cohomology of the topological space  $BG_{\mathbf{C}}$  localized at  $p$  is known to be a polynomial ring on generators in the same degrees, by Borel [7, Proposition 4.2, Théorème 4.5], [25, Theorem VII.2.12].

From here on, let  $k = \mathbf{F}_p$ , and write  $G$  for  $G_k$ . By definition of the Weyl group  $W$  as  $W = N_G(T)/T$ , the image of  $H_{\mathbb{H}}^*(BG/k)$  in  $H_{\mathbb{H}}^*(BT/k) = S(X(T) \otimes k)$  is contained in the subring of  $W$ -invariants. We now use that  $p$  is not a torsion prime for  $G$ . By Demazure, except in the case where  $p = 2$  and  $G$  has an  $Sp(2n)$  factor, the ring of  $W$ -invariants in  $S(X(T) \otimes k)$  is a polynomial algebra over  $k$ , with the degrees of generators equal to the fundamental degrees of  $G$  [12, Théorème].

By Theorem 9.1,  $H_{\mathbb{H}}^*(BG/k)$  is equal to the ring of invariants  $O(\mathfrak{g})^G$ . By Theorem 8.1 (due to Chaput and Romagny), for any simple group  $G$  over a field  $k$  of characteristic  $p$  with  $p$  not a torsion prime, except for  $G = Sp(2n)$  with  $p = 2$ , the restriction  $O(\mathfrak{g})^G \rightarrow O(\mathfrak{t})^W$  is an isomorphism, and hence  $O(\mathfrak{g})^G$  is the polynomial ring we want.

The case of  $Sp(2n)$  in characteristic 2 (including  $SL(2) = Sp(2)$ ) is a genuine exception: here  $O(\mathfrak{g})^G$  is a subring of  $O(\mathfrak{t})^W$ , not equal to it. However, it is still true in this case that  $O(\mathfrak{g})^G$  is a polynomial ring with generators in the fundamental degrees of  $G$ , that is,  $2, 4, \dots, 2n$ , by Chaput and Romagny [9, Theorem 6.6].  $\square$

## 10 $\mu_p$

**Proposition 10.1.** *Let  $k$  be a field of characteristic  $p > 0$ . Let  $G$  be the group scheme  $\mu_p$  of  $p$ th roots of unity over  $k$ . Then*

$$H_{\mathbb{H}}^*(B\mu_p/k) \cong k[c_1]\langle v_1 \rangle,$$

where  $c_1$  is in  $H^1(B\mu_p, \Omega^1)$  and  $v_1$  is in  $H^0(B\mu_p, \Omega^1)$ . Likewise,  $H_{\text{dR}}^*(B\mu_p/k) \cong k[c_1]\langle v_1 \rangle$  with  $|v_1| = 1$  and  $|c_1| = 2$ .

Here  $R\langle v \rangle$  denotes the exterior algebra over a graded-commutative ring  $R$  with generator  $v$ ; that is,  $R\langle v \rangle = R \oplus R \cdot v$ , with product  $v^2 = 0$ . See section 1 for the definitions of Hodge and de Rham cohomology we are using for a non-smooth group scheme such as  $\mu_p$ . Proposition 10.1 can help to compute Hodge cohomology of  $BG$  for smooth group schemes  $G$ , as we will see in the proof of Theorem 11.1 for  $G = SO(n)$ .

Proposition 10.1 is roughly what the topological analogy would suggest. Indeed, the group scheme  $\mu_p$  of  $p$ th roots of unity is defined over  $\mathbf{Z}$ , with  $(\mu_p)_{\mathbf{C}}$  isomorphic to the group  $\mathbf{Z}/p$ . For  $k$  of characteristic  $p$ , the ring  $H^*((B\mu_p)_{\mathbf{C}}, k)$  is a polynomial ring  $k[x]$  with  $|x| = 1$  if  $p = 2$ , or a free graded-commutative algebra  $k\langle x, y \rangle$  with  $|x| = 1$  and  $|y| = 2$  if  $p$  is odd. So  $H_{\text{dR}}^*(B\mu_p/k)$  is isomorphic to  $H^*((B\mu_p)_{\mathbf{C}}, k)$  additively for any prime  $p$ , and as a graded ring if  $p > 2$ .

*Proof.* Let  $G = \mu_p$  over  $k$ . The co-Lie complex  $l_G$  in the derived category of  $G$ -modules, discussed in section 3, has  $H^0(l_G) \cong \mathfrak{g}^* \cong k$  and also  $H^{-1}(l_G) \cong k$ , with other cohomology groups being zero. (In short, this is because  $G$  is a complete intersection in the affine line, defined by the one equation  $x^p = 1$ .)

Since representations of  $G$  are completely reducible, we have  $\text{Ext}_G^{>0}(M, N) = 0$  for all  $G$ -modules  $M$  and  $N$  [21, Lemma I.4.3]. The isomorphism class of  $l_G$  is described by an element of  $\text{Ext}_G^2(k, k)$ , which is zero. So  $l_G \cong k \oplus k[1]$  in the derived category of  $G$ -modules.

By Theorem 3.1, we have

$$H^i(BG, \Omega^j) \cong H^{i-j}(G, S^j(l_G)).$$

Here

$$\begin{aligned} S^j(l_G) &\cong \bigoplus_{m=0}^j S^m(k) \otimes S^{j-m}(k[1]) \\ &\cong \bigoplus_{m=0}^j S^m(k) \otimes \Lambda^{j-m}(k)[j-m], \end{aligned}$$

which is isomorphic to  $k \oplus k[1]$  if  $j \geq 1$  and to  $k$  if  $j = 0$ . Therefore,  $H^i(BG, \Omega^j)$  is isomorphic to  $k$  if  $0 \leq i = j$  or if  $0 \leq i = j - 1$ , and is otherwise zero.

Write  $c_1$  for the generator of  $H^1(BG, \Omega^1)$ , which is pulled back from the Chern class  $c_1$  in  $H^1(BG_m, \Omega^1)$  via the inclusion  $G \hookrightarrow G_m$ . Write  $v_1$  for the generator of

$H^0(BG, \Omega^1)$ . We have  $v_1^2 = 0$  because  $H^0(BG, \Omega^2) = 0$ . Theorem 3.1 also describes the ring structure on the Hodge cohomology of  $BG$ . In particular,  $\bigoplus_i H^i(BG, \Omega^i)$  is the ring of invariants of  $G$  acting on  $O(\mathfrak{g})$ , which is the polynomial ring  $k[c_1]$ . Finally, the description of  $S^j(l_G)$  also shows that  $\bigoplus_i H^i(BG, \Omega^{i+1})$  is the free module over  $k[c_1]$  on the generator  $v_1$ . This completes the proof that

$$H_{\mathbb{H}}^*(BG/k) \cong k[c_1]\langle v_1 \rangle.$$

Finally, consider the Hodge spectral sequence for  $BG$  from section 1. The element  $v_1$  is a permanent cycle because  $H^0(BG, \Omega^2) = 0$ , and  $c_1$  is a permanent cycle because it is pulled back from a permanent cycle on  $BG_m$ . Therefore, the Hodge spectral sequence degenerates at  $E_1$ . We have  $v_1^2 = 0$  in de Rham cohomology as in Hodge cohomology, because  $\bigoplus_i H^0(BG, \Omega^i)$  is a subring of de Rham cohomology, using degeneration of the Hodge spectral sequence. Therefore, the de Rham cohomology of  $BG$  is isomorphic to  $k[c_1]\langle v_1 \rangle$  as a graded ring.  $\square$

**Lemma 10.2.** *Let  $G$  be a discrete group, considered as a group scheme over a field  $k$ . Then the Hodge cohomology of the algebraic stack  $BG$  is the group cohomology of  $G$ :*

$$H^i(BG, \Omega^j) \cong \begin{cases} H^i(G, k) & \text{if } j = 0 \\ 0 & \text{otherwise.} \end{cases}$$

It follows that  $H_{\text{dR}}^*(BG/k) \cong H^*(G, k)$ .

*Proof.* Since  $G$  is smooth over  $k$ , we can compute the Hodge cohomology of the stack  $BG$  as the étale cohomology of the simplicial scheme  $B_{\text{simp}}G$  with coefficients in  $\Omega^j$ . Since  $G$  is discrete, the sheaf  $\Omega^j$  is zero for  $j > 0$ . For  $j = 0$ , the spectral sequence

$$E_1^{ab} = H^b(G^a, O) \Rightarrow H^{a+b}(BG, O)$$

reduces to a single row, since  $H^b(G^a, O) = 0$  for  $b > 0$ . That is,  $H^*(BG, O)$  is the cohomology of the standard complex that computes the cohomology of the group  $G$  with coefficients in  $k$ .  $\square$

Although Lemma 10.2 applies to any discrete group  $G$ , it is probably most meaningful in the case where  $G$  is finite (so that the associated group scheme over a field  $k$  is affine). A generalization of that case is the following ‘‘Hochschild-Serre’’ spectral sequence for the Hodge cohomology of a non-connected group scheme:

**Lemma 10.3.** *Let  $G$  be an affine group scheme of finite type over a field  $k$ . Let  $G^0$  be the identity component of  $G$ , and suppose that the finite group scheme  $G/G^0$  is the  $k$ -group scheme associated to a finite group  $Q$ . Then there is a spectral sequence*

$$E_2^{ij} = H^i(Q, H^j(BG^0, \Omega^a)) \Rightarrow H^{i+j}(BG, \Omega^a).$$

for any  $a \geq 0$ .

*Proof.* By Theorem 3.1,  $H^r(BG, \Omega^a)$  is isomorphic to  $H^{r-a}(G, S^a(l_G))$ . The lemma then becomes a special case of the Hochschild-Serre spectral sequence for the cohomology of  $G$  as an algebraic group, Theorem 6.4:

$$E_2^{ij} = H^i(Q, H^j(G^0, S^a(l_G))) \Rightarrow H^{i+j}(G, S^a(l_G)).$$

(Strictly speaking, move the Hochschild-Serre spectral sequence up by  $a$  rows to obtain the spectral sequence of the lemma.)  $\square$

## 11 The orthogonal groups

**Theorem 11.1.** *Let  $G$  be the split group  $SO(n)$  (also called  $O^+(n)$ ) over a field  $k$  of characteristic 2. Then the Hodge cohomology ring of  $BG$  is a polynomial ring  $k[u_2, u_3, \dots, u_n]$ , where  $u_{2a}$  is in  $H^a(BG, \Omega^a)$  and  $u_{2a+1}$  is in  $H^{a+1}(BG, \Omega^a)$ . Also, the Hodge spectral sequence degenerates at  $E_1$ , and so  $H_{\text{dR}}^*(BG/k)$  is also isomorphic to  $k[u_2, u_3, \dots, u_n]$ .*

*Likewise, the Hodge and de Rham cohomology rings of  $BO(2r)$  are isomorphic to the polynomial ring  $k[u_1, u_2, \dots, u_{2r}]$ . Finally, the Hodge and de Rham cohomology rings of  $BO(2r+1)$  are isomorphic to  $k[v_1, c_1, u_2, \dots, u_{2r+1}]/(v_1^2)$ , where  $v_1$  is in  $H^0(BO(2r+1), \Omega^1)$  and  $c_1$  is in  $H^1(BO(2r+1), \Omega^1)$ .*

Thus the de Rham cohomology ring of  $BSO(n)_{\mathbf{F}_2}$  is isomorphic to the mod 2 cohomology ring of the topological space  $BSO(n)_{\mathbf{C}}$  as a graded ring:

$$H^*(BSO(n)_{\mathbf{C}}, \mathbf{F}_2) \cong \mathbf{F}_2[w_2, w_3, \dots, w_n],$$

where the classes  $w_i$  are the Stiefel-Whitney classes. Theorem 11.1 gives a new analog of the Stiefel-Whitney classes for quadratic bundles in characteristic 2. (Note that the  $k$ -group scheme  $O(2r+1)$  is not smooth in characteristic 2. Indeed, it is isomorphic to  $SO(2r+1) \times \mu_2$ . By contrast,  $O(2r)$  is smooth but not connected, and we write  $SO(2r)$  for the kernel of the Dickson determinant  $O(2r) \rightarrow \mathbf{Z}/2$ , which describes the action of  $O(2r)$  on the center  $k \times k$  of the even Clifford algebra.)

The proof is inspired by topology. In particular, it involves some hard work with spectral sequences, related to Borel's transgression theorem and Zeeman's comparison theorem. The method should be useful for other reductive groups.

The formula for the classes  $u_i$  of a direct sum of two quadratic bundles is not the same as for the Stiefel-Whitney classes in topology. To state this, define a quadratic form  $(q, V)$  over a field  $k$  to be *nondegenerate* if the radical  $V^\perp$  of the associated bilinear form is zero, and *nonsingular* if  $V^\perp$  has dimension at most 1 and  $q$  is nonzero on any nonzero element of  $V^\perp$ . (In characteristic not 2, nonsingular and nondegenerate are the same.) The orthogonal group is defined as the automorphism group scheme of a nonsingular quadratic form; the precise group over  $k$  depends on the choice of form [22, section VI.23]. For example, over a field  $k$  of characteristic 2, the quadratic form

$$x_1x_2 + x_3x_4 + \dots + x_{2r-1}x_{2r}$$

is nonsingular of even dimension  $2r$ , while the form

$$x_1x_2 + x_3x_4 + \dots + x_{2r-1}x_{2r} + x_{2r+1}^2$$

is nonsingular of odd dimension  $2r+1$ , with  $V^\perp$  of dimension 1. (These particular forms define the *split* orthogonal groups over  $k$ .) Let  $u_0 = 1$ .

**Proposition 11.2.** *Let  $X$  be a scheme of finite type over a field  $k$  of characteristic 2. Let  $E$  and  $F$  be vector bundles with nondegenerate quadratic forms over  $X$  (hence of even rank). Write  $u_i$  for the characteristic classes from Theorem 11.1. Then, for any  $a \geq 0$ , in either Hodge cohomology or de Rham cohomology,*

$$u_{2a}(E \oplus F) = \sum_{j=0}^a u_{2j}(E)u_{2a-2j}(F)$$

and

$$u_{2a+1}(E \oplus F) = \sum_{l=0}^{2a+1} u_l(E)u_{2a+1-l}(F).$$

Thus the *even*  $u$ -classes of  $E \oplus F$  depend only on the even  $u$ -classes of  $E$  and  $F$ . By contrast, Stiefel-Whitney classes in topology satisfy

$$w_m(E \oplus F) = \sum_{l=0}^m w_l(E)w_{m-l}(F)$$

for all  $m$  [25, Theorem III.5.11].

Theorem 12.1 gives an example of a reductive group  $G$  for which the de Rham cohomology of  $BG_{\mathbf{F}_p}$  and the mod  $p$  cohomology of  $BG_{\mathbf{C}}$  are not isomorphic. It is a challenge to find out how close these rings are, in other examples.

Via Corollary 2.2, Theorem 11.1 can be viewed as a calculation in the representation theory of the algebraic group  $G = SO(n)$  for any  $n$ , over a field  $k$  of characteristic 2. For example, when  $G = SO(3) = PGL(2)$  over  $k$  of characteristic 2, we find (what seems to be new):

$$H^i(G, S^j(\mathfrak{g}^*)) \cong \begin{cases} k & \text{if } 0 \leq i \leq j \\ 0 & \text{otherwise.} \end{cases}$$

*Proof.* (Theorem 11.1) We will assume that  $k = \mathbf{F}_2$ . This implies the theorem for any field of characteristic 2.

We begin by computing the ring  $\oplus_i H^i(BG, \Omega^i)$  for  $G = SO(n)$ . By Corollary 2.2, this is equal to the ring of  $G$ -invariant polynomial functions on the Lie algebra  $\mathfrak{g}$  over  $k$ . By Theorem 8.1, since  $G$  is not a symplectic group, the restriction  $O(\mathfrak{g})^G \rightarrow O(\mathfrak{t})^W$  is an isomorphism.

Let  $r = \lfloor n/2 \rfloor$ . For  $n = 2r + 1$ , the Weyl group  $W$  is the semidirect product  $S_r \ltimes (\mathbf{Z}/2)^r$ . There is a basis  $e_1, \dots, e_r$  for  $\mathfrak{t}$  on which  $(\mathbf{Z}/2)^r$  acts by changing the signs, and so that action is trivial since  $k$  has characteristic 2. The group  $S_r$  has its standard permutation action on  $e_1, \dots, e_r$ . Therefore, the ring of invariants  $O(\mathfrak{t})^W$  is the ring of symmetric functions in  $r$  variables. Let  $u_2, u_4, \dots, u_{2r}$  denote the elementary symmetric functions. By the isomorphisms mentioned, we can view  $u_{2a}$  as an element of  $H^a(BSO(2r+1), \Omega^a)$  for  $1 \leq a \leq r$ , and  $\oplus_i H^i(BSO(2r+1), \Omega^i)$  is the polynomial ring  $k[u_2, u_4, \dots, u_{2r}]$ .

For  $n = 2r$ , the Weyl group  $W$  of  $SO(2r)$  is the semidirect product  $S_r \ltimes (\mathbf{Z}/2)^{r-1}$ . Again, the subgroup  $(\mathbf{Z}/2)^{r-1}$  acts trivially on  $\mathfrak{t}$ , and  $S_r$  acts by permutations as usual. So  $\oplus_i H^i(BSO(2r), \Omega^i)$  is also the polynomial ring  $k[u_2, u_4, \dots, u_{2r}]$ , with  $u_{2a}$  in  $H^a(BSO(2r), \Omega^a)$  for  $1 \leq a \leq r$ .

For the smooth  $k$ -group  $G = O(2r)$ , we can also compute the ring  $\oplus_i H^i(BG, \Omega^i)$ . By Corollary 2.2, this is the ring of  $G$ -invariant polynomial functions on the Lie algebra  $\mathfrak{g} = \mathfrak{so}(2r)$ . This is contained in the ring of  $SO(2r)$ -invariant functions on  $\mathfrak{g}$ , and I claim that the two rings are equal. It suffices to show that an  $SO(2r)$ -invariant function on  $\mathfrak{g}$  is also invariant under the normalizer  $N$  in  $O(2r)$  of a maximal torus  $T$  in  $SO(2r)$ , since that normalizer meets both connected components of  $O(2r)$ . Here  $N = S_r \ltimes (\mathbf{Z}/2)^r$ , which acts on  $\mathfrak{t}$  in the obvious way; in particular,  $(\mathbf{Z}/2)^r$  acts

trivially on  $\mathfrak{t}$ . Therefore, an  $SO(2r)$ -invariant function on  $\mathfrak{g}$  (corresponding to an  $S_r$ -invariant function on  $\mathfrak{t}$ ) is also  $O(2r)$ -invariant. Thus we have  $\oplus_i H^i(BO(2r), \Omega^i) = k[u_2, u_4, \dots, u_{2r}]$ .

For a smooth group scheme  $G$  over  $R = \mathbf{Z}/4$ , define the *Bockstein*

$$\beta: H^i(BG_k, \Omega^j) \rightarrow H^{i+1}(BG_k, \Omega^j)$$

on the Hodge cohomology of  $BG_k$  (where  $k = \mathbf{Z}/2$ ) to be the boundary homomorphism associated to the short exact sequence of sheaves

$$0 \rightarrow \Omega_k^j \rightarrow \Omega_R^j \rightarrow \Omega_k^j \rightarrow 0$$

on  $BG_R$ . (The Bockstein on Hodge cohomology is also defined for group schemes  $G$  such as  $\mu_2$  which are flat but not smooth over  $R = \mathbf{Z}/4$ , because the Hodge cohomology of  $BG$  can be described using smooth schemes (Lemma 1.2).)

Next, define elements  $u_1, u_3, \dots, u_{2r-1}$  of  $H_{\mathbb{H}}^*(BO(2r)/k)$  as follows. First, let  $u_1 \in H^1(BO(2r), \Omega^0)$  be the pullback of the generator of  $H^1(\mathbf{Z}/2, k) = k$  via the surjection  $O(2r) \rightarrow \mathbf{Z}/2$  (Lemma 10.2). Next, use that the split group  $O(2r)$  over  $k = \mathbf{F}_2$  lifts to a smooth group  $O(2r)$  over  $\mathbf{Z}$ . As a result, we have a Bockstein homomorphism on the Hodge cohomology of  $BO(2r)$ . For  $0 \leq a \leq r-1$ , let  $u_{2a+1} = \beta u_{2a} + u_1 u_{2a} \in H^{a+1}(BO(2r), \Omega^a)$ . This agrees with the previous formula for  $u_1$ , if we make the convention that  $u_0 = 1$ . (The definition of  $u_{2a+1}$  is suggested by the formula for odd Stiefel-Whitney classes in topology:  $w_{2a+1} = \beta w_{2a} + w_1 w_{2a}$  [25, Theorem III.5.12].)

I claim that the homomorphism

$$k[u_1, u_2] \rightarrow H_{\mathbb{H}}^*(BO(2)/k)$$

is an isomorphism. To see this, consider the Hochschild-Serre spectral sequence of Lemma 10.3,

$$E_2^{ij} = H^i(\mathbf{Z}/2, H^j(BSO(2)_k, \Omega^*)) \Rightarrow H^{i+j}(BO(2)_k, \Omega^*).$$

Here  $SO(2)$  is isomorphic to  $G_m$ , and so we know the Hodge cohomology of  $BSO(2)$  by Theorem 4.1:  $H_{\mathbb{H}}^*(BSO(2)/k) \cong k[c_1]$  with  $c_1$  in  $H^1(BSO(2), \Omega^1)$ . We read off that the  $E_2$  page of the spectral sequence is the polynomial ring  $k[u_1, u_2]$ , with  $u_1$  in  $H^1(\mathbf{Z}/2, H^0(BSO(2), \Omega^0))$  and  $u_2$  in  $H^0(\mathbf{Z}/2, H^1(BSO(2), \Omega^1))$ . Here  $u_1$  is a permanent cycle, because all differentials send  $u_1$  to zero groups. Also, because the surjection  $O(2) \rightarrow \mathbf{Z}/2$  of  $k$ -groups is split, there are no differentials into the bottom row of the spectral sequence; so  $u_2$  is also a permanent cycle. It follows that the spectral sequence degenerates at  $E_2$ , and hence that  $H_{\mathbb{H}}^*(BO(2)/k) \cong k[u_1, u_2]$ .

We also need to compute the Bockstein on the Hodge cohomology of  $BO(2)$ , which is defined because  $O(2)$  lifts to a smooth group scheme over  $R := \mathbf{Z}/4$ . The Bockstein is related to the Hodge cohomology of  $BO(2)_R$  by the exact sequence

$$H^i(BO(2)_R, \Omega^j) \rightarrow H^i(BO(2)_k, \Omega^j) \xrightarrow{\beta} H^{i+1}(BO(2)_k, \Omega^j).$$

Consider the Hochschild-Serre spectral sequence of Lemma 10.3 for  $BO(2)_R$ :

$$E_2^{ij} = H^i(\mathbf{Z}/2, H^j(BSO(2)_R, \Omega^*)) \Rightarrow H^{i+j}(BO(2)_R, \Omega^*).$$

Here  $H^1(BO(2)_R, \Omega^1)$  is isomorphic to  $H^0(\mathbf{Z}/2, H^1(BSO(2)_R, \Omega^1))$ , where  $\mathbf{Z}/2$  acts by  $-1$  on  $H^1(BSO(2)_R, \Omega^1) \cong \mathbf{Z}/4$ . So the generator of  $H^1(BO(2)_R, \Omega^1) \cong \mathbf{Z}/2$  maps to zero in  $H^1(BO(2)_k, \Omega^1) = k \cdot u_2$ . Therefore,  $\beta(u_2) \neq 0$ . Since  $k = \mathbf{F}_2$ , the element  $\beta(u_2)$  in  $H^2(BO(2)_k, \Omega^1) = k \cdot u_1 u_2$  must be equal to  $u_1 u_2$ . A similar analysis shows that  $\beta(u_1) = u_1^2$ .

Finally, think of  $O(2)$  as the isometry group of the quadratic form  $q(x, y) = xy$  on  $V = A_k^2$ . There is an inclusion  $H = \mathbf{Z}/2 \times \mu_2 \subset O(2)$ , where  $\mathbf{Z}/2$  switches  $x$  and  $y$  and  $\mu_2$  acts by scalars on  $V$ . For later use, it is convenient to say something about the restriction from  $BO(2)$  to  $BH$  on Hodge cohomology. By Lemma 10.2, the Hodge cohomology of  $B(\mathbf{Z}/2)$  over  $k$  is the cohomology of  $\mathbf{Z}/2$  as a group, namely the polynomial ring  $k[s]$  with  $s \in H^1(B(\mathbf{Z}/2), O)$ . Also, by Proposition 10.1, the Hodge cohomology of  $B\mu_2$  is  $k[t, v]/(v^2)$  with  $t \in H^1(B\mu_2, \Omega^1)$  and  $v \in H^0(B\mu_2, \Omega^1)$ . Thus we have a homomorphism from  $H_{\mathbb{H}}^*(BO(2)/k) = k[u_1, u_2]$  to  $H_{\mathbb{H}}^*(BH/k) \cong k[s, t, v]/(v^2)$  (by the Künneth theorem, Proposition 5.1). Here  $u_1$  restricts to  $s$ , since both elements are pulled back from the generator of  $H^1(B\mathbf{Z}/2, O)$ . Also,  $u_2$  restricts to either  $t$  or  $t + sv$ , because  $u_2$  restricts to the generator  $c_1$  of  $H^1(BG_m, \Omega^1)$  and hence to  $t$  in  $H^1(B\mu_2, \Omega^1)$ . Thus the homomorphism from  $H_{\mathbb{H}}^*(BO(2)/k)$  to  $H_{\mathbb{H}}^*(BH/k)/\text{rad} = k[s, t]$  is an isomorphism. (Here the *radical* of a commutative ring means the ideal of nilpotent elements.) A direct cocycle computation shows that  $u_2$  restricts to  $t + sv$  in  $H^1(BH, \Omega^1)$ , but we do not need that fact in this paper.

We now return to the group  $O(2r)$  over  $k = \mathbf{F}_2$  for any  $r$ . To formulate the following lemma, let  $s_1, \dots, s_r \in H^1(BO(2)^r, \Omega^0)$  be the pullbacks of  $u_1$  from the  $r$   $BO(2)$  factors, and let  $t_1, \dots, t_r$  be the pullbacks of  $u_2$  from those  $r$  factors. By the Künneth theorem (Proposition 5.1), the Hodge cohomology of  $BO(2)^r$  is the polynomial ring  $k[s_1, \dots, s_r, t_1, \dots, t_r]$ .

**Lemma 11.3.** *The homomorphism*

$$\psi: k[u_1, u_2, \dots, u_{2r}] \rightarrow H_{\mathbb{H}}^*(BO(2r)/k)$$

*is injective. Also, the composition of  $\psi$  with the restriction  $\rho$  to the Hodge cohomology of  $BO(2)^r$  is given by*

$$u_{2a} \mapsto e_a(t_1, \dots, t_r) = \sum_{1 \leq i_1 < \dots < i_a \leq r} t_{i_1} \cdots t_{i_a}$$

*and*

$$u_{2a+1} \mapsto \sum_{m=1}^r s_m \sum_{\substack{1 \leq i_1 < \dots < i_a \leq r \\ \text{none equal to } m}} t_{i_1} \cdots t_{i_a}.$$

*Finally, the same formulas hold in de Rham cohomology as well as Hodge cohomology.*

*Proof.* The formula for the restriction  $\rho\psi(u_{2a})$  on Hodge cohomology follows from the definition of  $u_{2a}$ . Likewise, it is immediate that

$$u_1 \mapsto s_1 + \dots + s_r.$$

The inclusion  $O(2)^2 \subset O(2r)$  lifts to an inclusion of smooth groups over  $\mathbf{Z}$ , and so the restriction homomorphism commutes with the Bockstein. Therefore, for



$$0 \leq a \leq r-1,$$

$$\begin{aligned} u_{2a+1} &= \beta u_{2a} + u_1 u_{2a} \\ &\mapsto \beta \left( \sum_{1 \leq i_1 < \dots < i_a \leq r} t_{i_1} \cdots t_{i_a} \right) + (s_1 + \dots + s_r) \left( \sum_{1 \leq i_1 < \dots < i_a \leq r} t_{i_1} \cdots t_{i_a} \right) \\ &= \sum_{1 \leq i_1 < \dots < i_a \leq r} \left( \sum_{j=1}^a s_{i_j} + \sum_{m=1}^r s_m \right) t_{i_1} \cdots t_{i_a} \\ &= \sum_{m=1}^r s_m \sum_{\substack{1 \leq i_1 < \dots < i_a \leq r \\ \text{none equal to } m}} t_{i_1} \cdots t_{i_a}, \end{aligned}$$

as we want.

These formulas remain true in de Rham cohomology as well as in Hodge cohomology, using Lemma 8.2: for a smooth affine  $k$ -group  $G$  whose identity component is reductive, there are *canonical* maps  $H^a(BG, \Omega^a) \rightarrow H_{\text{dR}}^{2a}(BG/k)$  and  $H^{a+1}(BG, \Omega^a) \rightarrow H_{\text{dR}}^{2a+1}(BG/k)$ . These maps are compatible with products and with pullback under a homomorphism of smooth affine  $k$ -groups.

To show that the homomorphism  $\psi: k[u_1, \dots, u_{2r}] \rightarrow H_{\mathbb{H}}^*(BO(2r)/k)$  is injective, it suffices to show that the composition  $\rho\psi: k[u_1, \dots, u_{2r}] \rightarrow k[s_1, \dots, s_r, t_1, \dots, t_r]$  is injective. We can factor this homomorphism through  $k[u_1, u_3, \dots, u_{2r-1}, t_1, \dots, t_r]$ , by the homomorphism  $\mu$  sending  $u_2, u_4, \dots, u_{2r}$  to the elementary symmetric polynomials in  $t_1, \dots, t_r$ . Since  $\mu$  is injective, it remains to show that

$$\sigma: k[u_1, u_3, \dots, u_{2r-1}, t_1, \dots, t_r] \rightarrow k[s_1, \dots, s_r, t_1, \dots, t_r]$$

is injective.

More strongly, we will show that  $\sigma$  is generically étale; that is, its Jacobian determinant is not identically zero. Because  $\sigma$  is the identity on the  $t_i$  coordinates, it suffices to show that the determinant of the matrix of derivatives of  $u_1, u_3, \dots, u_{2r-1}$  with respect to  $s_1, \dots, s_r$  is nonzero for  $s_1, \dots, s_r, t_1, \dots, t_r$  generic. This matrix of derivatives in fact only involves  $t_1, \dots, t_r$ , because  $u_1, u_3, \dots, u_{2r-1}$  have degree 1 in  $s_1, \dots, s_r$ . For example, for  $r = 3$ , this matrix of derivatives is

$$\begin{pmatrix} 1 & t_2 + t_3 & t_2 t_3 \\ 1 & t_1 + t_3 & t_1 t_3 \\ 1 & t_1 + t_2 & t_1 t_2 \end{pmatrix},$$

where the  $a$ th column gives the derivatives of  $u_{2a-1}$  with respect to  $s_1, \dots, s_r$ . For any  $r$ , column 1 consists of 1s, while entry  $(j, a)$  for  $a \geq 2$  is

$$\sum_{\substack{1 \leq i_1 < \dots < i_{a-1} \leq r \\ \text{none equal to } j}} t_{i_1} \cdots t_{i_{a-1}}.$$

This determinant is equal to the Vandermonde determinant  $\delta := \prod_{i < j} (t_i - t_j)$ , and in particular it is not identically zero [14, Theorem 1]. (The reference works over  $\mathbf{C}$ , but it amounts to an identity of polynomials over  $\mathbf{Z}$ , which therefore holds over any field.)

Thus we have shown that the composition  $\psi: k[u_1, \dots, u_{2r}] \rightarrow H_{\mathbb{H}}^*(BO(2r)/k)$  is injective, because the composition  $\rho\psi$  to  $H_{\mathbb{H}}^*(BO(2)^r/k)$  is injective.  $\square$

To avoid an excess of notation, let us also write  $\psi$  for the homomorphism  $k[u_2, u_3, \dots, u_n] \rightarrow H_{\mathbb{H}}^*(BSO(n)/k)$ .

**Lemma 11.4.** *The homomorphism*

$$\psi: k[u_2, u_3, \dots, u_n] \rightarrow H_{\mathbb{H}}^*(BSO(n)/k)$$

is injective. Also, in the case  $n = 2r + 1$ , the composition of  $\psi$  with the restriction  $\rho$  to the Hodge cohomology of  $BO(2)^r$  is given by

$$u_{2a} \mapsto e_a(t_1, \dots, t_r) = \sum_{1 \leq i_1 < \dots < i_a \leq r} t_{i_1} \cdots t_{i_a},$$

$$u_{2a+1} \mapsto \sum_{m=1}^r s_m \sum_{\substack{1 \leq i_1 < \dots < i_a \leq r \\ \text{one equal to } m}} t_{i_1} \cdots t_{i_a}$$

for  $1 \leq a \leq r$ . Finally, the same formulas hold in de Rham cohomology as well as Hodge cohomology.

*Proof.* For  $n = 2r + 1$ , this is an easy consequence of Lemma 11.3, using the inclusions  $O(2)^r \subset O(2r) \subset SO(2r + 1)$ . Write  $u_2, u_3, \dots, u_{2r+1}$  for the elements of the Hodge cohomology of  $BSO(2r + 1)$  defined by the same formulas as used above for  $BO(2r)$  (which simplify to  $u_{2a+1} = \beta u_{2a}$ , since there is no element  $u_1$  for  $BSO(2r + 1)$ ). Also, let  $v_1, \dots, v_{2r}$  be the elements of the Hodge cohomology of  $BO(2r)$  that were called  $u_1, \dots, u_{2r}$  above. Then restricting from  $BSO(2r + 1)$  to  $BO(2r)$  sends  $u_{2a} \mapsto v_{2a}$  and  $u_{2a+1} = \beta u_{2a} \mapsto \beta v_{2a} = v_{2a+1} + v_1 v_{2a}$  for  $1 \leq a \leq r - 1$ . It is not immediate how to compute the restriction of the remaining element  $u_{2r+1}$  to  $BO(2r)$ , but we can compute its restriction to  $BO(2)^r$ :

$$\begin{aligned} u_{2r+1} &= \beta u_{2r} \\ &\mapsto \beta v_{2r} \\ &= \beta(t_1 \cdots t_r) \\ &= (s_1 + \cdots + s_r)(t_1 \cdots t_r). \end{aligned}$$

Thus we have proved the desired formulas for the restriction on Hodge cohomology from  $BSO(2r + 1)$  to  $BO(2)^r$ . Since the generators are in  $H^i(BSO(2r + 1), \Omega^i)$  or  $H^{i+1}(BSO(2r + 1), \Omega^i)$ , the same formulas hold in de Rham cohomology.

Thus, the restriction from  $BSO(2r + 1)$  to  $BO(2)^r$  sends  $k[u_2, \dots, u_{2r+1}]$  into the subring

$$k[v_1, \dots, v_{2r}] \subset k[s_1, \dots, s_r, t_1, \dots, t_r],$$

by  $u_{2a} \mapsto v_{2a}$  for  $1 \leq a \leq r$ ,  $u_{2a+1} \mapsto v_{2a+1} + v_1 v_{2a}$  for  $1 \leq a \leq r - 1$ , and  $u_{2r+1} \mapsto v_1 v_{2r}$ . This homomorphism is injective, because the corresponding morphism  $A^{2r} \rightarrow A^{2r}$  is birational (for  $u_{2r} \neq 0$ , one can solve for  $v_1, \dots, v_{2r}$  in terms of  $u_2, \dots, u_{2r+1}$ ). So the homomorphism  $\psi: k[u_2, \dots, u_{2r+1}] \rightarrow H_{\mathbb{H}}^*(BSO(2r + 1)/k)$  is injective (because its composition to  $H_{\mathbb{H}}^*(BO(2)^r/k)$  is injective).

For  $SO(2r)$ , we argue a bit differently. As discussed above, there is a subgroup  $\mathbf{Z}/2 \times \mu_2 \subset O(2)$ . Therefore, we have a  $k$ -subgroup scheme  $(\mathbf{Z}/2 \times \mu_2)^r \subset O(2)^r \subset O(2r)$ . Since  $SO(2r)$  is the kernel of a homomorphism from  $O(2r)$  onto  $\mathbf{Z}/2$ ,  $SO(2r)$

contains a  $k$ -subgroup scheme  $H \cong (\mathbf{Z}/2)^{r-1} \times (\mu_2)^r$ . By Lemma 10.2, the Hodge cohomology of  $B(\mathbf{Z}/2)$  over  $k$  is the cohomology of  $\mathbf{Z}/2$  as a group, namely the polynomial ring  $k[x]$  with  $x \in H^1(B(\mathbf{Z}/2), \Omega^0)$ . Also, by Proposition 10.1, the Hodge cohomology of  $B\mu_2$  is  $k[t, v]/(v^2)$  with  $t \in H^1(B\mu_2, \Omega^1)$  and  $v \in H^0(B\mu_2, \Omega^1)$ . Thus we have a homomorphism  $\psi$  from  $k[u_2, u_3, \dots, u_{2r}]$  to  $H_{\mathbf{H}}^*(BSO(2r)/k)$ , and a homomorphism from there to  $H_{\mathbf{H}}^*(BH/k) \cong k[x_1, \dots, x_{r-1}, t_1, \dots, t_r, v_1, \dots, v_r]/(v_i^2)$  (by the Künneth theorem, Proposition 5.1). We want to show that this composition is injective. For convenience, we will prove the stronger statement that the composition  $\rho\psi$  from  $k[u_2, u_3, \dots, u_{2r}]$  to

$$H_{\mathbf{H}}^*(BH/k)/\text{rad} = k[x_1, \dots, x_{r-1}, t_1, \dots, t_r]$$

is injective.

We compare the restriction from  $O(2r)$  to  $(\mathbf{Z}/2)^r \times (\mu_2)^r$  with that from  $SO(2r)$  to  $H$ :

$$\begin{array}{ccc} k[u_1, \dots, u_{2r}] & \longrightarrow & k[u_2, u_3, \dots, u_{2r}] \\ \downarrow & & \downarrow \\ H_{\mathbf{H}}^*(BO(2r)/k) & \longrightarrow & H_{\mathbf{H}}^*(BSO(2r)/k) \\ \downarrow & & \downarrow \\ k[s_1, \dots, s_r, t_1, \dots, t_r] & \longrightarrow & k[x_1, \dots, x_{r-1}, t_1, \dots, t_r] \end{array}$$

The bottom homomorphism is given (for a suitable choice of generators  $x_1, \dots, x_{r-1}$ ) by  $s_i \mapsto x_i$  for  $1 \leq i \leq r-1$  and  $s_r \mapsto x_1 + \dots + x_{r-1}$  (agreeing with the fact that  $u_1 \mapsto s_1 + \dots + s_r \mapsto 0$  in the Hodge cohomology of  $BH$ ). By the formulas for  $O(2r)$ , we know how the elements  $u_2, \dots, u_{2r}$  restrict to  $k[s_1, \dots, s_r, t_1, \dots, t_r]$ , and hence to  $k[x_1, \dots, x_{r-1}, t_1, \dots, t_r]$ . Namely,

$$u_{2a} \mapsto e_a(t_1, \dots, t_r) = \sum_{1 \leq i_1 < \dots < i_a \leq r} t_{i_1} \cdots t_{i_a},$$

and, for  $1 \leq a \leq r-1$ ,

$$\begin{aligned} u_{2a+1} &\mapsto \sum_{1 \leq i_1 < \dots < i_a \leq r} \left( \sum_{j=1}^a s_{i_j} + \sum_{m=1}^r s_m \right) t_{i_1} \cdots t_{i_a} \\ &\mapsto \sum_{1 \leq i_1 < \dots < i_a \leq r-1} \left( \sum_{j=1}^a x_{i_j} \right) t_{i_1} \cdots t_{i_a} \\ &\quad + \sum_{1 \leq i_1 < \dots < i_{a-1} \leq r-1} \left( x_1 + \dots + x_{r-1} + \sum_{j=1}^{a-1} x_{i_j} \right) t_{i_1} \cdots t_{i_{a-1}} t_r \\ &= \sum_{j=1}^{r-1} x_j (t_j + t_r) \sum_{\substack{1 \leq i_1 < \dots < i_{a-1} \leq r-1 \\ \text{none equal to } j}} t_{i_1} \cdots t_{i_{a-1}}. \end{aligned}$$

We want to show that this homomorphism  $\rho\psi: k[u_2, u_3, \dots, u_{2r}] \rightarrow k[x_1, \dots, x_{r-1}, t_1, \dots, t_r]$  is injective. It can be factored through  $k[u_3, u_5, \dots, u_{2r-1}, t_1, \dots, t_r]$ , by the homomorphism  $\mu$  sending  $u_2, u_4, \dots, u_{2r}$  to the elementary symmetric polynomials in  $t_1, \dots, t_r$ . Since  $\mu$  is injective, it remains to show that  $\sigma: k[u_3, u_5, \dots, u_{2r-1}, t_1, \dots, t_r] \rightarrow k[x_1, \dots, x_{r-1}, t_1, \dots, t_r]$  is injective.

As in the argument for  $O(2r)$ , we will show (more strongly) that  $\sigma$  is generically étale; that is, its Jacobian determinant is not identically zero. Because  $\sigma$  is the identity on the  $t_i$  coordinates, it suffices to show that the determinant of the matrix of derivatives of  $u_3, u_5, \dots, u_{2r-1}$  with respect to  $x_1, \dots, x_{r-1}$  is nonzero for  $x_1, \dots, x_{r-1}, t_1, \dots, t_r$  generic. This matrix of derivatives in fact only involves  $t_1, \dots, t_r$ , because  $u_3, u_5, \dots, u_{2r-1}$  have degree 1 as polynomials in  $x_1, \dots, x_{r-1}$ . For example, for  $r = 3$ , this  $(r-1) \times (r-1)$  matrix of derivatives is

$$\begin{pmatrix} t_1 + t_3 & (t_1 + t_3)(t_2) \\ t_2 + t_3 & (t_2 + t_3)(t_1) \end{pmatrix},$$

where the  $a$ th column gives the derivatives of  $u_{2a+1}$  with respect to  $x_1, \dots, x_{r-1}$ . For any  $r$ , the entry  $(j, a)$  of the matrix (with  $j, a \in \{1, \dots, r-1\}$ ) is  $(t_j + t_r)e_{ja}$ , where

$$e_{ja} = \sum_{\substack{1 \leq i_1 < \dots < i_{a-1} \leq r-1 \\ \text{none equal to } j}} t_{i_1} \cdots t_{i_{a-1}}.$$

Since row  $j$  is a multiple of  $(t_j + t_r)$  for each  $r$ , the determinant is  $(t_1 + t_r)(t_2 + t_r) \cdots (t_{r-1} + t_r)$  times the determinant of the  $(r-1) \times (r-1)$  matrix  $E = (e_{ja})$ . So it suffices to show that the determinant of  $E$  is not identically zero. Indeed, the determinant of  $E$  is the same determinant shown to be nonzero in the calculation above for  $O(2r)$ , but with  $r$  replaced by  $r-1$ .

Thus we have shown that  $\psi: k[u_2, \dots, u_n] \rightarrow H_{\mathbb{H}}^*(BSO(n)/k)$  is injective for  $n$  even as well as for  $n$  odd.  $\square$

Having shown that  $\psi: k[u_2, \dots, u_n] \rightarrow H_{\mathbb{H}}^*(BSO(n)/k)$  is injective, we now show that it is an isomorphism.

Let  $r = \lfloor n/2 \rfloor$  and  $s = \lfloor (n-1)/2 \rfloor$ . Let  $P$  be the parabolic subgroup of  $G = SO(n)$  that stabilizes a maximal isotropic subspace (that is, an isotropic subspace of dimension  $r$ ). Then the quotient of  $P$  by its unipotent radical is isomorphic to  $GL(r)$ . By Proposition 9.3, we have a spectral sequence

$$E_2^{ij} = H_{\mathbb{H}}^i(BG/k) \otimes H_{\mathbb{H}}^j((G/P)/k) \Rightarrow H_{\mathbb{H}}^{i+j}(BGL(r)/k).$$

The Chow ring of  $G/P$  is isomorphic to

$$\mathbf{Z}[e_1, \dots, e_s]/(e_i^2 - 2e_{i-1}e_{i+1} + 2e_{i-2}e_{i+2} - \cdots + (-1)^i e_{2i}),$$

where  $e_i \in CH^i(G/P)$  is understood to mean zero if  $i > s$  [25, III.6.11]. (This uses the theorem, discussed in section 7, that the Chow ring of  $G/P$  for a split group  $G$  is independent of the characteristic of  $k$ , and is isomorphic to the integral cohomology ring of  $G_{\mathbb{C}}/P_{\mathbb{C}}$ . The reference assumes that  $n$  is even, but that is enough, because the obvious map  $SO(2r+1)/P \rightarrow SO(2r+2)/P$  is an isomorphism.) By Proposition 7.1, it follows that the Hodge cohomology ring of  $G/P$  is isomorphic to

$$k[e_1, \dots, e_s]/(e_i^2 = e_{2i}),$$

where  $e_i$  is in  $H^i(G/P, \Omega^i)$ . For any list of variables  $x_1, \dots, x_m$ , write  $\Delta(x_1, \dots, x_m)$  for the  $k$ -vector space with basis consisting of all products  $x_{i_1} \cdots x_{i_j}$  with  $1 \leq i_1 < \cdots < i_j \leq m$  and  $0 \leq j \leq m$ . Then we can say that

$$H_{\mathbb{H}}^*((G/P)/k) = \Delta(e_1, \dots, e_s).$$

The spectral sequence converges to  $H_{\mathbb{H}}^*(BGL(r)/k) = k[c_1, \dots, c_r]$ , by Theorem 9.2. Write  $\Phi: H_{\mathbb{H}}^*(BG/k) \rightarrow H_{\mathbb{H}}^*(BGL(r)/k)$  for the restriction homomorphism, which is the “edge map” associated to the 0th row in the spectral sequence. The restriction  $\Phi$  takes the elements  $u_2, u_4, \dots, u_{2r}$  (where  $u_{2i}$  is in  $H^i(BG, \Omega^i)$ ) to  $c_1, c_2, \dots, c_r$ . So the  $E_{\infty}$  term of the spectral sequence is concentrated on the 0th row and consists of the polynomial ring  $k[u_2, u_4, \dots, u_{2r}]$ .

To analyze the structure of the spectral sequence further, we use Zeeman’s comparison theorem, which he used to simplify the proof of the Borel transgression theorem [25, Theorem VII.2.9]. The key point is to show that the elements  $e_i$  (possibly after adding decomposable elements) are transgressive. (By definition, an element  $u$  of  $E_2^{0,q}$  in a first-quadrant spectral sequence is *transgressive* if  $d_2 = \dots = d_q = 0$  on  $u$ ; then  $u$  determines an element  $\tau(u) := d_{q+1}(u)$  of  $E_{q+1}^{q+1,0}$ , called the transgression of  $u$ .)

In order to apply Zeeman’s comparison theorem, we define a model spectral sequence that maps to the spectral sequence we want to analyze. (To be precise, we consider spectral sequences of  $k$ -vector spaces, not of  $k$ -algebras.) As above, let  $k = \mathbf{F}_2$ . For a positive integer  $q$ , define a spectral sequence  $G_*$  with  $E_2$  page given by  $G_2 = \Delta(y) \otimes k[u]$ ,  $y$  in bidegree  $(0, q)$ ,  $u$  in bidegree  $(q+1, 0)$ , and  $d_{q+1}(yu^j) = u^{j+1}$ .

$$\begin{array}{ccccccc} k \cdot y & & k \cdot yu & & k \cdot yu^2 & & \dots \\ & \searrow \cong & & \searrow \cong & & & \\ k \cdot 1 & & k \cdot u & & k \cdot u^2 & & \dots \end{array}$$

Suppose that, for some positive integer  $a$ , we have found elements  $y_i$  of  $H_{\mathbb{H}}^{2i}((G/P)/k)$  for  $1 \leq i \leq a$  which are transgressive in the spectral sequence  $E_*$  above. Because  $y_i$  is transgressive, there is a map of spectral sequences  $G_* \rightarrow E_*$  that takes the element  $y$  (in degree  $q = 2i$ ) to  $y_i$ . Since  $E_*$  is a spectral sequence of algebras, tensoring these maps gives a map of spectral sequences

$$\alpha: F_* := G_*(y_1) \otimes \dots \otimes G_*(y_a) \otimes k[u_2, u_4, \dots, u_{2r}] \rightarrow E_*.$$

(Here we are using that the elements  $u_2, u_4, \dots, u_{2r}$  are in  $H_{\mathbb{H}}^*(BG/k)$ , which is row 0 of the  $E_2$  page on the right, and so they are permanent cycles.) Although we do not view the domain as a spectral sequence of algebras, its  $E_2$  page is the tensor product of row 0 and column 0, and the map  $\alpha: F_2 \rightarrow E_2$  of  $E_2$  pages is the tensor product of the maps on row 0 and column 0.

Using these properties, we have the following version of Zeeman’s comparison theorem, as sharpened by Hilton and Roitberg [25, Theorem VII.2.4]:

**Theorem 11.5.** *Let  $N$  be a natural number. Suppose that the homomorphism  $\alpha: F_* \rightarrow E_*$  of spectral sequences is bijective on  $E_{\infty}^{i,j}$  for  $i+j \leq N$  and injective for  $i+j = N+1$ , and that  $\alpha$  is bijective on row 0 of the  $E_2$  page in degrees  $\leq N+1$  and injective in degree  $N+2$ . Then  $\alpha$  is bijective on column 0 of the  $E_2$  page in degree  $\leq N$  and injective in degree  $N+1$ .*

The inductive step for computing the Hodge cohomology of  $BSO(n)$  is as follows.

**Lemma 11.6.** *Let  $G$  be  $SO(n)$  over  $k = \mathbf{F}_2$ ,  $P$  the parabolic subgroup above,  $r = \lfloor n/2 \rfloor$ ,  $s = \lfloor (n-1)/2 \rfloor$ . Let  $N$  be a natural number, and let  $a = \min(s, \lfloor N/2 \rfloor)$ . Then, for each  $1 \leq i \leq a$ , there is an element  $y_i$  in  $H^i(G/P, \Omega^i)$  with the following*

properties. First,  $y_i$  is equal to  $e_i$  modulo polynomials in  $e_1, \dots, e_{i-1}$  with exponents  $\leq 1$ . Also, each element  $y_i$  is transgressive, and any lift  $v_{2i+1}$  to  $H^{i+1}(BG, \Omega^i)$  of the element  $\tau(y_i)$  has the property that

$$k[u_2, u_4, \dots, u_{2r}; v_3, v_5, \dots, v_{2a+1}] \rightarrow H_{\mathbb{H}}^*(BG/k)$$

is bijective in degree  $\leq N + 1$  and injective in degree  $N + 2$ . Finally, each element  $v_{2i+1}$  is equal to  $u_{2i+1}$  modulo polynomials in  $u_2, u_3, \dots, u_{2i}$ .

More precisely, if this statement holds for  $N - 1$ , then it holds for  $N$  with the same elements  $y_i$ , possibly with one added.

We will apply Lemma 11.6 with  $N = \infty$ , but the formulation with  $N$  arbitrary is convenient for the proof.

*Proof.* As discussed earlier, the  $E_\infty$  page of the spectral sequence

$$E_2^{ij} = H_{\mathbb{H}}^i(BG/k) \otimes H_{\mathbb{H}}^j((G/P)/k) \Rightarrow H_{\mathbb{H}}^{i+j}(BGL(r)/k)$$

is isomorphic to  $k[u_2, u_4, \dots, u_{2r}]$ , concentrated on row 0.

We prove the lemma by induction on  $N$ . For  $N = 0$ , it is true, using that  $H_{\mathbb{H}}^0(BG/k) = k$  and  $H_{\mathbb{H}}^1(BG/k) = 0$ , as one checks using our knowledge of the  $E_\infty$  term.

We now assume the result for  $N - 1$ , and prove it for  $N$ . By the inductive assumption, for  $b := \min(s, \lfloor (N - 1)/2 \rfloor)$ , we can choose  $y_1, \dots, y_b$  such that  $y_i \in H^i(G/P, \Omega^i)$  is equal to  $e_i$  modulo polynomials in  $e_1, \dots, e_{i-1}$  with exponents  $\leq 1$ ,  $y_i$  is transgressive for the spectral sequence, and, if we define  $v_{2i+1} \in H^{i+1}(BG, \Omega^i)$  to be any lift (from the  $E_{2i+1}$  page to the  $E_2$  page) of the transgression  $\tau(y_i)$  for  $1 \leq i \leq b$ , the homomorphism

$$k[u_2, u_4, \dots, u_{2r}; v_3, v_5, \dots, v_{2b+1}] \rightarrow H_{\mathbb{H}}^*(BG/k)$$

is bijective in degree  $\leq N$  and injective in degree  $N + 1$ . Finally, the element  $v_{2i+1}$  for  $1 \leq i \leq b$  is equal to  $u_{2i+1}$  modulo polynomials in  $u_2, u_3, \dots, u_{2i}$ .

Also, by the injectivity in degree  $N + 1$  (above), it follows that there is a  $k$ -linearly independent set (possibly empty) of elements  $z_i$  in  $H_{\mathbb{H}}^{N+1}(BG/k)$  such that

$$\varphi: k[u_2, u_4, \dots, u_{2r}; v_3, v_5, \dots, v_{2b+1}; z_i] \rightarrow H_{\mathbb{H}}^*(BG/k)$$

is bijective in degrees at most  $N + 1$ . (Recall that  $b = \min(s, \lfloor (N - 1)/2 \rfloor)$ .) The elements  $z_i$  do not affect the domain of  $\varphi$  in degree  $N + 2$  (because that ring is zero in degree 1). Therefore,  $\varphi$  is injective in degree  $N + 2$ , because

$$k[u_2, u_4, \dots, u_{2r}; v_3, v_5, \dots, v_{2b+1}] \rightarrow H_{\mathbb{H}}^*(BG/k)$$

is injective. (This uses that  $v_{2i+1}$  is equal to  $u_{2i+1}$  modulo polynomials in  $u_2, u_3, \dots, u_{2i}$ , together with the injectivity of  $k[u_2, u_3, \dots, u_n] \rightarrow H_{\mathbb{H}}^*(BG/k)$ , shown in Lemma 11.4.)

The elements  $z_i$  can be chosen to become zero in the  $E_\infty$  page, because the  $E_\infty$  page is just  $k[u_2, u_4, \dots, u_{2r}]$  on row 0. Therefore, there are transgressive elements  $w_i \in H_{\mathbb{H}}^N((G/P)/k)$  with  $z_i = \tau(w_i)$  in the  $E_{N+1}$  page. (If  $z_i$  is killed before  $E_{N+1}$ , we can simply take  $w_i = 0$ .)

Consider the map of spectral sequences

$$\alpha: F_* := \Delta(y_1, \dots, y_b; w_i) \otimes k[u_2, u_4, \dots, u_{2r}, v_3, v_3, \dots, v_{2b+1}; z_i] \rightarrow E_*.$$

The map on  $E_\infty$  terms is an isomorphism (to  $k[u_2, u_4, \dots, u_{2r}]$ ), and we showed two paragraphs back that the map on column 0 of the  $E_2$  terms is bijective in degrees at most  $N+1$  and injective in degree  $N+2$ . Therefore, Zeeman's comparison theorem (Theorem 11.5) gives that the homomorphism

$$\psi: \Delta(y_1, \dots, y_b; w_i) \rightarrow H_{\mathbb{H}}^*((G/P)/k)$$

is bijective in degrees  $\leq N$  and injective in degree  $N+1$ .

Let  $a = \min(s, \lfloor N/2 \rfloor)$ . We know that  $\Delta(e_1, \dots, e_a) \rightarrow H_{\mathbb{H}}^*((G/P)/k)$  is bijective in degrees  $\leq N$ . Since the elements  $w_i$  are in degree  $N$ , while  $b = \min(s, \lfloor (N-1)/2 \rfloor)$ , we deduce that there is no element  $w_i$  if  $N$  is odd or  $N > 2s$ , and there is exactly one  $w_i$  if  $N$  is even and  $N \leq 2s$ . In the latter case, we have  $a = N/2$ ; in that case, let  $y_a$  denote the single element  $w_i$ . Since we know that  $H_{\mathbb{H}}^*((G/P)/k) = \Delta(e_1, \dots, e_s)$ ,  $y_a$  must be equal to  $e_a$  modulo polynomials in  $e_1, \dots, e_{a-1}$  with exponents  $\leq 1$ . By construction,  $y_a$  is transgressive. Also, in the case where  $N$  is even and  $N \leq 2s$ , let  $v_{2a+1}$  in  $H^{a+1}(BG, \Omega^a)$  be a lift to the  $E_2$  page of the element  $\tau(y_a)$  (formerly called  $z_i$ ). Then we know that

$$\varphi: k[u_2, u_4, \dots, u_{2r}; v_3, v_5, \dots, v_{2a+1}] \rightarrow H_{\mathbb{H}}^*(BG/k)$$

is bijective in degree  $\leq N+1$ . In the case where  $N$  is even and  $N \leq 2s$  (where we have added one element  $v_{2a+1}$  to those constructed before), this bijectivity in degree  $N+1 = 2a+1$  together with the injectivity of  $k[u_2, u_3, \dots, u_n] \rightarrow H_{\mathbb{H}}^*(BG/k)$  in all degrees implies that  $v_{2a+1}$  must be equal to  $u_{2a+1}$  modulo polynomials in  $u_2, u_3, \dots, u_{2a}$ . By the same injectivity, it follows that  $\varphi$  is injective in degree  $N+2$ .  $\square$

We can take  $N = \infty$  in Lemma 11.6, because the elements  $y_1, \dots, y_s$  do not change as we increase  $N$ . This gives that  $k[u_2, u_3, \dots, u_n] \rightarrow H_{\mathbb{H}}^*(BSO(n)/k)$  is an isomorphism. (The element  $v_{2i+1}$  produced by Lemma 11.6 need not be the element  $u_{2i+1}$  defined earlier, but  $v_{2i+1}$  is equal to  $u_{2i+1}$  modulo decomposable elements, which gives this conclusion.)

Using the Hodge cohomology of  $BSO(2r)$ , we can compute the Hodge cohomology of  $BO(2r)$  over  $k$  using the Hochschild-Serre spectral sequence of Lemma 10.3:

$$E_2^{ij} = H^i(\mathbf{Z}/2, H^j(BSO(2r), \Omega^*)) \Rightarrow H^{i+j}(BO(2r), \Omega^*).$$

We have a homomorphism  $k[u_1, u_2, \dots, u_{2r}] \rightarrow BO(2r)$  whose composition to  $BSO(2r)$  is surjective. Therefore,  $\mathbf{Z}/2$  acts trivially on the Hodge cohomology of  $BSO(2r)$ , and all differentials are zero on column 0 of this spectral sequence. It follows that the spectral sequence degenerates at  $E_2$ , and hence

$$\begin{aligned} H_{\mathbb{H}}^*(BO(2r)/k) &\cong H^*(\mathbf{Z}/2, k) \otimes H_{\mathbb{H}}^*(BSO(2r)/k) \\ &\cong k[u_1, u_2, \dots, u_{2r}]. \end{aligned}$$

Finally, we show that the Hodge spectral sequence

$$E_1^{ij} = H^j(BG, \Omega^i) \Rightarrow H_{\text{dR}}^{i+j}(BG/k)$$

degenerates for  $G = SO(n)$  over  $k$ . Indeed, by restricting to a maximal torus  $T = (G_m)^r$  of  $G$ , the elements  $u_2, u_4, \dots, u_{2r}$  restrict to the elementary symmetric polynomials in the generators of  $H_{\text{dR}}^*(BT/k) = k[t_1, \dots, t_r]$ . Therefore, the ring  $k[u_2, u_4, \dots, u_{2r}]$  injects into  $H_{\text{dR}}^*(BG/k)$ . So all differentials into the main diagonal  $\oplus_i H^{i,i}$  of the Hodge spectral sequence for  $BG$  are zero.

$$\begin{array}{ccccc}
H^2(BG, \Omega^0) & \xrightarrow{d_1} & H^2(BG, \Omega^1) & \xrightarrow{d_1} & H^2(BG, \Omega^2) \\
& & \text{---} & \text{---} & \text{---} \\
& & & & \text{---} \\
H^1(BG, \Omega^0) & \xrightarrow{d_1} & H^1(BG, \Omega^1) & \longrightarrow & 0 \\
& & \text{---} & \text{---} & \text{---} \\
& & & & \text{---} \\
H^0(BG, \Omega^0) & \longrightarrow & 0 & \longrightarrow & 0
\end{array}$$

It follows that all differentials are zero on the elements  $u_{2i+1} \in H^{i+1}(BG, \Omega^i)$ : only  $d_1$  maps  $u_{2i+1}$  into a nonzero group, and that is on the main diagonal. Also, all differentials are zero on the elements  $u_{2i}$  in the main diagonal (since they map into zero groups). This proves the degeneration of the Hodge spectral sequence. Therefore,  $H_{\text{dR}}^*(BSO(n)/k)$  is isomorphic to  $k[u_2, u_3, \dots, u_n]$ .

The same argument proves the degeneration of the Hodge spectral sequence for  $BO(2r)$ . Therefore,  $H_{\text{dR}}^*(BO(2r)/k)$  is isomorphic to  $k[u_1, u_2, \dots, u_{2r}]$ .

Finally,  $O(2r+1)$  is isomorphic to  $SO(2r+1) \times \mu_2$ , and so the calculation for  $BO(2r+1)$  follows from those for  $BSO(2r+1)$  (above) and  $B\mu_2$  (Proposition 10.1), by the Künneth theorem (Proposition 5.1). Theorem 11.1 is proved.  $\square$

*Proof.* (Proposition 11.2) Let  $2r$  and  $2s$  be the ranks of the quadratic bundles  $E$  and  $F$ . The problem amounts to computing the restriction from  $BO(2r+2s)$  to  $BO(2r) \times BO(2s)$  on Hodge cohomology or de Rham cohomology. We first compute  $u(E \oplus F)$  in Hodge cohomology. The formula for  $u_{2a}(E \oplus F)$  follows from the definition of  $u_{2a}$  in  $H^a(BO(2r+2s), \Omega^a)$ . (Since  $u_{2a}$  is in  $H^a(BO(2r+2s), \Omega^a)$ , its restriction to the Hodge cohomology of  $BO(2r) \times BO(2s)$  must be in  $H^a(BO(2r) \times BO(2s), \Omega^a)$ , which explains why only the even  $u$ -classes of  $E$  and  $F$  appear in the formula.) The formula for  $u_{2a+1}(E \oplus F)$  follows from the formula for  $u_{2a}(E \oplus F)$ , using that  $u_{2a+1} = \beta u_{2a} + u_1 u_{2a}$ .

In de Rham cohomology, the same formulas hold for  $u(E \oplus F)$ . This uses that the subring  $\oplus_i H^i(BG, \Omega^i)$  of Hodge cohomology canonically maps into de Rham cohomology (Lemma 8.2).  $\square$

## 12 The spin groups

In contrast to the other calculations in this paper, we now exhibit a reductive group  $G$  such that the mod 2 cohomology of the topological space  $BG_{\mathbf{C}}$  is not isomorphic to the de Rham cohomology of the algebraic stack  $BG_{\mathbf{F}_2}$ , even additively. The example was suggested by the observation of Feshbach, Benson, and Wood that the restriction  $H^*(BG_{\mathbf{C}}, \mathbf{Z}) \rightarrow H^*(BT_{\mathbf{C}}, \mathbf{Z})^W$  fails to be surjective for  $G = \text{Spin}(n)$  if  $n \geq 11$  and  $n \equiv 3, 4, 5 \pmod{8}$  [4]. For simplicity, we work out the case of  $\text{Spin}(11)$ . It would be interesting to make a full computation of the de Rham cohomology of  $B\text{Spin}(n)$  in characteristic 2.



**Theorem 12.1.**

$$\dim_{\mathbf{F}_2} H_{\text{dR}}^{32}(B \text{Spin}(11)/\mathbf{F}_2) > \dim_{\mathbf{F}_2} H^{32}(B \text{Spin}(11)_{\mathbf{C}}, \mathbf{F}_2).$$

*Proof.* Let  $k = \mathbf{F}_2$ . Let  $n$  be an integer at least 6; eventually, we will restrict to the case  $n = 11$ . Let  $G$  be the split group  $\text{Spin}(n)$  over  $k$ , and let  $T$  be a maximal torus in  $G$ . Let  $r = \lfloor n/2 \rfloor$ . The Weyl group  $W$  of  $G$  is  $S_r \rtimes (\mathbf{Z}/2)^r$  for  $n = 2r + 1$ , and the subgroup  $S_r \rtimes (\mathbf{Z}/2)^{r-1}$  for  $n = 2r$ . We start by computing the ring  $O(\mathfrak{t})^W$  of  $W$ -invariant functions on the Lie algebra  $\mathfrak{t}$  of  $T$ .

First consider the easier case where  $n$  is odd,  $n = 2r + 1$ . The element  $-1$  in  $(\mathbf{Z}/2)^r \subset W$  acts as the identity on  $\mathfrak{t}$ , since we are in characteristic 2. The ring  $O(\mathfrak{t})^W$  can also be viewed as  $S(X^*(T) \otimes k)^W$ . Computing this ring is similar to, but simpler than, Benson and Wood's calculation of  $S(X^*(T))^W = H^*(BT_{\mathbf{C}}, \mathbf{Z})^W$  [4]. We follow their notation.

We have

$$S(X^*(T)) \cong \mathbf{Z}[x_1, \dots, x_r, A]/(2A = x_1 + \dots + x_r),$$

by thinking of  $T$  as the double cover of a maximal torus in  $SO(2r + 1)$ . The symmetric group  $S_r$  in  $W$  permutes  $x_1, \dots, x_r$  and fixes  $A$ . The elementary abelian group  $E_r = (\mathbf{Z}/2)^r$  in  $W$ , with generators  $\epsilon_1, \dots, \epsilon_r$ , acts by:  $\epsilon_i$  changes the sign of  $x_i$  and fixes  $x_j$  for  $j \neq i$ , and  $\epsilon_i(A) = A - x_i$ . So

$$S(X^*(T) \otimes k) \cong k[x_1, \dots, x_r, A]/(x_1 + \dots + x_r).$$

Note that  $-1 := \epsilon_1 \cdots \epsilon_r$  in  $W$  acts as the identity on  $S^*(X^*(T) \otimes k)$ .

We first compute the invariants of the subgroup  $E_r$  on  $S(X^*(T) \otimes k)$ , using the following lemma.

**Lemma 12.2.** *Let  $R$  be an  $\mathbf{F}_2$ -algebra which is a domain,  $S$  the polynomial ring  $R[x]$ , and  $a$  a nonzero element of  $R$ . Let  $G = \mathbf{Z}/2$  act on  $S$  by fixing  $R$  and sending  $x$  to  $x + a$ . Then the ring of invariants is*

$$S^G = R[u],$$

where  $u = x(x + a)$ .

*Proof.* Clearly  $u = x(x + a)$  in  $S$  is  $G$ -invariant. Since  $u$  is a monic polynomial of degree 2 in  $x$ , we have  $S = R[u] \oplus x \cdot R[u]$ . Let  $\sigma$  be the generator of  $G = \mathbf{Z}/2$ . Any element of  $S$  can be written as  $f + xg$  for some (unique) elements  $f, g \in R[u]$ . If  $f + xg$  is  $G$ -invariant, then  $0 = \sigma(f + xg) - (f + xg) = (x + a)g - xg = ag$ . Since  $a$  is a non-zero-divisor in  $R$ , it is a non-zero-divisor in  $R[u]$ ; so  $g = 0$ . Thus  $S^G = R[u]$ .  $\square$

Let  $E_j \cong (\mathbf{Z}/2)^j$  be the subgroup of  $W$  generated by  $\epsilon_1, \dots, \epsilon_j$ . Let

$$\eta_j = \prod_{I \subset \{1, \dots, j\}} \left( A - \sum_{i \in I} x_i \right),$$

which is  $E_j$ -invariant. Here  $\eta_j$  has degree  $2^j$  in  $S^*(X^*(T) \otimes k)$ . By Lemma 12.2 (with  $R = k[x_1, \dots, x_r]/(x_1 + \dots + x_r)$ ) and induction on  $j$ , we have

$$S^*(X^*(T) \otimes k)^{E_j} = k[x_1, \dots, x_r, \eta_j]/(x_1 + \dots + x_r = 0)$$

for  $1 \leq j \leq r-1$ . Since  $-1 = \epsilon_1 \cdots \epsilon_r$  acts as the identity on these rings, we also have

$$S^*(X^*(T) \otimes k)^{E_r} = k[x_1, \dots, x_r, \eta_{r-1}]/(x_1 + \cdots + x_r = 0).$$

The symmetric group  $S_r$  permutes  $x_1, \dots, x_r$ , and it fixes  $\eta_{r-1}$ . Therefore, computing the invariants of the Weyl group on  $S^*(X^*(T) \otimes k)$  reduces to computing the invariants of the symmetric group  $S_r$  on  $k[x_1, \dots, x_r]/(x_1 + \cdots + x_r)$ . That is known, by the following result [26, Proposition 4.1]. Write  $c_1, \dots, c_r$  for the elementary symmetric polynomials in  $k[x_1, \dots, x_r]$ .

**Lemma 12.3.** *Let  $k$  be a field of characteristic 2. If  $r \geq 3$ , then the ring of invariants of  $S_r$  on  $R = k[x_1, \dots, x_r]/(x_1 + \cdots + x_r)$  is equal to  $k[c_1, \dots, c_n]/(c_1) = k[c_2, \dots, c_r]$ . If  $r = 2$ , on the other hand, then  $S_2$  acts trivially on  $R = k[x_1, x_2]/(x_1 + x_2)$ , and so  $R^{S_2} = R = k[x_1]$ .*

Combining Lemma 12.3 with the calculations above, we have found the invariants for the Weyl group  $W$  of  $G = \text{Spin}(2r+1)$ : for  $r \geq 1$ ,

$$S^*(X^*(T) \otimes k)^W = \begin{cases} k[c_2, \dots, c_r, \eta_{r-1}] & \text{if } r \neq 2, \\ k[x_1, \eta_1] & \text{if } r = 2. \end{cases}$$

Here  $|c_i| = i$  for  $2 \leq i \leq r$ ,  $|x_1| = 2$ , and  $|\eta_{r-1}| = 2^{r-1}$ .

We now compute  $S^*(X^*(T) \otimes k)^W$  for  $G = \text{Spin}(2r)$ . Note that a maximal torus in  $\text{Spin}(2r)$  is also a maximal torus in  $\text{Spin}(2r+1)$ . So we have again

$$S^*(X^*(T) \otimes k) \cong k[x_1, \dots, x_r, A]/(x_1 + \cdots + x_r).$$

The Weyl group  $W = S_r \rtimes (\mathbf{Z}/2)^{r-1}$  acts on this ring by:  $S_r$  permutes  $x_1, \dots, x_r$ , and fixed  $A$ , and  $(\mathbf{Z}/2)^{r-1}$  is the subgroup  $\langle \epsilon_1 \epsilon_2, \dots, \epsilon_1 \epsilon_r \rangle$  in the notation above. Thus  $\epsilon_1 \epsilon_j$  fixes each  $x_j$  (since we are working modulo 2) and sends  $A$  to  $A - x_1 - x_j$ .

For  $1 \leq j \leq r$ , let  $F_j$  be the subgroup  $\langle \epsilon_1 \epsilon_2, \dots, \epsilon_1 \epsilon_j \rangle \cong (\mathbf{Z}/2)^{j-1} \subset W$ . Let

$$\mu_j = \prod_{\substack{I \subset \{1, \dots, j\} \\ |I| \text{ even}}} \left( A - \sum_{i \in I} x_i \right).$$

Then  $|\mu_j| = 2^{j-1}$  and  $\mu_1 = A$ . Clearly  $\mu_j$  is  $F_j$ -invariant. Benson and Wood observed (or one can check directly) that if  $r$  is even and  $r \geq 4$ , then  $\mu_{r-1}$  is in fact  $W$ -invariant, while if  $r$  is odd and  $r \geq 3$ , then  $\mu_r$  is  $W$ -invariant [4, Proposition 4.1].

For  $1 \leq j \leq r-1$ , an induction on  $j$  using Lemma 12.2 gives that

$$S^*(X^*(T) \otimes k)^{F_j} = k[x_1, \dots, x_r, \mu_j]/(x_1 + \cdots + x_r).$$

If  $r$  is even, then  $-1 := \epsilon_1 \cdots \epsilon_r$  is in  $F_r \subset W$ , and it acts trivially on  $S^*(X^*(T) \otimes k)$ . Therefore, for  $r$  even, we have

$$S^*(X^*(T) \otimes k)^{F_r} = k[x_1, \dots, x_r, \mu_{r-1}]/(x_1 + \cdots + x_r).$$

If  $r$  is odd, then we can apply Lemma 12.2 one more time, yielding that

$$S^*(X^*(T) \otimes k)^{F_r} = k[x_1, \dots, x_r, \mu_r]/(x_1 + \cdots + x_r).$$

The subgroup  $S_r \subset W$  permutes  $x_1, \dots, x_r$ , and fixes  $\mu_{r-1}$ , resp.  $\mu_r$ . We showed above that

$$k[x_1, \dots, x_r]/(x_1 + \dots + x_r)^{S_r} = k[c_2, \dots, c_r].$$

Therefore, for  $G = \text{Spin}(2r)$ , we have

$$S^*(X^*(T) \otimes k)^W = \begin{cases} k[c_2, \dots, c_r, \mu_{r-1}] & \text{if } r \text{ is even} \\ k[c_2, \dots, c_r, \mu_r] & \text{if } r \text{ is odd.} \end{cases}$$

Here  $|c_i| = i$  for  $2 \leq i \leq r$  and  $|\mu_{r-1}| = 2^{r-2}$ , resp.  $|\mu_r| = 2^{r-1}$ .

Thus we have determined  $S^*(X^*(T) \otimes k)^W$  for  $G = \text{Spin}(n)$  for all  $n$ , even or odd. Now think of  $G = \text{Spin}(n)$  as a split reductive group over  $k$ . By Theorem 8.1 (due to Chaput and Romagny), the ring  $S^*(X^*(T) \otimes k)^W = O(\mathfrak{t})^W$  can be identified with  $O(\mathfrak{g})^G$  for all  $n \geq 6$ . (The exceptional cases  $\text{Spin}(3)$ ,  $\text{Spin}(4)$ ,  $\text{Spin}(5)$  are the spin groups that have a factor isomorphic to a symplectic group:  $\text{Spin}(3) \cong Sp(2)$ ,  $\text{Spin}(4) \cong Sp(2) \times Sp(2)$ , and  $\text{Spin}(5) \cong Sp(4)$ .) We deduce that for  $n \geq 6$ ,

$$O(\mathfrak{g})^G = \begin{cases} k[c_2, \dots, c_r, \eta_{r-1}] & \text{if } n = 2r + 1 \\ k[c_2, \dots, c_r, \mu_{r-1}] & \text{if } n = 2r \text{ and } r \text{ is even} \\ k[c_2, \dots, c_r, \mu_r] & \text{if } n = 2r \text{ and } r \text{ is odd.} \end{cases}$$

For  $G = \text{Spin}(n)$  and any  $n \geq 6$ , we have homomorphisms

$$O(\mathfrak{g})^G \rightarrow H_{\text{dR}}^*(BG/k) \rightarrow H_{\text{dR}}^*(BT/k)^W = O(\mathfrak{t})^W,$$

whose composition is the obvious inclusion. (The first homomorphism comes from the isomorphism of  $O(\mathfrak{g})^G$  with  $\bigoplus_i H^i(BG_k, \Omega^i)$ , using that  $H^i(BG_k, \Omega^j) = 0$  for  $i < j$ .) In this case, the restriction  $O(\mathfrak{g})^G \rightarrow O(\mathfrak{t})^W$  is a bijection. So  $H_{\text{dR}}^*(BG/k)$  contains the ring computed above (with degrees multiplied by 2), and retracts onto it. It follows that for all  $n \geq 6$ ,  $H_{\text{dR}}^*(BG/k)$  has an indecomposable generator in degree  $2^r$  if  $n = 2r + 1$ , in degree  $2^{r-1}$  if  $n = 2r$  and  $r$  is even, and in degree  $2^r$  if  $n = 2r$  and  $r$  is odd. (For this argument, we do not need to find all the indecomposable generators of  $H_{\text{dR}}^*(BG/k)$ .) For our application to  $\text{Spin}(11)$ , we note the following information:

**Lemma 12.4.** *The image of  $H_{\text{dR}}^*(B \text{Spin}(2r + 1)/k) \rightarrow H_{\text{dR}}^*(BT/k)$  is the polynomial ring  $k[c_2, \dots, c_r, \eta_{r-1}]$ , where  $|c_i| = 2i$  and  $|\eta_{r-1}| = 2^r$ .*

Compare this with Quillen's calculation of the cohomology of the classifying space of the complex reductive group  $\text{Spin}(n)_{\mathbb{C}}$ , or equivalently of the compact Lie group  $\text{Spin}(n)$  [28, Theorem 6.5]:

$$H^*(B \text{Spin}(n)_{\mathbb{C}}, k) \cong H^*(BSO(n)_{\mathbb{C}}, k)/J \otimes k[w_{2^h}(\Delta_\theta)].$$

Here  $\Delta_\theta$  is a faithful orthogonal representation of  $\text{Spin}(n)_{\mathbb{C}}$  of minimal dimension, and  $J$  is the ideal generated by the regular sequence

$$w_2, Sq^1 w_2, \dots, Sq^{2^{h-2}} \dots Sq^2 Sq^1 w_2$$

in the polynomial ring  $H^*(BSO(n)_{\mathbb{C}}, k) = k[w_2, w_3, \dots, w_n]$ , where  $|w_i| = i$ . Finally, the number  $h$  is given by the following table:

$n$	$h$
$8l + 1$	$4l + 0$
$8l + 2$	$4l + 1$
$8l + 3$	$4l + 2$
$8l + 4$	$4l + 2$
$8l + 5$	$4l + 3$
$8l + 6$	$4l + 3$
$8l + 7$	$4l + 3$
$8l + 8$	$4l + 3$

The Steenrod operations on the mod 2 cohomology of  $BSO(n)_{\mathbf{C}}$ , as used in the formula above, are known, by Wu's formula [25, Theorem III.5.12]:

$$Sq^i w_j = \sum_{l=0}^i \binom{j-l-1}{i-l} w_l w_{i+j-l}$$

for  $0 \leq i \leq j$ , where by convention  $\binom{-1}{0} = 1$ .

Write  $r = \lfloor n/2 \rfloor$ . If  $n = 2r + 1$ , then the generator  $w_{2h}(\Delta_\theta)$  is in degree  $2^r$  if  $r \equiv 0, 3 \pmod{4}$  and in degree  $2^{r+1}$  if  $r \equiv 1, 2 \pmod{4}$ . If  $n = 2r$ , then the generator  $w_{2h}(\Delta_\theta)$  is in degree  $2^{r-1}$  if  $r \equiv 0 \pmod{4}$  and in degree  $2^r$  if  $r \equiv 1, 2, 3 \pmod{4}$ . Therefore, for  $n \geq 11$ ,  $H^*(B\text{Spin}(n)_{\mathbf{C}}, k)$  has no indecomposable generator in degree  $2^r$  if  $n \equiv 3, 5 \pmod{8}$ , and no indecomposable generator in degree  $2^{r-1}$  if  $n \equiv 4 \pmod{8}$ . But  $H_{\text{dR}}^*(BG/k)$  does have an indecomposable generator in the indicated degree  $2^a$ , as shown above. Thus, for  $G = \text{Spin}(n)$ ,  $H^*(BG_{\mathbf{C}}, k)$  is not isomorphic to  $H_{\text{dR}}^*(BG/k)$  as a graded ring when  $n \geq 11$  and  $n \equiv 3, 4, 5 \pmod{8}$ .

We want to show, more precisely, that for  $n = 11$ ,  $H_{\text{dR}}^{32}(BG/k)$  is bigger than  $H^{32}(BG_{\mathbf{C}}, k)$ . We know the cohomology of  $BG_{\mathbf{C}}$  by Quillen (above), and so it remains to give a lower bound for the de Rham cohomology of  $BG$  over  $k$ .

We do this by restricting to a suitable abelian  $k$ -subgroup scheme of  $G = \text{Spin}(n)$ . Assume that  $n \not\equiv 2 \pmod{4}$ ; this includes the case  $\text{Spin}(11)$  that we are aiming for. Then the Weyl group  $W$  of  $\text{Spin}(n)$  contains  $-1$ . So  $\text{Spin}(n)$  contains an extension of  $\mathbf{Z}/2$  by a split maximal torus  $T \cong (G_m)^r$ , where  $\mathbf{Z}/2$  acts by inversion on  $T$ . Let  $L$  be the subgroup of the form  $1 \rightarrow T[2] \rightarrow L \rightarrow \mathbf{Z}/2 \rightarrow 1$ ; then  $L$  is abelian (because inversion is the identity on  $T[2] \cong (\mu_2)^r$ ). Since the field  $k = \mathbf{F}_2$  is perfect, the reduced subscheme of  $L$  is a  $k$ -subgroup scheme (isomorphic to  $\mathbf{Z}/2$ ) [24, Corollary 1.39], and so the extension splits. That is,  $L \cong (\mu_2)^r \times \mathbf{Z}/2$ .

Let us compute the pullbacks of the generators  $u_i$  of  $H_{\text{dR}}^*(BSO(n)/k)$  (Theorem 11.1) to the subgroup  $L$  of  $G = \text{Spin}(n)$ . It suffices to compute the restrictions of the classes  $u_i$  to the image  $K$  of  $L$  in  $SO(n)$ ; clearly  $K \cong (\mu_2)^{r-1} \times \mathbf{Z}/2$ . In notation similar to that used earlier in this proof, the ring of polynomial functions on the Lie algebra of the subgroup  $(\mu_2)^{r-1}$  here is

$$k[t_1, \dots, t_r]/(t_1 + \dots + t_r).$$

This ring can be viewed as the Hodge cohomology ring of  $B(\mu_2)^{r-1}$  modulo its radical, with the generators  $t_i$  in  $H^1(B(\mu_2)^{r-1}, \Omega^1)$  (by Propositions 10.1 and 5.1). Using Lemma 10.2, we conclude that

$$H_{\mathbb{H}}^*(BK/k)/\text{rad} \cong k[s, t_1, \dots, t_r]/(t_1 + \dots + t_r),$$

where  $s$  is pulled back from the generator of  $H^1(B(\mathbf{Z}/2), O)$ . The Hodge spectral sequence for  $BK$  degenerates at  $E_1$ , since we know this degeneration for  $B\mathbf{Z}/2$  and  $B(\mu_2)^{r-1}$ . Therefore,

$$H_{\text{dR}}^*(BK/k)/\text{rad} \cong k[s, t_1, \dots, t_r]/(t_1 + \dots + t_r),$$

Note that the surjection  $L \rightarrow K$  is split. So if we compute that an element of  $H_{\text{dR}}^*(BSO(n)/k)$  has nonzero restriction to  $K$ , then it has nonzero restriction to  $L$ , hence a fortiori to  $G = \text{Spin}(n)$ .

Now strengthen the assumption  $n \not\equiv 2 \pmod{4}$  to assume that  $n$  is odd and  $n \geq 7$ . In Lemma 11.4, we computed the restriction of  $u_2, u_3, \dots, u_{2r+1}$  in de Rham cohomology from  $SO(2r+1)$  to its subgroup  $O(2)^r$ , and hence to its subgroup  $(\mu_2)^r \times (\mathbf{Z}/2)^r$ . We now want to restrict to the smaller subgroup  $K = (\mu_2)^{r-1} \times \mathbf{Z}/2$ . This last step sends  $H_{\text{dR}}^*(B((\mu_2)^r \times (\mathbf{Z}/2)^r)/k)/\text{rad} = k[s_1, \dots, s_r, t_1, \dots, t_r]$  to  $H_{\text{dR}}^*(BK/k)/\text{rad} = k[s, t_1, \dots, t_r]/(t_1 + \dots + t_r)$  by  $s_i \mapsto s$  for all  $i$  and  $t_i \mapsto t_i$ . By Lemma 11.4, the element  $u_{2a}$  (for  $1 \leq a \leq r$ ) restricts to the elementary symmetric polynomial

$$c_a = \sum_{1 \leq i_1 < \dots < i_a \leq r} t_{i_1} \cdots t_{i_a}.$$

Thus  $u_2$  restricts to 0 on  $K$ , but  $u_4, u_6, \dots, u_{2r}$  restrict to generators of the polynomial ring

$$(k[t_1, \dots, t_r]/(t_1 + \dots + t_r))^{S_r} \subset H_{\text{dR}}^*(BK/k)/\text{rad},$$

using that  $r \geq 3$ , by Lemma 12.3.

Next, by Lemma 11.4, for  $1 \leq a \leq r$ , the restriction of  $u_{2a+1}$  to  $H_{\text{dR}}^*(BK/k)/\text{rad}$  is (first restricting from  $SO(2r+1)$  to its subgroup  $(\mu_2)^r \times (\mathbf{Z}/2)^r$ , and then to  $K = (\mu_2)^{r-1} \times \mathbf{Z}/2$ ):

$$\begin{aligned} u_{2a+1} &\mapsto \sum_{m=1}^r s_m \sum_{\substack{1 \leq i_1 < \dots < i_a \leq r \\ \text{one equal to } m}} t_{i_1} \cdots t_{i_a} \\ &\mapsto as \sum_{1 \leq i_1 < \dots < i_a \leq r} t_{i_1} \cdots t_{i_a} \\ &= asu_{2a}. \end{aligned}$$

Thus, for all  $1 \leq a \leq r$ ,  $u_{2a+1}$  restricts in  $H_{\text{dR}}^*(BK/k)/\text{rad}$  to  $su_{2a}$  if  $a$  is odd, and otherwise to zero. (But  $u_2$  restricts to 0, and so this also means that  $u_3$  restricts to 0.)

This gives a lower bound for the image of  $H_{\text{dR}}^*(BSO(n)/k) \rightarrow H_{\text{dR}}^*(B\text{Spin}(n)/k)$  for  $n$  odd. In particular, for  $n = 11$ , this image has Hilbert series at least that of the ring

$$k[u_4, u_6, u_7, u_8, u_{10}, u_{11}]/(u_{11}u_6 + u_{10}u_7),$$

since the latter ring is isomorphic to the image of restriction from  $SO(11)$  to  $H_{\text{dR}}^*(BL/k)/\text{rad}$ , where  $L \subset \text{Spin}(11)$ .

We now compare this to Quillen's computation (above) in the case of  $\text{Spin}(11)$ :

$$\begin{aligned} H^*(B\text{Spin}(11)_{\mathbf{C}}, k) &= k[w_4, w_6, w_7, w_8, w_{10}, w_{11}, w_{64}(\Delta_\theta)]/(w_{11}w_6 + w_{10}w_7, \\ &\quad w_{11}^3 + w_{11}^2w_7w_4 + w_{11}w_8w_7^2). \end{aligned}$$

Since the last generator  $w_{64}(\Delta_\theta)$  is in degree 64 and the last relation is in degree 33, the degree-32 component of this ring has the same dimension as the degree-32 component of the lower bound above for  $H_{\text{dR}}^*(B\text{Spin}(11)/k)$ . However, Lemma 12.4 shows that  $H_{\text{dR}}^*(B\text{Spin}(11)/k)$  has an extra generator  $\eta_4$  in degree 32. This is linearly independent of the image of restriction from  $SO(11)$ , as we see by restricting to a maximal torus  $T$  in  $\text{Spin}(11)$ . Indeed, by Lemma 12.4, the image of  $H_{\text{dR}}^*(B\text{Spin}(11)/k) \rightarrow H_{\text{dR}}^*(BT/k)$  is the polynomial ring  $k[c_2, \dots, c_5, \eta_4]$ , whereas the image of the pullback from  $SO(11)$  to  $T \subset \text{Spin}(11)$  is just  $k[c_2, \dots, c_5]$  ( $= k[w_4, w_6, w_8, w_{10}]$ ). Thus we have shown that

$$\dim_k H_{\text{dR}}^{32}(B\text{Spin}(11)/k) > \dim_k H^{32}(B\text{Spin}(11)_{\mathbf{C}}, k).$$

□

## References

- [1] M. Artin, A. Grothendieck, and J.-L. Verdier, eds. *Théorie des topos et cohomologie étale des schémas* (SGA 4), v. 2. Lecture Notes in Mathematics 270. Springer (1972). 8
- [2] M. Atiyah. Complex analytic connections in fibre bundles. *Trans. Amer. Math. Soc.* **85** (1957), 181–207. 6
- [3] K. Behrend. On the de Rham cohomology of differential and algebraic stacks. *Adv. Math.* **198** (2005), 583–622. 3
- [4] D. Benson and J. Wood. Integral invariants and cohomology of  $B\text{Spin}(n)$ . *Topology* **34** (1995), 13–28. 32, 33, 34
- [5] B. Bhatt. Completions and derived de Rham cohomology. [arXiv:1207.6193](https://arxiv.org/abs/1207.6193) 3, 5
- [6] B. Bhatt, M. Morrow, and P. Scholze. Integral  $p$ -adic Hodge theory. [arXiv:1602.03148](https://arxiv.org/abs/1602.03148) 2
- [7] A. Borel. Sous-groupes commutatifs et torsion des groupes de Lie compacts connexes. *Tohoku Math. J.* **13** (1961), 216–240. 16, 18
- [8] R. Bott. On the Chern-Weil homomorphism and the continuous cohomology of Lie groups. *Adv. Math.* **11** (1973), 289–303. 1, 5
- [9] P.-E. Chaput and M. Romagny. On the adjoint quotient of Chevalley groups over arbitrary base schemes. *J. Inst. Math. Jussieu* **9** (2010), 673–704. 15, 19
- [10] C. Chevalley. Sur les décompositions cellulaires des espaces  $G/B$ . *Algebraic groups and their generalizations: classical methods* (University Park, PA, 1991), 1–23. Proc. Sympos. Pure Math. 56, Part 1, Amer. Math. Soc. (1994). 15
- [11] P. Deligne. Théorie de Hodge III. *Publ. Math. IHES* **44** (1974), 5–77. 4
- [12] M. Demazure. Invariants symétriques entiers des groupes de Weyl et torsion. *Invent. Math.* **21** (1973), 287–301. 15, 18

- [13] S. Donkin. On conjugating representations and adjoint representations of semisimple groups. *Invent. Math.* **91** (1988), 137–145. 1, 11
- [14] A. Dresden. On the generalized Vandermonde determinant and symmetric functions. *Bull. Amer. Math. Soc.* **39** (1933), 443–449. 25
- [15] F. El Zein. Complexe dualisant et applications à la classe fondamentale d’un cycle. *Bull. Soc. Math. France Mém.* No. 58 (1978), 93 pp. 15
- [16] M. Gros. Classes de Chern et classes de cycles en cohomologie de Hodge-Witt logarithmique. *Mém. Soc. Math. France* **21** (1985), 87 pp. 15
- [17] J. Humphreys. *Reflection groups and Coxeter groups*. Cambridge (1990). 2
- [18] L. Illusie. *Complexe cotangent et déformations. I*. Lecture Notes in Mathematics 239, Springer (1971). 7, 10
- [19] L. Illusie. Cotangent complex and deformations of torsors and group schemes. *Toposes, algebraic geometry and logic*, 159–189. Lecture Notes in Mathematics 274, Springer (1972). 9
- [20] L. Illusie. *Complexe cotangent et déformations. II*. Lecture Notes in Mathematics 283, Springer (1972). 6, 7, 9, 10
- [21] J. Jantzen. *Representations of algebraic groups*. Amer. Math. Soc. (2003). 1, 6, 8, 11, 13, 14, 15, 19
- [22] M.-A. Knus, A. Merkurjev, M. Rost, and J.-P. Tignol. *The book of involutions*. Amer. Math. Soc. (1998). 21
- [23] J. P. May. *Simplicial objects in algebraic topology*. Chicago (1992). 12
- [24] J. Milne. *Algebraic groups*. Cambridge (2017). 11, 36
- [25] M. Mimura and H. Toda. *Topology of Lie groups*. Amer. Math. Soc. (1991). 18, 22, 23, 28, 29, 36
- [26] H. Nakajima. Invariants of finite groups generated by pseudoreflections in positive characteristics. *Tsukuba J. Math.* **3** (1979), 109–122. 34
- [27] M. Olsson. Sheaves on Artin stacks. *J. Reine Angew. Math.* **603** (2007), 55–112. 5
- [28] D. Quillen. The mod 2 cohomology rings of extra-special 2-groups and the spinor groups. *Math. Ann.* **194** (1971), 197–212. 35
- [29] C. Simpson and C. Teleman. De Rham’s theorem for  $\infty$ -stacks. <https://math.berkeley.edu/~teleman/lectures.html> 1, 5
- [30] T. Springer and R. Steinberg. Conjugacy classes. *Seminar in algebraic groups and related finite groups*, 167–266. Lecture Notes in Mathematics 131, Springer (1970). Addendum to second corrected printing (1986), 322–323. Also in *Robert Steinberg: Collected Papers*, Amer. Math. Soc. (1997), 293–394. 15

- [31] V. Srinivas. Gysin maps and cycle classes in Hodge cohomology. *Proc. Indian Acad. Sci. (Math. Sci.)* **103** (1993), 209–247. 15
- [32] The Stacks Project Authors. *Stacks Project* (2017). <http://stacks.math.columbia.edu> 3, 4, 6, 7, 9, 12
- [33] C. Teleman. The quantization conjecture revisited. *Ann. Math.* **152** (2000), 1–43. 3

UCLA MATHEMATICS DEPARTMENT, BOX 951555, LOS ANGELES, CA 90095-1555  
TOTARO@MATH.UCLA.EDU