

Rationality does not specialize among terminal varieties

Burt Totaro

An algebraic variety is *rational* if it becomes isomorphic to projective space after removing lower-dimensional subvarieties from both sides. Little is known about how rationality behaves in families. In particular, given a family of projective varieties for which the geometric generic fiber is rational, is every fiber geometrically rational? (“Geometric” refers to properties of a variety after extending its base field to be algebraically closed.)

Matsusaka proved that the analogous question for geometric ruledness has a positive answer [5, Theorem IV.1.6]. (By definition, a variety is ruled if it is birational to the product of the projective line with some variety.) That is, ruledness specializes in families of varieties. For example, Kollár used Matsusaka’s theorem to show that a large class of Fano hypersurfaces are not ruled and therefore not rational [5, Theorem V.5.14]. By contrast, rationality does not specialize in this generality, as shown by a family of cubic surfaces over the complex numbers \mathbf{C} with most fibers smooth and one fiber the projective cone over a smooth cubic curve. Every smooth cubic surface is rational, but the cone over a smooth cubic curve E is birational to $E \times \mathbf{P}^1$, which is not rational because it has a nonzero holomorphic 1-form.

Note, however, that the cone over a cubic curve has a fairly bad singularity: it is log canonical but not klt (Kawamata log terminal). This suggests the question of whether rationality specializes among varieties with milder singularities. Indeed, it follows from de Fernex and Fusi [2, Theorem 1.3] and Hacon and McKernan [3, Corollary 1.5] that rationality specializes among klt complex varieties of dimension at most 3.

Extending work of Voisin [9] and Colliot-Thélène and Pirutka [1], [8, Theorem 2.1] showed that a large class of Fano hypersurfaces X are not stably rational. (That is, no product of X with projective space is rational.) As an application, suggested by de Fernex and Fusi [2, Example 3.4], [8, Corollary 4.1] showed that rationality does not specialize among klt varieties of dimension 4 or higher.

In this paper, we find that the results of [8] are strong enough to imply that rationality does not specialize even among *terminal* varieties. Terminal singularities form the narrowest class of singularities that comes up in the minimal model program. The examples are in any dimension at least 5.

Some natural remaining questions are: Does rationality specialize among terminal 4-folds? Does rationality specialize among smooth varieties?

Here we are asking whether rationality is a closed condition in families. At this writing, it is also unknown whether rationality is an open condition in families of smooth complex projective varieties.

1 The example

Theorem 1.1. *There is a flat projective morphism $f: X \rightarrow C$ with C a Zariski open subset of the complex affine line such that 0 is in C , all fibers of f have terminal singularities, all fibers of f over $C - 0$ are rational, and the fiber F over 0 is not rational.*

Such examples exist with F of any dimension at least 5. There is also a family of 4-folds with canonical singularities over a Zariski open subset C of $A_{\mathbf{C}}^1$ such that all fibers over $C - 0$ are rational and the fiber F over 0 is not rational.

In other words, rationality does not specialize among terminal varieties of dimension at least 5, or among canonical varieties of dimension at least 4. (Throughout, we are talking about families of projective varieties.)

Remark 1.2. Theorem 1.1 was inspired by an example proposed by de Fernex and Fusi [2, Example 3.4], before the technology of [8] was available. Note, however, that the degeneration considered in [2, Example 3.4] has general fiber \mathbf{P}^n and special fiber the projective cone S over $(W, O(n))$, where W is a smooth hypersurface of degree n in \mathbf{P}^n . The special fiber is not the projective cone $T \subset \mathbf{P}^{n+1}$ over $(W, O(1))$, as one sees by a computation of degrees. Here T has canonical singularities, while S is klt but not canonical. As a result, we need a different degeneration (with general fiber not just projective space) in order to prove Theorem 1.1.

Proof. (Theorem 1.1)

We start with the following old observation.

Lemma 1.3. *If X is a hypersurface of degree d in \mathbf{P}^{n+1} over a field k such that X has multiplicity equal to $d - 1$ at some k -rational point p , and if the singular locus of X has codimension at least 2, then X is rational over k .*

Proof. The assumption on the singular locus ensures that X is irreducible. The assumption on the multiplicity of X at p implies that a general line through p meets X in exactly one other point. That gives a birational map over k from the projective space \mathbf{P}^n of lines through p to X . \square

We return to the proof of Theorem 1.1. By [8, Theorem 2.1], a very general quartic 4-fold in $\mathbf{P}_{\mathbf{C}}^5$ is not stably rational. Choose one smooth quartic 4-fold Y over \mathbf{C} which is not stably rational. Let X_0 be the projective cone over Y in \mathbf{P}^6 . Then X_0 is a quartic 5-fold, and X_0 is not rational because it is birational to $\mathbf{P}^1 \times Y$. Also, X_0 is terminal, because Y has Fano index 2 which is greater than 1, meaning that the anticanonical bundle $-K_Y$ is given by $-K_Y \cong -(K_{\mathbf{P}^5} + Y)|_Y = O(6-4)|_Y = O(2)|_Y$ [6, Lemma 3.1].

Let Y be defined by the equation $f_4(x_0, \dots, x_5) = 0$. Then X_0 is defined by the same equation in $\mathbf{P}^6 = \{[x_0, \dots, x_6]\}$. Let $g_3(x_0, \dots, x_5)$ be a nonzero cubic form over \mathbf{C} . Consider the pencil of quartics in \mathbf{P}^6 given by the equation

$$f_4(x_0, \dots, x_5) + ag_3(x_0, \dots, x_5)x_6 = 0$$

for a in the affine line $A_{\mathbf{C}}^1$. This gives a flat family $f: X \rightarrow A^1$ of hypersurfaces, and the fiber over 0 is the cone X_0 . Since “terminal” is a Zariski-open condition in families [7, Corollary VI.5.3], there is a Zariski open neighborhood C of 0 in A^1

such that all fibers of the restricted family $f: X_C \rightarrow C$ are terminal. In particular, the fibers are normal and hence have singular locus of codimension at least 2.

Finally, for all $a \neq 0$ in C , the fiber X_a is a hypersurface of degree 4 in \mathbf{P}^6 with multiplicity equal to 3 at the point $[0, \dots, 0, 1]$. By Lemma 1.3, it follows that X_a is rational for all $a \neq 0$ in C . Since X_0 is not rational, this completes the proof that rationality does not specialize among terminal varieties.

The example given is a family of 5-folds. Multiplying the family with any projective space \mathbf{P}^m shows that rationality does not specialize among terminal varieties of any dimension at least 5. (Here again, it is important that Y is not stably rational, so that $X_0 \times \mathbf{P}^m$ is not rational.)

Finally, replace the 4-fold Y by a smooth quartic 3-fold (again called Y) in $\mathbf{P}_{\mathbb{C}}^4$ which is not stably rational. Such a variety exists, by Colliot-Thélène and Pirutka [1]. It follows that the projective cone X_0 over Y in \mathbf{P}^5 (rather than \mathbf{P}^6) is not rational. Since Y has Fano index 1, X_0 has canonical but not terminal singularities. Also, “canonical” is a Zariski open condition in families [4]. A pencil of hypersurfaces in \mathbf{P}^5 given by the same formula as above shows that rationality does not specialize among 4-folds with canonical singularities. \square

Acknowledgements. I thank the referee for useful suggestions. This work was supported by NSF grant DMS-1303105.

References

- [1] J.-L. Colliot-Thélène and A. Pirutka. Hypersurfaces quartiques de dimension 3: non rationalité stable. *Ann. Sci. École Normale Supérieure*, to appear. 1, 3
- [2] T. de Fernex and D. Fusi. Rationality in families of threefolds. *Rend. Circ. Mat. Palermo* **62** (2013), 127–135. 1, 2
- [3] C. Hacon and J. McKernan. On Shokurov’s rational connectedness conjecture. *Duke Math. J.* **138** (2007), 119–136. 1
- [4] Y. Kawamata. Deformations of canonical singularities. *J. Amer. Math. Soc.* **12** (1999), 85–92. 3
- [5] J. Kollár. *Rational curves on algebraic varieties*. Springer (1996). 1
- [6] J. Kollár. *Singularities of the minimal model program*. Cambridge (2013). 2
- [7] N. Nakayama. *Zariski-decomposition and abundance*. Mathematical Society of Japan (2004). 2
- [8] B. Totaro. Hypersurfaces that are not stably rational. *J. Amer. Math. Soc.*, to appear. 1, 2
- [9] C. Voisin. Unirational threefolds with no universal codimension 2 cycle. *Invent. Math.* **201** (2015), 207–237. 1

UCLA MATHEMATICS DEPARTMENT, BOX 951555, LOS ANGELES, CA 90095-1555
TOTARO@MATH.UCLA.EDU