

Non-injectivity of the map from the Witt group of a variety to the Witt group of its function field

Burt Totaro

For a regular noetherian scheme X with 2 invertible in X , let $W(X)$ denote the Witt group of X [9]. By definition, the Witt group is the quotient of the Grothendieck group of vector bundles with quadratic forms over X by the subgroup generated by quadratic bundles V which admit a Lagrangian subbundle E , meaning that E is its own orthogonal complement in V . A fundamental question is whether the homomorphism from the Witt group of X to the Witt group of its function field is injective. (When X does not have this injectivity property, and X is a variety over a field so that Ojanguren's injectivity theorem for local rings applies [15], then X has an even-dimensional quadratic bundle which is Zariski-locally trivial but which has no Lagrangian subbundle.) Ojanguren [16] and Pardon [17] proved this injectivity when the regular noetherian scheme X is affine of dimension at most 3, and this was extended to all regular noetherian schemes X of dimension at most 3 by Balmer and Walter [1]. There are simple examples of 4-dimensional varieties where injectivity fails, for example the affine quadric $S_{\mathbf{R}}^4$ over the real numbers defined by $x_0^2 + \cdots + x_4^2 = 1$ and the smooth projective quadric of dimension 4 over the complex numbers. But Pardon showed that injectivity holds for all smooth complex affine varieties of dimension 4 [17]. There remained the question of whether Pardon's result could hold in a larger range of dimensions. The only known counterexamples to injectivity for smooth complex affine varieties began in dimension 8, for example $S_{\mathbf{C}}^4 \times S_{\mathbf{C}}^4$, as pointed out by Knus, Ojanguren, and Sridharan ([10], section 5), as well as $S_{\mathbf{C}}^8$. More generally, using Karoubi's calculation of the hermitian K -group of $S_{\mathbf{C}}^n$ for all n ([8], p. 141), one can check that $W(S_{\mathbf{C}}^n) \rightarrow W(\mathbf{C}(S_{\mathbf{C}}^n))$ fails to be injective if and only if $n \equiv 0, 1 \pmod{8}$ and $n \geq 8$.

In this paper, we complete the story as follows.

Theorem 0.1 *There is a smooth complex affine 5-fold U with $W(U) \rightarrow W(\mathbf{C}(U))$ not injective.*

The main technical tools are the spectral sequences of Balmer and Walter [1] and Pardon [18], which together make the relation between Witt groups and Chow groups modulo 2 fairly explicit. For example, these spectral sequences easily imply Parimala's theorem (Theorem 1.4, below, originally stated only for affine varieties) that the Witt group of a smooth complex 3-fold X is finite if and only if the Chow group $CH^2(X)/2$ is finite [19]. Schoen recently found smooth complex 3-folds X with $CH^2(X)/l$ infinite for some primes l [21], and I expect that his method can be extended to give such examples with $l = 2$. That would give the first known examples of smooth complex varieties with infinite Witt group.

For Theorem 0.1, we need to identify the first differential in Pardon's spectral sequence with the Steenrod operation Sq^2 on Chow groups modulo 2, as defined

by Voevodsky [24] and Brosnan [3]. Given that, the main problem is to exhibit a smooth complex affine variety of high dimension n whose Chow group of curves modulo 2 is nonzero. (Notice that the Chow group of zero-cycles modulo 2 is zero in this situation, as is the cohomology group $H^{2n-2}(X, \mathbf{F}_2)$ to which the Chow group of curves maps.) Probably one could construct such an example by the methods of Kollár and van Geemen [12] or Schoen [21]. But we use a more elementary method, based on the phenomenon that a fibration of a “sufficiently nonsingular” variety which has a multiple fiber cannot have a section.

I would like to thank Paul Balmer, Philippe Gille, and Manuel Ojanguren for useful discussions.

1 The Gersten-Witt complex and Chow groups

Let X be a regular noetherian integral separated scheme of finite dimension with 2 invertible in X . Balmer and Walter [1] and Pardon [18] defined the Gersten-Witt complex W_X of X as a cochain complex

$$0 \rightarrow W(k(X)) \rightarrow \bigoplus_{x \in X^{(1)}} W(k(x)) \rightarrow \cdots \rightarrow \bigoplus_{x \in X^{(n)}} W(k(x)) \rightarrow 0,$$

where $X^{(i)}$ denotes the set of codimension- i points of the scheme X , or equivalently the set of codimension- i subvarieties of X . More canonically, as explained in Balmer and Walter’s equation (6) [1] or Pardon’s Corollary 0.11 [18], this sequence can be written

$$0 \rightarrow W(k(X)) \rightarrow \bigoplus_{x \in X^{(1)}} W(k(x); \omega_{x/X}) \rightarrow \cdots \rightarrow \bigoplus_{x \in X^{(n)}} W(k(x); \omega_{x/X}) \rightarrow 0.$$

Here $\omega_{x/X}$ denotes the 1-dimensional $k(x)$ -vector space $\det(m_x/m_x^2)$, where m_x is the maximal ideal in the local ring $O_{X,x}$, and $W(F; L)$ is the Witt group of quadratic forms over a field F that take values in a given 1-dimensional vector space L over F . Let $H^*(W_X)$ denote the cohomology groups of this cochain complex, with $W(k(X))$ placed in degree 0. The group $H^0(W_X)$ is also known as the unramified Witt group $W_{nr}(X)$. The groups $H^*(W_X)$ are related to the 4-periodic Witt groups $W^*(X)$ by Balmer and Walter’s “Gersten-Witt spectral sequence” [1], which reduces to the following exact sequence for X of dimension at most 7:

$$\begin{aligned} 0 \rightarrow H^4(W) \rightarrow W^0 \rightarrow H^0(W) \rightarrow H^5(W) \rightarrow W^1 \rightarrow H^1(W) \\ \rightarrow H^6(W) \rightarrow W^2 \rightarrow H^2(W) \rightarrow H^7(W) \rightarrow W^3 \rightarrow H^3(W) \rightarrow 0. \end{aligned}$$

In particular, this exact sequence shows that for X of dimension at most 7, the group $H^4(W_X)$ is isomorphic to the kernel of the homomorphism from the Witt group of X to the Witt group of its function field. The group $H^4(W_X)$ is obviously zero for X of dimension at most 3, which gives Balmer and Walter’s theorem that the homomorphism is injective for X of dimension at most 3. In this paper we will prove that the homomorphism is not injective for some smooth complex affine 5-fold U ; equivalently, $H^4(W_U)$ can be nonzero for such a U .

To compute the groups $H^*(W_X)$, we can filter the complex W_X by the subgroups I^n in the Witt groups of fields. This gives Pardon’s spectral sequence ([18], 0.13), as follows. Write $H_{\text{Zar}}^i(X, H_{\mathbf{F}_2}^j)$, or $H^i(X, H^j)$ for short, to mean the Zariski cohomology of X with coefficients in the presheaf $H_{\mathbf{F}_2}^j(V) := H_{\text{et}}^j(V, \mathbf{F}_2)$, for open

subsets $V \subset X$. Remarkably, these are the same groups that form the E_2 term of the Bloch-Ogus spectral sequence (Theorem 1.2, below), although the following spectral sequence is different.

Theorem 1.1 *For any smooth variety X over a field k of characteristic not 2, there is a spectral sequence $E_2^{ij} = H_{Zar}^i(X, H_{\mathbf{F}_2}^j) \Rightarrow H^i(W_X)$. The differentials are in an unusual direction: d_r has bidegree $(1, r-1)$ for $r \geq 2$. The groups $H^i(X, H^j)$ are zero unless $0 \leq i \leq j$. The groups on the main diagonal are the Chow groups modulo 2: $H^i(X, H^i) = CH^i(X)/2$. Finally, the differential on this main diagonal, $d_2 : CH^i(X)/2 \rightarrow CH^{i+1}(X)/2$, is the Steenrod operation Sq^2 , as defined on Chow groups by Voevodsky [24] and Brosnan [3].*

$$\begin{array}{cccc}
 H^0(X, H^3) & H^1(X, H^3) & H^2(X, H^3) & H^3(X, H^3) \\
 \nearrow & \nearrow & \nearrow & \nearrow \\
 H^0(X, H^2) & H^1(X, H^2) & H^2(X, H^2) & \\
 \nearrow & \nearrow & \nearrow & \\
 H^0(X, H^1) & H^1(X, H^1) & & \\
 \nearrow & \nearrow & & \\
 H^0(X, H^0) & & &
 \end{array}$$

The Chow group $CH^i X$ is defined as the group of codimension- i algebraic cycles on X modulo rational equivalence. A general reference on Chow groups is Fulton's book ([5], 1.3).

Proof. Most of this is Pardon's Corollary 0.13 [18], specialized to the case where the sheaf C is the trivial line bundle O_X . Also, Pardon's grading of the spectral sequence differs from that used here. The vanishing of $H^i(X, H^j)$ for $i > j$ and the isomorphism $H^i(X, H_{\mathbf{F}_2}^i) = CH^i(X)/2$ were proved by Bloch and Ogus ([2], 6.2 and proof of 7.7). The new point is the identification of the differential $d_2 : CH^i(X)/2 \rightarrow CH^{i+1}(X)/2$ with Sq^2 .

To prove this, it is convenient to reduce to the case where the field k is perfect. Indeed, for any field extension E of k , there is a natural flat pullback map of Chow groups, $CH^i(X) \rightarrow CH^i(X_E)$. Both the Pardon differential d_2 and the Steenrod operation Sq^2 are compatible with this pullback map. Furthermore, when E is a finite extension of k of odd degree, the map $CH^i(X)/2 \rightarrow CH^i(X_E)/2$ is injective, by Fulton [5], 15.1.5. Since k does not have characteristic 2, any inseparable extension of k has odd degree. By taking direct limits, it follows that the natural map $CH^i(X)/2 \rightarrow CH^i(X_E)/2$ is injective when E is the perfect closure of k (that is, $E = \cup_r k^{1/p^r}$ for k of characteristic $p > 0$, or $E = k$ in characteristic zero). Therefore, if we prove that the Pardon differential d_2 is equal to Sq^2 for varieties over a perfect field, then the same is true for varieties over any field. Thus, we can now assume that the base field k is perfect.

Let n be the dimension of the smooth variety X . Let Y be a smooth $(n-i)$ -dimensional variety with a proper map $f : Y \rightarrow X$. Then the Steenrod operation

$Sq^2 : CH^i(X)/2 \rightarrow CH^{i+1}(X)/2$ is defined in such a way that

$$\begin{aligned} Sq^2(f_*[Y]) &= f_*(c_1 f^*(TX) - c_1 TY) \\ &= c_1(TX) f_*[Y] - f_* c_1(TY), \end{aligned}$$

by 8.10, 8.11, 9.4 in Brosnan [3]. This property defines Sq^2 uniquely. In fact, for any codimension- i subvariety Z of X , the normalization of Z is regular in codimension 1, since a normal noetherian local ring of dimension 1 is regular ([13], 11.2). Therefore, we can remove a codimension-2 subset of Z from Z and from X , so as to make the normalization Y of Z regular. Since the base field k is perfect, it follows that Y is smooth over k ([13], 26.3, Lemma 1 in section 28, 30.3). Removing that subset does not change the group $CH^{i+1}(X)/2$, since the algebraic cycles and rational equivalences that define this group are unchanged. Having thus modified X , we can define $Sq^2[Z]$ by the above formula applied to the proper map $Y \rightarrow X$.

Furthermore, the variety Z is Cohen-Macaulay in codimension 1, since any reduced noetherian local ring of dimension 1 is Cohen-Macaulay ([13], Exercise 17.1(b)). So, after removing a suitable closed subset of codimension 2 in Z from Z and from X , we can assume that Z is Cohen-Macaulay, as well as having smooth normalization. As above, removing this subset does not change the group $CH^{i+1}(X)/2$ we are concerned with. With this change made, Z has a dualizing sheaf ω_Z (a short notation for $\omega_{Z/k}$). Indeed, any variety Z over k has a dualizing complex, defined as $\pi^1(k)$ where π is the map from Z to $\text{Spec } k$, and Cohen-Macaulayness of Z means that this complex is a single sheaf ω_Z placed in degree $-\dim Z$. This aspect of Grothendieck-Verdier duality is described in Hartshorne's book [7], V, Exercise 9.7, or in more detail in Conrad's book [4], 3.5.1.

Having described Sq^2 of the codimension- i subvariety Z of X , we now compare it to the differential d_2 . To compute $d_2[Z]$, we first lift the nonzero element of $W/I(k(z), \omega_{z/X}) \cong \mathbf{F}_2$ to an element of $W(k(z); \omega_{z/X})$. For example, we can do this by choosing a 1-dimensional $k(z)$ -vector space L together with an isomorphism $L \otimes L \cong \omega_{z/X}$. We can choose a basis element for the 1-dimensional vector space L ; then we get an isomorphism $k(z) \cong \omega_{z/X}$. Since Z is Cohen-Macaulay, we can view the isomorphism $k(z) \cong \omega_{z/X}$ as a nonzero rational section of the relative dualizing sheaf $\omega_{Z/X} := \omega_Z \otimes_{O_Z} \omega_X^*|_Z$.

The differential in the Gersten-Witt complex is defined by Balmer and Walter's Proposition 8.5 [1], as a sum over codimension-1 points d in Z of elements M in the Witt group of finite-length $O_{X,d}$ -modules, which can be identified with $W(k(d), \omega_{d/X})$. To compute $d_2[Z]$ in the Pardon spectral sequence, we only need to know the class of these elements in $W/I(k(d), \omega_{d/X}) \cong \mathbf{F}_2$, that is, the length modulo 2 of the finite-length $O_{X,d}$ -module M . Thus $d_2[Z]$ is a divisor with \mathbf{F}_2 coefficients on Z . Balmer and Walter's exact sequence (35) shows that this divisor has the form

$$d_2[Z] = \omega_{Z/X} - O_Z$$

in the first graded piece of the Grothendieck group of coherent sheaves on Z , filtered by codimension of support, $(\text{gr}^1 G_0 Z)/2$. Here inspection of Quillen's spectral sequence that converges to this filtration of $G_0 Z$ shows that $\text{gr}^1 G_0 Z$ is isomorphic to the divisor class group of Z ([20], section 7, Theorem 5.4). Also, to obtain the above formula, we use that the module $\text{Ext}_O^{e-1}(A, O)$ which Balmer and Walter call

ω_A in their exact sequence (35), apparently corresponding to ω_Z in the notation here, is really the relative dualizing sheaf $\omega_{Z/X}$.

Since $\omega_{Z/X}$ is the tensor product of the sheaf ω_Z with the line bundle $\omega_X^*|_Z$, it is easy to rewrite the formula for $d_2[Z]$ as:

$$\begin{aligned}\omega_{Z/X} - O_Z &= -(\omega_X|_Z - O_Z) + (\omega_Z - O_Z) \\ &= c_1(TX)|_Z + (\omega_Z - O_Z)\end{aligned}$$

in the divisor class group $(\text{gr}^1 G_0 Z)/2$. To show that this class equals $Sq^2[Z]$ in $CH^{i+1}(X)/2$, as defined by Brosnan's formula above, it remains to identify the class $\omega_Z - O_Z$ in $(\text{gr}^1 G_0 Z)/2$ with $f_*c_1(TY)$, where $f : Y \rightarrow Z$ is the normalization of Z . Recall that we have removed a codimension-2 subset of Z in such a way that the normalization Y is smooth. Here we can write $f_*(c_1 TY)$ as $f_*\omega_Y - f_*O_Y$ in $(\text{gr}^1 G_0 Z)/2$.

Let A and B denote the cokernels of the natural maps of sheaves on Z :

$$0 \rightarrow O_Z \rightarrow f_*O_Y \rightarrow A \rightarrow 0$$

and

$$0 \rightarrow f_*\omega_Y \rightarrow \omega_Z \rightarrow B \rightarrow 0.$$

The equality we want, saying that $\omega_Z - O_Z$ equals $f_*\omega_Y - f_*O_Y$ in $(\text{gr}^1 G_0 Z)/2$, will clearly follow if we can show that the sheaves A and B are equal in $(\text{gr}^1 G_0 Z)/2$. Since the sheaves A and B are zero at the generic point of Z , it suffices to show that A and B have the same coefficient in \mathbf{F}_2 at each codimension-1 point d in Z . Let $O_{Z,d}$ denote the local ring of Z at d . In these terms, we need to know that the $O_{Z,d}$ -modules A and B have the same length modulo 2.

Using Grothendieck-Verdier duality, we find an isomorphism of O_Z -modules $B \cong \text{Ext}_{O_Z}^1(A, \omega_Z)$. In more detail, consider the long exact sequence of O_Z -modules obtained from the first exact sequence above by applying $\text{Hom}_{O_Z}(\cdot, \omega_Z)$:

$$\text{Hom}_{O_Z}(f_*O_Y, \omega_Z) \rightarrow \text{Hom}_{O_Z}(O_Z, \omega_Z) \rightarrow \text{Ext}_{O_Z}^1(A, \omega_Z) \rightarrow \text{Ext}_{O_Z}^1(f_*O_Y, \omega_Z).$$

We have $\text{Ext}_{O_Z}^i(Rf_!M, N) \cong f_*\text{Ext}_{O_Y}^i(M, f^!N)$ for all i , by Theorem 1 in Verdier [23]. In the case at hand, $Rf_!O_Y$ is just the sheaf f_*O_Y because f is finite. Also, since Z is Cohen-Macaulay, we have already mentioned that the natural map $\pi_Z : Z \rightarrow \text{Spec } k$ has the property that the complex $\pi_Z^!(k)$ is simply a sheaf ω_Z placed in degree $-\dim Z$. The same goes for the smooth variety Y . In view of the commutative diagram

$$\begin{array}{ccc} Y & \xrightarrow{f} & Z \\ & \searrow \pi_Y & \downarrow \pi_Z \\ & & \text{Spec } k, \end{array}$$

it follows that $f^!\omega_Z$ is isomorphic to the sheaf ω_Y . So the above exact sequence can be rewritten as

$$f_*\omega_Y \rightarrow \omega_Z \rightarrow \text{Ext}_{O_Z}^1(A, \omega_Z) \rightarrow f_*\text{Ext}_{O_Y}^1(O_Y, \omega_Y) = 0.$$

This proves that $B \cong \text{Ext}_{O_Z}^1(A, \omega_Z)$, as we claimed.

Finally, for each codimension-1 point d in Z , the functor $A \mapsto \text{Ext}_{O_{Z,d}}^1(A, \omega_Z)$ is a duality on the category of finite-length $O_{Z,d}$ -modules, preserving the length of such modules. This is a special case of Grothendieck's local duality theorem, using that the local ring $O_{Z,d}$ is Cohen-Macaulay of dimension 1 ([7], IV.6.3). Thus A and B have the same length at each divisor in Z . As mentioned above, this completes the proof that the differential $d_2 : CH^i(X)/2 \rightarrow CH^{i+1}(X)/2$ is the Steenrod operation Sq^2 . QED

In order to understand the groups $H^i(X, H^j)$, and compute them in some cases, one can use the Bloch-Ogus spectral sequence ([2], 6.3). It has the same E_2 term as Pardon's spectral sequence, but with differentials in the usual direction:

Theorem 1.2 (Bloch-Ogus) *For any smooth variety X over a field of characteristic not 2, there is a spectral sequence $E_2^{i,j} = H_{\text{Zar}}^i(X, H_{\mathbf{F}_2}^j) \Rightarrow H_{\text{et}}^{i+j}(X, \mathbf{F}_2)$. The groups $H^i(X, H^j)$ in the E_2 term are zero unless $0 \leq i \leq j$. The groups on the main diagonal are the Chow groups modulo 2: $H^i(X, H^i) = CH^i(X)/2$. Finally, the differentials are in the usual direction: d_r has bidegree $(r, 1 - r)$, $r \geq 2$.*

$$\begin{array}{ccccccc}
 H^0(X, H^3) & & H^1(X, H^3) & & H^2(X, H^3) & & H^3(X, H^3) \\
 & \searrow & & & & & \\
 H^0(X, H^2) & & H^1(X, H^2) & & H^2(X, H^2) & & \\
 & & & & & & \\
 H^0(X, H^1) & & H^1(X, H^1) & & & & \\
 & & & & & & \\
 H^0(X, H^0) & & & & & &
 \end{array}$$

For completeness, let me explain how the groups $H_{\text{Zar}}^i(X, H_{\mathbf{F}_2}^j)$ in the E_2 term of the Bloch-Ogus spectral sequence are related to motivic cohomology groups. This will not be used in the rest of the paper.

Let X be a smooth variety over a field k of characteristic not 2. There is a natural sequence of homomorphisms relating the motivic cohomology groups of X with different weights,

$$\dots \rightarrow H_M^i(X, \mathbf{F}_2(j)) \rightarrow H_M^i(X, \mathbf{F}_2(j+1)) \rightarrow \dots,$$

given by cup product with the nonzero element of $H_M^0(k, \mathbf{F}_2(1)) \cong \mu_2(k) \cong \mathbf{F}_2$. The motivic cohomology groups in this sequence are initially zero, namely for $i > 2j$. The first group to appear is the Chow group $CH^j(X)/2$, when $i = 2j$. Finally, the groups in the sequence are eventually isomorphic to the etale cohomology $H_{\text{et}}^i(X, \mathbf{F}_2)$, namely when $j \geq i$. Now we can explain the groups $H_{\text{Zar}}^i(X, H_{\mathbf{F}_2}^j)$ in the E_2 term of the Bloch-Ogus spectral sequence: they are the differences between one motivic cohomology group and the next one, in the above sequence. More precisely:

Theorem 1.3 *For each $j \geq 0$, there is a long exact sequence:*

$$\rightarrow H_M^{i+j}(X, \mathbf{F}_2(j-1)) \rightarrow H_M^{i+j}(X, \mathbf{F}_2(j)) \rightarrow H_{\text{Zar}}^i(X, H_{\mathbf{F}_2}^j) \rightarrow H_M^{i+j+1}(X, \mathbf{F}_2(j-1)) \rightarrow$$

Proof. The ingredients here are deep. First, by Voevodsky, we know the Milnor conjecture, also known as the Bloch-Kato conjecture with \mathbf{F}_2 coefficients ([25], Theorem 1.3). Namely, for any field k of characteristic not 2, the natural map from the Milnor K -theory of k modulo 2 to the Galois cohomology of k with \mathbf{F}_2 coefficients is an isomorphism.

Next, by Geisser and Levine ([6], Theorem 1.1), generalizing the work of Suslin and Voevodsky in characteristic zero [22], we know that the Bloch-Kato conjecture with \mathbf{F}_p coefficients implies the Beilinson-Lichtenbaum conjecture with \mathbf{F}_p coefficients. As a result, we know that the Beilinson-Lichtenbaum conjecture with \mathbf{F}_2 coefficients is true.

This means that the natural map

$$\mathbf{F}_2(j) \rightarrow \tau_{\leq j} R\alpha_*(\mathbf{F}_2)$$

is an isomorphism for all $j \geq 0$. This is a map in the derived category of sheaves in the Zariski topology on any smooth variety X over a field of characteristic not 2. The object $\mathbf{F}_2(j)$ is defined so that $H_{\text{Zar}}^i(X, \mathbf{F}_2(j))$ is the motivic cohomology of X with \mathbf{F}_2 coefficients, also written $H_M^i(X, \mathbf{F}_2(j))$.

On the right, $\tau_{\leq j}$ is a truncation, α is the projection from the étale site to the Zariski site, and \mathbf{F}_2 is a sheaf in the étale topology on X . More concretely, this means that the object $\tau_{\leq j} R\alpha_*(\mathbf{F}_2)$ in the derived category of Zariski sheaves on X has cohomology sheaf in degree i given as follows. For $i > j$, this sheaf is zero; for $i \leq j$, it is the sheaf $H_{\mathbf{F}_2}^i$ of étale cohomology groups.

In particular, there is a distinguished triangle in the derived category of Zariski sheaves on X :

$$\tau_{\leq j-1} R\alpha_*(\mathbf{F}_2) \rightarrow \tau_{\leq j} R\alpha_*(\mathbf{F}_2) \rightarrow H_{\mathbf{F}_2}^j[-j],$$

where the shift $[-j]$ just means that the sheaf $H_{\mathbf{F}_2}^j$ is placed in degree j . The associated long exact sequence of Zariski cohomology groups is the result we want. QED

As an application of the Balmer-Walter and Pardon spectral sequences, not needed for the rest of the paper, we can give an easy proof of the following theorem of Parimala (originally stated only for affine varieties) [19].

Theorem 1.4 *Let X be a smooth complex 3-fold. Then the Witt group $W(X)$ is finite if and only if the Chow group $CH^2(X)/2$ is finite.*

Proof of Theorem 1.4. Consider the Bloch-Ogus spectral sequence for X . Since X has dimension 3 over the complex numbers, the groups $H^i(X, H^j)$ in the E_2 term are zero unless $j \leq 3$. Combining this with the other properties listed in Theorem 1.2, we see that the only possible differential in the Bloch-Ogus spectral sequence is $d_2 : H^0(X, H^3) \rightarrow H^2(X, H^2) = CH^2(X)/2$. The spectral sequence converges to $H^*(X, \mathbf{F}_2)$, which is finite. So all the groups $H^i(X, H^j)$ are finite except possibly for $H^0(X, H^3)$ and $H^2(X, H^2) = CH^2(X)/2$, and $H^0(X, H^3)$ is finite if and only if $CH^2(X)/2$ is finite.

Thus, if $CH^2(X)/2$ is finite, then all the groups $H^i(X, H^j)$ are finite. By the Pardon spectral sequence, it follows that the groups $H^i(W_X)$ are finite, and by the Balmer-Walter spectral sequence, it follows that the Witt groups $W^i(X)$ are finite. In particular, the classical Witt group $W^0(X)$ is finite.

Conversely, if $CH^2(X)/2$ is infinite, then $H^0(X, H^3)$ is infinite. There are no possible differentials involving this group in the Pardon spectral sequence, and so $H^0(W_X)$ is infinite. Also, for dimension reasons, there are no differentials in the Balmer-Walter spectral sequence, and so $W^0(X)$ is infinite. QED

2 The example

We come to the problem of finding a smooth complex affine 5-fold U with $W(U) \rightarrow W(\mathbf{C}(U))$ not injective. As explained in section 1, it is equivalent to make $H^4(W_U)$ nonzero. Using Pardon's spectral sequence, Theorem 1.1, for any smooth complex affine 5-fold U , we can ask when an element α of $H^4(U, H^4) = CH^4(U)/2$ corresponds to some nonzero element of $H^4(W_U)$. First, the only possible differential on α maps it to $H^5(U, H^5) = CH^5(U)/2$, but this group is zero. Indeed, more generally, the Chow group of zero-cycles on any noncompact complex variety is a divisible abelian group, by Fulton's Example 1.6.6 [5]. As a result, $CH_0(U)/2 = 0$. (Note that $CH^5 U = CH_0 U$ for U of dimension 5.)

The only differential in Pardon's spectral sequence that might hit α is $Sq^2 : CH^3(U)/2 \rightarrow CH^4(U)/2$. Thus we have shown:

Lemma 2.1 *Let U be a noncompact smooth complex 5-fold. If $Sq^2 : CH^3(U)/2 \rightarrow CH^4(U)/2$ is not surjective, then $H^4(W_U)$ is not zero.*

We will choose U to be a suitable open subset of $Q^4 \times (A^1 - 0)$, where Q^4 denotes the smooth complex 4-dimensional quadric. Any open subset U of $Q^4 \times (A^1 - 0)$ has the convenient property that the Steenrod operation $Sq^2 : CH^3(U)/2 \rightarrow CH^4(U)/2$ is identically zero, as we now show. The pullback map $CH^*(Q^4) \rightarrow CH^*(Q^4 \times A^1)$ is an isomorphism ([5], 3.3(a)), and the restriction to an open subset $CH^*(Q^4 \times A^1) \rightarrow CH^*U$ is surjective ([5], 1.8). So it suffices to show that

$$Sq^2 : CH^3(Q^4)/2 \rightarrow CH^4(Q^4)/2$$

is zero. The Chow ring of a complex quadric maps isomorphically to its cohomology ([5], 19.1.11), and so it suffices to show that

$$Sq^2 : H^6(Q^4, \mathbf{F}_2) \rightarrow H^8(Q^4, \mathbf{F}_2)$$

is zero. For this, we use Wu's formula (see Milnor and Stasheff [14], 11.14): to show that Sq^2 is zero when mapping into the top degree of $H^*(Q^4, \mathbf{F}_2)$, it is equivalent to show that the Wu class $v_2(Q^4) = w_2(Q^4) + w_1^2(Q^4)$ is zero in $H^2(Q^4, \mathbf{F}_2)$. Since Q^4 is a complex manifold, the Wu class is the reduction modulo 2 of $c_1(Q^4)$. But $c_1(Q^4)$ is 4 times the hyperplane class, and so its reduction modulo 2 is zero, as we want. (In other words, we are using that an even-dimensional complex quadric is a spin manifold.)

Thus $Sq^2 : CH^3(U)/2 \rightarrow CH^4(U)/2$ is zero. Combining this with Lemma 2.1 gives:

Lemma 2.2 *Let U be an open subset of $Q^4 \times (A^1 - 0)$. If $CH^4(U)/2$ is not zero, then $H^4(W_U)$ is not zero.*

It remains to find an affine open subset U of $Q^4 \times (A^1 - 0)$ such that the Chow group $CH^4(U)/2 = CH_1(U)/2$ of curves modulo 2 is not zero. Let $Q^4 \subset \mathbf{P}^5$ be defined by

$$x_0x_1 + x_2x_3 + x_4x_5 = 0.$$

We define U to be the complement of the hypersurface Y in $Q^4 \times (A^1 - 0)$ defined by

$$x_0^8 + tx_1^8 + t^2x_2^8 + t^3x_3^8 + t^4x_4^8 + t^5x_5^8 = 0.$$

(Equations of this form were used in a different context by Kollár [11], IV.6.4.3.1.) The complement of the hypersurface defined by this equation in $\mathbf{P}^5 \times (A^1 - 0)$ is clearly affine, and U is the closed subset of this affine variety defined by the quadric. Therefore the smooth complex 5-fold U is affine.

It remains to show that $CH_1(U)/2$ is not zero. By the exact sequence

$$CH_1(Y)/2 \rightarrow CH_1(Q^4 \times (A^1 - 0))/2 \rightarrow CH_1(U)/2 \rightarrow 0$$

from Fulton ([5], 1.8), it is equivalent to show that $CH_1(Y)/2 \rightarrow CH_1(Q^4 \times (A^1 - 0))/2$ is not surjective. The proper pushforward map

$$CH_1(Q^4 \times (A^1 - 0))/2 \rightarrow CH_1(A^1 - 0)/2 = \mathbf{F}_2$$

is surjective, since the curve $p \times (A^1 - 0)$ has degree 1 over $A^1 - 0$ for any point p in Q^4 . Therefore we will be done if we can show that the pushforward map $CH_1(Y)/2 \rightarrow CH_1(A^1 - 0)/2 = \mathbf{F}_2$ is not surjective, in other words that every curve in Y has even degree over $A^1 - 0$.

We can in fact forget about the quadric at this point, and only consider the hypersurface Z^5 in $\mathbf{P}^5 \times (A^1 - 0)$ defined by

$$x_0^8 + tx_1^8 + t^2x_2^8 + t^3x_3^8 + t^4x_4^8 + t^5x_5^8 = 0.$$

We will show that every curve in Z has even degree over $A^1 - 0$, which obviously implies the same statement for curves in $Y^4 \subset Z^5$. The point is that if there was a curve on Z of odd degree over $A^1 - 0$, then we could restrict this curve over a neighborhood of 0 in A^1 . It could fall into several pieces, but at least one would have odd degree over $\mathbf{C}((t))$. Thus, we are finished if we can show that the morphism $t : Z \rightarrow A^1 - 0$ has no section over $\mathbf{C}((t^{1/r}))$ for any odd r .

This is immediate by a power series calculation. Let $u = t^{1/r}$, r odd, and suppose that $Z \rightarrow A^1 - 0$ has a section over $\text{Spec } \mathbf{C}((u)) \rightarrow A^1 - 0$. This means that there are Laurent series $x_i(u)$ which satisfy the equation

$$x_0^8 + u^r x_1^8 + u^{2r} x_2^8 + u^{3r} x_3^8 + u^{4r} x_4^8 + u^{5r} x_5^8 = 0.$$

But the lowest degrees in u of these 6 terms are congruent to $0, r, 2r, \dots, 5r \pmod{8}$. Since r is odd, these lowest degrees are all different. So the only way the 6 terms can add up to zero is if all are identically zero. That would imply that $x_i(u) = 0$ for all i , but this does not correspond to a point in projective space. Thus $Z \rightarrow A^1 - 0$ has no section over $\mathbf{C}((t^{1/r}))$ for any odd r .

This completes the proof that the smooth complex affine 5-fold U has $H^4(W_U)$ nonzero, and so the homomorphism $W(U) \rightarrow W(\mathbf{C}(U))$ is not injective. QED (Theorem 0.1).

References

- [1] P. Balmer and C. Walter. A Gersten-Witt spectral sequence for regular schemes. *Ann. Sci. ENS* **35** (2002), 127–152.
- [2] S. Bloch and A. Ogus. Gersten’s conjecture and the homology of schemes. *Ann. Sci. ENS* **7** (1974), 181–201.
- [3] P. Brosnan. Steenrod operations in Chow theory. *Trans. AMS* **355** (2003), 1869–1903.
- [4] B. Conrad. *Grothendieck duality and base change*. Lecture Notes in Mathematics **1750**, Springer (2000).
- [5] W. Fulton. *Intersection theory*. Springer (1984).
- [6] T. Geisser and M. Levine. The Bloch-Kato conjecture and a theorem of Suslin-Voevodsky. *J. Reine Angew. Math.* **530** (2001), 55–103.
- [7] R. Hartshorne. *Residues and duality*. Lecture Notes in Mathematics **20**, Springer (1966).
- [8] M. Karoubi. Localisation de formes quadratiques II. *Ann. Sci. ENS* **8** (1975), 99–155.
- [9] M.-A. Knus. *Quadratic and hermitian forms over rings*. Springer (1991).
- [10] M.-A. Knus, M. Ojanguren, and R. Sridharan. Quadratic forms and Azumaya algebras. *J. Reine Angew. Math.* **303/304** (1978), 231–248.
- [11] J. Kollár. *Rational curves on algebraic varieties*. Springer (1996).
- [12] J. Kollár and B. van Geemen. Trento examples. *Classification of irregular varieties* (Trento, 1990), 134–135. Lecture Notes in Mathematics **151**, Springer (1992).
- [13] H. Matsumura. *Commutative ring theory*. Cambridge (1986).
- [14] J. Milnor and J. Stasheff. *Characteristic classes*. Princeton (1974).
- [15] M. Ojanguren. Quadratic forms over regular rings. *J. Ind. Math. Soc.* **44** (1980), 109–116.
- [16] M. Ojanguren. A splitting theorem for quadratic forms. *Comment. Math. Helv.* **57** (1982), 145–157.
- [17] W. Pardon. A relation between Witt groups and zero-cycles in a regular ring. *Algebraic K-theory, number theory, geometry and analysis* (Bielefeld, 1982), 261–328. Lecture Notes in Mathematics **1046**, Springer (1984).
- [18] W. Pardon. The filtered Gersten-Witt complex for regular schemes. <http://www.math.uiuc.edu/K-theory/0419>
- [19] R. Parimala. Witt groups of affine 3-folds. *Duke Math. J.* **57** (1988), 947–954.

- [20] D. Quillen. Higher algebraic K -theory I. *Algebraic K-theory I* (Seattle, 1972), 85–147. Lecture Notes in Mathematics **341**, Springer (1973).
- [21] C. Schoen. Complex varieties for which the Chow group mod n is not finite. *J. Alg. Geo.* **11** (2001), 41–100.
- [22] A. Suslin and V. Voevodsky. Bloch-Kato conjecture and motivic cohomology with finite coefficients. *The arithmetic and geometry of algebraic cycles* (Banff, 1998), 117–189. Kluwer (2000).
- [23] J.-L. Verdier. Base change for twisted inverse image of coherent sheaves. *Algebraic geometry* (Bombay, 1968), 393–408. Oxford (1969).
- [24] V. Voevodsky. Reduced power operations in motivic cohomology. *Publ. Math. IHES*, to appear. <http://www.math.uiuc.edu/K-theory/0487>
- [25] V. Voevodsky. On 2-torsion in motivic cohomology. *Publ. Math. IHES*, to appear. <http://www.math.uiuc.edu/K-theory/0502>

DPMMS, UNIVERSITY OF CAMBRIDGE, WILBERFORCE ROAD, CAMBRIDGE CB3
0WB, ENGLAND
B.TOTARO@DPMMS.CAM.AC.UK