

# MUTUAL STATIONARITY AND COMBINATORICS AT $\aleph_\omega$

WILLIAM ADKISSON AND DIMA SINAPOVA

ABSTRACT. Mutual stationarity for  $\langle \kappa_n \mid n < \omega \rangle$  says that for any stationary sequence  $S_n \subset \kappa_n$  and any algebra on  $\sup_n \kappa_n$ , there is a simultaneous witness for stationarity i.e. an elementary substructure  $M$  such that for all  $n$ ,  $\sup(M \cap \kappa_n) \in S_n$ . We prove that mutual stationarity for  $\langle \aleph_n \cap \text{cof}(\omega_k) \mid k < n < \omega \rangle$  is consistent with the tree property at  $\aleph_{\omega+1}$ . Our second theorem is that mutual stationarity for  $\langle \aleph_n \cap \text{cof}(\omega_k) \mid k < n < \omega \rangle$  is consistent with the failure of SCH at  $\aleph_\omega$ . Both theorems use large cardinal hypotheses.

## 1. INTRODUCTION

Stationary sets are a fundamental notion in modern set theory. They are related to elementary substructures by identifying clubs as algebras as follows. For a regular cardinal  $\kappa$  and  $\kappa < \lambda$ ,  $S \subset \kappa$  is stationary iff for every algebra  $\mathfrak{A}$  on  $\lambda$ , there is an elementary  $N \prec \mathfrak{A}$ , such that  $\sup(N \cap \kappa) \in S$ . This notion has an analogue for singular cardinals, called *mutual stationarity*.

Mutual stationarity was introduced in 2001 by Foreman and Magidor in [7], and was used to show the nonsaturation of the nonstationary ideal on  $\mathcal{P}_{\omega_1}(\lambda)$ . Here is the definition:

**Definition 1.1.** Let  $R$  be a set of uncountable regular cardinals and  $S = \langle S_\kappa \mid \kappa \in R \rangle$  be a sequence of stationary sets with  $S_\kappa \subseteq \kappa$ . The sequence  $S$  is mutually stationary if for every algebra  $\mathfrak{A}$  on  $\sup(R)$  there is  $M \prec \mathfrak{A}$  such that  $\sup(M \cap \kappa) \in S_\kappa$  for every  $\kappa \in R \cap M$ .

Suppose now that  $R$  consists of an increasing sequence of cardinals  $\langle \kappa_n \mid n < \omega \rangle$  with limit  $\kappa$ . Given  $A_n \subset \kappa_n$ , we say that **mutual stationarity holds at**  $\langle A_n \mid n < \omega \rangle$  if every sequence of stationary sets  $S_n \subset A_n$  is mutually stationary.

Restricting to countable cofinality, in [7] Foreman and Magidor showed that mutual stationarity holds for  $\langle \kappa_n \cap \text{cof}(\omega) \mid n < \omega \rangle$ . On the other hand, they showed that this result does not generalize to higher fixed cofinality. In particular, in  $L$  there is a sequence of stationary sets  $S_n \subset \aleph_n \cap \text{cof}(\omega_1)$ ,  $n > 1$ , which is not mutually stationary. This prompted the question of whether it is consistent to have mutual stationarity at the  $\aleph_n$ 's for higher fixed cofinality.

Since then there has been a long line of results on this topic. It turns out that mutual stationarity for uncountable cofinality both follows from large cardinals and has large cardinal strength. Here are some highlights:

- (1) If each  $\kappa_n$  is supercompact, then every sequence of stationary sets  $S_n \subset \kappa_n$  is mutually stationary.

---

The second author was partially supported by the National Science Foundation under Grant No. DMS - 1954117.

- (2) (Cummings-Foreman-Magidor) [4] If  $\mathbb{P}$  is the Prikry forcing to singularize  $\kappa$  and  $\langle \kappa_n \mid n < \omega \rangle$  is the corresponding Prikry sequence, then in  $V[\mathbb{P}]$  every sequence of stationary sets  $S_n \subset \kappa_n$  is mutually stationary.
- (3) (Koepke) [9] From a measurable cardinal, one can force mutual stationarity for  $\langle \aleph_{2n+1} \cap \text{cof}(\omega_1) \mid 1 < n < \omega \rangle$ .
- (4) (Koepke-Welch) [10] A measurable cardinal is necessary to obtain mutual stationarity for  $\langle \kappa_n \cap \text{cof}(\omega_1) \mid n < \omega \rangle$ , where  $\langle \kappa_n \mid n < \omega \rangle$  are increasing regular cardinals.

Still, for a long time Foreman and Magidor's original question remained open. Then in 2019, Ben Neria [2] gave a positive answer. More precisely, he showed that from  $\omega$  supercompacts it is consistent that every sequence of stationary sets  $S_n \subseteq \omega_n$  of some fixed cofinality is mutually stationary. His model is obtained by forcing with Levy collapses to make the supercompacts be the  $\aleph_n$ 's. In Ben Neria's model, SCH holds at  $\aleph_\omega$  (and actually GCH is true). Moreover, approachability at  $\aleph_\omega$  holds, and all known models of the tree property at  $\aleph_{\omega+1}$  have failure of approachability at  $\aleph_\omega$ . This raises some natural questions: whether mutual stationarity at the  $\aleph_n$ 's for a fixed uncountable cofinality is consistent with the failure of SCH; and whether it is consistent with the tree property. In this paper we show the answer to both questions is yes.

**Theorem 1.2.** *Suppose that  $\langle \kappa_n \mid n < \omega \rangle$  are  $\nu^+$ -supercompact cardinals, where  $\nu = \sup_n \kappa_n$ . Then for all  $0 < l < \omega$ , there is a forcing extension where for all  $l \neq k < \omega$ , mutual stationarity holds for  $\langle \aleph_n \cap \text{cof}(\omega_k) \mid k < n < \omega \rangle$  and the tree property holds at  $\aleph_{\omega+1}$ .*

**Theorem 1.3.** *Suppose that  $\kappa < \mu < \lambda$  are supercompact cardinals, and let  $0 < l < \omega$ . Then there is a forcing extension where SCH fails at  $\aleph_\omega$  and mutual stationarity holds for  $\langle \aleph_n \cap \text{cof}(\omega_k) \mid k < n < \omega \rangle$  both for all  $k < l$  and for unboundedly many  $k > l$ .*

**Remark 1.4.** The unbounded set of  $k$ 's in the theorem above is  $\omega \setminus \{l+3i \mid i < \omega\}$ .

The first theorem, together with Ben Neria's model, shows that mutual stationarity and the tree property are in a sense orthogonal.

The motivation for the second theorem is that the failure of SCH is an instance of incompleteness, since it requires small powerset below a singular  $\kappa$  and large powerset at  $\kappa$ . In contrast, mutual stationarity can be viewed as a compactness type principle, as it is similar in spirit to stationary reflection and follows from large cardinals. In addition, a corollary of this theorem is that one can reduce the large cardinal assumption of Ben-Neria's result in [2] at the price of losing some cofinalities.

The paper is organized as follows. In section 2 we go over some preliminaries and facts which will be used to prove mutual stationarity. In section 3 we prove Theorem 1.2. Then in section 4 we prove Theorem 1.3.

## 2. OBTAINING MUTUAL STATIONARITY FROM IDEALS

In this section we summarize techniques due to Ben Neria [2] we will use throughout this paper to prove mutual stationarity. For a more detailed exposition of these techniques, see [2, Section 2]. Note that [2] uses the Jerusalem forcing convention, which this paper does not.

**Definition 2.1.** Suppose  $M \prec \mathfrak{A}$ . We call an extension  $N$  of  $M$  an *end-extension* of  $M$  above  $\lambda$  if  $M \prec N \prec \mathfrak{A}$  and  $N \cap \lambda = M \cap \lambda$ .

To show that a sequence is mutually stationary, we will work inductively, starting with  $M_n$  and producing an end-extension  $M_{n+1}$ . The following standard result shows that it is enough to verify mutual stationarity on a tail, so we can start this process at any finite stage  $n$ .

**Fact 2.2.** [7, Lemma 23] Let  $\nu$  be a regular cardinal less than the least element of a set of regular cardinals  $K$ . If  $\{S_\kappa \mid \kappa \in K\}$  is mutually stationary, and for all  $\kappa$ ,  $S_\kappa \subseteq \text{cof}(\leq \nu)$ , then for all  $\lambda_1, \dots, \lambda_n$  greater than  $\nu$  and not in  $K$ , and all sequences of stationary sets  $S_{\lambda_i} \subseteq \lambda_i \cap \text{cof}(\leq \nu)$ , the sequence  $\{S_\kappa \mid \kappa \in K\} \cup \{S_{\lambda_i}, \dots, S_{\lambda_n}\}$  is mutually stationary.

End extensions will be constructed via ideals.

**Definition 2.3.** A nonprincipal  $\kappa$ -complete ideal  $I$  on  $\kappa$  is  $\mu + 1$ -closed if  $I^+$  has a  $\leq_I$ -dense subset  $D$  such that the  $\leq_I \upharpoonright D$  is closed under  $\mu$ -sequences.

An ideal on  $\kappa$  is *nonstationary* if it extends the nonstationary ideal.

**Definition 2.4.** Suppose that  $\mathbb{P}$  is a forcing with  $|\mathbb{P}| \leq \mu$ , and  $G$  is generic for  $\mathbb{P}$  over  $V$ . A nonprincipal  $\kappa$ -complete ideal  $I \in V[G]$  is *formerly  $\mu + 1$ -closed* if  $I$  is the downwards closure of a  $\mu + 1$ -closed ideal  $I_0 \in V$ .

Note that  $\mu$ -closed ideals are formerly  $\mu$ -closed, where  $\mathbb{P}$  is the trivial forcing.

**Definition 2.5.** Let  $\kappa \in M$  be a regular cardinal and  $A \subseteq \kappa$  an unbounded set. We say  $A$  is  $\lambda$ -homogeneous for  $M$  if  $f \upharpoonright A$  is constant for every function  $f : \kappa \rightarrow \lambda$  in  $M$ .  $A$  is  $\lambda$ -homogeneous and approximated if for every function  $f : \kappa \rightarrow \lambda$  in  $M$  there is a set  $A_f \in M$  so that  $A \subseteq A_f \subseteq \kappa$  and  $f \upharpoonright A_f$  is constant.

To build these homogeneous and approximated subsets, we will make use of the following fact.

**Remark 2.6.** [2] Let  $\lambda < \kappa$ . Suppose  $I$  is a  $\kappa$ -complete ideal on  $\kappa$  and  $f : \kappa \rightarrow \lambda$  is a function. Then if  $I, f \in M$  and  $B \in I^+ \cap M$ , there is  $\nu^* < \lambda$  such that  $f^{-1}(\nu^*) \cap B \in I^+$ .

We will also need the following lemma, a slight strengthening of [2, Lemma 2.6].

**Lemma 2.7.** Let  $\mathbb{P}$  be a forcing with  $|\mathbb{P}| \leq \nu$ , and let  $G$  be generic for  $\mathbb{P}$ . In  $V[G]$ , let  $M \prec \mathfrak{A}$  be a  $|M|$ -closed model, with  $|M| = \nu$  and  $\mathbb{P} \subset M$ ,  $\mathbb{P}, G \in M$ . Suppose  $\kappa > \nu$  is a regular cardinal. Suppose  $I \in M$  is a  $\kappa$ -complete formerly  $(\nu + 1)$ -closed ideal on  $\kappa$ , generated by a  $\kappa$ -complete  $(\nu + 1)$ -closed ideal  $I_0$  contained in  $V$ . Then for every  $A \in I^+ \cap M$  and  $\lambda \in M \cap \kappa$ , there exists a subset  $B \subseteq A$  in  $I^+$  which is  $\lambda$ -homogeneous and approximated in  $M$ .

*Proof.* By our assumptions,  $M$  is of the form  $\bar{M}[G]$ , where  $\bar{M} = M \cap V$ .

Let  $<_\theta$  be a definable well-order of  $\mathfrak{A}$ . Let  $D \in \bar{M}$  be a  $\leq_{I_0}$ -dense set of  $I_0^+$  containing lower bounds to all  $\leq_I$ -decreasing sequences of length  $\leq \nu$ . Enumerate all functions from  $\kappa$  to  $\lambda$  in  $M$  by  $\langle f_i \mid i < \nu \rangle$ . While the full sequence may not be in  $M$ , since  $M$  is  $\nu$ -closed, every initial segment will be. In  $V$ , we obtain a corresponding sequence  $\langle \dot{f}_i \mid i < \nu \rangle$  of names, forced by the empty condition to enumerate all such functions.

Working in  $V$ , we wish to build a  $\leq_{I_0}$ -decreasing sequence of sets  $A_i \in D \cap \bar{M}$  such that  $1_{\mathbb{P}} \Vdash \dot{f}_i \restriction A_{i+1}$  has constant value for each  $i < \nu$ . Choose  $A_0 \subset A$  to be some element of  $I_0^+$  contained in  $D \cap \bar{M}$ .

Suppose that we have defined  $A_i$ . First we show the following claim (in  $V$ ).

**Claim 2.8.** *There is  $B \subset A_i$ ,  $B \in I_0^+ \cap D$ , such that  $1_{\mathbb{P}} \Vdash \dot{f}_i \restriction B$  is constant.*

*Proof.* Enumerate the conditions in  $\mathbb{P}$  by  $\langle p_\alpha \mid \alpha < |\mathbb{P}| \rangle$ . We will build a  $\leq_{I_0}$ -decreasing sequence of sets  $\langle A'_\alpha \mid \alpha < |\mathbb{P}| \rangle$  in  $I_0^+ \cap D$ , with  $A'_0 \subset A_i$ , so that for all  $\alpha$ ,  $p_\alpha$  does not force that  $\dot{f}_i \restriction A'_{\alpha+1}$  is not constant. We do so inductively.

At successor stages  $\alpha + 1$ , we note that by Remark 2.6, the empty condition forces that  $\dot{f}_i$  must be constant on some subset of  $A'_\alpha$  that is in  $I_0^+$ .

We claim that any set  $B \in I_0^+$  must have a subset  $A' \subseteq B$  in  $I_0^+$ . Let  $\dot{B}$  be a name for  $B$ . For each  $p \in \mathbb{P}$ , let  $B_p = \{\alpha \mid p \Vdash \alpha \in \dot{B}\}$ ; then  $B = \bigcup_{p \in G} B_p$ . Since  $I$  is  $\kappa$ -complete and  $|\mathbb{P}| < \kappa$ , if every  $B_p$  is in  $I$ ,  $B$  itself would be in  $I$ . We conclude that some  $B_p$  is in  $I^+$ . Since this  $B_p$  is in  $V$ , it must also be in  $I_0^+$ .

So in fact the empty condition forces that  $\dot{f}_i$  is constant on a subset of  $A'_\alpha$  that is in  $I_0^+ \cap D$ . It follows that  $p_\alpha$  cannot force that  $\dot{f}_i$  is nonconstant on every subset of  $A'_\alpha$  in  $I_0^+ \cap D$ , so we can find  $A'_{\alpha+1}$  as desired.

At limit stages  $\gamma$ , let  $A'_\gamma$  be a  $\leq_{I_0}$ -lower bound of  $\langle A'_\alpha \mid \alpha < \gamma \rangle$  contained in  $D$ .

Now that we have built the sequence  $\langle A'_\alpha \mid \alpha < |\mathbb{P}| \rangle$ , we let  $B$  be a  $\leq_{I_0}$ -lower bound of  $\langle A'_\alpha \mid \alpha < |\mathbb{P}| \rangle$  in  $D$ . Then, by construction,  $B$  is as desired.  $\square$

By elementarity,  $\bar{M}$  also satisfies the statement of the claim, so we obtain a set  $A_{i+1} \in I_0^+ \cap D \cap \bar{M}$  as desired.

When  $i$  is a limit ordinal, we let  $A_i$  be the  $<_\theta$ -least  $\leq_{I_0}$ -lower bound of the sequence  $\langle A_j \mid j < i \rangle$ .

Finally, let  $A' \in D$  be the  $<_\theta$ -least  $\leq_{I_0^+}$ -lower bound of the sequence  $\langle A_i \mid i < \nu \rangle$ .

Let us pause to ensure that at each stage our construction is definable in  $\bar{M}$ . We first verify that  $\bar{M}$  is  $\nu$ -closed in  $V$ . Since  $\bar{M} = M \cap V$  and  $M$  is  $\nu$ -closed, it follows that  $\bar{M}^{<\nu} \cap V \subseteq M^{<\nu} \cap V \subseteq M \cap V = \bar{M}$ .

Since the full sequence of names  $\langle \dot{f}_i \mid i < \nu \rangle$  is in  $V$ , and  $\bar{M}$  is  $\nu$ -closed in  $V$ , all initial segments of  $\langle \dot{f}_i \mid i < \nu \rangle$  are in  $\bar{M}$ . The sets  $A_i$  are defined from the sequences  $\langle \dot{f}_j, A_j \mid j < i \rangle$ , the poset  $\mathbb{P}$ , the dense set  $D$ , and the definable well-order  $<_\theta$ . It follows that each  $A_i$  is contained in  $\bar{M}$ , and thus in  $M$ .

By construction, for each  $i < \nu$ , there is some  $B_i \in I_0$  such that the empty condition forces that  $A' \setminus B_i \subseteq A_i$ . In particular, this means that  $\dot{f}_i$  is forced to be constant on  $A' \setminus B_i$ . Let  $B' = \bigcup_{i < \nu} B_i$ ; this is in  $I_0$  by  $\kappa$ -completeness. Then  $A' \setminus B'$  is in  $I_0^+$ . Let  $A'' = A' \setminus B'$ . Since  $B' \in I_0$  and  $A' \leq_{I_0} A$ , we see that  $A'' \in I_0^+$  and thus in  $I^+$ . Moreover,  $A''$  is forced by the empty condition to be  $\lambda$ -homogeneous and approximated in  $V[G]$ .  $\square$

The following lemma follows [2, Proposition 2.12], weakening the closure assumption to former closure.

**Lemma 2.9.** *Suppose  $\mu < \kappa$  are regular cardinals and  $\mathfrak{A}$  is an algebra extending  $\langle H_\theta, \in, <_\theta \rangle$  for some regular cardinal  $\theta > 2^\kappa$ . Let  $M \prec \mathfrak{A}$  be a substructure of size  $\mu$  closed under sequences of size  $< \mu$ , and let  $S \subseteq \kappa \cap \text{cof}(\mu)$  be a stationary subset of  $\kappa$  in  $M$ . Suppose also that at least one of the following holds:*

- (1)  $S$  consists of approachable points<sup>1</sup> or  
 (2) either  $\kappa$  is inaccessible or  $\kappa = \tau^+$  and  $\tau^{<\tau} = \tau$ .

If  $S$  is positive with respect to some nonstationary  $\kappa$ -complete formerly  $(\mu + 1)$ -closed ideal on  $\kappa$ , then for every regular cardinal  $\lambda \in M \cap \kappa$ , there is a  $\mu$ -closed substructure  $N \prec \mathfrak{A}$  of size  $\mu$  which is an end-extension of  $M$  above  $\lambda$  and satisfies  $\sup(N \cap \kappa) \in S$ .

*Proof.* The proof is identical to the proof of [2, Lemma 2.12], using Lemma 2.7 in place of [2, Lemma 2.6].  $\square$

**Remark 2.10.** For the proof in the case of the approachability assumption, see [2, Remark 2.9] for details.

To check that every sequence of stationary sets is mutually stationary, it suffices to show that these hypotheses are satisfied at each stage of the induction. Next, we define a principle that captures the key hypothesis of Lemma 2.9.

**Definition 2.11.** Let  $\nu < \theta$  be uncountable cardinals. We say  $\dagger_\theta^\nu$  holds if for all stationary  $S \subset \theta$ , there is a nonstationary  $\theta$ -complete,  $(\nu + 1)$ -closed ideal, for which  $S$  is a positive set. Given a poset  $\mathbb{Q}$ , we say that  $\dagger_{\theta, \mathbb{Q}}^\nu$  holds if  $1_{\mathbb{Q}}$  forces that for all stationary  $\dot{S} \subset \theta$ , there is a nonstationary  $\theta$ -complete,  $(\nu + 1)$ -closed ideal, for which  $\dot{S}$  is a positive set.

**Lemma 2.12.** Suppose that  $\nu < \kappa$  are cardinals, and  $\dagger_\kappa^\nu$  holds in  $V$ . Let  $\mathbb{P}$  be a poset in  $V$ , such that  $|\mathbb{P}| \leq \nu$  and  $\mathbb{P}$  preserves  $\nu$ , and let  $G$  be generic for  $\mathbb{P}$ . Let  $S$  be a stationary subset of  $\kappa$  in  $V[G]$ . Then in  $V[G]$ , there is a  $\kappa$ -complete nonstationary formerly- $(\nu + 1)$ -closed ideal  $I$  such that  $S \in I^+$ .

*Proof.* Since  $\mathbb{P}$  has size less than  $\kappa$ , there is a generic condition  $p \in G$ , such that  $S_p = \{\alpha \mid p \Vdash_{\mathbb{P}}^\nu \alpha \in \dot{S}\}$  is stationary.

Let  $I_1$  be the ideal given by  $\dagger_\kappa^\nu$  in  $V$  applied to  $S_p$ . Now, going back to  $V[G]$ , let  $I$  be the ideal obtained from  $I_1$ . More precisely,  $I = \{X \subset \kappa \mid \exists \bar{X} \in I_1, X \subset \bar{X}\}$ .

First we show that  $I$  is a  $\kappa$ -complete ideal. Suppose that for some  $\tau < \kappa$ ,  $\langle X_i \mid i < \tau \rangle \in V[G]$  is a sequence of sets in  $I$ . Working in  $V$ , let  $D_i = \{p \in \mathbb{P} \mid \exists \bar{X} \in I_1, p \Vdash \dot{X}_i \subset \bar{X}\}$ . This is a dense subset of  $\mathbb{P}$ . For each  $p \in D_i$ , let  $\bar{X}_{i,p} \in I_1$  witness membership. By  $\kappa$ -completeness of  $I_1$ , we have that  $X := \bigcup_{i,p} \bar{X}_{i,p} \in I_1$ . Since  $\bigcup_i X_i \subset X$ , we get that  $\bigcup_i X_i \in I$ .

Also, since  $S_p \in I_1^+$ , and  $S_p \subset S$ , we have that  $S \in I^+$ . Also if  $A \subset \kappa$  is a nonstationary set in  $V[G]$ , since  $|\mathbb{P}| < \kappa$ , there is a nonstationary  $A_1 \in V$  with  $A \subset A_1$ , and so  $A \in I$ .  $\square$

By the previous two lemmas, to ensure that mutual stationarity holds below  $\aleph_\omega$  for sets of points of cofinality  $\aleph_k$ , it suffices to check that  $\dagger_{\aleph_n}^{\aleph_k}$  holds for cofinitely many  $n < \omega$  (possibly in an inner model by a small forcing) and that all relevant stationary sets are approachable. More precisely:

**Lemma 2.13.** Suppose that  $\mathbb{P}$  is a forcing in  $V$  of size  $\leq \nu$ , such that for some  $k < \omega$ ,  $1_{\mathbb{P}} \Vdash \nu = \aleph_k$ , and let  $G$  be  $\mathbb{P}$ -generic over  $V$ . Suppose that for all large  $n$ ,  $\dagger_{(\aleph_n)^{V[G]}}^\nu$  holds in  $V$ , and that in  $V[G]$ , all stationary sets of  $\aleph_n$  are approachable or *GCH* holds. Then in  $V[G]$ , mutual stationarity holds for  $\langle \aleph_n \cap \text{cof}(\aleph_k) \mid k < n < \omega \rangle$ .

<sup>1</sup>For the definition of approachable points see for example [3, Section 8]

*Proof.* Work in  $V[G]$ . Fix a stationary sequence  $S_n \subset \aleph_n \cap \text{cof}(\aleph_k)$ , for  $n > k$ . Suppose that  $\mathfrak{A}$  is an algebra on  $\aleph_\omega$ . Construct a sequence of elementary substructures of  $\mathfrak{A}$ ,  $\langle M_n \mid k < n < \omega \rangle$  by induction on  $n$ , as follows. Let  $M_{k+1}$  be such that  $\sup(M_{k+1} \cap \aleph_{k+1}) \in S_{k+1}$ . Now, suppose  $n > k+1$  and we have defined  $M_{n-1}$ . Since  $\dot{\upharpoonright}_{\aleph_n}^{\aleph_k}$  held in  $V$ , by Lemma 2.12 there is a nonstationary  $\aleph_n$ -complete formerly  $(\aleph_k + 1)$ -closed ideal  $I$  on  $\aleph_n$  such that  $S_n \in I^+$ . Then by Lemma 2.9, there is an elementary substructure  $M_n$  of  $\mathfrak{A}$ , such that  $M_n$  is an end extension of  $M_{n-1}$  above  $\aleph_{n-1}$  and  $\sup(M_n \cap \aleph_n) \in S_n$ . Finally, let  $M = \bigcup_n M_n \prec \mathfrak{A}$ . Then for all  $n > k$ ,  $\sup(M \cap \aleph_n) = \sup(M_n \cap \aleph_n) \in S_n$ .  $\square$

Ideals as above are obtained from large cardinal embeddings.

**Lemma 2.14.** [2, Fact 2.14] *Let  $j : V \rightarrow M$  be an elementary embedding with  $\text{crit}(j) = \kappa$  and  ${}^\kappa M \subseteq M$ . Let  $\mathbb{P} \in V$  be a poset and let  $G$  be generic for  $\mathbb{P}$ . Suppose that  $j(\mathbb{P})$  projects to  $\mathbb{P}$ , so that every  $j(\mathbb{P})/G$  generic contains  $j''G$ . Working in  $V[G]$ , for every  $\gamma \in j(\kappa) \setminus \kappa$  and  $r \in j(\mathbb{P})/G$ , define an ideal  $I_{\gamma,r}$  by*

$$I_{\gamma,r} = \{\dot{X}_G \mid r \Vdash_{j(\mathbb{P})/G} \gamma \notin j(\dot{X})\}.$$

*Then this ideal is well defined and has the following properties:*

- $I_{\gamma,r}$  is  $\kappa$ -complete and nonprincipal.
- $I_{\gamma,r}$  is nonstationary iff  $r \Vdash \gamma \in j(\dot{C})$  for every  $\mathbb{P}$ -name  $\dot{C}$  for a club subset of  $\kappa$ .
- If  $j(\mathbb{P})/G$  is  $(\mu + 1)$ -closed for some  $\mu < \kappa$ , then  $I_{\gamma,r}$  is a  $(\mu + 1)$ -closed ideal.

*Proof.* We only briefly outline the proof. More details can be found in Foreman's chapter in the handbook [6]; see the interlude to Section 7, "The Basic Idea".

First, note that if  $\dot{X}_G = \dot{X}'_G$ , then some condition in  $j''G$  will force that  $j(\dot{X}) = j(\dot{X}')$ . Since any generic extension by  $j(\mathbb{P})/G$  must contain  $j''G$ , any condition in  $j(\mathbb{P})/G$  will force that  $j(\dot{X}) = j(\dot{X}')$ . It follows that the ideal is well-defined.

$I_{\gamma,r}$  is  $\kappa$ -complete because  $\kappa$  is the critical point of the embedding, and it is non-principal, because  $\gamma \geq \kappa$ . The second assertion of the lemma is clear.

The last claim follows from the fact that  $j(\mathbb{P})/G$  induces a generic for the poset  $(I_{\gamma,r}^+, \leq_{I_{\gamma,r}})$ . For example, if  $j$  is derived from a  $\kappa$ -complete measure  $U$ , one can consider the following projection  $\pi$  from  $j(\mathbb{P})/G$  below  $r$ , to the poset  $(I_{\gamma,r}^+, \leq_{I_{\gamma,r}})$ . Let  $\gamma = [f_\gamma]_U$ ; for  $q = [f_q]_U \in j(\mathbb{P})$ ,  $q \leq r$ , set  $\pi(q) = \{f_\gamma(x) \mid f_q(x) \in G\}$ . Clearly,  $\pi(q)$  is a positive  $I_{\gamma,r}$  set, since  $q$  forces that it is in the dual filter. In particular,  $q \Vdash_{j(\mathbb{P})/G} \gamma \in j(\pi(\dot{q})) := j(\{f_\gamma(x) \mid f_q(x) \in \dot{G}\})$ ; since  $q \leq r$ ,  $r$  certainly can't force  $\gamma$  to not be in  $j(\pi(\dot{q}))$ , so  $\pi(q) \notin I_{\gamma,r}$ . Also, if  $q' \leq q$ , then  $\pi(q') \subset \pi(q)$ , so the map is order preserving.

Finally, we verify that  $\pi$  is indeed a projection. Suppose  $Y = \dot{Y}_G \leq_{I_{\gamma,r}^+} \pi(q)$ . We claim that (in  $V$ ),  $A := \{x \mid f_q(x) \not\Vdash f_\gamma(x) \notin j(\dot{Y})\} \in U$ . If  $A$  is not in  $U$ , then its complement must be, so  $q \Vdash_{j(\mathbb{P})/G} \gamma \notin j(\dot{Y})$ . Note that the empty condition of  $j(\mathbb{P})/G$  forces  $\gamma \in j(\pi(\dot{q})) \Leftrightarrow q \in j(\dot{G})$ . It follows that the empty condition forces  $\gamma \in j(\pi(\dot{q})) \implies \gamma \notin j(\dot{Y})$ . We conclude that  $r \Vdash \gamma \notin j(\dot{Y}) \cap j(\pi(\dot{q}))$ , so  $Y \cap \pi(q) \in I_{\gamma,r}$ . But  $Y \leq_{I_{\gamma,r}^+} \pi(q)$  by assumption, so  $Y \cap \pi(q) \in I_{\gamma,r}^+$ , a contradiction. Since  $A \in U$ , we can define a condition  $q' = [x \rightarrow q'_x]_U \leq q$ , such that for all  $x \in A$ ,  $q'_x \Vdash f_\gamma(x) \in j(\dot{Y})$  and if  $x \notin A$ ,  $q'_x \perp q_x$ . By density, one can find such a condition

in  $j(\mathbb{P})/G$ . Then  $\pi(q') = Y$ . We conclude that  $\pi$  is a projection from  $j(\mathbb{P})/G$  to  $(I_{\gamma,r}^+, \leq_{I_{\gamma,r}})$ , so a generic for  $j(\mathbb{P})/G$  will induce a generic for  $(I_{\gamma,r}^+, \leq_{I_{\gamma,r}})$ .  $\square$

To verify  $\dagger_{\aleph_n}^{\aleph_n}$ , we will need to use embeddings that give sufficiently closed quotients, and we will need to check that the ideals we produce are nonstationary and meet the requisite stationary set. To do so we will use the following lemma, which is implicit in [2].

**Lemma 2.15.** *Let  $\lambda \geq 2^\kappa$ , and let  $j : V \rightarrow M$  be a  $\lambda$ -supercompactness embedding with critical point  $\kappa$ . Suppose  $\mathbb{P}$  is a  $\lambda$ -cc poset with  $(|\mathbb{P}|^{<\lambda})^\kappa = \lambda$ , such that  $\mathbb{P}$  and  $j$  meet the hypotheses of Lemma 2.14, and  $\Vdash_{\mathbb{P}} j(\mathbb{P})/\dot{G}$  is  $(\mu+1)$ -closed. Let  $G$  be generic for  $\mathbb{P}$  over  $V$ . Let  $S \subset \kappa$  be a stationary set in  $V[G]$ . Then there is a condition  $r$  and ordinal  $\gamma$  such that the ideal  $I_{\gamma,r}$  given by Lemma 2.14 is  $(\mu+1)$ -closed and nonstationary, and  $S \in I_{\gamma,r}^+$ .*

*Proof.* Since  $\mathbb{P}$  is  $\lambda$ -cc, we can enumerate (possibly with repetitions) all  $\mathbb{P}$ -names for clubs in  $\kappa$  by  $\vec{C} = \langle \dot{C}_i \mid i < \lambda \rangle$ . Since  $j$  is a  $\lambda$ -supercompactness embedding, this sequence is contained in  $M$ . The sequence  $j''\vec{C} = \langle j(\dot{C}_i) \mid i < \lambda \rangle$  will also be in  $M$ , and is a sequence of  $j(\mathbb{P})$ -names for clubs in  $j(\kappa)$ . It follows that the empty condition of  $j(\mathbb{P})$  forces that  $\dot{C}^* = \bigcap_{i < \lambda} j(\dot{C}_i)$  is a club in  $j(\kappa)$ .

Let  $\dot{S}$  be a  $\mathbb{P}$ -name for  $S$ ;  $\dot{S}$  is forced to be stationary by some condition  $p \in G$ . Then the empty condition of  $j(\mathbb{P})/G$  forces that  $j(\dot{S})$  is stationary in  $j(\kappa)$ . Then there is a condition  $r \in j(\mathbb{P})/G$  and an ordinal  $\gamma \geq \kappa$  such that  $r \Vdash \dot{\gamma} \in j(\dot{S}) \cap \dot{C}^*$ . Let  $I = I_{\gamma,r}$ . By Lemma 2.14, we conclude that  $I$  is  $\kappa$ -complete, nonprincipal, nonstationary, and  $(\mu+1)$ -closed. Since  $r \Vdash \dot{\gamma} \in j(\dot{S})$ , by the definition of  $I_{\gamma,r}$ , we have that  $S \in I^+$  (and actually in the dual filter).  $\square$

We end this section with a lemma about forcings that preserve  $\dagger$ .

**Lemma 2.16.** *Suppose that  $\nu < \kappa$  are cardinals, and  $\dagger_\kappa^\nu$  holds in  $V$ . Let  $\mathbb{Q}$  be a poset in  $V$ , such that  $|\mathbb{Q}| < \kappa$ , and  $\mathbb{Q}$  is  $\leq \nu$ -distributive (i.e. does not add sequences of size  $\nu$ ). Then if  $H$  is  $\mathbb{Q}$ -generic,  $\dagger_\kappa^\nu$  still holds in  $V[H]$ .*

*Proof.* Suppose that  $S \subset \kappa$  is a stationary set in  $V[H]$ . Since  $\mathbb{Q}$  has size less than  $\kappa$ , there is a generic condition  $q \in H$ , such that  $S_q = \{\alpha \mid q \Vdash_{\mathbb{Q}} \alpha \in \dot{S}\}$  is stationary.

Let  $I_1$  be the ideal given by  $\dagger_\kappa^\nu$  in  $V$  applied to  $S_q$ . Now, going back to  $V[H]$ , let  $I$  be the ideal obtained from  $I_1$ . More precisely,  $I = \{X \subset \kappa \mid \exists \bar{X} \in I_1, X \subset \bar{X}\}$ .

First we show that  $I$  is a  $\kappa$ -complete ideal. Suppose that for some  $\tau < \kappa$ ,  $\langle X_i \mid i < \tau \rangle \in V[H]$  is a sequence of sets in  $I$ . Working in  $V$ , let  $D_i = \{q \in \mathbb{Q} \mid \exists \bar{X} \in I_1, q \Vdash \dot{X}_i \subset \bar{X}\}$ . This is a dense subset of  $\mathbb{Q}$ . For each  $q \in D_i$ , let  $\bar{X}_{i,q} \in I_1$  witness membership. By  $\kappa$ -completeness of  $I_1$ , we have that  $X := \bigcup_{i,q} \bar{X}_{i,q} \in I_1$ . Since  $\bigcup_i X_i \subset X$ , we get that  $\bigcup_i X_i \in I$ .

Also, since  $S_q \in I_1^+$ , and  $S_q \subset S$ , we have that  $S \in I^+$ . Also if  $A \subset \kappa$  is a nonstationary set in  $V[H]$ , since  $|\mathbb{Q}| < \kappa$ , there is a nonstationary  $A_1 \in V$  with  $A \subset A_1$ , and so  $A \in I$ .

It remains to show the existence of a closed dense set. Let  $D$  be the  $(\nu+1)$ -closed dense subset of  $I_1^+$ . For the density, we use that if  $X$  is an  $I$ -positive set in  $V[H]$ , by the  $\kappa$ -completeness of  $I_1$  and  $|\mathbb{Q}| < \kappa$ , there is  $p \in H$ , such that  $X_p := \{\alpha \mid p \Vdash \alpha \in \dot{X}\} \in I_1^+$ . Since  $D$  is dense there is  $A \in D$  with  $A \subset X_p \subset X$ . For the closure, suppose that  $\langle Y_i \mid i < \nu \rangle \in V[H]$  is a decreasing sequence of sets

in  $D$ . Since  $\mathbb{Q}$  is  $\leq \nu$ -distributive, this sequence is in  $V$ , and by closure of  $D$ ,  $\bigcap Y_i \in D$ .  $\square$

### 3. MUTUAL STATIONARITY AND THE TREE PROPERTY

To obtain the tree property at  $\aleph_{\omega+1}$  along with mutual stationarity below  $\aleph_\omega$ , we use the arguments of [12, Section 3]. The main complication is that to use these techniques, we cannot determine the cardinal that will become  $\aleph_1$  in advance.

We will use the following lemma to obtain the tree property. The proof is almost identical to [12, Lemma 3.6], except that we allow a finite gap in the definition of  $\mathbb{L}_\mu$ .

**Lemma 3.1.** (*[12, Lemma 3.6]*) *Let  $\langle \kappa_n \mid n < \omega \rangle$  be a strictly increasing sequence of regular cardinals with supremum  $\kappa_\omega$ , and let  $0 < l < \omega$ . Suppose that the following holds:*

- $\kappa_0$  is  $\kappa_\omega^+$ -supercompact.
- For each  $n > 0$ , there is a generic  $\kappa_\omega^+$ -supercompactness embedding with domain  $V$  and critical point  $\kappa_n$ , added by a  $\kappa_{n-1}$ -closed forcing.

*For each strong limit cardinal  $\mu < \kappa_0$  with  $\text{cf}(\mu) = \omega$ , let  $\mathbb{L}_\mu$  be the poset  $\text{Col}(\omega, \mu) \times \text{Col}(\mu^{+l}, < \kappa_0)$ . Then there is  $\mu < \kappa_0$  such that in the extension by  $\mathbb{L}_\mu$ , the tree property holds at  $\kappa_\omega^+$ .*

**Theorem 3.2.** *Let  $\langle \kappa_n \mid n < \omega \rangle$  be an increasing sequence of  $\kappa_\omega^+$ -supercompact cardinals, with supremum  $\kappa_\omega$  and  $0 < l < \omega$ . Then there is a forcing extension in which the tree property holds at  $\aleph_{\omega+1}$  and for all  $l \neq k < \omega$ , mutual stationarity holds for  $\langle \aleph_n \cap \text{cof}(\omega_k) \mid k < n < \omega \rangle$ .*

*Proof.* Let  $\langle \kappa_n \mid n < \omega \rangle$  be an increasing sequence of supercompact cardinals with supremum  $\kappa_\omega$ . Assume also that  $\kappa_0$  is indestructibly supercompact. Let  $\mathbb{H} = \langle \mathbb{H}_n, \dot{\mathbb{H}}(n) \mid n < \omega \rangle$  be the full support iteration where each  $\dot{\mathbb{H}}(n)$  is a  $\mathbb{H}_n$ -name of  $\text{Col}(\kappa_n, < \kappa_{n+1})$ . Let  $H$  be generic for  $\mathbb{H}$ . Note that in  $V[H]$ ,  $\kappa_0$  remains supercompact and  $\kappa_{n+1} = \kappa_n^+$  for all  $n < \omega$ .

Fix  $n < \omega$  and let  $j$  be a  $\nu^+$  supercompact embedding in  $V$  with critical point  $\kappa_n$ . Recall that  $\mathbb{H}$  decomposes into  $\mathbb{H} = \mathbb{H}_{n-1} * \text{Col}(\kappa_{n-1}, < \kappa_n) * (\mathbb{H}/\mathbb{H}_n)$ ;  $\mathbb{H}_{n-1}$  is below the critical point, while  $\text{Col}(\kappa_{n-1}, < \kappa_n) * (\mathbb{H}/\mathbb{H}_n)$  is  $\kappa_{n-1}$ -closed. Note that the poset  $j(\mathbb{H})$  projects to  $\mathbb{H}$ ; this projection is the identity on  $\mathbb{H}_{n-1}$ , and the induced quotient is  $\kappa_{n-1}$ -closed. It follows that for all  $n < \omega$ , in  $V[H]$  there is a generic  $\kappa_\omega^+$ -supercompactness embedding with critical point  $\kappa_n$ , added by a  $\kappa_{n-1}$ -closed forcing.

Applying Lemma 3.1, we see that there exists some strong limit cardinal  $\mu$  with cofinality  $\omega$  so that in the extension of  $V[H]$  by a generic  $L$  for  $\mathbb{L}_\mu := \text{Col}(\omega, \mu) \times \text{Col}(\mu^{+l}, < \kappa_0)$ , the tree property holds at  $\kappa_\omega^+$ . In  $V[H][L]$ ,  $\aleph_{n+l+1} = \kappa_n$  and  $\aleph_\omega = \kappa_\omega$ , so the tree property holds at  $\aleph_{\omega+1}$ . Note that GCH holds in this model below  $\aleph_\omega$ .

**Remark 3.3.** It follows from work of the first author [1] that when  $l = 1$  in the model above, the strong tree property holds at  $\kappa_\omega^+ = \aleph_{\omega+1}$ . Defining  $\mathbb{L}_\mu$  to be  $\text{Col}(\omega, \mu) \times \text{Col}(\mu^{+l}, < \kappa_0)$  for  $l > 1$ , we could even obtain the super tree property.

Now we turn our attention to proving mutual stationarity. Let  $l > 0$ , and  $L$  be generic for  $\mathbb{L}_\mu := \text{Col}(\omega, \mu) \times \text{Col}(\mu^{+l}, < \kappa_0)$  over  $V[H]$ . We claim that in  $V[H][L]$ ,



for all  $k \neq l$  we have mutual stationarity for  $\langle \aleph_n \cap \text{cof}(\aleph_k) \mid k < n < \omega \rangle$ . Fix  $0 < k < \omega$ ,  $k \neq l$ , and denote  $\nu = (\aleph_k)^{V[H][L]}$ . We divide into two cases.

Case 1.  $k > l$ . Then  $\nu \geq \kappa_0$  and since  $|\mathbb{L}_\mu| \leq \nu$  and  $\kappa_n = (\aleph_{n+l+1})^{V[H][L]}$ , by Lemma 2.13, it is enough to show  $\dagger_{\kappa_n}^\nu$  in  $V[H]$  for all large  $n$  in order to get mutual stationarity in  $V[H][L]$ . To that end, let  $n > k$  and in  $V[H]$ , let  $S \subseteq \kappa_n$  be stationary. In  $V$ , let  $j$  be a  $\kappa_{n+1}$ -supercompactness embedding with critical point  $\kappa_n$ . As before,  $j(\mathbb{H})$  projects to  $\mathbb{H}$  with a  $\kappa_{n-1}$ -closed quotient. In particular, we meet the hypotheses of Lemma 2.14.

Now consider the decomposition  $\mathbb{H} = \mathbb{H}_{n+1} * \mathbb{H}/\mathbb{H}_{n+1}$ . Note that the quotient  $\mathbb{H}/\mathbb{H}_{n+1}$  is  $< \kappa_{n+1}$ -distributive, and so  $S$  is a stationary set in  $V[\mathbb{H}_{n+1}]$ . Since  $\mathbb{H}_{n+1}$  is  $\kappa_{n+1}$ -cc, we apply Lemma 2.15 in  $V[\mathbb{H}_{n+1}]$  to conclude that there is some condition  $r \in \mathbb{H}_{n+1}$  and ordinal  $\gamma$  such that in  $V[\mathbb{H}_{n+1}]$ ,  $I_{\gamma,r}$  is nonstationary and  $\nu + 1$ -closed and  $S \in I_{\gamma,r}^+$ . Since the rest of the forcing is  $< \kappa_{n+1}$ -distributive,  $I_{\gamma,r}$  still has the desired properties in the full extension. We have verified that  $\dagger_{\kappa_n}^\nu$  holds, as desired.

Case 2.  $k < l$ . Write  $L = L_0 \times L_1$ , where  $L_0$  is  $\text{Col}(\omega, \mu)$ -generic and  $L_1$  is  $\text{Col}(\mu^{+l}, < \kappa_0)$ -generic. Note that the collapses are computed the same way in  $V$  and in  $V[H]$ . Then  $(\mu^+)^V = (\mu^+)^{V[H][L_1]} = \aleph_1^{V[H][L]} \leq \nu < \mu^{+l}$ . Fix  $n > k$ . Since  $|\text{Col}(\omega, \mu)| < \nu$ , it is enough to show  $\dagger_{\kappa_n}^\nu$  in  $V[H][L_1]$  for all  $n$  in order to get mutual stationarity in  $V[H][L]$ . Again, we verify the hypotheses of Lemmas 2.14 and 2.15.

Let  $n > 0$  and in  $V[H][L_1]$ , let  $S \subseteq \kappa_n$  be stationary. In  $V$ , let  $j$  be a  $\kappa_{n+1}$ -supercompactness embedding with critical point  $\kappa_n$ . As before,  $j(\mathbb{H})$  projects to  $\mathbb{H}$  with a  $\kappa_{n-1}$ -closed quotient. Also, since  $n > 0$ ,  $j(\text{Col}(\mu^{+l}, < \kappa_0)) = \text{Col}(\mu^{+l}, < \kappa_0)$ . So,  $j(\mathbb{H} \times \text{Col}(\mu^{+l}, < \kappa_0))$  projects to  $\mathbb{H} \times \text{Col}(\mu^{+l}, < \kappa_0)$ , and with a  $\mu^{+l}$ -closed quotient. In particular, the quotient is  $(\nu + 1)$ -closed, and we meet the hypotheses of Lemma 2.14.

As before, since the quotient  $\mathbb{H}/\mathbb{H}_{n+1}$  is  $< \kappa_{n+1}$ -distributive,  $S$  is a stationary set in  $V[\text{Col}(\mu^{+l}, < \kappa_0) \times \mathbb{H}_{n+1}]$ . Since  $\text{Col}(\mu^{+l}, < \kappa_0) \times \mathbb{H}_{n+1}$  is  $\kappa_{n+1}$ -cc, we apply Lemma 2.15 in  $V[\text{Col}(\mu^{+l}, < \kappa_0) \times \mathbb{H}_{n+1}]$  to conclude that there is some condition  $r \in \text{Col}(\mu^{+l}, < \kappa_0) \times \mathbb{H}_{n+1}$  and ordinal  $\gamma$  such that in  $V[\text{Col}(\mu^{+l}, < \kappa_0) \times \mathbb{H}_{n+1}]$ ,  $I_{\gamma,r}$  is nonstationary and  $\nu + 1$ -closed and  $S \in I_{\gamma,r}^+$ . Since the rest of the forcing is  $< \kappa_{n+1}$ -distributive,  $I_{\gamma,r}$  still has the desired properties in the full extension. We have verified that  $\dagger_{\kappa_n}^\nu$  holds, completing the proof.  $\square$

#### 4. MUTUAL STATIONARITY AND THE FAILURE OF SCH

Suppose that in  $V_0$ ,  $\kappa < \mu < \lambda$  are all supercompact cardinals, with  $\kappa$  indestructibly supercompact. Let  $H$  be  $\text{Col}(\kappa, < \mu) * \dot{\text{Col}}(\mu, < \lambda) * \dot{\text{Add}}(\kappa, \lambda)$ -generic and let  $V = V_0[H]$ . Fix  $l < \omega$ .

**Lemma 4.1.**<sup>2</sup> *There is a normal measure  $U^* \in V$  on  $\mathcal{P}_\kappa(\lambda^+)$ , such that if  $U_\mu$  is the projected measure to  $\mathcal{P}_\kappa(\mu)$ , then for every  $\gamma < j_{U_\mu}(\kappa)$ ,  $\gamma = j_{U^*}(f)(\kappa)$  for some  $f : \kappa \rightarrow \kappa$ .*

*Proof.* Let  $j_{\lambda^+} : V \rightarrow M^*$  be a  $\lambda^+$ -supercompact embedding with critical point  $\kappa$ . Let  $j_\mu : V \rightarrow M$  be the projected ultrapower to a normal measure on  $\mathcal{P}_\kappa(\mu)$ . Let  $V = \bar{V}[E]$ , where  $E$  is the  $\text{Add}(\kappa, \lambda)$ -generic. Let  $\bar{j}_\mu : \bar{V} \rightarrow \bar{M}$  be the restriction

<sup>2</sup>We do not need this lemma, if we assume a slightly stronger large cardinal hypothesis. See Remark 4.13.

of  $j_\mu$  to  $\bar{V}$  and  $\bar{j}_{\lambda^+} : \bar{V} \rightarrow \bar{M}^*$  be the restriction of  $j_{\lambda^+}$  to  $\bar{V}$ . Since  $|\bar{j}_\mu(\kappa)|^V = |j_\mu(\kappa)|^V = 2^\kappa = \lambda$ , enumerate (in  $V$ ) the interval  $[\kappa, \bar{j}_\mu(\kappa)) = \langle u_\alpha \mid \alpha < \lambda \rangle$ . We can view the  $Add(\kappa, \lambda)$ -generic  $E$  as a function from  $\lambda \times \kappa \rightarrow \kappa$ , and let  $E_\alpha : \kappa \rightarrow \kappa$  be  $E_\alpha(\delta) = E(\alpha, \delta)$ . Let  $E^* = j_{\lambda^+}(E)$ ; a function from  $\bar{j}_{\lambda^+}(\lambda) \times \bar{j}_{\lambda^+}(\kappa) \rightarrow \bar{j}_{\lambda^+}(\kappa)$ . Next we make small changes to  $E^*$  to obtain a generic  $F^*$  for  $\bar{j}_{\lambda^+}(Add(\kappa, \lambda))$ . Set  $F^*$  to be such that for all  $\alpha < \lambda$ ,  $F^*(\bar{j}_{\lambda^+}(\alpha), \kappa) = u_\alpha$ , otherwise  $F^*$  coincides with  $E^*$ . Since the change is captured by a condition,  $F^*$  is still generic, and by construction,  $\bar{j}_{\lambda^+}''E \subset F^*$ . So now we can lift  $\bar{j}_{\lambda^+}$  to  $j'_{\lambda^+} : V = \bar{V}[E] \rightarrow \bar{M}^*[F^*]$ .

**Claim 4.2.** *For every  $u_\alpha$ , there is a function  $f : \kappa \rightarrow \kappa$ , such that  $j'_{\lambda^+}(f)(\kappa) = u_\alpha$ .*

*Proof.* Take  $f = E_\alpha$ . Then  $j'_{\lambda^+}(f)(\kappa) = F^*_{j'_{\lambda^+}(\alpha)}(\kappa) = F^*(j'_{\lambda^+}(\alpha), \kappa) = u_\alpha$ .  $\square$

We make the analogous change to  $j_\mu(E)$  to obtain a generic  $F$  for  $\bar{j}_\mu(Add(\kappa, \lambda))$ , such that for all  $\alpha < \lambda$ ,  $F(\bar{j}_\mu(\alpha), \kappa) = u_\alpha$ . Here although the change is not quite captured by a condition, all of its initial segments are, so we still have that  $F$  is generic. This argument is due to Gitik-Sharon ([8], see Lemma 2.26).

Lift  $\bar{j}_\mu$  to  $j'_\mu$  with respect to  $F$ . As in [5, Section 4.1],  $j'_{\lambda^+}$  is obtained by a normal measure  $U_{\lambda^+}$  on  $\mathcal{P}_\kappa(\lambda^+)$ , and  $j'_\mu$  is obtained from its projection to a normal measure on  $\mathcal{P}_\kappa(\mu)$ .  $\square$

Let  $U_{\lambda^+}$  be the normal measure on  $\mathcal{P}_\kappa(\lambda^+)$  from the above lemma, and let  $U_\lambda$  be its projection to  $\mathcal{P}_\kappa(\lambda)$ . Also let  $U_\mu$  be its projection to  $\mathcal{P}_\kappa(\mu)$  and let  $U$  be its projection to a normal measure on  $\kappa$ . Set  $j_{\lambda^+} := j_{U_{\lambda^+}} : V \rightarrow M_{\lambda^+}$ ,  $j_\lambda := j_{U_\lambda} : V \rightarrow M_\lambda$ ,  $j_\mu := j_{U_\mu} : V \rightarrow M_\mu$ , and  $j := j_U : V \rightarrow M$ .

Let  $k : M \rightarrow M_\mu$  be  $k([f]_U) = j_\mu(f)(\kappa)$ . Then  $j_\mu = k \circ j$ , and by construction each  $u_\alpha$  is in the range of  $k$ . It follows that  $\text{crit}(k) \geq j_\mu(\kappa)$ . And actually, since  $j_\mu(\kappa)$  is also in the range of  $k$ ,  $\text{crit}(k) > j_\mu(\kappa)$ , and so  $j(\kappa) = j_\mu(\kappa)$ .

Similarly, let  $k^* : M \rightarrow M_{\lambda^+}$  be  $k^*([f]_U) = j_{\lambda^+}(f)(\kappa)$ . Then  $j_{\lambda^+} = k^* \circ j$ , and by construction each  $u_\alpha$  is in the range of  $k^*$ . It follows that  $\text{crit}(k^*) \geq j_\mu(\kappa) = j(\kappa)$ . Since  $|j(\kappa)|^V = \lambda < |j_{\lambda^+}(\kappa)|^V = \lambda^{++}$ , we must have  $\text{crit}(k^*) = j(\kappa)$ .

Let  $\mathbb{P}$  be the Prikry forcing with respect to  $U$  with interleaved collapses and guiding generics to make  $\kappa = \aleph_\omega$  and preserve cardinals above  $\kappa$ , with the Prikry sequence starting above  $\omega_l$ . More precisely, conditions in  $\mathbb{P}$  are of the form  $p = \langle d, \alpha_0, c_0, \dots, \alpha_{n-1}, c_{n-1}, A, C \rangle$ , where  $\text{lh}(p) = n$  and:

- (1)  $\langle \alpha_i \mid i < n \rangle$  is an increasing sequence in  $\kappa$  with  $\alpha_0 > \omega_l$ , and  $A \in U$ ;
- (2)  $d \in \text{Col}(\omega_l, < \alpha_0)$  if  $n > 0$ ; otherwise  $d \in \text{Col}(\omega_l, < \kappa)$ .
- (3)  $c_i \in \text{Col}(\alpha_i^{++}, < \alpha_{i+1})$  if  $i < n-1$ , and  $c_{n-1} \in \text{Col}(\alpha_{n-1}^{++}, < \kappa)$ ;
- (4)  $\text{dom}(C) = A$ , for each  $\alpha \in A$ ,  $C(\alpha) \in \text{Col}(\alpha^{++}, < \kappa)$ ,  $[C] \in K$ , where  $K$  is a guiding generic for  $\text{Col}(\kappa^{++}, < j(\kappa))^{U_{\text{lt}}(V, U)}$ .

Let us briefly describe how we get  $K$ . The number of antichains in  $\mathbb{C} := \text{Col}(\kappa^{++}, < j(\kappa))^{U_{\text{lt}}(V, U)}$  is  $\kappa^{++}$ ; enumerate them by  $\langle A_i \mid i < \kappa^{++} \rangle$ . By the high critical point of  $k$ , we have that  $k(\mathbb{C}) = \mathbb{C}$  and for each  $i$ ,  $k(A_i) = k''A_i = A_i$ . So working in  $M_\mu$ , which is closed under sequences of length  $\kappa^+$ , and satisfies that  $\mathbb{C}$  is  $< \kappa^{++}$ -closed, build a decreasing sequence of conditions meeting these antichains. Then use them to define  $K$ .

Let  $G$  be  $\mathbb{P}$ -generic. We have the following standard properties about  $V[G]$ :

- (1)  $\kappa$  is preserved by the Prikry lemma, and becomes  $\aleph_\omega$ .

- (2)  $\mathbb{P}$  has the  $\kappa^+$  chain condition, so cardinals above  $\kappa$  are preserved, and  $2^{\aleph_\omega} = \lambda = \aleph_{\omega+2}$ .
- (3)  $G$  adds a Prikry sequence  $\langle \kappa_n \mid n < \omega \rangle$ , with limit  $\kappa$ , such that for all  $A \in U$ , for all large  $n$ ,  $\kappa_n \in A$ ;
- (4)  $G$  adds a sequence  $\langle c_n^* \mid n < \omega \rangle$  and  $d^*$ , such that for each  $n$ ,  $c_n^*$  is generic for  $Col^V(\kappa_n^{++}, < \kappa_{n+1})$ , and  $d^*$  is generic for  $Col^V(\omega_l, < \kappa_0)$ .

We will show that  $V[G]$  is the desired model for theorem 1.3. Note that when showing mutual stationarity for the  $\aleph_i$ 's, as computed in  $V[G]$ , we only have to worry about cardinals of one of the following three types:  $\kappa_n$ ,  $\kappa_n^+$ , and  $\kappa_n^{++}$  for  $n < \omega$ .

More precisely, we will prove that for all  $k \in \omega \setminus \{l + 3i \mid i < \omega\}$ , mutual stationarity for  $\langle \aleph_n \cap \text{cof}(\aleph_k) \mid k < n < \omega \rangle$  holds in  $V[G]$ . Note that the first Prikry point becomes  $\aleph_{l+1}$  in  $V[G]$ , and for all  $k \leq l$ ,  $(\aleph_k)^V = (\aleph_k)^{V[G]}$ . Since the arguments when  $k > l$  is a little different than when  $k < l$ , we will argue in parallel for the two cases.

To address the former case, for the duration of this section, fix  $\bar{k} > l, \bar{k} \in \omega \setminus \{l + 3i \mid i < \omega\}$ . Then in  $V[G]$ ,  $\aleph_{\bar{k}}$  is either a Prikry point or its successor. Let  $\bar{\nu} < \kappa$  and  $\bar{n} < \omega$  be such that some condition in  $G$  forces that  $\bar{\nu} = \dot{\aleph}_{\bar{k}} \in \{\kappa_{\bar{n}}, \kappa_{\bar{n}}^+\}$ .

Now, set  $W_{\bar{k}} := V[\langle \kappa_n \mid n < \omega \rangle, \langle c_n^* \mid \bar{n} \leq n < \omega \rangle]$ , i.e. the generic extension obtained when we take  $\mathbb{P}$  but omit collapses below the Prikry point  $\kappa_{\bar{n}}$  (and so all the collapses used are  $\bar{\nu}^+$ -closed). Then  $V[G] = W_{\bar{k}}[\langle d^*, c_n^* \mid n < \bar{n} \rangle]$ , i.e.  $V[G]$  is an extension of  $W_{\bar{k}}$  by a forcing of size  $\kappa_{\bar{n}} \leq \bar{\nu}$ . So, by Lemma 2.13, to show the desired mutual stationarity, it is enough to prove the following:

- (1) In  $V[G]$ , for all  $k < l$ ,  $\dagger_{\kappa_n}^{\omega_k}$ ,  $\dagger_{\kappa_n^+}^{\omega_k}$ , and  $\dagger_{\kappa_n^{++}}^{\omega_k}$  hold for all large  $n$ ;
- (2) In  $W_{\bar{k}}$ , the properties  $\dagger_{\kappa_n}^{\bar{\nu}}$ ,  $\dagger_{\kappa_n^+}^{\bar{\nu}}$ , and  $\dagger_{\kappa_n^{++}}^{\bar{\nu}}$  hold for all large  $n$ ;
- (3) All relevant stationary sets consists of approachable points.

**4.1. The Prikry points.** Let  $\nu \in \{\omega_k \mid 0 < k < l\} \cup \{\bar{\nu}\}$ .

**Lemma 4.3.** *In  $V$ , for all regular  $\tau$  with  $\nu < \tau < \kappa$ , we have that  $\dagger_{\kappa, Col(\tau^{++}, < \kappa)}^\nu$  holds. Moreover, there is a measure one set  $A_\tau \in U$ , such that for all  $\alpha \in A_\tau$ ,  $\dagger_{\alpha, Col(\tau^{++}, < \alpha)}^\nu$  holds.*

*Proof.* Note that  $\dagger_{\kappa, Col(\tau^{++}, < \kappa)}^\nu$  asserts the existence of certain ideals on  $\kappa$ , which are subsets of  $2^\kappa$ . We will construct these ideals from the supercompactness of  $\kappa$ , using Lemma 2.14.

Fix  $\tau$ . Recall that  $j_\lambda : V \rightarrow M_\lambda$  is the  $\lambda$ -supercompactness embedding with critical point  $\kappa$ , projecting to  $U$ . I.e.  $U = \{A \mid \kappa \in j_\lambda(A)\}$  is the normal measure used in the definition of the Prikry forcing. We have that there exists a projection from  $j_\lambda(Col(\tau^{++}, < \kappa))$  onto  $Col(\tau^{++}, < \kappa)$  with a  $\tau^{++}$ -closed quotient, so by Lemma 2.14 every ideal  $I_{\gamma, r}$  will be  $(\nu + 1)$ -closed. It remains to verify that for any name for a stationary set  $\dot{S}$ , there is some choice of  $(\gamma, r)$  such that the ideal  $I_{\gamma, r}$  is nonstationary and  $\dot{S}$  is a positive set with respect to this ideal. This follows from Lemma 2.15, noting that  $Col(\tau^{++}, < \kappa)$  is  $\kappa$ -cc. So,  $\dagger_{\kappa, Col(\tau^{++}, < \kappa)}^\nu$  holds in  $V$ .

Since  $M_\lambda^\lambda = M_\lambda^{2^\kappa} \subseteq M_\lambda$ , in  $M_\lambda$ ,  $\dagger_{\kappa, Col(\tau^{++}, < \kappa)}^\nu$  also holds. It follows that for  $U$ -many  $\alpha$ ,  $\dagger_{\alpha, Col(\tau^{++}, < \alpha)}^\nu$  holds in  $V$ .  $\square$

Now, let  $A_\tau$  be given by the above lemma for each  $\tau > \nu$  and set  $A^* = \bigtriangleup_{\tau < \kappa} A_\tau$ . By forcing below  $A^*$ , we may assume that each Prikry point  $\kappa_n \in A^*$ .

**Lemma 4.4.** *For all large  $n$ , in  $V$ ,  $\dagger_{\kappa_n, Col(\kappa_{n-1}^{++}, < \kappa_n)}^\nu$  holds.*

*Proof.* Fix  $n$  such that it is forced that  $\nu < \kappa_{n-1}$ . By choice of  $A^*$ , we have that for all  $\tau$ , for all  $\alpha \in A^* \setminus (\tau + 1)$ ,  $\dagger_{\alpha, Col(\tau^{++}, < \alpha)}^\nu$  holds in  $V$ . In particular, for all  $\alpha \in A^*$  with  $\alpha > \kappa_{n-1}$ ,  $\dagger_{\alpha, Col(\kappa_{n-1}^{++}, < \alpha)}^\nu$  holds in  $V$ . Since  $\kappa_n \in A^*$  with  $\kappa_n > \kappa_{n-1}$ , we have that  $\dagger_{\kappa_n, Col(\kappa_{n-1}^{++}, < \kappa_n)}^\nu$  holds in  $V$ .  $\square$

As a corollary, by definition of  $\dagger$ , we have that:

**Lemma 4.5.** *For all large  $n$ ,  $\dagger_{\kappa_n}^\nu$  holds in  $V[c_{n-1}^*]$ .*

**Lemma 4.6.** *For all  $k < l$ , for all large  $n$ ,  $\dagger_{\kappa_n}^{\omega_k}$  holds in  $V[\langle d^*, c_i^* \mid i < n \rangle]$ . Also, for all large  $n$ ,  $\dagger_{\kappa_n}^{\bar{\nu}}$  holds in  $V[\langle c_i^* \mid \bar{n} \leq i < n \rangle]$ .*

*Proof.* The first statement follows from Lemma 2.16, since the poset to get from  $V[c_{n-1}^*]$  to  $V[\langle d^*, c_i^* \mid i < n \rangle]$  has size  $< \kappa_n$  and is  $\omega_l$ -closed.

The second statement also follows from Lemma 2.16, since the poset to get from  $V[c_{n-1}^*]$  to  $V[\langle c_i^* \mid \bar{n} \leq i < n \rangle]$  has size  $< \kappa_n$  and is  $\kappa_n^{++}$ -closed, and  $\bar{\nu} < \kappa_n^{++}$ . (Because recall that  $\bar{\nu} \in \{\kappa_n, \kappa_n^+\}$ .) So in particular, it is  $\bar{\nu}^+$  closed and we can apply the lemma.  $\square$

**Lemma 4.7.** *In  $V[G]$ , for all  $k < l$ , for all large  $n$ ,  $\dagger_{\kappa_n}^{\omega_k}$  holds. And in  $W_{\bar{k}}$ , for all large  $n$ ,  $\dagger_{\kappa_n}^{\bar{\nu}}$  holds.*

*Proof.* First note that  $V[G]$  projects to  $V[\langle d^*, c_i^* \mid i \leq n \rangle]$  by a quotient that does not add subsets of  $\kappa_{n+1}$  (this is [11, Theorem 3.2]), and  $\dagger_{\kappa_n}^\nu$  is a statement about subsets of  $\mathcal{P}(\kappa_n)$ . So if  $\dagger_{\kappa_n}^\nu$  holds in  $V[\langle d^*, c_i^* \mid i \leq n \rangle]$ , then it also holds in  $V[G]$ . Similarly, each  $W_{\bar{k}}$  projects to  $V[\langle c_i^* \mid \bar{n} \leq i \leq n \rangle]$  by a quotient that does not add subsets of  $\kappa_{n+1}$ , and so if  $\dagger_{\kappa_n}^{\bar{\nu}}$  holds in  $V[\langle c_i^* \mid \bar{n} \leq i \leq n \rangle]$ , then it also holds in  $W_{\bar{k}}$ .

It follows, by the preceding lemma, that in both cases it is enough to show that  $c_n^*$  preserves  $\dagger_{\kappa_n}^\nu$ . Recall that  $c_n^*$  is generic for  $Col(\kappa_n^{++}, < \kappa_{n+1})$ , and let  $W'$  denote some ground model where  $\dagger_{\kappa_n}^\nu$  holds. We claim that  $\dagger_{\kappa_n}^\nu$  still holds in  $W'[c_n^*]$ .

Suppose that  $S \subset \kappa_n$  is a stationary set in  $W'[c_n^*]$ . By closure of  $Col(\kappa_n^{++}, < \kappa_{n+1})$ ,  $S \in W'$ . Let  $I \in W'$  be the nonstationary  $\kappa_n$ -complete,  $\nu + 1$ -closed ideal on  $\kappa_n$ , with  $S \in I^+$ , given by  $\dagger_{\kappa_n}^\nu$  in that model. Since  $Col(\kappa_n^{++}, < \kappa_{n+1})$  does not add new subsets of  $\kappa_n$ ,  $I$  is still a non stationary ideal in the bigger model  $W'[c_n^*]$ . Moreover, since  $Col(\kappa_n^{++}, < \kappa_{n+1})$  is  $\kappa_n^{++}$ -closed,  $I$  is still  $\kappa_n$ -complete and  $\nu + 1$ -closed. So  $I$  is as desired.  $\square$

#### 4.2. The first successors, $\kappa_n^+$ .

**Lemma 4.8.** *In  $V[G]$ , for all  $k < l$ , for all large  $n$ ,  $\dagger_{\kappa_n^+}^{\omega_k}$  holds. And in  $W_{\bar{k}}$ , for all large  $n$ ,  $\dagger_{\kappa_n^+}^{\bar{\nu}}$  holds.*

*Proof.* As before, let  $\nu \in \{\omega_k \mid 0 < k < l\} \cup \{\bar{\nu}\}$ .

**Claim 4.9.** *For all large  $n$ ,  $\dagger_{\kappa_n^+}^\nu$  holds in  $V$ .*

*Proof.* Recall that  $V$  is the extension of  $V_0$  by the poset  $Col(\kappa, < \mu) * Col(\mu, < \lambda) * Add(\kappa, \lambda)$ . Let  $i : V_0 \rightarrow M_0$  be a  $2^\mu = \lambda$ -supercompactness embedding with critical point  $\mu$ . Note that  $i(Col(\kappa, < \mu) * Col(\mu, < \lambda) * Add(\kappa, \lambda))$  absorbs  $Col(\kappa, < \mu) * Col(\mu, < \lambda) * Add(\kappa, \lambda)$  and the quotient is  $\kappa$ -closed.

By Lemmas 2.14 and 2.15, noting that  $Col(\kappa, < \mu) * Col(\mu, < \lambda) * Add(\kappa, \lambda)$  is  $\lambda$ -cc and  $\kappa$ -closed, we conclude that  $\dagger_\mu^\nu$  holds in  $V$ .

Now we use the  $\lambda$ -supercompactness embedding with critical point  $\kappa$ ,  $j_\lambda : V \rightarrow M_\lambda$ . Since  $2^\mu = \lambda$  and  $M_\lambda^\lambda \subset M_\lambda$ , we also have that  $\dagger_\mu^\nu$  (i.e.  $\dagger_{\kappa^+}^\nu$ ) holds in  $M_\lambda$ . Then there is a measure one set  $A \in U$  such that for all  $\alpha \in A$ ,  $\dagger_{\alpha^+}^\nu$  holds in  $V$ . It follows that for all large  $n$ ,  $\kappa_n \in A$ , and so  $\dagger_{\kappa_n^+}^\nu$  holds in  $V$ .  $\square$

**Claim 4.10.** *For all large  $n$ ,  $\dagger_{\kappa_n^+}^\nu$  holds in  $V[c_n^*]$ .*

*Proof.* Let  $n$  be such that  $\dagger_{\kappa_n^+}^\nu$  holds in  $V$ . Suppose that  $S \subset \kappa_n^+$  is a stationary set in  $V[c_n^*]$ . Since  $Col(\kappa_n^{++}, < \kappa_{n+1})$  does not add any subsets of  $\kappa_n^+$ ,  $S$  is a stationary set in  $V$ . Let  $I \in V$  be a nonstationary,  $\kappa_n^+$ -complete,  $(\nu + 1)$ -closed ideal on  $\kappa_n^+$  with  $S \in I^+$ , given by  $\dagger_{\kappa_n^+}^\nu$  in  $V$ . Since  $Col(\kappa_n^{++}, < \kappa_{n+1})$  is  $\kappa_n^{++}$ -closed,  $I$  remains a nonstationary,  $\kappa_n^+$ -complete,  $(\nu + 1)$ -closed ideal in  $V[c_n^*]$ .  $\square$

Next, by Lemma 2.16 and the above claim we get that for  $k < l$ ,  $\dagger_{\kappa_n^+}^{\omega_k}$  holds in  $V[\langle d^*, c_i^* \mid i \leq n \rangle]$ .

Note that  $2^{\kappa_n^+} = \kappa_n^{++}$ . Since the quotient to get to  $V[G]$  from  $V[\langle d^*, c_i^* \mid i \leq n \rangle]$  does not add subsets of  $\kappa_{n+1}$ , we have that  $\dagger_{\kappa_n^+}^{\omega_k}$  holds in  $V[G]$  for  $k < l$ .

Similarly, by Lemma 2.16, we have  $\dagger_{\kappa_n^+}^{\bar{\nu}}$  in  $V[\langle c_i^* \mid \bar{n} \leq i \leq n \rangle]$ . And since the the quotient to get to  $W_{\bar{k}}$  from  $V[\langle c_i^* \mid \bar{n} \leq i \leq n \rangle]$  does not add subsets of  $\kappa_{n+1}$ , we get  $\dagger_{\kappa_n^+}^{\bar{\nu}}$  in  $W_{\bar{k}}$ .  $\square$

**4.3. The second successors,  $\kappa_n^{++}$ .** Again, we set  $\nu \in \{\omega_k \mid 0 < k < l\} \cup \{\bar{\nu}\}$ .

**Lemma 4.11.** *In  $V$ ,  $\dagger_{\lambda, Col(\lambda, < \tau)}^\nu$  holds for all  $\tau > \lambda$ .*

*Proof.* Let  $\tau > \lambda$ . Let  $V'$  be a generic extension of  $V_0$  by  $Col(\kappa, < \mu)$ , and let  $j : V' \rightarrow M$  be a  $\tau$ -supercompact embedding with critical point  $\lambda$ . Since  $j(Col(\mu, < \lambda) * Add(\kappa, \lambda) * Col(\lambda, < \tau))$  projects to  $Col(\mu, < \lambda) * Add(\kappa, \lambda) * Col(\lambda, < \tau)$ , we can lift  $j$  to  $j : V'[Col(\mu, < \lambda) * Add(\kappa, \lambda) * Col(\lambda, < \tau)] \rightarrow M^*$ . Moreover,  $\mathbb{P} := Col(\mu, < \lambda) * Add(\kappa, \lambda) * Col(\lambda, < \tau)$  is  $\tau$ -c.c.

Let  $S$  be stationary in  $V'[Col(\mu, < \lambda) * Add(\kappa, \lambda) * Col(\lambda, < \tau)]$ . By Lemma 2.15 that there is some condition  $r \in j(\mathbb{P})/\mathbb{P}$  and ordinal  $\gamma \in j(\kappa) \setminus \kappa$  so that the ideal  $I_{\gamma, r}$  is nonstationary and  $S \in I_{\gamma, r}^+$ . Note also that the quotient  $j(Col(\mu, < \lambda) * Add(\kappa, \lambda) * Col(\lambda, < \tau)) / (Col(\mu, < \lambda) * Add(\kappa, \lambda) * Col(\lambda, < \tau))$  is  $\kappa$ -closed. Since  $\nu < \kappa$ , from Lemma 2.14, we can conclude that  $I_{\gamma, r}$  is  $\lambda$ -complete, nonprincipal, and  $(\nu + 1)$ -closed.

Since  $V$  is the extension of  $V'$  by  $Col(\mu, < \lambda) * Add(\kappa, \lambda)$ , we conclude that in  $V$ ,  $\dagger_{\lambda, Col(\lambda, < \tau)}^\nu$  holds.  $\square$

**Remark 4.12.** By the same argument as above, we can get  $\dagger_{\lambda, Col(\lambda, < \gamma)}^\nu$  in  $V$  even if  $\gamma$  is not a cardinal. We just have to use a  $|\gamma|^+$ -supercompact embedding with critical point  $\lambda$ .

**Remark 4.13.** Next we will use Lemma 4.1. We note that we do not need it if we assume a slightly stronger large cardinal hypothesis that there is a normal measure on  $\mathcal{P}_\kappa(\lambda)$ , such that for measure one many  $\tau < \kappa$ ,  $\tau$  is  $< j(\kappa)$ -supercompact in the ultrapower.

**Lemma 4.14.** *In  $V[G]$ , for all  $k < l$ , for all large  $n$ ,  $\dagger_{\kappa_n^{++}}^{\omega_k}$  holds. And in  $W_k$ ,  $\dagger_{\kappa_n^{++}}^{\bar{\nu}}$  holds for all large  $n$ .*

*Proof.* As before, it is enough to show that  $\dagger_{\kappa_n^{++}}^{\nu}$  holds in  $V[c_n^*]$ .

Recall that we chose a  $\lambda^+$ -supercompact embedding with critical point  $\kappa$ ,  $j_{\lambda^+} : V \rightarrow M_{\lambda^+}$ , so that the corresponding  $k^* : M \rightarrow M_{\lambda^+}$  has critical point  $j(\kappa)$ . (Here  $M = \text{Ult}(V, U)$  where  $U$  is the projected normal measure on  $\kappa$ , used in the definition of the Prikry forcing).

**Claim 4.15.** *There is a measure one set  $A \in U$  such that for all  $\alpha \in A$  and all  $\tau$  with  $\alpha^{++} < \tau < \kappa$  we have  $\dagger_{\alpha^{++}, \text{Col}(\alpha^{++}, < \tau)}^{\nu}$  holds in  $V$ .*

*Proof.* Let  $\lambda < \gamma < j(\kappa)$ ,  $\gamma$  a cardinal in  $M$ . By Lemma 4.11 and the subsequent remark, we have that  $\dagger_{\lambda, \text{Col}(\lambda, < \gamma)}^{\nu}$  holds in  $V$ . Since  $|\gamma|^V \leq \lambda$ ,  $2^\lambda = \lambda^+$ , and  $(M_{\lambda^+})^{\lambda^+} \subset M_{\lambda^+}$ , we also have that,  $\dagger_{\lambda, \text{Col}(\lambda, < \gamma)}^{\nu}$  holds in  $M_{\lambda^+}$ .

By the high critical point of  $k^*$ ,  $k^*(\gamma) = \gamma$ , so by the elementarity of  $k^*$ ,  $M \models \dagger_{\lambda, \text{Col}(\lambda, < \gamma)}^{\nu}$ .

We have shown that in  $M$ , for all  $\tau$  with  $\lambda < \tau < j(\kappa)$ ,  $\dagger_{\lambda, \text{Col}(\lambda, < \tau)}^{\nu}$  holds. So there is  $A \in U$ , such that for all  $\alpha \in A$ , and all  $\tau$  with  $\alpha^{++} < \tau < \kappa$  we have  $\dagger_{\alpha^{++}, \text{Col}(\alpha^{++}, < \tau)}^{\nu}$  holds in  $V$ .  $\square$

It follows from the claim that for all large  $n$ ,  $V \models \dagger_{\kappa_n^{++}, \text{Col}(\kappa_n^{++}, < \kappa_{n+1})}^{\nu}$ . So for all large  $n$ ,  $\dagger_{\kappa_n^{++}}^{\nu}$  holds in  $V[c_n^*]$ .  $\square$

**4.4. Mutual stationarity in the final model.** We can finally prove the main theorem of the section:

**Theorem 4.16.** *In  $V[G]$ , we have the failure of SCH at  $\aleph_\omega$  and mutual stationarity for  $\langle \aleph_n \cap \text{cof}(\aleph_k) \mid k < n < \omega \rangle$  for every  $k < l$  and for all  $k > l$  with  $k \in \omega \setminus \{l + 3i \mid i < \omega\}$ .*

*Proof.* Clearly SCH at  $\aleph_\omega$  fails. Fix  $k < \omega$ . It is a well-known fact due to Shelah [13] that for all  $n > k + 1$ ,  $\aleph_n \cap \text{cof}(\aleph_k)$  is approachable. Mutual stationarity follows since in  $V[G]$ , for the specified values of  $k$ , we have  $\dagger_{\aleph_n}^{\aleph_k}$  for all large  $n$ .  $\square$

We end with the following open questions:

**Question.** Do the analogues of our two main theorems hold for singular cardinals of uncountable cofinality? In particular, for any countable  $\rho$ , can we obtain mutual stationarity for  $\langle \aleph_\eta \cap \text{cof}(\aleph_{\rho+1}) \mid \rho + 1 < \eta < \omega_1 \rangle$  together with the failure of SCH at  $\aleph_{\omega_1}$ ? What about together with the tree property at  $\aleph_{\omega_1+1}$ ?

**Question.** Can we obtain a model where mutual stationarity for  $\langle \aleph_n \cap \text{cof}(\aleph_k) \mid k < n < \omega \rangle$  holds together with reflection at  $\aleph_{\omega+1}$  and the failure of SCH at  $\aleph_\omega$ ?

## REFERENCES

- [1] William Adkisson. The strong and super tree property at successors of singular cardinals. *The Journal of Symbolic Logic*, 2023.
- [2] Omer Ben-Neria. On singular stationarity I (mutual stationarity and ideal-based methods). *Adv. Math.*, 356:106790, 25, 2019.
- [3] James Cummings. Notes on singular cardinal combinatorics. *Notre Dame J. Formal Logic*, 46(3):251–282, 2005.

- [4] James Cummings, Matthew Foreman, and Menachem Magidor. Canonical structure in the universe of set theory. II. *Ann. Pure Appl. Logic*, 142(1-3):55–75, 2006.
- [5] James Cummings, Yair Hayut, Menachem Magidor, Itay Neeman, Dima Sinapova, and Spencer Unger. The ineffable tree property and failure of the singular cardinals hypothesis. *Trans. Amer. Math. Soc.*, 373(8):5937–5955, 2020.
- [6] Matthew Foreman. Ideals and generic elementary embeddings. In *Handbook of set theory. Vols. 1, 2, 3*, pages 885–1147. Springer, Dordrecht, 2010.
- [7] Matthew Foreman and Menachem Magidor. Mutually stationary sequences of sets and the non-saturation of the non-stationary ideal on  $P_\kappa(\lambda)$ . *Acta Math.*, 186(2):271–300, 2001.
- [8] Moti Gitik and Assaf Sharon. On SCH and the approachability property. *Proc. Amer. Math. Soc.*, 136(1):311–320, 2008.
- [9] Peter Koepke. Forcing a mutual stationarity property in cofinality  $\omega_1$ . *Proc. Amer. Math. Soc.*, 135(5):1523–1533, 2007.
- [10] Peter Koepke and Philip Welch. On the strength of mutual stationarity. In *Set theory*, Trends Math., pages 309–320. Birkhäuser, Basel, 2006.
- [11] Menachem Magidor. On the singular cardinals problem. I. *Israel J. Math.*, 28(1-2):1–31, 1977.
- [12] Itay Neeman. The tree property up to  $\aleph_{\omega+1}$ . *J. Symbolic Logic*, 79(2):429–459, 2014.
- [13] Saharon Shelah. Reflecting stationary sets and successors of singular cardinals. *Arch. Math. Logic*, 31(1):25–53, 1991.