

NEAR-MAXIMA OF THE TWO-DIMENSIONAL DISCRETE GAUSSIAN FREE FIELD

MAREK BISKUP¹, STEPHAN GUFLER² AND OREN LOUIDOR²

¹ *Department of Mathematics, UCLA, Los Angeles, California, USA*

² *Faculty of Industrial Engineering and Management, Technion, Haifa, Israel*

October 26, 2020

Abstract: For the Discrete Gaussian Free Field (DGFF) in domains $D_N \subseteq \mathbb{Z}^2$ arising, via scaling by N , from nice domains $D \subseteq \mathbb{R}^2$, we study the statistics of the values order- $\sqrt{\log N}$ below the absolute maximum. Encoded as a point process on $D \times \mathbb{R}$, the scaled spatial distribution of these near-extremal level sets in D_N and the field values (in units of $\sqrt{\log N}$ below the absolute maximum) tends in law as $N \rightarrow \infty$ to the product of the critical Liouville Quantum Gravity (cLQG) Z^D and the Rayleigh law. The convergence holds jointly with the extremal process, for which Z^D enters as the intensity measure of the limiting Poisson point process, and that of the DGFF itself; the cLQG defined by the limit field then coincides with Z^D . While the limit near-extremal process is measurable with respect to the limit continuum GFF, the limit extremal process is not. In a follow up paper, our results are used to study the fractal properties of the cLQG.

1. INTRODUCTION

Recent years have witnessed considerable progress in the understanding of extremal behavior of logarithmically correlated spatial random processes. A canonical example among these is the two-dimensional Discrete Gaussian Free Field (DGFF). In its \mathbb{Z}^2 -based version, this is a centered Gaussian process $\{h_x^V : x \in V\}$, for non-empty $V \subsetneq \mathbb{Z}^2$, with covariances given by

$$\text{Cov}(h_x^V, h_y^V) = G^V(x, y), \quad x, y \in V, \quad (1.1)$$

where G^V is the Green function of the simple random walk; i.e., with $G^V(x, y)$ given as the expected number of visits to y of the walk started at x before exiting V .

We will focus on the asymptotic properties of h^{D_N} , where $\{D_N\}_{N \geq 1}$ is a sequence of lattice domains that arise, more or less through scaling by N , from a nice continuum domain $D \subseteq \mathbb{R}^2$. The existence of the limiting law of the centered maximum, $\max_{x \in D_N} h_x^{D_N} - m_N$, where the centering sequence is given by

$$m_N := 2\sqrt{g} \log N - \frac{3}{4}\sqrt{g} \log \log(N \vee e) \quad (1.2)$$

© 2020 M. Biskup, S. Gufler, O. Louidor. Reproduction, by any means, of the entire article for non-commercial purposes is permitted without charge.

for g a suitable constant (see (2.2)), was settled in Bramson and Zeitouni [13] and Bramson, Ding and Zeitouni [12]. In [8–10], two of the present authors gave a full description of the *extremal* process cast via the point process convergence

$$\sum_{x \in D_N} \delta_{x/N} \otimes \delta_{h_x^{D_N} - m_N} \xrightarrow[N \rightarrow \infty]{\text{law}} \sum_{i \geq 1} \sum_{z \in \mathbb{Z}^2} \delta_{x_i} \otimes \delta_{h_i - \phi_z^{(i)}}, \quad (1.3)$$

where the process on the right is defined as follows: $\{(x_i, h_i, \phi^{(i)}): i \geq 1\}$ enumerates the sample “points” of the Poisson point process

$$\text{PPP}\left(\bar{c} Z^D(dx) \otimes e^{-\alpha h} dh \otimes \nu(d\phi)\right) \quad (1.4)$$

on $D \times \mathbb{R} \times [0, \infty)^{\mathbb{Z}^2}$, in which $\bar{c} \in (0, \infty)$ is a constant, Z^D is a *random* a.s.-finite Borel measure on D normalized so that

$$\lim_{s \downarrow 0} \frac{E(Z^D(A) e^{-s Z^D(D)})}{\log(1/s)} = \int_A r^D(x)^2 dx, \quad (1.5)$$

where $r^D(x)$ is, for D simply connected, the conformal radius of D from x (see (3.1)), $\alpha := 2/\sqrt{g}$ and ν is a (deterministic) probability measure on $[0, \infty)^{\mathbb{Z}^2}$.

The measure Z^D encodes the macroscopic correlations of the DGFF that “survive” the scaling limit. As it turns out (see [9, Theorem 2.9]), Z^D can be independently characterized as a version of the *critical* Liouville Quantum Gravity (LQG). The LQG was introduced by Duplantier and Sheffield [18] (although the concept goes back to Kahane [21]) in order to give a rigorous meaning to the formal object $1_D(x) e^{\beta h(x)} dx$ for h denoting the continuum Gaussian Free Field (CGFF) on D . Since CGFF exists only in the sense of distributions, the constructions of LQG invariably proceed by regularization. This is particularly subtle in the critical case $\beta = \alpha$, which is the one relevant for the present paper. A number of different regularizations have been considered here (Duplantier, Rhodes, Sheffield and Vargas [16, 17]) which then raises the question of uniqueness; i.e., independence of the regularization used. This was addressed in Rhodes and Vargas [26] and ultimately settled in Junnila and Saksman [20] and Powell [29].

In another paper by two of the present authors [11], the subcritical ($0 < \beta < \alpha$) LQGs have been shown to describe the spatial part of the $N \rightarrow \infty$ limit of random measures

$$\frac{1}{K_N} \sum_{x \in D_N} \delta_{x/N} \otimes \delta_{h_x^{D_N} - 2\sqrt{g}\lambda \log N'} \quad (1.6)$$

where $\lambda := \beta/\alpha$ and $K_N := N^{2(1-\lambda^2)}/\sqrt{\log N}$. This gave a way to quantify the overall growth of the cardinality of the level set $\{x \in D_N: h_x^{D_N} \geq 2\sqrt{g}\lambda \log N'\}$, called “intermediate” in [11] but also known as the “DGFF thick points.” The goal of the present paper is to derive similar control for (what we call) the *near-extremal* level sets

$$\{x \in D_N: h_x^{D_N} \geq m_N - r\sqrt{\log N}\} \quad (1.7)$$

for $r > 0$. We will again glean this from a corresponding near-extremal point process,

$$\zeta_N^D := \frac{1}{\log N} \sum_{x \in D_N} e^{\alpha(h_x^{D_N} - m_N)} \delta_{x/N} \otimes \delta_{(m_N - h_x^{D_N})/\sqrt{\log N}}. \quad (1.8)$$

It is actually known that, modulo normalization, the spatial part of this measure,

$$\frac{1}{\log N} \sum_{x \in D_N} e^{\alpha(h_x^{D_N} - m_N)} \delta_{x/N} \quad (1.9)$$

to be referred to as the *lattice-cLQG measure* below, converges in law to Z^D as $N \rightarrow \infty$ (Rhodes and Vargas [26, Theorem 5.13] and Junnila and Saksman [20, Theorem 1.1]). The measure (1.8) extends (1.9) to include information about the statistics of the field values that carry the mass of the lattice-cLQG measure.

Our main findings are as follows: First, as anticipated in (1.7–1.8), the bulk of the lattice-cLQG measure is carried by the fields where $m_N - h_x = O(\sqrt{\log N})$ with the scaled difference asymptotically Rayleigh distributed. Second, the convergence of the near-extremal process holds *jointly* with that of the extremal process in (1.3) as well as the underlying DGFF. Third, the Z^D -measure arising in the limit of the near-extremal process coincides, in a path-wise sense, with the intensity in (1.4) as well as the cLQG defined by the limiting CGFF. In particular, this shows that, while the limit near-extremal process is a measurable function of the CGFF, the extremal process is not.

2. MAIN RESULT

We now move to the statements of the main theorem which requires introduction of necessary notation and definitions of some basic concepts. Throughout we rely on the notation of references [8–10] and the review [6].

Let \mathcal{D} denote the class of bounded open subsets of \mathbb{R}^2 whose boundary consists of a finite number of connected components each of which has a positive Euclidean diameter. We will approximate D by a sequence of $\{D_N\}_{N \geq 1}$ lattice domains defined by

$$D_N := \{x \in \mathbb{Z}^2 : d_\infty(x/N, D^c) > 1/N\}, \quad (2.1)$$

where d_∞ denotes the ℓ^∞ -distance on \mathbb{Z}^2 . For $A \subseteq \mathbb{Z}^2$, we will write ∂A for the set of vertices in A^c that have a neighbor in A . We will use $\lfloor x \rfloor$ to denote the unique $z \in \mathbb{Z}^2$ such that $x - z \in [0, 1)^2$. The set of all points in \mathbb{R}^2 whose ℓ^∞ -distance to x is less than r will be denoted by $B(x, r)$.

Much of what is to follow depends on a precise control of the Green function in domains of size (order) N . Two specific constants,

$$g := \frac{2}{\pi} \quad \text{and} \quad c_0 := \frac{2\gamma + \log 8}{\pi} \quad (2.2)$$

with γ denoting the Euler constant, appear throughout the derivations in the expansions of the diagonal Green function; see Lemma 3.1. These arise through an asymptotic formula of the potential kernel.

For points in D_N separated by distances of order N , the lattice Green function G^{D_N} is well approximated by its continuum counterpart $\widehat{G}^D : D \times D \rightarrow \mathbb{R} \cup \{+\infty\}$. One way to view \widehat{G}^D is as the integral kernel for the inverse of the (scaled negative) Laplacian $-\frac{1}{4}\Delta$ with Dirichlet boundary conditions on ∂D . (Other representations of \widehat{G}^D exist as well; see (3.2).) Let $C_0(D)$ be the set of continuous functions on \overline{D} that vanish on ∂D .

Definition 2.1 *The Continuum Gaussian Free Field (CGFF) in D is a family of random variables $\{h^D(f) : f \in C_0(D)\}$ such that $f \mapsto h^D(f)$ is linear and, for all $f \in C_0(D)$,*

$$h^D(f) = \mathcal{N}(0, \sigma_f^2) \quad \text{for} \quad \sigma_f^2 := \int_{D \times D} dx dy f(x) f(y) \widehat{G}^D(x, y). \quad (2.3)$$

Regarding $f \mapsto h^D(f)$ as a coordinate map, we may and will henceforth treat h^D as an element of $\mathbb{R}^{C_0(D)}$ endowed with the product topology.

We now move to a definition of the critical LQG measure. We start by a suitable representation of the CGFF. Given a domain $D \in \mathfrak{D}$, let $\{f_k : k \geq 1\}$ be an orthonormal basis (in L^2) of eigenfunctions of $-\frac{1}{4}\Delta$ on D and let λ_k be the eigenvalue corresponding to f_k . As $\lambda_k > 0$ and $f_k(x) = \lambda_k \int_D dy \widehat{G}^D(x, y) f_k(y)$, elementary estimates for \widehat{G}^D ensure that $f_k \in C_0(D)$. It follows that $h^D(f_k)$ is defined for all $k \geq 1$ and obeys $\text{Cov}(h^D(f_k), h^D(f_\ell)) = \delta_{k,\ell} \lambda_k^{-1}$, meaning that

$$\{h^D(f_k) : k \geq 1\} \text{ are independent with } h^D(f_k) = \mathcal{N}(0, \lambda_k^{-1}). \quad (2.4)$$

Moreover, for each $t \in \mathbb{R}$,

$$h_t^D(x) := \sum_{k \geq 1} e^{-\frac{1}{2}e^{-t}\lambda_k} f_k(x) h^D(f_k) \quad (2.5)$$

converges in L^2 and also uniformly in $x \in D$ almost surely, thanks to known estimates on the growth of $k \mapsto \lambda_k$ and the bound $|f_k(x)| \leq \lambda_k [\int_D dy \widehat{G}^D(x, y)^2]^{1/2}$. In particular, $x \mapsto h_t^D(x)$ is continuous a.s. The covariances

$$\text{Cov}(h_t^D(x), h_t^D(y)) = \sum_{k \geq 1} \lambda_k^{-1} e^{-\lambda_k e^{-t}} f_k(x) f_k(y), \quad (2.6)$$

viewed as a kernel of an integral operator, tend to \widehat{G}^D as $t \rightarrow \infty$. Hence, also $h_t^D \xrightarrow{\text{law}} h^D$ as $t \rightarrow \infty$ in the above product topology on $\mathbb{R}^{C_0(D)}$.

The field h_t^D is a measurable function of h^D . Since (unlike h^D) it exists pointwise, we may use it to define the (random) measure

$$Z_t^D(dx) := 1_D(x) \alpha \sqrt{t} e^{\alpha h_t^D(x) - \frac{1}{2} \alpha^2 \text{Var}(h_t^D(x))} r^D(x)^2 dx, \quad (2.7)$$

where $\alpha := 2/\sqrt{g} = \sqrt{2\pi}$ and $r^D(x)$ is the conformal radius of D from x ; see (3.1) for an explicit formula. By [17, Theorem 10] (and the proof of [9, Theorem 2.9] for precise normalization), there exists a random measure Z^D on D normalized as in (1.5) and such that $Z_t^D(A) \rightarrow Z^D(A)$ in probability as $t \rightarrow \infty$ for each measurable $A \subseteq D$.

Definition 2.2 *The critical Liouville Quantum Gravity (cLQG) on D associated with h^D is any version of Z^D that is measurable with respect to h^D .*

Let \overline{D} be the closure of D and denote by $\overline{\mathbb{R}} := \mathbb{R} \cup \{+\infty, -\infty\}$ the two-point compactification of \mathbb{R} . The topology on $\mathbb{R} \cup \{+\infty\}$ is defined so that sets of the form $(a, \infty]$, with $a \in \mathbb{R}$, are compact. We then have:

Theorem 2.3 (Near-critical process convergence) *Let $D \in \mathfrak{D}$ and let $\{D_N\}_{N \geq 1}$ be such that (2.1) holds. Given a sample h^{D_N} of the DGFF in D_N , define ζ_N^D by (1.8) with $\alpha := 2/\sqrt{g} = \sqrt{2\pi}$ and regard h^{D_N} as an element of $\mathbb{R}^{C_0(D)}$ via*

$$h^{D_N}(f) := \int_D dx f(x) h_{\lfloor xN \rfloor}^{D_N}, \quad f \in C_0(D). \quad (2.8)$$

Let η_N^D be the measure on the left of (1.3). Then, relative to the vague topology on the space of Radon measures on $\overline{D} \times \overline{\mathbb{R}}$, resp., $\overline{D} \times (\mathbb{R} \cup \{+\infty\})$ for ζ_N^D , resp., η_N^D and the product topology on $\mathbb{R}^{C_0(D)}$ for h^{D_N} ,

$$(\zeta_N^D, \eta_N^D, h^{D_N}) \xrightarrow[N \rightarrow \infty]{\text{law}} (\zeta^D, \eta^D, h^D), \quad (2.9)$$

where h^D is the CGFF in D and, for Z^D the cLQG measure on D associated with h^D ,

$$\zeta^D = c_* Z^D(dx) \otimes 1_{[0, \infty)}(t) t e^{-t^2/(2g)} dt, \quad (2.10)$$

with $c_* := \frac{1}{2\sqrt{g}} e^{2c_0/g}$ for g and c_0 as in (2.2), while η^D is the measure on the right of (1.3) defined from the above sample of Z^D via (1.4).

To further elucidate the structure of the joint law of the limiting triplet in (2.9), for each non-negative $f_1, f_2 \in C_c(D \times \mathbb{R})$ and arbitrary $f_3 \in C_0(D)$ we have

$$\begin{aligned} & E(e^{-\langle \zeta^D, f_1 \rangle - \langle \eta^D, f_2 \rangle + h^D(f_3)}) \\ &= e^{\frac{1}{2} \langle f_3, \widehat{G}^D f_3 \rangle} E \left(\exp \left\{ -c_* \int_D Z^D(dx) e^{\alpha(\widehat{G}^D f_3)(x)} \int_0^\infty dt t e^{-\frac{t^2}{2g}} f_1(x, t) \right. \right. \\ & \quad \left. \left. - \bar{c} \int_D Z^D(dx) e^{\alpha(\widehat{G}^D f_3)(x)} \int_{\mathbb{R}} dh e^{-\alpha h} \int_{\mathbb{R}^{\mathbb{Z}^2}} \nu(d\phi) (1 - e^{-\sum_{z \in \mathbb{Z}^2} f_2(x, h - \phi_z)}) \right\} \right), \quad (2.11) \end{aligned}$$

where the expectation is with respect to Z^D . Note that the limiting near extremal process ζ^D is a measurable function of the field h^D and that so through Z^D in (2.10). The extremal process η^D contains an additional degree of randomness due to Poisson sampling in (1.4) and is thus *not* measurable with respect to h^D .

Remark 2.4 The above conclusions assume a very specific (albeit canonical) discretization (2.1) of the continuum domain $D \in \mathfrak{D}$. This is stricter than (but included in) the discretizations considered in [8–11] where only

$$D_N \subseteq \{x \in \mathbb{Z}^2 : d_\infty(x/N, D^c) > 1/N\} \quad (2.12)$$

and, for each $\delta > 0$

$$\{x \in \mathbb{Z}^2 : d_\infty(x/N, D^c) > \delta\} \subseteq D_N \quad (2.13)$$

were assumed. As it turns out, the limit in Theorem 2.3 remains in effect provided that the convergence is claimed only for functions compactly supported in D , not \overline{D} . This is because of our inability to control the tightness of the near-extreme process when D_N has a large number of small “holes” near the boundary.

We note that our conclusions are used in a companion paper [7] to control the fractal properties of the cLQG and prove the validity of KPZ relation for the cLQG. We also expect that the control of the near-critical process will be relevant for the study of the law of the most visited site for random walks in planar domains; see [1, 2] for recent results in this direction.

The remainder of this paper is organized as follows. In the next section (Section 3) we recall a number of known facts about the DGFF, its extreme values and connection to the ballot theorems. Section 4 establishes useful conditions for identifying random measures with cLQG. Section 5 is devoted to calculations and estimates of moments of the measure ζ_N^D which are then used, in Section 6, to prove Theorem 2.3.

3. PRELIMINARIES

The exposition of the proofs starts by collecting various preliminary facts about the Green functions, DGFF and its extreme values and the ballot estimates for the DGFF. An uninterested (or otherwise informed) reader may consider skipping this section and returning to it only when the stated facts are used in later proofs.

3.1 Green function.

To address the covariance of the DGFF, we need the large- N behavior of the Green functions G^{D_N} in lattice domains D_N approximating a continuum domain $D \in \mathfrak{D}$ as detailed in (2.1). For this, let $\Pi^D(x, \cdot)$ be the harmonic measure on ∂D that we define using the exit distribution from D of the Brownian motion, $\Pi^D(x, A) := P^x(B_{\tau_{D^c}} \in A)$ where $\tau_{D^c} := \inf\{t \geq 0: B_t \notin D\}$. Writing $\|\cdot\|_2$ for the Euclidean norm on \mathbb{R}^2 , we then define

$$r^D(x) := \exp \left\{ \int_{\partial D} \Pi^D(x, dz) \log \|x - z\|_2 \right\}. \quad (3.1)$$

For D simply connected, this coincides with the notion of conformal radius of D from x defined as the modulus of $f'(0)$ for the conformal map f of the unit disc in \mathbb{C} centered at 0 onto D such that $f(0) = x$.

Closely related is the continuum Green function which is explicitly given by

$$\widehat{G}^D(x, y) := -g \log \|x - y\|_2 + g \int_{\partial D} \Pi^D(x, dz) \log \|y - z\|_2 \quad (3.2)$$

with g as in (2.2). We then have:

Lemma 3.1 (Green function asymptotic) *For all $D \in \mathfrak{D}$, domains $\{D_N\}_{N \geq 1}$ as in (2.12–2.13), all $\delta > 0$ and $D^\delta := \{x \in D: d_\infty(x, D^c) > \delta\}$,*

$$\lim_{N \rightarrow \infty} \sup_{x \in D^\delta} \left| G^{D_N}(\lfloor xN \rfloor, \lfloor xN \rfloor) - g \log N - c_0 - g \log r^D(x) \right| = 0, \quad (3.3)$$

where g and c_0 are the constants from (2.2). In addition, for all $\delta > 0$,

$$\lim_{N \rightarrow \infty} \sup_{\substack{x, y \in D^\delta \\ |x-y| > \delta}} \left| G^{D_N}(\lfloor xN \rfloor, \lfloor yN \rfloor) - \widehat{G}^D(x, y) \right| = 0. \quad (3.4)$$

Proof. This follows from e. g. from Theorem 4.4.4 and Proposition 4.6.2 of [24] in conjunction with [9, Lemma A.1]; see [6, Theorem 1.17] for a full proof. \square

We will also need the following hard estimates:

Lemma 3.2 *For each $D \in \mathfrak{D}$ there is $c = c(D) > 0$ such that for all $N \geq 1$, domains $\{D_N\}_{N \geq 1}$ as in (2.1) and all $x \in D_N$,*

$$g \log d_\infty(x, D_N^c) - c \leq G^{D_N}(x, x) \leq g \log d_\infty(x, D_N^c) + c. \quad (3.5)$$

Proof. Writing \mathfrak{a} for the potential kernel on \mathbb{Z}^2 and $H^U(x, z)$ for the probability that simple random walk started at $x \in U$ exits U at z , the diagonal Green function admits the representation

$$G^{D_N}(x, x) = \sum_{z \in \partial D_N} H^{D_N}(x, z) \mathfrak{a}(x - z). \quad (3.6)$$

Using the constants in (2.2), the potential kernel exhibits the asymptotic

$$\mathfrak{a}(x) = g \log \|x\|_2 + c_0 + O(\|x\|_2^{-2}), \quad \|x\|_2 \rightarrow \infty, \quad (3.7)$$

see, e.g., Lawler [23] or Kozma and Schreiber [22]. The assumptions on the component sizes of D along with our way to discretize D into D_N ensure that there is $\tilde{c} > 0$ such that for all $x \in D_N$ and with $A_k := \{z \in \mathbb{Z}^2 : e^k < d_\infty(x, z) \leq e^{k+1}\}$,

$$\sum_{z \in A_k \cap \partial D_N} H^{D_N}(x, z) \leq e^{-\tilde{c}k}, \quad k \geq \log d_\infty(x, D_N^c). \quad (3.8)$$

Plugging (3.7) into (3.6), the claim follows by a simple estimate. \square

We remark that Lemma 3.2 is the reason why we have to work with the discretization (2.1) instead of (2.12–2.13), for which the upper bound (3.5) fails in general.

3.2 Gibbs-Markov property.

One of the important properties of the DGFF is the behavior under the restriction to a subdomain. As this arises directly from the Gibbsian structure of the law of the DGFF and is similar to the Markov property for stochastic processes, we refer to this as the Gibbs-Markov property, although the term domain-Markov has been used as well.

Lemma 3.3 (Gibbs-Markov property) *Let $\emptyset \neq U \subseteq V \subsetneq \mathbb{Z}^2$ and let h^V be the DGFF in V . Define*

$$\varphi^{V,U}(x) := \mathbb{E}(h_x^V \mid \sigma(h_y^V : y \in V \setminus U)). \quad (3.9)$$

Then

$$h^V - \varphi^{V,U} \perp\!\!\!\perp \varphi^{V,U} \quad \text{and} \quad h^V - \varphi^{V,U} \stackrel{\text{law}}{=} h^U. \quad (3.10)$$

Moreover, a.e. sample of $\varphi^{V,U}$ is discrete harmonic on U .

Proof. See [6, Lemma 3.1]. \square

Consider now two sequences $\{D_N\}_{N \geq 1}$, resp., $\{\tilde{D}_N\}_{N \geq 1}$ of lattice domains related, via (2.1), to continuum domains D , resp., \tilde{D} subject to $\tilde{D} \subseteq D$. While the DGFFs h^{D_N} ,

resp., $h^{\tilde{D}_N}$ do not allow for pointwise limits, the field $\varphi^{D_N, \tilde{D}_N}$ does, due to controlled behavior of the variances:

Lemma 3.4 *Let $D, \tilde{D} \in \mathfrak{D}$ obey $\tilde{D} \subseteq D$. Define*

$$C^{D, \tilde{D}}(x, y) := g \int_{\partial D} \Pi^D(x, dz) \log \|y - z\|_2 - g \int_{\partial \tilde{D}} \Pi^{\tilde{D}}(x, dz) \log \|y - z\|_2. \quad (3.11)$$

Then $x, y \mapsto C^{D, \tilde{D}}(x, y)$, for $x, y \in \tilde{D}$, is symmetric, positive semi-definite, continuous and harmonic in each variable. There exists a centered Gaussian process $\Phi^{D, \tilde{D}} = \mathcal{N}(0, C^{D, \tilde{D}})$ with continuous and harmonic sample paths a.s.

Proof. See [9, Lemma 2.3]. □

The equicontinuity of the sample paths implied by harmonicity permits a locally-uniform coupling of the discrete and continuum processes:

Lemma 3.5 *Assume the setting of Lemma 3.4 and let $\{D_N\}_{N \geq 1}$, resp., $\{\tilde{D}_N\}_{N \geq 1}$ be lattice domains approximating D , resp., \tilde{D} in the sense (2.12–2.13). Suppose $\tilde{D}_N \subseteq D_N$ for all $N \geq 1$. Then for each $N \geq 1$ there is a coupling of $\varphi^{D_N, \tilde{D}_N}$ and $\Phi^{D, \tilde{D}}$ such that, for each $\delta > 0$,*

$$\sup_{x \in \tilde{D}^\delta} \left| \varphi^{D_N, \tilde{D}_N}(\lfloor xN \rfloor) - \Phi^{D, \tilde{D}}(x) \right| \longrightarrow 0 \quad (3.12)$$

in probability as $N \rightarrow \infty$, where $\tilde{D}^\delta := \{x \in \tilde{D} : \text{dist}(x, \tilde{D}^c) > \delta\}$.

Proof. See [9, Appendix] or [6, Lemma 4.4]. □

Another property we will use in some arguments of the proof deal with positive correlations. The following is a direct consequence of the positivity of the Green function:

Lemma 3.6 (strong FKG for DGFF) *Let $U \subsetneq \mathbb{Z}^2$. The law \mathbb{P} of DGFF h^U on \mathbb{R}^U is then strong FKG in the sense that, for any $V \subseteq U$ and any $f, g \in L^2(\mathbb{P})$ that admit a version which is non-decreasing in each coordinate,*

$$\mathbb{E}(f(h^U)g(h^U) \mid \mathcal{F}_V) \geq \mathbb{E}(f(h^U) \mid \mathcal{F}_V)\mathbb{E}(g(h^U) \mid \mathcal{F}_V) \quad (3.13)$$

where $\mathcal{F}_V := \sigma(h_x^U : x \in V)$. Moreover,

$$h \mapsto \mathbb{E}(f(h^U) \mid \mathcal{F}_V)(h) \quad (3.14)$$

admits a version that is non-decreasing in every coordinate.

Proof. For multivariate Gaussians, positivity of correlations implies the validity of the strong FKG condition; cf [6, Proposition 5.22]. The monotonicity (3.14) is validated easily using the Gibbs-Markov property. □

3.3 Ballot estimates.

Our proofs of the near-critical process convergence rely heavily on computations involving the tail asymptotic for the DGFF maximum. A starting point is the precise bound (to within a multiplicative constant) on the upper tail:

Lemma 3.7 *There exists a constant $c < \infty$ such that for all $N \geq 1$ and all $u \in [1, \sqrt{\log N})$,*

$$\mathbb{P}(\max_{x \in D_N} h_x^{D_N} > m_N + u) \leq cue^{-\alpha u}. \quad (3.15)$$

Proof. For $D_N := (0, N)^2 \cap \mathbb{Z}^2$, this is [15, Theorem 1.4]. A routine extension to general domains is performed in [9, Proposition 3.3]. \square

In addition, we will need to control the tail probability even under conditioning on the field at a vertex to be fairly large. We begin with an asymptotic statement:

Lemma 3.8 *There exists a function $\mathcal{R}: [0, \infty) \rightarrow (0, \infty)$ and, for each $x \in D$, a function $\mathcal{L}^x: [0, \infty) \rightarrow (0, \infty)$ such that for all $\delta > 0$, uniformly in $x \in D^\delta$,*

$$\lim_{s \rightarrow \infty} \frac{\mathcal{L}^x(s)}{s} = \lim_{s \rightarrow \infty} \frac{\mathcal{R}(s)}{s} = 1 \quad (3.16)$$

and such that for all $x \in D$ and all $u \geq 1$, the quantity $o(1)$ defined by

$$\mathbb{P}(h^{D_N} \leq m_N + u \mid h_{\lfloor Nx \rfloor}^{D_N} = m_N - s) = [2 + o(1)] \frac{\mathcal{L}^x(u) \mathcal{R}(u + s)}{g \log N} \quad (3.17)$$

tends to zero as $N \rightarrow \infty$ uniformly in $s \in [0, (\log N)^{1-\delta}]$.

Proof. The proof draws heavily on the conclusions of the technical paper [19] by two of the present authors so we give only the main steps. Given $N \in \mathbb{N}$ and $x \in D$, denote $D^{(N)} := -\lfloor Nx \rfloor / N + D$. Then $D^{(N)} \in \mathfrak{D}$ and $D_N^{(N)} = -\lfloor Nx \rfloor + D_N$ and so the left-hand side of (3.17) equals

$$\mathbb{P}(h^{D_N^{(N)}} - m_N - u \leq 0 \mid h_0^{D_N^{(N)}} - m_N - u = -s - u). \quad (3.18)$$

By Theorem 1.1 in [19], this probability is asymptotically equivalent to

$$2\mathcal{L}_{\log N, 0, D^{(N)}}(-u 1_{\partial D_N^{(N)}}) \mathcal{R}_{0,0,B(0,1/2)}(-u - s) (g \log N)^{-1} \quad (3.19)$$

in the limit as $N \rightarrow \infty$, uniformly in $s \in [0, (\log N)^{1-\delta}]$ for any fixed $u \geq 1$. Here the functions on the right-hand side are written in the notation of [19] with the discretization of the continuum domain $D^{(N)}$ for $\mathcal{L}_{\log N, 0, D^{(N)}}$ chosen as in the present paper (see [19, Remark 2]), while that for $\mathcal{R}_{0,0,B(0,1/2)}$ as in (1.3) of [19].

By the continuity of the \mathcal{L} -functional in the underlying domain (see [19, Proposition 1.6]), the quantity $\mathcal{L}_{\log N, 0, D^{(N)}}(-u 1_{\partial D_N^{(N)}})$ has the same $N \rightarrow \infty$ asymptotic as its continuum analogue $\mathcal{L}_{\log N, 0, -x+D}(-u 1_{\partial(-x+D)_N})$ which by [19, Proposition 1.3] converges, as $N \rightarrow \infty$, to some $\mathcal{L}^x(u)$. Setting $\mathcal{R}(s) := \mathcal{R}_{0,0,B(0,1/2)}(-s)$, we arrive at (3.17). Proposition 1.4 of [19] yields (3.16). \square

We remark that asymptotic expressions of the form (3.17) lie at the heart of [10] where they were derived using the so called concentric decomposition. In [6], the concentric decomposition was used to give alternative proofs of tightness and convergence in law of the centered DGFF maximum that avoid the use of the modified branching random walk which the earlier proofs by Bramson and Zeitouni [13] and Bramson, Ding and Zeitouni [12] relied on. (A key additional input on top of [10] was the first limit in (3.16);

see [6, Proposition 12.5].) Notwithstanding, Lemma 3.8 is stronger than any of these prior conclusions; e.g., compared with [10, Proposition 5.2]. Indeed, there $-s = u$ and s has to be confined to a finite interval.

In [19] the concentric decomposition has been developed further to yield uniform asymptotics as well as bounds for various ‘‘ballot’’ probabilities in more general setups. This includes the case where the field is conditioned on a subset of its domain which is not necessarily a single point, as needed in the next lemma:

Lemma 3.9 *There is $\delta_0 \in (0, 1)$ and, for each $\delta \in (0, \delta_0)$, there is $c = c(\delta) \in (0, \infty)$ such that the following holds for all $D, \tilde{D} \in \mathfrak{D}$ with D connected and $B(0, \delta) \subseteq D \cap \tilde{D} \subseteq B(0, 1/\delta)$: For all $1 \leq K \leq \frac{1}{3}\delta^2 N$, all $u \in \mathbb{R}^{\partial D_N}$ and all $v \in \mathbb{R}^{\partial(\tilde{D}^c)_K}$, abbreviating $U := D_N \cap (\tilde{D}^c)_K$,*

$$\begin{aligned} \mathbb{P}\left(h^U \Big|_{D_N^{\delta} \cap (\tilde{D}^c)_K^{\delta}} \leq 0 \mid h^U \Big|_{\partial D_N} = -m_N + u, h^U \Big|_{\partial(\tilde{D}^c)_K} = -m_K + v\right) \\ \leq c \frac{(1 + u_{\star}^{-}) (e^{-v_{\star}^{+}})^{3/21} \mathbf{1}_{\{u_{\star}^{-} \leq \sqrt{\log N/K}\}} + v_{\star}^{-}}{\log(N/K)}, \end{aligned} \quad (3.20)$$

where

$$u_{\star} := \min_{x \in D_N^{\delta}} \bar{u}(x) \quad \text{and} \quad v_{\star} := \min_{x \in (\tilde{D}^c)_K^{\delta}} \bar{v}(x) \quad (3.21)$$

for \bar{u} , resp., \bar{v} denoting the (unique) bounded harmonic extensions of u , resp., v onto D_N , resp., $(\tilde{D}^c)_K$. In addition, for all $u \geq 0$, all $v \in \mathbb{R}$, all $x \in D_N^{\delta}$ and all $N \geq 1$,

$$\mathbb{P}\left(h^{D_N} \leq m_N + u \mid h_x^{D_N} = m_N + v\right) \leq c \frac{(1 + u)(1 + (u - v)^+)}{\log N}. \quad (3.22)$$

Proof. This follows immediately from Theorem 1.8 of [19] and Remark 2 therein. \square

4. CHARACTERIZATIONS OF CLQG MEASURE

In order to identify the measure arising in the limit (2.9) with the cLQG measure constructed by the limit of the measures in (2.7), we will also need conditions that characterize the cLQG measure uniquely. For subcritical LQG measures, such conditions have been found by Shamov [30], but the argument does not extend to the critical case. We will thus rely on [9, Theorem 2.8] in the following specific form:

Theorem 4.1 *Suppose $\{M^D : D \in \mathfrak{D}\}$ is a family of random Borel measures (not necessarily realized on the same probability space) that obey:*

- (1) M^D is concentrated on D and $M^D(D) < \infty$ a.s.
- (2) $P(M^D(A) > 0) = 0$ for any Borel $A \subseteq D$ with $\text{Leb}(A) = 0$.
- (3) For any $a \in \mathbb{R}^2$,

$$M^{a+D}(a + dx) \stackrel{\text{law}}{=} M^D(dx). \quad (4.1)$$

- (4) If $D, \tilde{D} \in \mathfrak{D}$ are disjoint then

$$M^{D \cup \tilde{D}}(dx) \stackrel{\text{law}}{=} M^D(dx) + M^{\tilde{D}}(dx), \quad (4.2)$$

with M^D and $M^{\tilde{D}}$ on the right regarded as independent. If $D, \tilde{D} \in \mathfrak{D}$ instead obey $\tilde{D} \subseteq D$ and $\text{Leb}(D \setminus \tilde{D}) = 0$, then (for $\alpha := 2/\sqrt{g}$)

$$M^D(\mathrm{d}x) \stackrel{\text{law}}{=} e^{\alpha\Phi^{D,\tilde{D}}(x)} M^{\tilde{D}}(\mathrm{d}x), \quad (4.3)$$

where $\Phi^{D,\tilde{D}}$ is a centered Gaussian field with covariance $C^{D,\tilde{D}}$, independent of $M^{\tilde{D}}$.

(5) There is $c \in (0, \infty)$ such that for every open A with $\bar{A} \subseteq D$ and $\text{Leb}(\partial A) = 0$, we have

$$\lim_{\lambda \downarrow 0} \sup_{K \geq 1} \left| \frac{K^4 E(M^{K^{-1}D}(K^{-1}A) e^{-\lambda K^4 M^{K^{-1}D}(K^{-1}D)})}{\log(1/\lambda)} - c \int_A r^D(x)^2 \mathrm{d}x \right| = 0, \quad (4.4)$$

where $r^D(x)$ is as in (3.1).

Then

$$M^D \stackrel{\text{law}}{=} cZ^D, \quad D \in \mathfrak{D}, \quad (4.5)$$

where Z^D denotes cLQG measure on D .

Proof. With (3) replaced by the requirement of shift and dilation covariance; namely,

$$M^{a+\lambda D}(a + \lambda \mathrm{d}x) \stackrel{\text{law}}{=} \lambda^4 M^D(\mathrm{d}x), \quad a \in \mathbb{R}^2, \lambda > 0, \quad (4.6)$$

and (5) weakened to just the case of $K := 1$ in (4.4), i.e.,

$$\lim_{\lambda \downarrow 0} \frac{E(M^D(A) e^{-\lambda M^D(D)})}{\log(1/\lambda)} = c \int_A r^D(x)^2 \mathrm{d}x, \quad (4.7)$$

this becomes an exact restatement of [9, Theorem 2.8] (modulo the specific normalization of Z^D). That dilation covariance may be dropped at the expense of imposing the uniform bound (4.4) is justified in [9, Remark 6.9]. \square

As it turns out, not only does Theorem 4.1 determine the law of cLQG measures uniquely but it also offers simple criteria that identify two such families of measures in path-wise sense. This is the content of:

Theorem 4.2 *Let $\{M^D : D \in \mathfrak{D}\}$, resp., $\{\widehat{M}^D : D \in \mathfrak{D}\}$ be two families of random Borel measures satisfying conditions (1-3) and (4.2) in Theorem 4.1 and such that (4.4) holds with a constant $c > 0$ for $\{M^D : D \in \mathfrak{D}\}$ and a constant $\hat{c} > 0$ for $\{\widehat{M}^D : D \in \mathfrak{D}\}$. Assume that M^D and \widehat{M}^D are defined, for each $D \in \mathfrak{D}$, on the same probability space so that (4.3) holds jointly for both measures, i.e., for all $D, \tilde{D} \in \mathfrak{D}$ with $\tilde{D} \subseteq D$ and $\text{Leb}(D \setminus \tilde{D}) = 0$, we have*

$$(M^D(\mathrm{d}x), \widehat{M}^D(\mathrm{d}x)) \stackrel{\text{law}}{=} \left(e^{\alpha\Phi^{D,\tilde{D}}(x)} M^{\tilde{D}}(\mathrm{d}x), e^{\alpha\Phi^{D,\tilde{D}}(x)} \widehat{M}^{\tilde{D}}(\mathrm{d}x) \right), \quad (4.8)$$

where $\Phi^{D,\tilde{D}}$ is independent of $(M^{\tilde{D}}, \widehat{M}^{\tilde{D}})$ on the right-hand side. Then

$$c^{-1}M^D = \hat{c}^{-1}\widehat{M}^D \quad \text{a.s.} \quad (4.9)$$

for each $D \in \mathfrak{D}$.

Proof. The proof could in principle be referred to [9, Theorem 6.1] which states that any measure M^D satisfying conditions (1-5) can be represented, by sequential application of the Gibbs-Markov property (4.3), as a weak limit of approximating measures that depend only on the binding fields $\Phi^{D, \tilde{D}}$ used in (4.3). We will instead provide an independent argument that only uses the conclusion of Theorem 4.1.

Given $p \in (0, 1)$, consider the random measure

$$\tilde{M}_p^D := pM^D + (1-p)\hat{M}^D. \quad (4.10)$$

By our assumptions, the family $\{\tilde{M}_p^D : D \in \mathfrak{D}\}$ satisfies conditions (1-4) of Theorem 4.1. We claim that it satisfies also condition (5) with constant

$$c_p := pc + (1-p)\hat{c}. \quad (4.11)$$

For this pick $K \geq 1$, abbreviate

$$X(\cdot) := K^4 M^{K^{-1}D}(K^{-1}\cdot) \quad \text{and} \quad Y(\cdot) := K^4 \hat{M}^{K^{-1}D}(K^{-1}\cdot) \quad (4.12)$$

and consider the rewrite

$$\begin{aligned} E\left(K^4 \tilde{M}_p^{K^{-1}D}(K^{-1}A) e^{-\lambda K^4 \tilde{M}_p^{K^{-1}D}(K^{-1}D)}\right) = \\ pE(X(A)e^{-\lambda pX(D)}) + (1-p)E(Y(A)e^{-\lambda(1-p)Y(D)}) \\ - pE\left(X(A)e^{-\lambda pX(D)}(1 - e^{-\lambda(1-p)Y(D)})\right) \\ - (1-p)E\left(Y(A)e^{-\lambda(1-p)Y(D)}(1 - e^{-\lambda pX(D)})\right). \end{aligned} \quad (4.13)$$

Invoking (4.4) for the two families of measures, as $\lambda \downarrow 0$ (with p fixed), the sum of the first two terms on the right divided by $\log(1/\lambda)$ tends to $c_p \int_A r^D(x)^2 dx$, uniformly in $K \geq 1$. We claim that the two remaining terms are $o(\log(1/\lambda))$.

Focusing only on the third term by symmetry, the Cauchy-Schwarz inequality bounds this by the square root of

$$E\left(X(A)^2 e^{-2\lambda pX(D)}\right) E\left((1 - e^{-\lambda(1-p)Y(D)})^2\right). \quad (4.14)$$

Since Theorem 4.1 applied to X and Y separately ensures that X and Y have the (marginal) law of $K^4 Z^{K^{-1}D}$ which has the law of Z^D by exact scaling properties of Z^D under dilation (cf [9, Corollary 2.2]), the quantities in (4.14) are independent of $K \geq 1$. Dominating $X(A)^2 e^{-2\lambda pX(D)} \leq \frac{1}{\lambda p} X(D) e^{-\lambda pX(D)}$ shows, with the help of [9, Corollary 2.7] that the first term in (4.14) is $O(\lambda^{-1} \log(1/\lambda))$. From the known properties of Z^D (see again [9, Corollary 2.7]) we also get that the limit

$$\lim_{\lambda \downarrow 0} \frac{1}{\lambda \log(1/\lambda)} E(1 - e^{-\lambda Y(D)}) \quad (4.15)$$

exists and is finite. Then

$$E\left((1 - e^{-\lambda Y(D)})^2\right) = 2E(1 - e^{-\lambda Y(D)}) - E(1 - e^{-2\lambda Y(D)}) \quad (4.16)$$

shows that the second term in (4.14) is $o(\lambda \log(1/\lambda))$ as $\lambda \downarrow 0$. Hence, the quantity in (4.14) is $o(\log(1/\lambda)^2)$ and so the third (and by symmetry also the fourth) term on the right of (4.13) are $o(\log(1/\lambda))$, uniformly in $K \geq 1$.

Having verified the conditions of Theorem 4.1, we thus get that for each $p \in [0, 1]$ and each $D \in \mathfrak{D}$,

$$pM^D + (1-p)\widehat{M}^D \stackrel{\text{law}}{=} (pc + (1-p)\hat{c})Z^D \quad (4.17)$$

Pick non-negative $f \in C_c(\overline{D})$ and abbreviate $U := c^{-1}\langle M^D, f \rangle$, $T := \hat{c}^{-1}\langle \widehat{M}^D, f \rangle$ and $q := pc/c_p$. By (4.17) the law of the non-negative random variable $qU + (1-q)T$ does not depend on $q \in [0, 1]$. This implies $Ee^{-(qU+(1-q)T)} = qE(e^{-U}) + (1-q)E(e^{-T})$ and so, by Jensen's inequality, $e^{-(qU+(1-q)T)} = qe^{-U} + (1-q)e^{-T}$ a.s. By Jensen's inequality again, this is only possible if $U = T$ a.s. As this holds for all non-negative $f \in C_c(\overline{D})$, we get (4.9) as desired. \square

As a consequence of the above, we do get a characterization of cLQG by its behavior under the Cameron-Martin shifts of the underlying CGFF:

Theorem 4.3 *Let $\{M^D : D \in \mathfrak{D}\}$ be a family of random Radon measures satisfying the conditions of Theorem 4.1 with $c := 1$ in (4.4). Suppose also that, for each $D \in \mathfrak{D}$, there exists a coupling of M^D with the CGFF h^D such that for all $f \in C_0(D)$ and for \mathbb{P}_f defined by*

$$\mathbb{P}_f(A) := \mathbb{E}(e^{h^D(f) - \frac{1}{2}\text{Var}(h^D(f))} \mathbf{1}_A), \quad (4.18)$$

we have

$$M^D \text{ under } \mathbb{P}_f \stackrel{\text{law}}{=} e^{\alpha(\widehat{G}^D f)(x)} M^D(dx) \text{ under } \mathbb{P}, \quad (4.19)$$

where $\widehat{G}^D f(x) := \int_D \widehat{G}^D(x, y) f(y) dy$. Then

$$M^D = Z^D \text{ a.s.} \quad (4.20)$$

where Z^D is the cLQG associated with h^D .

Proof. By Theorem 4.2 it suffices to show (4.8) for $\widehat{M}^D := Z^D$. Let $D, \tilde{D} \in \mathfrak{D}$ be such that $\tilde{D} \subseteq D$ and $\text{Leb}(D \setminus \tilde{D}) = 0$ and let $\Phi^{D, \tilde{D}} = \mathcal{N}(0, C^{D, \tilde{D}})$ be a sample of the binding field independent of $M^{\tilde{D}}$ and $h^{\tilde{D}}$. Let $f_1, f_2 \in C_c(D)$ with $f_2 \geq 0$. Then

$$\begin{aligned} & \mathbb{E}\left(e^{h^{\tilde{D}}(f_1) + \Phi^{D, \tilde{D}}(f_1)} e^{-\langle e^{\alpha\Phi^{D, \tilde{D}}} M^{\tilde{D}}, f_2 \rangle}\right) \\ &= e^{\frac{1}{2}\langle f_1, C^{D, \tilde{D}} f_1 \rangle} \mathbb{E}\left(\exp\left\{h^{\tilde{D}}(f_1) - \langle e^{\alpha\Phi^{D, \tilde{D}} + \alpha C^{D, \tilde{D}}} f_1 M^{\tilde{D}}, f_2 \rangle\right\}\right) \\ &= e^{\frac{1}{2}\langle f_1, G^{\tilde{D}} f_1 \rangle + \frac{1}{2}\langle f_1, C^{D, \tilde{D}} f_1 \rangle} \mathbb{E}\left(\exp\left\{-\langle e^{\alpha\Phi^{D, \tilde{D}} + \alpha C^{D, \tilde{D}}} f_1 e^{\alpha G^{\tilde{D}}} f_1 M^{\tilde{D}}, f_2 \rangle\right\}\right) \end{aligned} \quad (4.21)$$

where we used (4.19) to get the second equality. The expectation on the right-hand side involves only $M^{\tilde{D}}$ and so is determined by the *marginal* law of $M^{\tilde{D}}$ only. By assumption, this law obeys the Gibbs-Markov property meaning that $e^{\alpha\Phi^{D, \tilde{D}}} M^{\tilde{D}} \stackrel{\text{law}}{=} M^D$. Using also

that $\widehat{G}^{\bar{D}} + C^{D, \bar{D}} = \widehat{G}^D$ and one more time (4.19), we get

$$\mathbb{E}\left(e^{h^{\bar{D}}(f_1) + \Phi^{D, \bar{D}}(f_1)} e^{-\langle e^{\alpha\Phi^{D, \bar{D}}} M^{\bar{D}}, f_2 \rangle}\right) = \mathbb{E}\left(e^{h^D(f_1)} e^{-\langle M^D, f_2 \rangle}\right). \quad (4.22)$$

This implies

$$(e^{\alpha\Phi^{D, \bar{D}}} M^{\bar{D}}, h^{\bar{D}} + \Phi^{D, \bar{D}}) \stackrel{\text{law}}{=} (M^D, h^D). \quad (4.23)$$

By (2.5) and (2.7), there exists a measurable function F with $F(h^D) = Z^D$ and $F(h^D + \varphi) = e^{\alpha\varphi} Z^D$ for all $D \in \mathfrak{D}$ and $\varphi \in \mathbb{R}^{C_0(D)}$. Hence we get

$$(e^{\alpha\Phi^{D, \bar{D}}} M^{\bar{D}}, e^{\alpha\Phi^{D, \bar{D}}} Z^{\bar{D}}) \stackrel{\text{law}}{=} (M^D, Z^D). \quad (4.24)$$

Theorem 4.2 now implies (4.20). \square

If it were not for the reliance on the conditions of Theorem 4.1, Theorem 4.3 would run very close to the characterization of the *subcritical* Gaussian Multiplicative Chaos measures by their behavior under Cameron-Martin shifts established by Shamov [30]. In order to get the critical case aligned with subcritical ones, one would have to prove the conditions of Theorem 4.1 directly from (4.19), a task we will consider returning to in future work.

5. MOMENT CALCULATIONS AND ESTIMATES

We are ready to move to the proofs of our main results. In this section we perform moment calculations for the near-critical measure ζ^D ; these will feed directly into the proofs of Theorem 2.3. Throughout, we will assume that a domain $D \in \mathfrak{D}$ and a set of approximating lattice domains $\{D_N\}_{N \geq 1}$ have been fixed.

5.1 Strategy and key lemmas.

In order to motivate the forthcoming derivations, let us first discuss the main steps of the proof of Theorem 2.3. Focussing only on the limit of the measures ζ_N^D , our aim is to show convergence in law of $\langle \zeta_N^D, f \rangle$ for a class of test functions f . We will proceed using a similar strategy as in [8–11]: First prove tightness, which permits extraction of subsequential limits, and then identify the limit uniquely by its properties.

The tightness will be resolved by a first-moment calculation. However, since Z^D is known to lack the first moment and the convergence of $\langle \zeta_N^D, f \rangle$ cannot thus hold in the mean, the moment calculation must be restricted by a suitable truncation. In order to identify the subsequential limits with the expression on the right of (2.9), we invoke Theorem 4.1. This requires checking the conditions in the statement of which the most subtle is the uniform Laplace-transform tail (4.4). It is here where we need the present version as opposed to [9, Theorem 2.8] because, due to our reliance on subsequential convergence, we do not have a direct argument for dilation covariance of the limit.

Both the tightness and the uniform Laplace-transform tail will be inferred from the following two propositions whose proof constitutes the bulk of this section:

Proposition 5.1 (Truncated first moment) *For all bounded continuous $f: D \times \mathbb{R} \rightarrow \mathbb{R}$,*

$$\lim_{u \rightarrow \infty} \limsup_{N \rightarrow \infty} \left| \frac{1}{u} \mathbb{E} \left(\langle \zeta_N^D, f \rangle; h^{D_N} \leq m_N + u \right) - \frac{2e^{2c_0/g}}{g\sqrt{2\pi g}} \int_{D \times (0, \infty)} s e^{-\frac{s^2}{2g}} r^D(x)^2 f(x, s) \, dx \, ds \right| = 0, \quad (5.1)$$

where the limit in u is uniform in $f \in C(D \times \mathbb{R})$ with $\|f\|_\infty \leq 1$. Moreover, there exists a constant $c' = c'(D) < \infty$ such that

$$\mathbb{E} \left(\zeta_N^D(D \times \mathbb{R}); h^{D_N} \leq m_N + u \right) \leq c'u \quad (5.2)$$

for all $N \geq 1$ and $u \geq 1$.

Proposition 5.2 (Truncated second moment) *For each $\delta > 0$ there exists a continuous function $\theta: [0, \infty) \rightarrow [0, \infty)$ with $\lim_{u \rightarrow \infty} \theta(u)/u = 0$ such that*

$$\limsup_{N \rightarrow \infty} \mathbb{E} \left(\zeta_N^D(D^\delta \times (-\infty, \delta^{-1}])^2; h^{D_N} \leq m_N + u \right) \leq e^{\alpha u + \theta(u)} \quad (5.3)$$

for all $u \geq 0$.

Before delving into the proof, let us see how tightness follows from this:

Lemma 5.3 (Tightness) *The family $\{\zeta_N^D(D \times \mathbb{R}): N \geq 1\}$ of \mathbb{R} -valued random variables is tight. Consequently, the measures $\{\zeta_N^D: N \geq 1\}$ are tight relative to the vague topology on $\overline{D} \times \overline{\mathbb{R}}$.*

Proof. Since ζ_N^D is supported on $D \times \mathbb{R}$, the second part of the claim follows from the first and so we just need to show that $\{\zeta_N^D(D \times \mathbb{R}): N \geq 1\}$ is tight. Let $u > 0$ and $c > 0$. By the union bound and Markov's inequality,

$$\begin{aligned} \mathbb{P}(\zeta_N^D(D \times \mathbb{R}) > a) &\leq \mathbb{P}\left(\max_{x \in D_N} h^{D_N} > m_N + u\right) + \frac{1}{a} \mathbb{E}\left(\zeta_N^D(D \times \mathbb{R}); h^{D_N} \leq m_N + u\right). \end{aligned} \quad (5.4)$$

Lemma 3.7 and Proposition 5.1 dominate the right-hand side by a quantity of order $ue^{-\alpha u} + a^{-1}u$, uniformly in $N \geq 1$. This can be made arbitrary small by choosing, e.g., $a := u^2$ and taking u sufficiently large. \square

The proof of the Laplace transform asymptotic is more involved. The bulk of the computation is the content of:

Lemma 5.4 (Laplace transform asymptotic) *For all bounded continuous $f_1: D \rightarrow [0, \infty)$ and all nonnegative $f_2 \in C_c(\mathbb{R})$,*

$$\lim_{\lambda \downarrow 0} \limsup_{N \rightarrow \infty} \left| \frac{\mathbb{E} \left(\langle \zeta_N^D, f_1 \otimes f_2 \rangle e^{-\lambda \langle \zeta_N^D, 1_D \otimes f_2 \rangle} \right)}{\log(1/\lambda)} - c_\star \int_{D \times (0, \infty)} s e^{-\frac{s^2}{2s}} r^D(x)^2 f_1(x) f_2(s) \, dx \, ds \right| = 0, \quad (5.5)$$

where c_\star is the constant from Theorem 2.3.

Proof. We will separately show that the *limes superior*, resp., *limes inferior* as $N \rightarrow \infty$ and $\lambda \downarrow 0$ of the ratio is bounded from above, resp., below by the integral expression in (5.5). For the *limes superior* we parametrize the $\lambda \downarrow 0$ limit as the $u \rightarrow \infty$ limit of $\lambda := ue^{-\alpha u}$. For this choice of λ , the expectation in (5.5) is bounded from above by

$$\mathbb{E} \left(\langle \zeta_N^D, f_1 \otimes f_2 \rangle e^{-ue^{-\alpha u} \langle \zeta_N^D, 1_D \otimes f_2 \rangle}; \max_{x \in D_N} h_x^{D_N} > m_N + u \right) + \mathbb{E} \left(\langle \zeta_N^D, f_1 \otimes f_2 \rangle; h^{D_N} \leq m_N + u \right). \quad (5.6)$$

Using that $\max_{t \geq 0} te^{-\lambda t} = (\lambda e)^{-1}$, the first term is at most

$$\|f_1\|_\infty u^{-1} e^{\alpha u - 1} \mathbb{P} \left(\max_{x \in D_N} h_x^{D_N} > m_N + u \right), \quad (5.7)$$

which upon division by $\log(1/\lambda) = \alpha u + \log(1/u)$ tends to zero in the limits as $N \rightarrow \infty$ followed by $u \rightarrow \infty$, thanks to (3.15). By Proposition 5.1, the second term in (5.6) divided by αu tends to the integral term in (5.5) under these limits.

For the *limes inferior*, pick $\delta > 0$ and parametrize the $\lambda \downarrow 0$ limit as the $u \rightarrow \infty$ limit of $\lambda := e^{-\alpha u - \theta(u)}$, for θ as in Proposition 5.2. (This exhausts all small values of λ because θ is continuous with $\theta(u) = o(u)$.) For this choice of λ , the expectation in (5.5) is at least

$$\mathbb{E} \left(\langle \zeta_N^D, f_1 \otimes f_2 \rangle; h^{D_N} \leq m_N + u \right) - \mathbb{E} \left(\langle \zeta_N^D, f_1 \otimes f_2 \rangle \left(1 - \exp \left\{ -e^{-\alpha u - \theta(u)} \langle \zeta_N^D, 1_D \otimes f_2 \rangle \right\} \right); h^{D_N} \leq m_N + u \right). \quad (5.8)$$

By Proposition 5.1, the first term in the last expression divided by $\log(1/\lambda) = \alpha u + o(u)$ converges to the desired limit as $N \rightarrow \infty$ and then $u \rightarrow \infty$. The second term (without the minus sign) is at most $\|f_1\|_\infty$ times

$$\begin{aligned} & \mathbb{E} \left(\langle \zeta_N^D, 1_{D \setminus D^\delta} \otimes f_2 \rangle; h^{D_N} \leq m_N + u \right) \\ & + (1 - e^{-\delta}) \mathbb{E} \left(\langle \zeta_N^D, 1_{D^\delta} \otimes f_2 \rangle; h^{D_N} \leq m_N + u \right) \\ & + \mathbb{E} \left(\langle \zeta_N^D, 1_{D^\delta} \otimes f_2 \rangle; h^{D_N} \leq m_N + u, \langle \zeta_N^D, 1_D \otimes f_2 \rangle > \delta e^{\alpha u + \theta(u)} \right). \end{aligned} \quad (5.9)$$

Proposition 5.1 ensures that, upon division by u , the *limes superior* as $N \rightarrow \infty$ and $u \rightarrow \infty$ of first two terms is at most order $\text{Leb}(D \setminus D^\delta) + \delta$. Using the Cauchy-Schwarz and Markov inequalities, the third term is in turn at most the square root of

$$\begin{aligned} & \delta^{-1} e^{-\alpha u - \theta(u)} \mathbb{E} \left(\langle \zeta_N^D, 1_D \otimes f_2 \rangle; h^{D_N} \leq m_N + u \right) \\ & \quad \times \mathbb{E} \left(\langle \zeta_N^D, 1_{D^\delta} \otimes f_2 \rangle^2; h^{D_N} \leq m_N + u \right). \end{aligned} \quad (5.10)$$

Thanks to our restriction on the support of f_2 , Proposition 5.2 bounds the second expectation by a constant times $e^{\alpha u + \theta(u)}$ while Proposition 5.1 shows that the first expectation is at most $O(u)$. The last expectation in (5.9) is thus at most $O(\sqrt{u})$ which upon dividing by u vanishes in the limits $N \rightarrow \infty$ and $u \rightarrow \infty$.

We conclude that the *limes inferior* of the expectation term in (5.5) is at least the integral term minus a quantity of order $\text{Leb}(D \setminus D^\delta) + \delta$. As $D^\delta \uparrow D$ implies $\text{Leb}(D \setminus D^\delta) \downarrow 0$ as $\delta \downarrow 0$, the claim follows. \square

5.2 Truncated first moment.

The remainder of this section is devoted to the proof of the above two key lemmas. When dealing with various Gaussian densities, we will frequently refer to:

Lemma 5.5 *Denote $v_N(y) := g \log N + gy$. Then for all $c \geq 1$ and all $\epsilon > 0$:*

(1) *For all $N \geq e^c$, all $y \in [-c, c]$ and all s with $|s| \leq (\log N)^{1-\epsilon}$,*

$$\exp \left(-\frac{(m_N - s)^2}{2v_N(y)} \right) = N^{-2} (\log N)^{3/2} e^{as} \exp \left(-\frac{s^2}{2g \log N} + 2y + \delta_N(s, y) \right), \quad (5.11)$$

where $\delta_N(s, y) \in \mathbb{R}$ satisfies

$$\lim_{N \rightarrow \infty} \sup_{|s| \leq (\log N)^{1-\epsilon}} \sup_{|y| \leq c} |\delta_N(s, y)| = 0. \quad (5.12)$$

(2) *There exist constants $c_1, c_2 < \infty$ such that*

$$\exp \left(-\frac{(m_N + r)^2}{2v_N(y)} \right) \leq c_1 N^{-2} v_N(y)^{3/2} e^{-ar} \exp \left(-\frac{r^2}{2v_N(y)} + |r| \frac{3\alpha \log \log N}{8 \log N} \right) \quad (5.13)$$

$$\leq c_2 N^{-2} v_N(y)^{3/2} e^{-ar}$$

for all $N \geq e^c$, all $y \in [-c, c]$ and all $r \in \mathbb{R}$.

Proof. Abbreviate $n := \log N$. Then (for $n \geq 1$) $m_N = 2\sqrt{gn} - \frac{3}{4}\sqrt{g} \log n$ and so

$$-\frac{1}{2}(m_N + r)^2 = -2gn^2 + \frac{3}{2}gn \log n - 2\sqrt{g}rn - \frac{9}{32}g \log(n)^2 + \frac{3}{4}\sqrt{g}r \log n - \frac{1}{2}r^2. \quad (5.14)$$

From $v_N(y) = gn + gy$ we get

$$\frac{1}{v_N(y)} = \frac{1}{gn} - \frac{y}{gn^2 + gny} = \frac{1}{gn} - \frac{1}{gn^2} (y + o(1)), \quad (5.15)$$

where $o(1) \rightarrow 0$ as $N \rightarrow \infty$ uniformly in $y \in [-c, c]$. Then (5.11) follows by combining (5.14) with (5.15) and noting that $|s| \leq n^{1-\epsilon}$. For (5.13) we bound the second to last term in (5.14) by its absolute value and then use that $1/v_N(y) \leq 1/(gn - c)$. For the second line in (5.13) we also need that the minimum of $x \mapsto Ax^2 + Bx$ is, for $A > 0$, equal to $-\frac{1}{4}B^2/A$. \square

We start by the first moment asymptotics:

Proof of Proposition 5.1. Let us again use the shorthand $n := \log N$ whenever convenient. We first show (5.1) for f with compact support in $D \times \mathbb{R}$ as this illustrates the main computation driving the proof. For any $u \geq 1$, $x \in D$ and $r \in \mathbb{R}$, let

$$Q_{N,x}(u, r) := \mathbb{P}\left(h^{D_N} \leq m_N + u \mid h_{\lfloor Nx \rfloor}^{D_N} = m_N - r\right). \quad (5.16)$$

The expectation in (5.1) then equals

$$\frac{1}{n} \sum_{x \in D_N} \int_{\mathbb{R}} f(x/N, s) e^{-\alpha s \sqrt{n}} Q_{N,x/N}(u, s\sqrt{n}) \mathbb{P}\left(\frac{m_N - h_x^{D_N}}{\sqrt{n}} \in ds\right). \quad (5.17)$$

By our restriction to f with compact support, there is $\delta > 0$ such that $\text{dist}(x, D_N^c) > \delta N$ holds for all x effectively contributing to the sum. Using (5.11) in conjunction with the Green function asymptotic (3.3), for the integrating measure in (5.17) we get

$$e^{-\alpha s \sqrt{n}} \mathbb{P}\left(\frac{m_N - h_x^{D_N}}{\sqrt{n}} \in ds\right) = \frac{1}{\sqrt{2\pi g}} N^{-2} n^{3/2} r^D(x/N)^2 e^{2c_0/g + o(1)} e^{-\frac{s^2}{2g}} ds, \quad (5.18)$$

where $o(1) \rightarrow 0$ as $N \rightarrow \infty$ uniformly in $(x, s) \in \text{supp } f$. This permits us to bring the $o(1)$ term outside the integral which turns (5.17) into

$$\begin{aligned} & \frac{e^{o(1)+2c_0/g}}{\sqrt{2\pi g}} \frac{1}{N^2} \sum_{x \in D_N} r^D(x/N)^2 \int_{\mathbb{R}} f(x/N, s) \sqrt{n} Q_{N,x/N}(u, s\sqrt{n}) e^{-\frac{s^2}{2g}} ds \\ &= \frac{e^{o(1)+2c_0/g}}{\sqrt{2\pi g}} \int_{D \times \mathbb{R}} r^D(x)^2 f_N(x, s) \sqrt{n} Q_{N,x}(u, s\sqrt{n}) e^{-\frac{s^2}{2g}} dx ds, \end{aligned} \quad (5.19)$$

where $f_N(x, s) := f(\lfloor Nx \rfloor/N, s)$ and where we noted that $Q_{N,x} = Q_{N, \lfloor Nx \rfloor/N}$. Henceforth we assume that N is so large that (2.1) ensures that $f_N(x, s) \neq 0$ implies $x \in D^{\delta/2}$.

Concerning the integrand, the second part of Lemma 3.9 bounds $Q_{N,x}(u, r)$ by

$$Q_{N,x}(u, r) \leq c \frac{(1+u)(1+(u+r)^+)}{n} \quad (5.20)$$

for all $x \in D^{\delta/2}$, and so $\sqrt{n} Q_{N,x}(u, s\sqrt{n})$ is bounded by a constant times $(1+u)(1+s^+)$ uniformly in $(x, s) \in \text{supp}(f_N)$ and $N \geq 1$. For an asymptotic expression, we first note that $Q_{N,x}(u, s\sqrt{n}) = 0$ whenever $s\sqrt{n} < -u$ for all $x \in D$. On the other hand, if $s \geq 0$, then by Lemma 3.8, as $N \rightarrow \infty$,

$$Q_{N,x}(u, s\sqrt{n}) = [2 + o(1)] \frac{\mathcal{L}^x(u) \mathcal{R}(u + s\sqrt{n})}{gn}. \quad (5.21)$$

As \mathcal{R} is asymptotic to the identity function at large values of its argument, we get

$$\sqrt{n} Q_{N,x}(u, s\sqrt{n}) \xrightarrow{N \rightarrow \infty} \frac{2}{g} \mathcal{L}^x(u) s 1_{[0,\infty)}(s), \quad (5.22)$$

for all $x \in D$ and $s \in \mathbb{R}$.

Since the assumptions on f ensure that $f_N(x, s) \rightarrow f(x, s)$, the Dominated Convergence Theorem shows that (5.17) converges as $N \rightarrow \infty$ to

$$e^{2c_0/g} \frac{2}{g\sqrt{2\pi g}} \int_{D \times [0,\infty)} r^D(x)^2 f(x, s) \mathcal{L}^x(u) s e^{-\frac{s^2}{2g}} dx ds. \quad (5.23)$$

Dividing by u , using (3.16) (which holds uniformly whenever the integrand is non-zero) and applying the Dominated Convergence Theorem then yield the claim for all f with compact support. Using Hölder's Inequality for the difference between (5.23) divided by u and its limiting value then shows that the convergence is uniform in f as desired.

To include f whose support is of the form $A \times \mathbb{R}$ for a compact set $A \subseteq D$, we argue as before but restrict the integral to $s \in [M, \infty)$ and use (5.13) and Lemma 3.9 instead of (5.11) and Lemma 3.8 to obtain

$$\begin{aligned} \frac{1}{1+u} \mathbb{E} \left(\zeta_N^D(A \times [M, \infty)); h^{D_N} \leq m_N + u \right) \\ \leq c \int_{s \geq M} ds e^{-\frac{s^2}{2g} + s \frac{3\alpha \log n}{8\sqrt{n}}} \left(1 + \frac{u}{\sqrt{n}} + s \right), \end{aligned} \quad (5.24)$$

where the constant on the right-hand side may depend on D . The right-hand side vanishes in the limit as $N \rightarrow \infty$, $u \rightarrow \infty$ and $M \rightarrow \infty$ (in this order). Hence, the contribution to (5.17) of the part of the integral corresponding to $s \in [M, \infty)$ tends to 0 in these limits, with the rate of convergence depending on f through its supremum norm only.

Finally, to include functions with arbitrary support, we need:

Lemma 5.6 *For each $D \in \mathfrak{D}$ there is $c \in (0, \infty)$ such that for all measurable $A \subseteq D$, all $u \geq 1$ and all $N \geq 3$,*

$$u^{-1} \mathbb{E} \left(\zeta_N^D(A \times \mathbb{R}); h^{D_N} \leq m_N + u \right) \leq c \frac{1}{N^2} |\{x \in \mathbb{Z}^2 : x/N \in A\}|. \quad (5.25)$$

Postponing the proof for a moment, (5.25) shows that restricting the support of f to $D^\delta := \{x \in D : d_\infty(x, D^c) > \delta\}$ with the help of a bounded mollifier that differs from one or zero only $D^\delta \setminus D^{2\delta}$, the expectation in (5.1) changes by at most a constant times $uN^{-2}|D_N \setminus D_N^{2\delta}|$, where D_N^δ is the discretization of D^δ via (2.1). The integral in (5.1) also changes by a constant times $\text{Leb}(D \setminus D^{2\delta})$. Both constants depend on f through its supremum norm only. The errors tend to zero in light of

$$N^{-2}|D_N \setminus D_N^\delta| \leq \text{Leb}(D \setminus D^{2\delta}), \quad (5.26)$$

and the fact that $\text{Leb}(D \setminus D^{2\delta}) \rightarrow 0$ as $\delta \downarrow 0$. \square

It remains to give:

Proof of Lemma 5.6. Fix $x \in D_N$ and let $B := \{z \in D_N : d_\infty(x, z) < d_\infty(x, (D_N)^c)\}$, $B' := \{z \in D_N : d_\infty(x, z) < \frac{1}{2}d_\infty(x, (D_N)^c)\}$. Using the Gibbs-Markov property (Lemma 3.3) to write $h^{D_N} = h^B + \varphi^{D_N, B}$ we then get

$$\begin{aligned} & \mathbb{E}\left(e^{\alpha(h_x^{D_N} - m_N)}; h^{D_N} \leq m_N + u\right) \\ & \leq \mathbb{E}\left(e^{\alpha(h_x^{D_N} - m_N)}; h^{D_N} \leq m_N + u \text{ in } B'\right) \\ & \leq \sum_{k=0}^{\infty} \mathbb{P}\left(\max_{z \in B'} |\varphi_z^{D_N, B}| \geq k\right) e^{\alpha(k+1)} \mathbb{E}\left(e^{\alpha(h_x^B - m_N)}; h^B \leq m_N + u + k \text{ in } B'\right). \end{aligned} \quad (5.27)$$

A key point is to estimate the expectation on the right-hand side. First, a Girsanov-type computation along with the FKG inequality (cf Lemma 3.6) yields, for all $u' \in \mathbb{R}$,

$$\begin{aligned} & \mathbb{E}\left(e^{\alpha h^B(x)}; h^B \leq m_N + u' \text{ in } B'\right) \\ & = e^{\frac{\alpha^2}{2} G^B(x, x)} P\left(h^B \leq m_N + u' - \alpha G^B(x, \cdot) \text{ in } B'\right) \\ & \leq e^{\frac{\alpha^2}{2} G^B(x, x)} \frac{P\left(h^B \leq m_N + u' - \alpha G^B(x, \cdot)\right)}{P\left(h^B \leq m_N + u' - \alpha G^B(x, \cdot) \text{ in } B \setminus B'\right)}. \end{aligned} \quad (5.28)$$

The fact that $G^B(x, y)$ is bounded by a constant uniformly $y \in B \setminus B'$ regardless of the radius of the ball B (and thus also regardless of the exact position of x in D_N) in conjunction with the tightness of the DGFF maximum shows that the probability in the denominator is positive uniformly in $u' \geq 1$ and $x \in D_N$ and $N \geq 1$. Reverting the Girsanov transformation, we get that, for some $c \in (0, \infty)$,

$$\mathbb{E}\left(e^{\alpha h^B(x)}; h^B \leq m_N + u + k \text{ in } B'\right) \leq c \mathbb{E}\left(e^{\alpha h^B(x)}; h^B \leq m_N + u + k\right) \quad (5.29)$$

holds for all $k \geq 0$, all $u \geq 1$ all $x \in D_N$ and all $N \geq 1$.

Since x is at the center of B , the calculation (5.17–5.20) applies and bounds the right-hand side by a constant times

$$(m_N - m_L + u + k) \frac{1 \vee \log L}{L^2} e^{\alpha m_L} \quad \text{for } L := d_\infty(x, D_N), \quad (5.30)$$

uniformly in $x \in D_N$ and $u \geq 1$. Using that (for $N \geq 3$)

$$\frac{1 \vee \log L}{L^2} e^{\alpha m_L} = \frac{L^2}{N^2} \sqrt{\frac{\log N}{1 \vee \log L}} \frac{\log N}{N^2} e^{\alpha m_N} \quad (5.31)$$

and that $m_N - m_L = O(1) + O(\log(N/L))$ and $L = O(N)$ uniformly in $x \in D_N$, we conclude

$$\mathbb{E}\left(e^{\alpha(h_x^B - m_N)}; h^B \leq m_N + u + k \text{ in } B'\right) \leq c(u + k) \frac{\log N}{N^2} \quad (5.32)$$

for some constant $c \in (0, \infty)$, uniformly in $x \in D_N$, $k \geq 0$ and $u \geq 1$.

Thanks to Lemma 3.2, $\max_{z \in B'} \text{Var}(\varphi^{D_N, B}(z))$ is bounded uniformly in (the centering point) $x \in D_N$ and $N \geq 1$. A standard argument based on the Fernique estimate along with Borell-TIS inequality then shows that, for some $\tilde{c} > 0$,

$$\mathbb{P}\left(\max_{z \in B'} |\varphi_z^{D_N, B}| \geq k\right) \leq e^{-\tilde{c}k^2}, \quad k \geq 0. \quad (5.33)$$

Plugging this, along with (5.32), into (5.27), the sum over $k \geq 0$ may be performed resulting in an expression bounded by a constant times $u(\log N)N^{-2}$, uniformly in $u \geq 1$ and $x \in D_N$. The claim follows by summing over $x \in D_N$ with $x/N \in A$ and invoking the definition of ζ_N^D . \square

We remark that it is exactly Lemma 5.6 that forces us to restrict attention to discretization via (2.1). This is in turn caused by our reliance on Lemma 3.2 which only applies when the ‘‘maximal’’ discretization (2.1) is used.

5.3 Truncated second moment.

Our next task is the proof of the upper bound in Proposition 5.2 on the truncated second moment of measures ζ_N^D . Given an integer $N \geq 1$, distinct vertices $x, y \in D_N$ and a real number $u \geq 1$, let $f_u^{N;x,y} : \mathbb{R}^2 \rightarrow [0, \infty)$ denote the density in

$$\mathbb{P}\left(h^{D_N} \leq m_N + u, h_x^{D_N} - m_N \in ds, h_y^{D_N} - m_N \in dt\right) = f_u^{N;x,y}(s, t) ds dt. \quad (5.34)$$

In light of the explicit form of the law of $(h_x^{D_N}, h_y^{D_N})$, we may and will assume that $f_u^{N;x,y}$ is left continuous in each variable which fixes this function uniquely.

Let D_N^δ be the discretization (via (2.1)) of $D^\delta := \{x \in \mathbb{R}^2 : d_\infty(x, D^c) > \delta\}$. Abbreviate $M := \delta^{-1}$. Then, thanks to the restriction on the maximum of h^{D_N} ,

$$\begin{aligned} & \mathbb{E}\left(\left(\zeta_N^D(D^\delta \times (-\infty, M])\right)^2; h^{D_N} \leq m_N + u\right) \\ &= \frac{1}{(\log N)^2} \sum_{x \in D_N^\delta} \int_{[-M\sqrt{\log N}, u]} e^{2\alpha s} \mathbb{P}(h_x^{D_N} - m_N \in ds, h^{D_N} \leq m_N + u) \\ & \quad + \frac{1}{(\log N)^2} \sum_{\substack{x, y \in D_N^\delta \\ x \neq y}} \int_{[-M\sqrt{\log N}, u]^2} e^{\alpha(s+t)} f_u^{N;x,y}(s, t) ds dt. \end{aligned} \quad (5.35)$$

To control the second integral we will need a good estimate on $f_u^{N;x,y}(s, t)$:

Lemma 5.7 (2-density estimate) *Let $\ell_N(x, y)$ be the largest integer for which the ℓ^∞ -balls $B(x, e^\ell + 1)$ and $B(y, e^\ell + 1)$ are disjoint. For each $\delta, \epsilon > 0$ there exist $C, p \geq 1$ such that, with $\ell := \ell_N(x, y)$ and $n := \log N$,*

$$e^{\alpha(s+t)} f_u^{N;x,y}(s, t) \leq CN^{-2} e^{-2\ell} \left(\frac{n}{\ell(n-\ell)}\right)^{3/2} (1+s^-)(1+t^-) e^{(\alpha+\epsilon)u} \quad (5.36)$$

holds for all $N \geq 3$, all $x, y \in D_N^\delta$ with $1 \leq \ell_N(x, y) \leq n - p$, all $s, t \in \mathbb{R}$ and all $u \geq 1$.

Before we delve into the proof, let us show how this implies:

Proof of Proposition 5.2. Fix $\delta, \epsilon > 0$ and let p be as in Lemma 5.7. Abbreviate $n := \log N$ and $M := \delta^{-1}$. We start by bounding the short-range part of the double sum in (5.35) using that $e^{\alpha(s+t)} \leq e^{\alpha s + \alpha u}$ on the integration domain. This gives

$$\begin{aligned} & \frac{1}{(\log N)^2} \sum_{x \in D_N^\delta} \int_{[-M\sqrt{\log N}, u]} e^{2\alpha s} \mathbb{P}(h_x^{D_N} - m_N \in ds, h^{D_N} \leq m_N + u) \\ & + \frac{1}{(\log N)^2} \sum_{\substack{x, y \in D_N^\delta \\ \ell_N(x, y) < 1 \\ x \neq y}} \int_{[-M\sqrt{\log N}, u]^2} e^{\alpha(s+t)} f_u^{N; x, y}(s, t) ds dt \\ & \leq \frac{1}{n^2} \sum_{\substack{x, y \in D_N^\delta \\ \ell_N(x, y) < 1}} e^{\alpha u} \int e^{\alpha s} \mathbb{P}(h_x^{D_N} - m_N \in ds, h^{D_N} \leq m_N + u). \end{aligned} \quad (5.37)$$

Since $\ell_N(x, y) < 1$ implies $y \in B(x, 2e^1)$, using $|B(x, r) \cap D_N^\delta| \leq 4r^2$ we can estimate the right-hand side of (5.37) by

$$\mathbb{E} \left(\zeta_N^D(D^\delta \times \mathbb{R}); h^{D_N} \leq m_N + u \right) \frac{16e^{2+\alpha u}}{n}. \quad (5.38)$$

Lemma 5.6 ensures the expectation is bounded uniformly in $N \geq 1$. Taking $N \rightarrow \infty$, (5.38) tends to zero.

We will handle the remaining portion of the sum in (5.35) by a covering argument. For this let us consider a general measurable set $A \subseteq D^\delta$ with $\text{diam}(A) < \delta e^{-p}$ and denote $A_N := \{x \in D_N : x/N \in A\}$. Then, for δ sufficiently small, the definition of $\ell_N(x, y)$ ensures $\ell_N(x, y) \leq n - p$ for all $x, y \in A_N$. Lemma 5.7 then gives, for $N \geq N_0$,

$$\begin{aligned} & \frac{1}{(\log N)^2} \sum_{\substack{x, y \in A_N \\ \ell_N(x, y) \geq 1}} \int_{[-M\sqrt{\log N}, u]^2} e^{\alpha(s+t)} f_u^{N; x, y}(s, t) ds dt \\ & \leq C \frac{1}{N^2 (\log N)^2} \sum_{\ell=1}^{n-p} \sum_{\substack{x, y \in A_N \\ \ell_N(x, y) = \ell}} e^{-2\ell} \left(\frac{n}{\ell(n-\ell)} \right)^{3/2} \\ & \quad \times \int_{[-M\sqrt{\log N}, u]^2} (1+s^-)(1+t^-) e^{(\alpha+\epsilon)u} ds dt. \end{aligned} \quad (5.39)$$

For N sufficiently large (depending on u) the integral is at most order $M^4 (\log N)^2 e^{(\alpha+\epsilon)u}$. Since, for each $x \in D_N$, there are at most a constant times $e^{2\ell}$ vertices y with $\ell_N(x, y) = \ell$, the expression in (5.39) is thus at most a constant times

$$M^4 e^{(\alpha+\epsilon)u} \frac{1}{N^2} |A_N| \sum_{\ell=1}^{n-p} \left(\frac{n}{\ell(n-\ell)} \right)^{3/2} \leq 2^{5/2} M^4 e^{(\alpha+\epsilon)u} \frac{1}{N^2} |A_N| \sum_{\ell \geq 1} \ell^{-3/2}. \quad (5.40)$$

All the above bounds are uniform in the choice of the set A . Covering D^δ by $K = O(\delta^2 e^{2p})$ of ℓ^∞ -balls $A^{(1)}, \dots, A^{(K)}$ of radius δe^{-p} , using

$$\begin{aligned} & \mathbb{E} \left((\zeta_N^D(D^\delta \times (-\infty, M]))^2; h^{D_N} \leq m_N + u \right) \\ & \leq K \sum_{i=1}^K \mathbb{E} \left((\zeta_N^D(A^{(i)} \times (-\infty, M]))^2; h^{D_N} \leq m_N + u \right) \end{aligned} \quad (5.41)$$

and noting that the right-hand side of (5.40) is additive in $|A_N|$ we conclude

$$\limsup_{N \rightarrow \infty} \mathbb{E} \left((\zeta_N^D(D^\delta \times (-\infty, M]))^2; h^{D_N} \leq m_N + u \right) \leq c_\epsilon e^{(\alpha + \epsilon)u} \quad (5.42)$$

for a constant $c_\epsilon \in [1, \infty)$ depending on ϵ . As this holds for all $\epsilon \in (0, 1)$, we get (5.3) with $\theta(u) := \inf_{0 < \epsilon < 1} [\epsilon u + \log c_\epsilon]$. This is trivially $o(u)$ as $u \rightarrow \infty$ and is continuous in light of the bounds $\theta(u) \leq \theta(u + r) \leq \theta(u) + r$ for all $u, r \geq 0$. \square

The proof of Lemma 5.7 constitutes the remainder of this subsection. Given $x, y \in D_N$, let us abbreviate $\ell = \ell_N(x, y)$, set $L := \lfloor e^\ell \rfloor$ and define

$$B_1 := B(x, L) \cap \mathbb{Z}^2, \quad B_2 := B(y, L) \cap \mathbb{Z}^2, \quad W := B_1 \cup B_2, \quad B_3 := D_N \setminus (W \cup \partial_\infty W), \quad (5.43)$$

where $\partial_\infty W$ denotes the set of all vertices whose ℓ^∞ -distance to W is 1. Observe that ∂W is a subset of ∂B_3 which is the set of all vertices whose ℓ^∞ -distance to x or y is L . Also, pick $\epsilon' \in (0, 1/3)$, let $W' := \{z \in D_N^\delta : d_\infty(z, \partial_\infty W) > \epsilon' L\}$ and finally set

$$B'_1 := B_1 \cap W', \quad B'_2 := B_2 \cap W', \quad B'_3 := B_3 \cap W'. \quad (5.44)$$

To bound $f_u^{N; x, y}(s, t)$, we will first condition on the values of h^{D_N} on $\partial_\infty W$ and then use the Gibbs-Markov property, Lemma 3.9 and standard Gaussian estimates to bound this conditional density in terms of the unique bounded harmonic extension $\tilde{\varphi}^N$ of $h^{D_N}|_{\partial_\infty W}$ onto all of \mathbb{Z}^2 . Note that $\tilde{\varphi}^N = \varphi^{D_N, D_N \setminus \partial_\infty W}$ on $W \cup \partial_\infty W$.

For our purposes, sufficient control over $\tilde{\varphi}^N$ can be achieved by using its value at x , together with its oscillation on W' :

$$R_N(x, y) := \max_{z \in W'} |\tilde{\varphi}_z^N - \tilde{\varphi}_x^N| \quad (5.45)$$

which we control via:

Lemma 5.8 *Let $\delta, \epsilon' \in (0, 1/3)$. There exists $C > 0$ such that for all $N \geq 1$, all $x, y \in D_N^\delta$, all $r \geq 0$, and all $w \in \mathbb{R}$,*

$$\mathbb{P} \left(R_N(x, y) \geq r \mid \varphi_x^{D_N, D_N \setminus \partial_\infty W} = m_N - m_{\epsilon^\ell} + w \right) \leq C^{-1} \exp \left\{ -C \left[\left(r - \frac{|w|}{n - \ell} \right)^+ \right]^2 \right\}, \quad (5.46)$$

where $\ell = \ell_N(x, y)$.

Proof. This is a reformulation of [19, Proposition B.5]. We take $-x/N + D$ as U and $B(0, 1) \cup B((y - x)/L, 1)$ as W and use the discretization in [19] for the inner domain. \square

With this in hand, we are ready to give:

Proof of Lemma 5.7. We may assume p to be so large such that $\ell := \ell_N(x, y)$ obeys $\ell \leq n + 2 \log \delta - \log 3$. Then, in particular, $x, y \in D_N^\delta$ implies $W \subseteq D_N^{\delta/2}$. Given $v \in \mathbb{R}^{\partial_\infty W}$, let $f_u^{N;x,y}(s, t|v)$ be the function defined exactly as $f_u^{N;x,y}(s, t)$ in (5.34) except that the probability is conditioned on $\{h^{D_N}|_{\partial_\infty W} = m_N - m_L + v\}$. Using the spatial Markov property of h^{D_N} , the measure $f_u^{N;x,y}(s, t|v) ds dt$ is now dominated by the product of two conditional measures

$$\mathbb{P}\left(h_x^{B_1} - m_N \in ds \mid h^{B_1}|_{\partial B_1} = m_N - m_L + v\right) \quad (5.47)$$

$$\mathbb{P}\left(h_y^{B_2} - m_N \in dt \mid h^{B_2}|_{\partial B_2} = m_N - m_L + v\right) \quad (5.48)$$

and three conditional probabilities

$$\mathbb{P}\left(h^{B_3} \leq m_N + u \text{ on } B'_3 \mid h^{B_3}|_{\partial D_N} = 0, h^{B_3}|_{\partial_\infty W} = m_N - m_L + v\right) \quad (5.49)$$

$$\mathbb{P}\left(h^{B_1} \leq m_N + u \text{ on } B'_1 \mid h^{B_1}|_{\partial B_1} = m_N - m_L + v, h_x^{B_1} = m_N + s\right) \quad (5.50)$$

$$\mathbb{P}\left(h^{B_2} \leq m_N + u \text{ on } B'_2 \mid h^{B_2}|_{\partial B_2} = m_N - m_L + v, h_y^{B_2} = m_N + t\right). \quad (5.51)$$

We will now bound each of these terms separately to give an estimate on $f_u^{N;x,y}(s, t|v)$ which can then be integrated into the desired bound.

Let us write \bar{v} for the unique bounded harmonic extension of v from $\partial_\infty W$ to \mathbb{Z}^2 . Abbreviate

$$v_* := \min_{W'} \bar{v}, \quad w := \bar{v}_x \quad \text{and} \quad r := \max_{z \in W'} |\bar{v}_z - w|. \quad (5.52)$$

The Gibbs-Markov decomposition then dominates (5.47) by

$$\mathbb{P}\left(h_x^{B_1} - m_L + \bar{v}_x \in ds\right) \leq c \ell e^{-2\ell + \alpha(-s+w)} ds, \quad (5.53)$$

where we have used Lemma 3.2 and (5.13) to bound the Gaussian density. Similarly, the second probability is at most

$$\mathbb{P}\left(h_x^{B_1} - m_L + \bar{v}_y \in dt\right) \leq c \ell e^{-2\ell + \alpha(-t+\bar{v}_y)} dt \leq c \ell e^{-2\ell + \alpha(-t+w+r)} dt, \quad (5.54)$$

where the second inequality is based on $\bar{v}_y \leq \bar{v}_x + r$.

For the probabilities in (5.49–5.51) we use the upper bounds in Lemma 3.9. By shifting h^{B_3} by $-m_N - u$, the boundary conditions in (5.49) become $-m_N - u$ on ∂D_N and $-m_L + v - u$ on $\partial_\infty W$, while the event whose probability we are after is now $\{h^{B_3}|_{B'_3} \leq 0\}$. As $N/L \geq \frac{1}{3}\delta^2$ by our assumption on p , the first part of Lemma 3.9 with $-x/N + D$, resp., $B(0, 1) \cup B((y-x)/(L-1), 1)$ as D , resp., \tilde{D} and with N , resp., $L-1$ as N , resp., K then dominates (5.49) by

$$\begin{aligned} c \frac{(1+u)((v_* - u)^- + e^{-((v_* - u)^+)^{3/2}})}{n - \ell} &\leq c' \frac{(1+u)((w - u - r)^- + e^{-\rho(w-u-r)^+})}{n - \ell} \\ &\leq c'' \frac{(1+u)(1+r)e^{\rho r}((w - u)^- + e^{-\rho(w-u)^+})}{n - \ell}, \end{aligned} \quad (5.55)$$

where we set $\rho := \alpha + 2\epsilon$ and then choose $c', c'' < \infty$ accordingly.

Similarly, subtracting $m_N + u$ from h^{B_1} in (5.50) and applying Lemma 3.9 with $D := B(0, 1)$ and $N := L + 1$ bounds the probability in (5.50) by

$$c \frac{(1 + (v_* - u)^-)(1 + (s - u)^-)}{\ell} \leq c' \frac{(1 + u)^2(1 + r)(1 + w^-)(1 + s^-)}{\ell}. \quad (5.56)$$

An analogous bound with s replaced by t applies to (5.51). Combining these observations, $f_u^{N;x,y}(s, t|v)$ is at most a constant times

$$\frac{(1 + u)^5(1 + s^-)(1 + t^-)}{n - \ell} e^{-4\ell - \alpha(s+t)} \times \left(e^{2\alpha w + 2\rho r} ((w - u)^- + e^{-\rho(w-u)^+})(1 + r)^3(1 + w^-)^2 \right) \quad (5.57)$$

Recalling the definition of w and r , the integral with respect to $\mathbb{P}(h^{D_N}|_{\partial_\infty W} - m_N + m_L \in dw)$ of the expression inside the large brackets above is at most

$$\int_{w \in \mathbb{R}} \sum_{r \geq 1} \mathbb{P}(\tilde{\varphi}_x^N - m_N + m_L \in dw) \mathbb{P}(R_N(x, y) \geq r - 1 \mid \tilde{\varphi}_x^N - m_N + m_L = w) \times e^{2\alpha w + 2\rho r} ((w - u)^- + e^{-\rho(w-u)^+})(1 + r)^3(1 + w^-)^2. \quad (5.58)$$

For the first probability we first note that, by Lemma 3.2,

$$\text{Var}(\tilde{\varphi}_x^N) \leq g \log(N/L) + O(1). \quad (5.59)$$

Abbreviating $m_{N,L} := m_N - m_{N/L} - m_L$, Lemma 5.5 shows

$$\begin{aligned} \mathbb{P}(\tilde{\varphi}_x^N - m_N + m_L \in dw) &= \mathbb{P}(\tilde{\varphi}_x^N - m_{N/L} - m_{N,L} \in dw) \\ &\leq cN^{-2} e^{2\ell - \alpha(w + m_{N,L})} \log(N/L) dw \\ &= cN^{-2} (\log N / \log L)^{3/2} (\log(N/L))^{-1/2} e^{2\ell - \alpha w} dw, \end{aligned} \quad (5.60)$$

where we used that $m_{N,L} = \frac{3}{4}\sqrt{g}(-\log \log N + \log^+ \log(N/L) + \log^+ \log^+ L)$ once $N, L \geq e$. The second probability in (5.58) is handled by Lemma 5.8.

Integrating $f_u^{N;x,y}(s, t|v)$ with respect to $\mathbb{P}(h^{D_N}|_{\partial_\infty W} - m_N + m_L \in dv)$ and combining all of the above we obtain

$$\begin{aligned} &\frac{f_u^{N;x,y}(s, t)}{N^{-2} e^{-2\ell - \alpha(s+t)} (1 + u)^5 (1 + s^-) (1 + t^-) \left(\frac{n}{\ell(n-\ell)}\right)^{3/2}} \\ &\leq c' \int dw e^{\alpha w} \left((w - u)^- + e^{-\rho(w-u)^+} \right) (1 + |w|)^2 \sum_{r=1}^{\infty} e^{-3\rho(r - \frac{|w|}{n-\ell})} e^{2\rho r} (1 + r)^3 \\ &\leq c'' \int dw e^{\alpha w + \epsilon|w|} \left((w - u)^- + e^{-\rho(w-u)^+} \right), \end{aligned} \quad (5.61)$$

for some constants $c', c'' \in (0, \infty)$, provided p is large enough.

Splitting the integration domain into $w \in (u, \infty)$, $w \in [0, u]$, and $w \in (-\infty, 0)$, the integral is at most a constant times $e^{(\alpha+2\epsilon)u}$. The claim follows with 3ϵ replacing ϵ in the desired upper bound. \square

We note that also the previous lemma relied on estimates from Lemma 3.2 and is thus constrained to the discretization (2.1).

6. PROOF OF THEOREM 2.3

In this section we will give a formal proof of Theorem 2.3. For easier exposition, and also since this is the main novelty in the present work, we will first prove the convergence of the near-extremal process ζ_N^D alone and deal with the joint convergence with the extremal process and the field itself later.

6.1 Near-extremal process convergence.

Thanks to Lemmas 5.6 and 3.7, the processes $\{\zeta_N^D: N \geq 1\}$ are tight in the space of Radon measures on $\bar{D} \times \bar{\mathbb{R}}$. (This uses that ζ_N^D is supported on $D \times \mathbb{R}$ for each $N \geq 1$.) The strategy of the proof is similar to that used in [8–10] for the extremal process and [11] for the intermediate DGFF level sets: We extract a subsequential limit ζ^D of these processes and identify a properties that determine its law uniquely.

Our first item of concern is the dependence of the subsequential limit on the underlying domain D . This is the content of:

Proposition 6.1 *Every sequence of N 's tending to infinity contains a subsequence $\{N_k\}_{k \geq 1}$ and, for all $D \in \mathfrak{D}$, there is a random measure ζ^D such that*

$$\zeta_{N_k}^D \xrightarrow[k \rightarrow \infty]{\text{law}} \zeta^D, \quad D \in \mathfrak{D}. \quad (6.1)$$

The same applies (with the same subsequence and the limit measure) even if ζ_N^D is defined using a sequence of volumes $\{D_N\}_{N \geq 1}$ satisfying (2.12–2.13).

The proof is based on two lemmas that will be of independent interest. We start by recording a consequence of the first moment estimates from Section 5:

Lemma 6.2 (Stochastic absolute continuity) *Let $D \in \mathfrak{D}$ and let ζ^D be a subsequential weak limit of the measures $\{\zeta_N^D: N \geq 1\}$. Then for all measurable $A \subseteq \bar{D}$,*

$$\text{Leb}(A) = 0 \quad \Rightarrow \quad \zeta^D(A \times \bar{\mathbb{R}}) = 0 \quad \text{a.s.} \quad (6.2)$$

The measure ζ^D is concentrated on $D \times [0, \infty)$ with $\zeta^D(D \times [0, \infty)) < \infty$ a.s.

Proof. Let $A \subseteq \bar{D}$ be measurable with $\text{Leb}(A) = 0$. The outer regularity of the Lebesgue measure implies the existence of functions $\{f_j: j \geq 1\} \subseteq C_c(\mathbb{R}^2)$ with support restricted to an open ball containing \bar{D} such that $1_A \leq f_j$ and $\|f_j\| \leq 1$ hold for all $j \geq 1$ and $f_j \downarrow 1_A$

Lebesgue-a.e. as $j \rightarrow \infty$. Given $\epsilon > 0$, we can thus estimate

$$\mathbb{P}(\zeta^D(A \times \bar{\mathbb{R}}) \geq \epsilon) \leq \inf_{j \geq 1} \mathbb{P}(\langle \zeta^D, f_j \otimes 1_{\bar{\mathbb{R}}} \rangle \geq \epsilon) \leq \inf_{j \geq 1} \limsup_{k \rightarrow \infty} \mathbb{P}(\langle \zeta_{N_k}^D, f_j \otimes 1_{\bar{\mathbb{R}}} \rangle \geq \epsilon), \quad (6.3)$$

where $\{N_k\}_{k \geq 1}$ is a subsequence such that $\zeta_{N_k}^D \xrightarrow{\text{law}} \zeta^D$.

Pick $u > 0$. Thanks to the Markov inequality, the probability on the right hand side of (6.3) is at most

$$\epsilon^{-1} \mathbb{E} \left(\langle \zeta_N^D, f_j \otimes 1_{\bar{\mathbb{R}}} \rangle; h^{D_N} \leq m_N + u \right) + \mathbb{P} \left(\max_{x \in D_N} h_x^{D_N} > m_N + u \right) \quad (6.4)$$

with $N := N_k$. Lemma 5.6 along with a routine approximation argument give

$$\limsup_{N \rightarrow \infty} \mathbb{E} \left(\langle \zeta_N^D, f_j \otimes 1_{\bar{\mathbb{R}}} \rangle; h^{D_N} \leq m_N + u \right) \leq cu \int_D f_j(x) dx \quad (6.5)$$

and so, by the Bounded Convergence Theorem, the expectation in (6.4) tends to zero as $N \rightarrow \infty$ followed by $j \rightarrow \infty$. Lemma 3.7 in turn shows that the probability on the right of (6.4) vanishes as $N \rightarrow \infty$ followed by $u \rightarrow \infty$ and so we get (6.2).

The same argument applied to $A := \partial D$ shows that ζ^D puts no mass on $\partial D \times \bar{\mathbb{R}}$ a.s. The bound (5.24) shows that ζ^D does not charge $D \times \{+\infty\}$. By Lemma 3.7, the maximum of h^{D_N} is at most $m_N + o(\sqrt{\log N})$ and so $\zeta^D(D \times [-\infty, 0)) = 0$ a.s. The a.s.-finiteness of $\zeta^D(D \times [0, \infty))$ then follows from Lemma 5.3 \square

Next we will state the consequence Gibbs-Markov property:

Lemma 6.3 (Gibbs-Markov property) *Let $D, \tilde{D} \in \mathfrak{D}$ obey $\tilde{D} \subseteq D$ and $\text{Leb}(D \setminus \tilde{D}) = 0$. Suppose that ζ^D and $\zeta^{\tilde{D}}$ be subsequential weak limits of $\{\zeta_N^D: N \geq 1\}$ and $\{\zeta_N^{\tilde{D}}: N \geq 1\}$ respectively, along the same subsequence. Then*

$$\zeta^D(dx dt) \stackrel{\text{law}}{=} e^{\alpha \Phi^{D, \tilde{D}}(x)} \zeta^{\tilde{D}}(dx dt), \quad (6.6)$$

where $\Phi^{D, \tilde{D}} = \mathcal{N}(0, C^{D, \tilde{D}})$ is independent of $\zeta^{\tilde{D}}$ on the right-hand side.

Proof. We remark that the fact that $\text{Leb}(D \setminus \tilde{D}) = 0$ ensures, via Lemma 6.2, that neither ζ^D nor $\zeta^{\tilde{D}}$ charge the set $(D \setminus \tilde{D}) \times \mathbb{R}$ a.s. and so it is immaterial that $\Phi^{D, \tilde{D}}$ is not defined there.

Pick $f \in C_c(\tilde{D} \times \mathbb{R})$ with $f \geq 0$. Noting that $\tilde{D}_N \subseteq D_N$, let $h^{\tilde{D}_N}$, resp., φ^{D, \tilde{D}_N} be independent fields with the law of DGFF in \tilde{D}_N , resp., the binding field between domains D_N and \tilde{D}_N . Lemma 3.3 then shows that $h^{D_N} := h^{\tilde{D}_N} + \varphi^{D_N, \tilde{D}_N}$ has the law of DGFF in D_N . Define the functions

$$e_N(x) := \exp \{ \alpha \varphi^{D_N, \tilde{D}_N}(\lfloor Nx \rfloor) \} \quad (6.7)$$

and

$$f_N(x, t) := f \left(x, t - \varphi^{D_N, \tilde{D}_N}(\lfloor Nx \rfloor) / \sqrt{\log N} \right). \quad (6.8)$$

Using h^{D_N} to define ζ_N^D and $h^{\tilde{D}_N}$ to define $\zeta_N^{\tilde{D}}$, we then have

$$\langle \zeta_N^D, f \rangle = \langle \zeta_N^{\tilde{D}}, e_N f_N \rangle, \quad (6.9)$$

where $\zeta_N^{\tilde{D}}$ is independent of $\varphi^{D_N, \tilde{D}_N}$ (implicitly contained in f_N) on the right-hand side. Hereby we get

$$\left| \langle \zeta_N^D, f \rangle - \langle \zeta_N^{\tilde{D}}, e^{\alpha \Phi^{D, \tilde{D}}} f \rangle \right| \leq \zeta_N^{\tilde{D}}(D \times \mathbb{R}) \|e_N f_N - e^{\alpha \Phi^{D, \tilde{D}}} f\|_\infty. \quad (6.10)$$

The first term on the right-hand side is tight thanks to Lemma 5.3. The uniform continuity of f and the coupling in Lemma 3.5 show that the second term tends to 0 in probability as $N \rightarrow \infty$. Since $(x, t) \mapsto e^{\alpha \Phi^{D, \tilde{D}}(x)} f(x, t)$ is uniformly continuous a.s., if $\{N_k\}_{k \geq 1}$ is a sequence along with $\zeta_{N_k}^{\tilde{D}}$ tends to $\zeta^{\tilde{D}}$, by conditioning on $\Phi^{D, \tilde{D}}(x)$ we get

$$\mathbb{E}(e^{-\langle \zeta_{N_k}^{\tilde{D}}, e^{\alpha \Phi^{D, \tilde{D}}} f \rangle}) \xrightarrow[k \rightarrow \infty]{} \mathbb{E}(e^{-\langle \zeta^{\tilde{D}}, e^{\alpha \Phi^{D, \tilde{D}}} f \rangle}). \quad (6.11)$$

Applying the subsequential convergence to the first term on the left of (6.10) as well, we infer

$$\mathbb{E}(e^{-\langle \zeta^{\tilde{D}}, e^{\alpha \Phi^{D, \tilde{D}}} f \rangle}) = \mathbb{E}(e^{-\langle \zeta^D, f \rangle}), \quad f \in C_c(\tilde{D} \times \mathbb{R}). \quad (6.12)$$

The equality extends to all $f \in C(\overline{D} \times \overline{\mathbb{R}})$ by approximating f on $\tilde{D} \times \mathbb{R}$ by an increasing sequence of functions in $C_c(\tilde{D} \times \mathbb{R})$ and invoking the Bounded Convergence Theorem along with Lemma 6.2. Hereby we get (6.6). \square

We are ready to give:

Proof of Proposition 6.1. Let \mathfrak{D}_0 be the collection of domains in \mathbb{R}^2 consisting of finite unions of non-degenerate rectangles with vertices in \mathbb{Q}^2 . Since \mathfrak{D}_0 is countable, given a sequence of N 's tending to infinity, Cantor's diagonal argument permits us to extract a strictly increasing subsequence $\{N_k\}_{k \geq 1}$ such that (6.1) holds for all $D \in \mathfrak{D}_0$. We thus have to show that the convergence extends (with the same subsequence) to all $D \in \mathfrak{D}$.

Let $D \in \mathfrak{D}$ and, for each $n \geq 1$, let D^n be the union of all open rectangles contained in D with vertices in $(2^{-n}\mathbb{Z})^2$. Since D is open, we have $D^n \neq \emptyset$ once n is sufficiently large and $D^n \uparrow D$ as $n \rightarrow \infty$. Moreover, if $f \in C_c(D \times \mathbb{R})$, then $\text{supp}(f) \subseteq \overline{D^n} \times [-n, n]$ for all n sufficiently large, say $n \geq n_f$. Hence, for every limit ζ^D of $(\zeta_{N_k}^D)$ along a subsubsequence (k_ℓ) and every $f \in C_c(D \times \mathbb{R})$,

$$\langle \zeta^D, f \rangle = \lim_{n \rightarrow \infty} \langle \zeta^{D^n}, e^{\Phi^{D, D^n}} f \rangle \quad (6.13)$$

by Lemma 3.3. As the right-hand side and hence ζ^D does not depend on the subsubsequence (k_ℓ) , it follows that $\zeta_{N_k}^D$ converges vaguely to ζ^D . \square

In order to show that the limit measure does not depend on the underlying sequence, we need to verify the conditions of Theorem 4.1. Most of these conditions have already been checked; one that still requires some work is the content of:

Lemma 6.4 *Let $D \in \mathfrak{D}$ and, for each $K \geq 1$ natural, let $\zeta^{K^{-1}D}$ be a subsequential weak limit of $\{\zeta_N^{K^{-1}D} : N \geq 1\}$, using the same subsequence as ensured by Proposition 6.1. Then for any*

open $A \subseteq D$ with $\text{Leb}(\partial A) = 0$ and $\bar{A} \subset D$, and for any $f \in C_c(\mathbb{R})$ satisfying $f \geq 0$,

$$\limsup_{\lambda \downarrow 0} \sup_{K \geq 1} \left| \frac{\mathbb{E} \left(K^4 \langle \zeta^{K^{-1}D}, 1_{K^{-1}A} \otimes f \rangle e^{-\lambda K^4 \langle \zeta^{K^{-1}D}, 1_{K^{-1}D} \otimes f \rangle} \right)}{\log(1/\lambda)} \right. \\ \left. - c_\star \int_0^\infty f(s) s e^{-\frac{s^2}{2\delta}} ds \int_A r^D(x)^2 dx \right| = 0, \quad (6.14)$$

where c_\star is the constant from Theorem 2.3.

Proof. Let $\tilde{f} = f_1 \otimes f_2$ for $f_1 \in C_c(\bar{D})$ and $f_2 \in C_c(\mathbb{R})$, and let $\{N_j\}_{j \geq 1}$ be the sequence defining the limiting measures $\{\zeta^{K^{-1}D} : K \geq 1\}$ in the statement. The proof hinges on the convergence statement

$$\langle \zeta_{[N_j/K]}^D, \tilde{f} \rangle \xrightarrow{j \rightarrow \infty} K^4 \langle \zeta^{K^{-1}D}, \tilde{f}_K \rangle, \quad (6.15)$$

where

$$\tilde{f}_K(x, t) := \tilde{f}(xK, t). \quad (6.16)$$

Indeed, noting that $(1_A \otimes f)_K = 1_{K^{-1}A} \otimes f$, we can rewrite the quantity in (6.14) with $1_A \otimes f$ replaced by \tilde{f} (while leaving the function in the exponent unchanged) as

$$\limsup_{\lambda \downarrow 0} \lim_{K \geq 1} \lim_{j \rightarrow \infty} \left| \frac{\mathbb{E} \left(\langle \zeta_{[N_j/K]}^D, \tilde{f} \rangle e^{-\langle \zeta_{[N_j/K]}^D, 1_D \otimes f \rangle} \right)}{\log(1/\lambda)} \right. \\ \left. - c_\star \int_{D \times [0, \infty)} s e^{-\frac{s^2}{2\delta}} r^D(x)^2 \tilde{f}(x, s) dx ds \right|, \quad (6.17)$$

The $j \rightarrow \infty$ limit can be bounded by the *limes superior* as $N \rightarrow \infty$ of the same expression with $[N_j/K]$ replaced by N . This *limes superior* no longer depends on K , rendering the supremum over $K \geq 1$ redundant. Moreover, the $N \rightarrow \infty$ limit vanishes by Lemma 5.4. Routine approximation arguments in conjunction with Lemma 5.6 then extend the limit to \tilde{f} replaced by $1_A \otimes f$ with $A \subseteq D$ open.

Moving to the proof of (6.15), we start by noting that, by Lemma 5.6, it suffices to do this for $f \in C_c(D \times \mathbb{R})$. Then there exists an open set B with $\bar{B} \subseteq D$ such that $\text{supp}(f) \subseteq B \times \mathbb{R}$. For all naturals $K \geq 1$ we then have

$$\{x \in D_N : xK/N \in B\} \subseteq D_{[N/K]} \subseteq (K^{-1}D)_N \quad (6.18)$$

once N is sufficiently large. Using the Gibbs-Markov property to write $h^{(K^{-1}D)_N} = h^{D_{\lfloor N/K \rfloor}} + \varphi^{(K^{-1}D)_N, D_{\lfloor N/K \rfloor}}$ and denoting

$$\begin{aligned} \tilde{f}_{N,K}(x, t) &:= K^4 e^{\alpha(m_{\lfloor N/K \rfloor} - m_N)} \frac{\log \lfloor N/K \rfloor}{\log N} \\ &\times e^{\alpha \varphi_x^{(K^{-1}D)_N, D_{\lfloor N/K \rfloor}}} \tilde{f} \left(x \frac{K \lfloor N/K \rfloor}{N}, t \sqrt{\frac{\log \lfloor N/K \rfloor}{\log N}} + \frac{m_N - m_{\lfloor N/K \rfloor}}{\sqrt{\log N}} - \frac{\varphi_x^{(K^{-1}D)_N, D_{\lfloor N/K \rfloor}}}{\sqrt{\log N}} \right), \end{aligned} \quad (6.19)$$

which depends on the sample of $\varphi^{(K^{-1}D)_N, D_{\lfloor N/K \rfloor}}$, then gives

$$\begin{aligned} K^4 \langle \zeta_N^{K^{-1}D}, \tilde{f}_K \rangle &= \frac{K^4}{\log N} \sum_{x \in (K^{-1}D)_N} e^{\alpha(h_x^{(K^{-1}D)_N} - m_N)} \tilde{f} \left(x \frac{K}{N}, \frac{m_N - h_x^{(K^{-1}D)_N}}{\sqrt{\log N}} \right) \\ &= \frac{1}{\log \lfloor N/K \rfloor} \sum_{x \in D_{\lfloor N/K \rfloor}} e^{\alpha(h_x^{D_{\lfloor N/K \rfloor}} - m_{\lfloor N/K \rfloor})} \tilde{f}_{N,K} \left(\frac{x}{\lfloor N/K \rfloor}, \frac{m_{\lfloor N/K \rfloor} - h_x^{D_{\lfloor N/K \rfloor}}}{\sqrt{\log \lfloor N/K \rfloor}} \right) \\ &= \langle \zeta_{\lfloor N/K \rfloor}^D, \tilde{f}_{N,K} \rangle \end{aligned} \quad (6.20)$$

provided N is so large that the left inclusion in (6.18) applies. Since $\alpha(m_N - m_{\lfloor N/K \rfloor}) \rightarrow 4 \log K$ as $N \rightarrow \infty$ and, thanks to $\max_{x \in B_N} \text{Var}(\varphi_x^{(K^{-1}D)_N, D_{\lfloor N/K \rfloor}}) \rightarrow 0$ as $N \rightarrow \infty$,

$$\max_{\substack{x \in D_N: \\ xK/N \in B}} |\varphi_x^{(K^{-1}D)_N, D_{\lfloor N/K \rfloor}}| \xrightarrow[N \rightarrow \infty]{} 0 \quad (6.21)$$

in probability, we now get that $\|\tilde{f}_{N,K} - \tilde{f}\|_\infty$ tends to zero in probability as $N \rightarrow \infty$. Since $\|\tilde{f}_{N,K} - \tilde{f}\|_\infty$ depends only on the binding field and not the underlying measure, this implies (6.15) as desired. \square

We are now ready for stating and proving the part of Theorem 2.3 dealing solely with the near-extremal process:

Theorem 6.5 (Near-extremal process convergence) *In the limit $N \rightarrow \infty$, the measures $\{\zeta_N^D: N \geq 1\}$ converge in law to the measure ζ^D in (2.10).*

Proof. For each $D \in \mathfrak{D}$, let ζ^D be a weak subsequential limit of $\{\zeta_N^D: N \geq 1\}$ along the same subsequence as ensured by Proposition 6.1. Pick $f \in C_c(\mathbb{R})$ non-negative with $\text{Leb}(f > 0) > 0$. Define the Borel measure

$$M^D(B) := \langle \zeta^D, 1_B \otimes f \rangle. \quad (6.22)$$

We will now check that M^D satisfies the assumptions of Theorem 4.1. That M^D is concentrated on D , of finite total mass and not charging Lebesgue null sets a.s. was shown in Lemma 6.2 (using that f is bounded). The factorization (4.2) over disjoint domains follows from the fact that if $D \cap \tilde{D} = \emptyset$ then $h^{D_N} \perp\!\!\!\perp h^{\tilde{D}_N}$. The Gibbs-Markov property

(4.3) was shown in Lemma 6.3. The uniform Laplace transform tail (4.4) was verified in Lemma 6.4 with constant c given by

$$c := c_\star \int_0^\infty f(t) t e^{-\frac{t^2}{2g}} dt. \quad (6.23)$$

Invoking (4.5) in Theorem 4.1 we conclude that, for each $f \in C_c(\mathbb{R})$ non-negative with non-zero total integral, the measure

$$W_f^D(B) := \left(c_\star \int_0^\infty f(t) t e^{-\frac{t^2}{2g}} dt \right)^{-1} \langle \zeta^D, 1_B \otimes f \rangle. \quad (6.24)$$

has the law of cLQG.

Moreover, since the Gibbs-Markov property (4.3) uses the same binding field for all f as above, Theorem 4.2 ensures that, for all $f_1, f_2 \in C_c(\mathbb{R})$ non-negative and not vanishing Lebesgue-a.e.,

$$W_{f_1}^D = W_{f_2}^D \quad \text{a.s.} \quad (6.25)$$

The separability of $C_c(\mathbb{R})$ permits us to choose the null event so that the equality holds for all $f_1, f_2 \in C_c(\mathbb{R})$ simultaneously a.s. Standard extension arguments then show that ζ^D takes the form (2.10). \square

6.2 Adding the extremal process.

Our next item of business is adding the convergence of the extremal process η_N^D to that of ζ_N^D . We start by recalling from [10] the structured version of the extremal process. We denote $\Lambda_r(x) := \{y \in \mathbb{Z}^2 : d_\infty(x, y) < r\}$. Given any positive sequence $\{r_N\}_{N \geq 1}$, define a measure on $D \times \mathbb{R} \times \mathbb{R}^{\mathbb{Z}^2}$ by

$$\tilde{\eta}_N^D := \sum_{x \in D_N} \mathbf{1}_{\{h_x^{D_N} = \max_{z \in \Lambda_{r_N}(x)} h_z^{D_N}\}} \delta_{x/N} \otimes \delta_{h_x^{D_N} - m_N} \otimes \delta_{\{h_x^{D_N} - h_{x+z}^{D_N} : z \in \mathbb{Z}^2\}}. \quad (6.26)$$

This measure records the positions, centered values and configuration near and relative to the r_N -local maxima of h^{D_N} . Our aim is to show:

Theorem 6.6 (Joint convergence of near-extremal and extremal process) *For any $D \in \mathfrak{D}$ and any $\{r_N\}_{N \geq 1}$ with $r_N \rightarrow \infty$ and $N/r_N \rightarrow \infty$,*

$$(\zeta_N^D, \tilde{\eta}_N^D) \xrightarrow[N \rightarrow \infty]{\text{law}} (\zeta^D, \tilde{\eta}^D), \quad (6.27)$$

where ζ^D is derived from a sample Z^D of cLQG in D as in (2.10) and, conditional on Z^D ,

$$\tilde{\eta}^D = \text{PPP} \left(\bar{c} Z^D(dx) \otimes e^{-\alpha h} dh \otimes \nu(d\phi) \right) \quad (6.28)$$

where \bar{c} and ν are as in (1.4). More precisely, the joint law of $(\zeta^D, \tilde{\eta}^D)$ is such that, for each non-negative $f \in C_c(D \times \mathbb{R})$ and $\tilde{f} \in C_c(D \times \mathbb{R} \times \mathbb{R}^{\mathbb{Z}^2})$,

$$E\left(e^{-\langle \zeta^D, f \rangle - \langle \tilde{\eta}^D, \tilde{f} \rangle}\right) = E\left(\exp\left\{-c_* \int_D Z^D(dx) \int_0^\infty dt t e^{-\frac{t^2}{2s}} f(x, t) - \bar{c} \int_D Z^D(dx) \int_{\mathbb{R}} dh e^{-\alpha h} \int_{\mathbb{R}^{\mathbb{Z}^2}} \nu(d\phi) (1 - e^{-\tilde{f}(x, h, \phi)})\right\}\right), \quad (6.29)$$

where the expectation on the right is over the law of Z^D .

Thanks to Theorem 2.1 of [10] and to Lemma 5.3, the family of measure-valued pairs $\{(\zeta_N^D, \tilde{\eta}_N^D) : N \geq 1\}$ is tight. We may thus extract a (joint) subsequential weak limit to be called, with some abuse of notation, $(\zeta^D, \tilde{\eta}^D)$ for the remainder of this subsection. The marginal distributions are known from Theorem 2.1 of [10] and from Theorem 6.5. The main task is to characterize the joint law of this pair uniquely.

One way to achieve this is to follow the proofs of [8–10] while keeping track of the behavior of the near-extremal limit process throughout the manipulations. We will instead proceed by a direct argument that consists of two steps: We extract the spatial part of the intensity measure of $\tilde{\eta}^D$ by way of a suitable limit and then equate it with the spatial part of the measure ζ^D . The first step is the content of:

Lemma 6.7 *For all $D \in \mathfrak{D}$, the vague limit in probability (under the joint law of ζ^D and $\tilde{\eta}^D$)*

$$\vartheta^D(\cdot) := \lim_{\epsilon \rightarrow 0} \lim_{s \rightarrow \infty} \frac{1}{\epsilon} e^{-\alpha s} \tilde{\eta}^D(\cdot \times (-s - \epsilon, -s) \times \mathbb{R}^{\mathbb{Z}^2}) \quad (6.30)$$

exists and is equal in law to $\bar{c}Z^D$. Moreover, the conditional distribution of $\tilde{\eta}^D$ given ϑ^D is that of a Poisson point process with intensity $\vartheta^D \otimes e^{-\alpha h} dh \otimes \nu(d\phi)$ and, for any $f \in C(D)$,

$$\lim_{\epsilon \rightarrow 0} \lim_{s \rightarrow \infty} \frac{1}{\epsilon} e^{-\alpha s} \tilde{\eta}^D(dx \times (-s - \epsilon - f(x), -s - f(x)) \times \mathbb{R}^{\mathbb{Z}^2}) = e^{\alpha f(x)} \vartheta^D(dx) \quad (6.31)$$

vaguely in probability.

Proof. By [10, Theorem 2.1], $\tilde{\eta}^D$ is the Cox process $\text{PPP}(W^D(dx) \otimes e^{-\alpha h} dh \otimes \nu(d\phi))$, where W^D is a random Radon measure on \bar{D} with $W^D \stackrel{\text{law}}{=} \bar{c}Z^D$. Conditionally on W^D , for each $s > 0$, each $\epsilon > 0$ and each $B \subseteq D$ measurable,

$$\begin{aligned} \tilde{\eta}^D\left(\left\{(x, h, \phi) \in B \times \mathbb{R} \times \mathbb{R}^{\mathbb{Z}^2} : h \in (-s - \epsilon - f(x), -s - f(x))\right\}\right) \\ = \text{Poisson}\left(\int_B W^D(dx) \alpha^{-1} e^{\alpha(s+f(x))} (e^{\alpha\epsilon} - 1)\right). \end{aligned} \quad (6.32)$$

The claim follows from Fubini-Tonelli and the fact that $\lambda^{-1} \text{Poisson}(a\lambda) \rightarrow a$ in probability as $\lambda \rightarrow \infty$. \square

Let $\tilde{\zeta}^D := \zeta^D(\cdot \times \mathbb{R})$. In order to identify the joint law of $(\tilde{\zeta}^D, \vartheta^D)$, where ϑ^D is as in (6.30), we will rely on Theorem 4.2 for which we need:

Lemma 6.8 (Joint Gibbs-Markov property) *For any $D, \tilde{D} \in \mathfrak{D}$ satisfying $\tilde{D} \subseteq D$ and $\text{Leb}(D \setminus \tilde{D}) = 0$, and any subsequential limits along the same sequence of N 's,*

$$\left(\tilde{\zeta}^D(\mathrm{d}x), \vartheta^D(\mathrm{d}\tilde{x}) \right) \stackrel{\text{law}}{=} \left(e^{\alpha\Phi^{D,\tilde{D}}(x)} \tilde{\zeta}^{\tilde{D}}(\mathrm{d}x), e^{\alpha\Phi^{D,\tilde{D}}(\tilde{x})} \vartheta^{\tilde{D}}(\mathrm{d}\tilde{x}) \right), \quad (6.33)$$

where $\Phi^{D,\tilde{D}} = \mathcal{N}(0, C^{D,\tilde{D}})$ is independent of $(\tilde{\zeta}^{\tilde{D}}, \vartheta^{\tilde{D}})$ on the right-hand side.

Proof. By the Gibbs-Markov property (Lemma 3.3) for the DGFF h^{D_N} , there is a coupling of h^{D_N} and $h^{\tilde{D}_N}$ such that $h^{D_N} = \varphi^{D_N, \tilde{D}_N} + h^{\tilde{D}_N}$, with the fields on the right-hand side being independent. Let $\tilde{\zeta}_N^D, \tilde{\eta}_N^D$ be derived from h^{D_N} , and $\tilde{\zeta}_N^{\tilde{D}}, \tilde{\eta}_N^{\tilde{D}}$ from $h^{\tilde{D}_N}$, using any $\{r_N\}_{N \geq 1}$ with $r_N \rightarrow \infty$ and $N/r_N \rightarrow \infty$. We abbreviate $\varphi^N := \varphi^{D_N, \tilde{D}_N}$.

For any $f \in C_c(\tilde{D})$, we have

$$\langle \tilde{\zeta}_N^D, f \rangle = \langle \tilde{\zeta}_N^{\tilde{D}}, e^{\alpha\varphi^N} f \rangle \quad (6.34)$$

where we used that f does not depend on the field coordinate and so is unaffected by the shift by φ^N . For the extremal process, we will focus on quantities of the form

$$\langle \tilde{\eta}_N^D, \tilde{f} \otimes 1_{(-s-\epsilon, -s)} \otimes 1 \rangle \quad (6.35)$$

where $\tilde{f} \in C_c(\tilde{D})$. Our aim is to relate (6.35) to

$$\langle \tilde{\eta}_N^{\tilde{D}}, \tilde{f} \otimes 1_{(-s-\epsilon-\Phi^{D,\tilde{D}}, -s-\Phi^{D,\tilde{D}})} \otimes 1 \rangle \quad (6.36)$$

where $\Phi^{D,\tilde{D}}$ is the continuum binding field (regarded as independent of $\eta_N^{\tilde{D}}$ on the right hand side. Besides the fact that the test function is not continuous in the field variable, here we are faced with the additional difficulty that the additive term φ^N may change the spatial positions of relevant r_N -local extrema.

Let $a > 0$ be such that $\text{supp}(f) \subset \tilde{D}^a := \{x \in \tilde{D} : \text{dist}(x, D^c) > a\}$. To control the fluctuation of the binding field at scale $2r_N$, we define

$$L_N := \max_{\substack{x, y \in \tilde{D}_N^a \\ d_\infty(x, y) \leq 2r_N}} |\varphi_x^N - \varphi_y^N|. \quad (6.37)$$

Given $\epsilon > 0$ and $s > 0$, for each $N \geq 1$ and each $\delta > 0$ consider the event

$$F_N := \bigcap_{x \in \tilde{D}_N^a} \left\{ h_x^{D_N} \geq m_N - s - \epsilon \Rightarrow \max_{y \in \Lambda_{2r_N}(x) \setminus \Lambda_{r_N/2}(x)} h_y^{D_N} \leq h_x^{D_N} - \delta \right\}. \quad (6.38)$$

By the ‘‘gap estimate’’ in Theorem 9.2 of [6], we have

$$\lim_{N \rightarrow \infty} \mathbb{P}(F_N^c) = 0 \quad (6.39)$$

while

$$\lim_{N \rightarrow \infty} \mathbb{P}(L_N > \delta) = 0 \quad (6.40)$$

by the coupling to the continuum binding field in Lemma 3.5.

On F_N , for each $x \in \tilde{D}_N$ with $h_x^{D_N} \geq m_N - s - \epsilon$ there exists a.s.-unique $y_x \in \Lambda_{r_N/2}(x)$ where $h^{\tilde{D}_N}$ achieves its maximum on $\Lambda_{r_N}(y_x)$. On $F_N \cap \{L_N \leq \delta\}$,

$$h_x^{D_N} = h_x^{\tilde{D}_N} + \varphi_x^N \leq h_{y_x}^{\tilde{D}_N} + \varphi_{y_x}^N + \delta = h_{y_x}^{D_N} + \delta \leq h_x^{D_N} + \delta \quad (6.41)$$

and thus $|h_x^{\tilde{D}_N} - h_{y_x}^{\tilde{D}_N} - \varphi_{y_x}^N| \leq \delta$ hold for all $x \in \tilde{D}_N^a$ with $h_x^{D_N} \geq m_N - s - \epsilon$. Denoting $\text{osc}_u(\tilde{f}) := \sup\{|\tilde{f}(y) - \tilde{f}(x)| : |y - x| < u\}$ and assuming $\tilde{f} \geq 0$, we thus have

$$\begin{aligned} \langle \tilde{\eta}_N^D, \tilde{f} \otimes 1_{(-s-\epsilon, -s)} \otimes 1 \rangle &\leq \text{osc}_{r_N/N}(\tilde{f}) \eta_N(D \times (-s - \epsilon, -s) \times \mathbb{R}) \\ &\quad + \langle \tilde{\eta}_N^{\tilde{D}}, \tilde{f} \otimes 1_{(-s-\epsilon-2\delta-\Phi^{D,\tilde{D}}, -s+2\delta-\Phi^{D,\tilde{D}})} \otimes 1 \rangle, \end{aligned} \quad (6.42)$$

where we also used Lemma 3.5 to couple φ^N to a continuum binding field so that $E_N := \max_{x \in D_N^a} |\varphi_{[xN]}^N - \Phi_x^{D,\tilde{D}}| \leq \delta$.

We now take $N \rightarrow \infty$ along the subsequence such that (taking further subsequence if necessary) the pair $(\tilde{\eta}_N^D, \tilde{\eta}_N^{\tilde{D}})$ converges to $(\tilde{\eta}^D, \tilde{\eta}^{\tilde{D}})$ in law. Relying on (6.39–6.40) and also Lemma 3.5 to eliminate the contribution of $F_N^c \cup E_N^c \cup \{L_N > \delta\}$, and using that the limit measure $\tilde{\eta}^{\tilde{D}}$ is continuous in the second coordinate we then take also $\delta \downarrow 0$ and get the pointwise comparison

$$\langle \tilde{\eta}^D, \tilde{f} \otimes 1_{(-s-\epsilon, -s)} \otimes 1 \rangle \leq \langle \tilde{\eta}^D, \tilde{f} \otimes 1_{(-s-\epsilon-\Phi^{D,\tilde{D}}, -s-\Phi^{D,\tilde{D}})} \otimes 1 \rangle \quad \text{a.s.} \quad (6.43)$$

A completely analogous argument, which we skip for brevity, shows that “ \geq ” holds as well. Combining with (6.34), we conclude

$$\begin{aligned} &\left(\tilde{\zeta}^D(dx), \tilde{\eta}^D(d\tilde{x} \times (-s - \epsilon, -s) \times \mathbb{R}^{\mathbb{Z}^2}) \right)_{s \in \mathbb{Q}_+} \\ &\stackrel{\text{law}}{=} \left(e^{\alpha \Phi^{D,\tilde{D}}(x)} \tilde{\zeta}^{\tilde{D}}(dx), \tilde{\eta}^D(d\tilde{x} \times (-s - \Phi^{D,\tilde{D}}(\tilde{x}) - \epsilon, -s - \Phi^{D,\tilde{D}}(\tilde{x})) \times \mathbb{R}^{\mathbb{Z}^2}) \right)_{s \in \mathbb{Q}_+} \end{aligned} \quad (6.44)$$

where $\Phi^{D,\tilde{D}}$ is independent of $(\tilde{\zeta}^{\tilde{D}}, \tilde{\eta}^{\tilde{D}})$ on the right-hand side. The assertion now follows from Lemma 6.7. \square

We are ready to give:

Proof of Theorem 6.6. Let $(\tilde{\zeta}^D, \vartheta^D)$ be related to the subsequential weak-limit pair of measures $(\zeta^D, \tilde{\eta}^D)$ by letting $\tilde{\zeta}^D(\cdot) := \zeta^D(\cdot \times \mathbb{R})$ and defining ϑ^D from $\tilde{\eta}^D$ is as in (6.30). Since the law of ϑ^D is known from Lemma 6.7, ϑ^D satisfies all of the conditions of Theorem 4.1 with $c := \bar{c}$ in (4.4). The same has been shown with $c := gc_\star$ for $\tilde{\zeta}^D$ in the proof of Theorem 6.5. The joint validity of the Gibbs-Markov property holds true by Lemma 6.8. Theorem 4.2 thus implies

$$\bar{c}^{-1} \vartheta^D = (gc_\star)^{-1} \tilde{\zeta}^D \quad \text{a.s.} \quad (6.45)$$

Denote $Z^D := (gc_\star)^{-1} \tilde{\zeta}^D$ and note that, by Lemma 6.7, this measure has the law of the cLQG. Theorem 6.5 then shows that ζ^D is determined completely by the sample of Z^D and, in fact,

$$\zeta^D(dxdt) = c_\star Z^D(dx) \otimes 1_{(0,\infty)}(t) t e^{-\frac{t^2}{2g}} dt. \quad (6.46)$$

Lemma 6.7 in turn gives (6.28), with the Poisson process conditionally independent of Z^D . Hence, the subsequential limit (ζ^D, η^D) obeys (6.29). The limit object does not depend on the chosen subsequence, so we also get full convergence. \square

6.3 Including the field itself.

As our last item left to do, we need to include the DGFF itself in the distributional convergence. This is based on the following observation:

Lemma 6.9 *The weak limit*

$$(\zeta_N^D, \tilde{\eta}_N^D, h^{D_N}) \xrightarrow[N \rightarrow \infty]{\text{law}} (\zeta^D, \tilde{\eta}^D, h^D) \quad (6.47)$$

exists with the law of the right-hand side determined by

$$\begin{aligned} & E(e^{-\langle \zeta^D, f_1 \rangle - \langle \tilde{\eta}^D, f_2 \rangle + h^D(f_3)}) \\ &= e^{\frac{1}{2}\langle f_3, \widehat{G}^D f_3 \rangle} E\left(\exp\left\{-c_\star \int_D Z^D(dx) e^{\alpha(\widehat{G}^D f_3)(x)} \int_0^\infty dt t e^{-\frac{t^2}{2s}} f_1(x, t) \right. \right. \\ &\quad \left. \left. - \bar{c} \int_D Z^D(dx) e^{\alpha(\widehat{G}^D f_3)(x)} \int_{\mathbb{R}} dh e^{-\alpha h} \int_{\mathbb{R}^{\mathbb{Z}^2}} \nu(d\phi) (1 - e^{-f_2(x, h, \phi)})\right\}\right), \quad (6.48) \end{aligned}$$

for each non-negative $f_1 \in C_c(D \times \mathbb{R})$ and $f_2 \in C_c(D \times \mathbb{R} \times \mathbb{R}^{\mathbb{Z}^2})$ and arbitrary $f_3 \in C_0(D)$.

Proof. Under the measure

$$A \mapsto \mathbb{P}(e^{h^{D_N}(f_3) - \frac{1}{2}\langle f_3, G^{D_N} f_3 \rangle} 1_A) \quad (6.49)$$

the DGFF h^{D_N} has the law of $h^{D_N} + G^{D_N} f_3$ under \mathbb{P} . Since $G^{D_N} f_3$, with its argument scaled by N , tends uniformly to $\widehat{G}^D f_3$ as $N \rightarrow \infty$, we get

$$\mathbb{E}(e^{-\langle \zeta_N^D, f_1 \rangle - \langle \tilde{\eta}_N^D, f_2 \rangle + h^{D_N}(f_3)}) = o(1) + e^{\frac{1}{2}\langle f_3, \widehat{G}^D f_3 \rangle} \mathbb{E}(e^{-\langle \zeta_N^D, e^{\alpha \widehat{G}^D f_3} f_1 \rangle - \langle \tilde{\eta}_N^D, f_2 \circ \theta_{\widehat{G}^D f_3} \rangle}), \quad (6.50)$$

where $(f_2 \circ \theta_s)(x, h, \phi) := f_2(x, h(x) + s(x), \phi)$ and $o(1) \rightarrow 0$ as $N \rightarrow \infty$. The joint convergence of $(\zeta_N^D, \tilde{\eta}_N^D)$ to $(\zeta^D, \tilde{\eta}^D)$ then shows

$$E(e^{-\langle \zeta_N^D, f_1 \rangle - \langle \tilde{\eta}_N^D, f_2 \rangle + h^{D_N}(f_3)}) \xrightarrow[N \rightarrow \infty]{} e^{\frac{1}{2}\langle f_3, \widehat{G}^D f_3 \rangle} E(e^{-\langle \zeta^D, e^{\alpha \widehat{G}^D f_3} f_1 \rangle - \langle \tilde{\eta}^D, f_2 \circ \theta_{\widehat{G}^D f_3} \rangle}). \quad (6.51)$$

This proves the existence of the joint limit in (6.47). Invoking (6.29) along with a routine change of variables, we then get (6.48) as well. \square

We are finally ready to give:

Proof of Theorem 2.3. Lemma 6.9 shows the weak convergence of $(\zeta_N^D, \tilde{\eta}_N^D, h^{D_N})$ as $N \rightarrow \infty$. Moreover, (6.48) with $f_2 := 0$ and $f_1(x, t) := cg(x)$ for a suitable constant $c > 0$ shows that the law of Z^D underlying the limit processes ζ^D and $\tilde{\eta}^D$ obeys

$$E(e^{-\langle Z^D, g \rangle + h^D(f)}) = e^{\frac{1}{2}\langle f, \widehat{G}^D f \rangle} \mathbb{E}(e^{-\langle e^{\alpha \widehat{G}^D f} Z^D, g \rangle}) \quad (6.52)$$

for all $f \in C_0(D)$ all non-negative $g \in C_c(D)$. Theorem 4.3 implies that Z^D is, up to a modification on a set of vanishing probability, the cLQG associated with h^D .

This proves the claim for the structured extremal process; the reduction to the unstructured process is then the same as in the proof of [10, Corollary 2.2]. \square

ACKNOWLEDGMENTS

This project has been supported in part by the NSF award DMS-1712632, ISF grant No. 1382/17 and BSF award 2018330. The second author has also been supported in part by a Zeff Fellowship and a Minerva Fellowship at the Technion.

REFERENCES

- [1] Y. Abe and M. Biskup, Exceptional points of two-dimensional random walks at multiples of the cover time. arXiv: 1903.04045
- [2] Y. Abe, M. Biskup and S. Lee, Exceptional points of discrete-time random walks in planar domains. arXiv:1911.11810
- [3] J. Aru, E. Powell and A. Sepúlveda (2019). Critical Liouville measure as a limit of subcritical measures. *Electron. Commun. Probab.* **24**, paper no. 18, 16 pp.
- [4] J. Barral, A. Kupiainen, M. Nikula, E. Saksman, C. Webb (2015). Basic properties of critical lognormal multiplicative chaos. *Ann. Probab.* **43**, no. 5, 2205–2249
- [5] N. Berestycki, C. Garban, R. Rhodes and V. Vargas (2016). KPZ formula derived from Liouville heat kernel. *J. Lond. Math. Soc.* **94**, no. 1, 186–208.
- [6] M. Biskup (2020). Extrema of the two-dimensional Discrete Gaussian Free Field. In: M. Barlow and G. Slade (eds.): *Random Graphs, Phase Transitions, and the Gaussian Free Field*. SSPROB 2017. Springer Proceedings in Mathematics & Statistics, vol 304, pp 163–407. Springer, Cham.
- [7] M. Biskup, S. Gufler and O. Louidor (2020). Support sets and KPZ relation for critical Liouville Quantum Gravity. In preparation.
- [8] M. Biskup and O. Louidor (2016). Extreme local extrema of two-dimensional discrete Gaussian free field. *Commun. Math. Phys.* **345** 271-304.
- [9] M. Biskup and O. Louidor (2020). Conformal symmetries in the extremal process of two-dimensional discrete Gaussian free field. *Commun. Math. Phys.* **375**, no. 1, 175–235.
- [10] M. Biskup and O. Louidor (2018). Full extremal process, cluster law and freezing for two-dimensional discrete Gaussian free field. *Adv. Math.* **330** 589-687.
- [11] M. Biskup and O. Louidor (2019). On intermediate level sets of two-dimensional discrete Gaussian free field. *Ann. Inst. Henri Poincaré* **55**, no. 4, 1948–1987.
- [12] M. Bramson, J. Ding and O. Zeitouni (2016). Convergence in law of the maximum of the two-dimensional discrete Gaussian free field. *Commun. Pure Appl. Math* **69**, no. 1, 62–123.
- [13] M. Bramson and O. Zeitouni (2011). Tightness of the recentered maximum of the two-dimensional discrete Gaussian free field. *Comm. Pure Appl. Math.* **65**, 1–20.
- [14] F. David, A. Kupiainen, R. Rhodes and V. Vargas (2016). Liouville Quantum Gravity on the Riemann sphere. *Commun. Math. Phys.* **342**, no. 3, 869–907.
- [15] J. Ding and O. Zeitouni (2014). Extreme values for two-dimensional discrete Gaussian free field. *Ann. Probab.* **42**, no. 4, 1480–1515
- [16] B. Duplantier, R. Rhodes, S. Sheffield and V. Vargas (2014). Critical Gaussian multiplicative chaos: Convergence of the derivative martingale. *Ann. Probab.* **42**, no. 5, 1769–1808.
- [17] B. Duplantier, R. Rhodes, S. Sheffield and V. Vargas (2014). Renormalization of critical Gaussian multiplicative chaos and KPZ formula. *Commun. Math. Phys.* **330**, no. 1, 283–330.
- [18] B. Duplantier and S. Sheffield (2011). Liouville quantum gravity and KPZ. *Invent. Math.* **185**, 333-393.
- [19] S. Gufler and O. Louidor (2020). Ballot theorems for the two-dimensional discrete Gaussian free field. In preparation.
- [20] J. Junnila and E. Saksman (2017). Uniqueness of critical Gaussian chaos. *Elect. J. Probab* **22**, 1–31.
- [21] J.-P. Kahane (1985). Sur le chaos multiplicatif. *Ann. Sci. Math. Québec* **9**, no.2, 105–150.

- [22] G. Kozma and E. Schreiber (2004). An asymptotic expansion for the discrete harmonic potential. *Electron. J. Probab.* **9**, Paper no. 1, pages 10–17.
- [23] G.F. Lawler (1991). *Intersections of Random Walks*. Birkhäuser Boston, Inc., Boston, MA.
- [24] G.F. Lawler and V. Limic (2010). *Random walk: a modern introduction*. Cambridge Studies in Advanced Mathematics, vol. 123. Cambridge University Press, Cambridge, xii+364.
- [25] T.M. Liggett (1978). Random invariant measures for Markov chains, and independent particle systems. *Z. Wahrsch. Verw. Gebiete* **45** 297–313.
- [26] R. Rhodes and V. Vargas (2014). Gaussian multiplicative chaos and applications: A review. *Probab. Surveys* **11**, 315–392
- [27] R. Rhodes and V. Vargas (2011). KPZ formula for log-infinitely divisible multifractal random measures. *ESAIM: Probability and Statistics* **15**, 358–371.
- [28] R. Robert and V. Vargas (2010). Gaussian multiplicative chaos revisited. *Ann. Probab.* **38**, no. 2, 605–631.
- [29] E. Powell (2018). Critical Gaussian chaos: convergence and uniqueness in the derivative normalisation. *Electron. J. Probab.* **23**, paper no. 31, 26 pp.
- [30] A. Shamov (2016). On Gaussian multiplicative chaos. *J. Funct. Anal.* **270**, no. 9, 3224–3261