

Math 181

Lecture 9

Pricing Options for the Random Walk with Discrete Steps

Consider a single step in a random walk

$$S_0 \text{ given at } t = 0$$
$$S_1 = \begin{cases} uS_0 & \text{probability } p' \\ dS_0 & \text{probability } q' = 1-p' \end{cases}$$

in which

$$u = e^{\mu dt + \sigma\sqrt{dt}}$$
$$d = e^{\mu dt - \sigma\sqrt{dt}}$$

Also consider an option where value is

$$f_0 \quad \text{at } t = 0$$
$$f_u \quad \text{at } t = 1, \text{ if step is up}$$
$$f_d \quad \text{at } t = 1, \text{ if step is down.}$$

This is summarized in in Figure 1.

For example, if the option is a call with a strike price X such that $dS_0 < X < uS_0$, and if t_1 is the expiration time, then

$$f_u = uS_0 - X$$
$$f_d = 0.$$

It is tempting to think that the option price f_0 at t_0 is the expected value of the future option price, suitably discounted. Discounting the expected value f_1 back to $t = 0$ gives

$$\tilde{f}_0 = e^{-rdt} E(f_1) = e^{-rdt} (p' f_u + (1 - p') f_d). \quad (1)$$

As shown below, however, this price for f would not be arbitrage free!

In order to correctly price f , we perform an arbitrage-free analysis by considering two portfolios




Figure 1: A single step of a binary random walk, with stock price S , option price f and probabilities p' and q' of going up (u) or down (d).

A: long Δ shares of stock, each with value S_0 ; short one option of value f_0

B: cash K

at time t_0 . We will choose Δ so that there is no risk in the portfolio A. Then we'll choose K to make B equivalent to A . The value of A at t_0 and t_1 is

$$A(t_0) = \Delta S_0 - f_0$$

$$A(t_1) = \begin{cases} \Delta u S_0 - f_u & \text{if up} \\ \Delta d S_0 - f_d & \text{if down} \end{cases}$$

The profile will be riskless if the two outcomes are identical, i.e. if

$$\Delta u S_0 - f_u = \Delta d S_0 - f_d$$

i.e. if

$$\Delta = \frac{f_u - f_d}{u S_0 - d S_0}. \quad (2)$$

Now for portfolio B, take an amount of cash K equal to the initial value of A , i.e. at $t = t_0$,

$$B(t_0) = K = \Delta S_0 - f_0.$$

Then at $t = t_1$

$$B(t_1) = K e^{r dt}.$$

Since portfolio A is riskless and its initial value is the same as B's, then its value at t_1 is also the same; otherwise there would be an arbitrage opportunity. This shows that

$$\begin{aligned} A(t_1) &= B(t_1) \\ &= e^{rdt} B(t_0) \\ &= e^{rdt} A(t_0). \end{aligned}$$

In the future, we will dispense with the second portfolio. We will summarize this argument by saying that since A is riskfree, its value must grow at the riskfree rate r .

Now insert the values of $A(t_0)$, $A(t_1)$ and Δ to obtain

$$(\Delta u S_0 - f_u) = e^{rdt} (\Delta S_0 - f_0)$$

which leads to

$$f_0 = e^{-rdt} (p f_u + (1 - p) f_d) \tag{3}$$

in which

$$p = \frac{e^{rdt} - d}{u - d}. \tag{4}$$

The formula (3) is a simplified version of the Black-Scholes equation for pricing an option. It has some surprising properties:

- (i) The formula (3) for f_0 looks like a discounted average, as in (1). But the probability p in (3) is not the probability p' in for the random walk to move up. In fact p is determined just from u, d, r and dt .
- (ii) The formula is the same no what type of option is involved. It could be a call or put. The values of f_u and f_d would be different, but the relation between f_0 and these values would be the same.
- (iii) The probabilities p' and q' , for the stock to move up or down, do not influence the option price! They do not appear in (3) or (4). This says that the likelihood of the stock to move up or down does not affect the value of the option.

Although this is surprising, it has an explanation. The argument above shows that the risk in the option value is entirely due to risk in the stock, since a portfolio of stock and options can be riskless. Since the risks in the stock are already accounted for in its price, they do not enter into the price of the option.

This pricing method for the option is called “risk-neutral valuation”.




Figure 2: Two steps of a binary random walk, with stock price S , option price f and probabilities p' and q' of going up (u) or down (d).

Math 181

Lecture 10

Pricing Options on a Binomial Tree

The price of an option on a binomial tree can be determined from its values at expiration, by repeated application of the single step formulas from Lecture 9. First we perform this for a 2 step tree, then we generalize to an arbitrary sized tree.

Consider the two step tree in Figure 2.

In this formulation, the values S, u, d are known. They are the initial stock value, and the proportional increase or decrease in its value. We take the final time to be the exercise time for the option; so that f_{uu}, f_{ud}, f_{dd} are also known. (Note that $f_{du} = f_{ud}$.) We wish to compute f , and along the way we will compute f_u and f_d .

Computation of f_u and f_d performed on one step trees, as shown in Figure 3.

The value f_u is computed as a suitable discounted average of f_{uu} and f_{du} as in (3), (4) by the substitution (f_{uu}, f_{du}, f_n) for (f_u, f_d, f_0) . The result is

$$f_u = e^{-rdt}(pf_{uu} + (1-p)f_{du})$$




Figure 3: One step trees used for computation of f_u (left) and f_d (right).

$$p = \frac{e^{rdt} - d}{u - d}.$$

In a similar way

$$f_d = e^{-rdt}(pf_{ud} + (1 - p)f_{dd})$$

with the same p .

Now we compute f using the tree shown in Figure 4.

$$\begin{aligned} f &= e^{-rdt}(pf_u + (1 - p)f_d) \\ &= e^{-r2dt} \{p(pf_{uu} + (1 - p)f_{ud}) + (1 - p)(pf_{ud} + (1 - p)f_{dd})\} \\ &= e^{-r2dt} \{p^2 f_{uu} + 2p(1 - p)f_{ud} + (1 - p)^2 f_{dd}\}. \end{aligned}$$

Note that

$$\begin{aligned} p^2 + 2p(1 - p) + (1 - p)^2 &= (p + (1 - p))^2 \\ &= 1 \end{aligned}$$

so that this formula is still a discounted average.

Risk-Neutral Valuation

We now give a tree-interpretation of these formulas. The real random walk has probabilities $p', q' = (1 - p')$ of stepping up or down. We replace these




Figure 4: One step tree used for computation of f from f_u and f_d .

probabilities with the probabilities that occur in the option pricing formula, as shown in Figure 5.

The resulting random walk is called the “risk-neutral process”. This is not the real process. It has the same states S, uS, dS , etc., but they occur with different probabilities.

The usefulness of the risk-neutral process is that it can be directly used to calculate the option price as an average (with discounting).

For a single step the tree is drawn in Figure 6.

and the price is

$$\begin{aligned} f_0 &= e^{-rdt} \bar{E}(f_1) \\ &= e^{-rdt} (pf_u + (1-p)f_d) \end{aligned}$$

in which \bar{E} denotes average with respect to the risk-neutral process.

For the two step process the same formulas work:

$$\begin{aligned} f_0 &= e^{-r2dt} \bar{E}(f_2) \\ &= e^{-r2dt} (p^2 f_{uu} + 2p(1-p)f_{ud} + (1-p)^2 f_{dd}). \end{aligned}$$

More generally its true that for $m < n$

$$f_m = e^{-r(t_n - t_m)} \bar{E}(f_n). \quad (5)$$

This is the risk-neutral valuation formula.




Figure 5: One step of tree with risk-neutral probabilities p and $q = 1 - p$.




Figure 6: One step tree, with risk-neutral probabilities used for computation of f from f_u and f_d .