

PROBLEM SET 4

DUE FRIDAY, MAY 3

**Problem 1.** For each of the following inner product spaces  $V$  and linear operators  $T$  on  $V$ , evaluate  $T^*$  at the given vector in  $V$ .

- (1)  $V = \mathbb{R}^2$ ,  $T(a, b) = (2a + b, a - 3b)$  and  $x = (3, 5)$ .
- (2)  $V = \mathbb{C}^2$ ,  $T(z_1, z_2) = (2z_1 + iz_2, (1 - i)z_1)$  and  $x = (3 - i, 1 + 2i)$ .
- (3)  $V = P_1(\mathbb{R})$  with the inner product  $\langle f, g \rangle = \int_{-1}^1 f(t)g(t)dt$ ,  $T(f) = f' + 3f$ ,  $f(t) = 4 - 2t$ .

**Problem 2.** Let  $T$  be a linear operator on an inner product space  $V$ .

- (1) Let  $U_1 = T + T^*$  and  $U_2 = TT^*$ . Prove that  $U_1 = U_1^*$  and  $U_2 = U_2^*$ .
- (2) Prove that  $T^*T = T_0$  implies  $T = T_0$ .

Is the same result true if we assume that  $TT^* = T_0$ ?

**Problem 3.** Give an example of a linear operator  $T$  on an inner product space  $V$  such that  $N(T) \neq N(T^*)$ .

**Problem 4.** Let  $V$  be a finite-dimensional inner product space, and let  $T$  be a linear operator on  $V$ . Prove that if  $T$  is invertible, then  $T^*$  is invertible and  $(T^*)^{-1} = (T^{-1})^*$ .

**Problem 5.** Let  $V$  be a finite-dimensional vector space.

**Definition.** If  $V = W_1 \oplus W_2$ , then a linear operator  $T$  on  $V$  is the *projection on  $W_1$  along  $W_2$*  if, whenever  $x = x_1 + x_2$  with  $x_1 \in W_1$  and  $x_2 \in W_2$ , then we have  $T(x) = x_1$ .

We say that  $T$  is a *projection* if it is a projection on  $W_1$  along  $W_2$  for *some* subspaces  $W_1$  and  $W_2$  of  $V$ .

- (1) Show that  $R(T) = W_1$  and  $N(T) = W_2$ .  
Hence  $V = R(T) \oplus N(T)$ .
- (2) Prove that  $T \in \mathcal{L}(V)$  is a projection if and only if  $T = T^2$ .

**Problem 6.** Let  $V$  be a finite dimensional inner product space, and let  $W$  be a subspace.

- (1) Prove that  $V = W \oplus W^\perp$ .
- (2) Show that if  $T$  is a projection on  $W$  along  $W^\perp$ , then  $T = T^*$ .

**Problem 7.** Let  $T$  be a linear operator on an inner product space  $V$ . Prove that  $\|T(x)\| = \|x\|$  for all  $x \in V$  if and only if  $\langle T(x), T(y) \rangle = \langle x, y \rangle$  for all  $x, y \in V$ .

**Problem 8.** Let  $V$  be a finite-dimensional inner product space, and let  $T$  be a linear operator on  $V$ . Prove the following results.

- (1)  $R(T^*)^\perp = N(T)$  and  $R(T^*) = N(T)^\perp$ .
- (2)  $N(T^*T) = N(T)$ , and deduce from it that  $\text{rank}(T^*T) = \text{rank}(T)$ .
- (3)  $\text{rank}(T) = \text{rank}(T^*)$ , and deduce from (2) that  $\text{rank}(TT^*) = \text{rank}(T)$ .
- (4) For any  $n \times n$  matrix  $A$ ,  $\text{rank}(A^*A) = \text{rank}(AA^*) = \text{rank}(A)$ .