

MATH 33A Worksheet 2

TA: Emil Geisler and Caleb Partin

April 9, 2024

Exercise 1. Describe all solutions to the following linear systems.

(a)

$$x + 3y - 2z = 4$$

$$2x - y + 3z = 15$$

$$x - z = 3$$

(b)

$$x + y - 2z = 1$$

$$2x - 3y + z = 1$$

$$x - z = 2$$

(c)

$$x - 2y = 2$$

$$2y + 3z = 4$$

Exercise 2. Write down what it means for a matrix to be in row reduced echelon form (RREF). Which of the following matrices are in RREF? For each matrix, write its rank.

(a) $\begin{bmatrix} 1 & -2 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$

(b) $\begin{bmatrix} 1 & 2 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}$

(c) $\begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$

(d) $\begin{bmatrix} 1 & 2 & 0 & 3 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \end{bmatrix}$

(e) $\begin{bmatrix} 1 & 2 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$

Exercise 3. Find values a and b so that the ellipse $ax^2 + by^2 = 1$ goes through the points $(3, 2)$ and $(17, 12)$.

Exercise 4. Compute the following in the order of the parentheses, and observe they are the same. This is an example of the rule $A \cdot (\vec{v} + \vec{w}) = A \cdot \vec{v} + A \cdot \vec{w}$ for any $m \times n$ matrix A and $\vec{v}, \vec{w} \in \mathbb{R}^n$, i.e., A is a linear transformation.

$$(a) \begin{bmatrix} 2 & -1 & 3 \\ 0 & 1 & 1 \\ 1 & 0 & -1 \end{bmatrix} \left(\begin{bmatrix} 0 \\ 1 \\ -1 \end{bmatrix} + \begin{bmatrix} 1 \\ 0 \\ -2 \end{bmatrix} \right)$$

$$(b) \begin{bmatrix} 2 & -1 & 3 \\ 0 & 1 & 1 \\ 1 & 0 & -1 \end{bmatrix} \begin{bmatrix} 0 \\ 1 \\ -1 \end{bmatrix} + \begin{bmatrix} 2 & -1 & 3 \\ 0 & 1 & 1 \\ 1 & 0 & -1 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \\ -2 \end{bmatrix}$$

Exercise 5.

Compute the following in the order of the parentheses, and observe they are the same. This is an example of the rule $B \cdot (A \cdot \vec{v}) = (B \cdot A) \cdot \vec{v}$ for any $m \times n$ matrix B , $n \times q$ matrix A , and \vec{v} a vector in \mathbb{R}^q .

$$(a) \begin{bmatrix} 2 & -1 \\ 0 & 1 \\ 1 & 0 \end{bmatrix} \left(\begin{bmatrix} 0 & 1 & 2 \\ 3 & 1 & 0 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} \right)$$

$$(b) \left(\begin{bmatrix} 2 & -1 \\ 0 & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 0 & 1 & 2 \\ 3 & 1 & 0 \end{bmatrix} \right) \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$$
