## L-value formula

Haruzo Hida Department of Mathematics, UCLA, Lecture no.4 at NCTS, April 9, 2024

Lecture 4: We sketch the proof of the L-value formula for a division quaternion algebra  $D_{\mathbb{O}}$ . The algebra D can be definite or indefinite, though we describe mainly the details only for the indefinite case. Let  $B = D \otimes_{\mathbb{O}} E$  for a real semi-simple quadratic extension E. The non-trivial automorphism  $\sigma \in \text{Gal}(E/\mathbb{Q})$  acts on B through the factor E. Since the case  $E = \mathbb{Q} \times \mathbb{Q}$  is easier, we mainly assume that E is a field. A key point is the use of the see-saw principle for the decomposition  $D_{\sigma} = Z \oplus D_0$ , where  $D_{\sigma} := \{v \in B | v^{\iota} = v^{\sigma}\}$  with the reduced norm  $N : D_{\sigma} \to \mathbb{Q}$  and  $Z = D_{\sigma} \cap E$  and  $D_0 = \{v \in D_{\sigma} | \operatorname{Tr}(v) = 0\}$ . We need to use the Siegel–Weil formula for  $D_0$ . For simplicity, we assume  $M = \partial$ . The details are in Chapter 5, and the case  $M = M_2(\mathbb{Q})$  is dealt with in Section 5.5 of the notes.

§0. An idea of Waldspurger. For an elliptic cusp form f, an idea of Waldspurger of computing the period of a theta lift of f for a quadratic space  $V = W \oplus W^{\perp}$  over an orthogonal Shimura subvariety  $S_W \times S_{W^{\perp}} \subset S_V$  is two-folds:

(S) Split  $\theta(\phi')(\tau, h, h^{\perp}) = \theta(\phi)(\tau, h) \cdot \theta(\tau, \phi^{\perp})(h^{\perp})$   $(h^? \in O_{W?}(\mathbb{A}))$ for a decomposition  $\phi' = \phi \otimes \phi^{\perp}$   $(\phi \text{ and } \phi^{\perp} \text{ Schwartz-Bruhat}$ functions on  $W_{\mathbb{A}}$  and  $W_{\mathbb{A}}^{\perp}$ );

(R) For the theta lift  $\theta^*(\phi)(f)(h) = \int_X f(\tau)\theta(\phi)(\tau,h)d\mu$  with an SL(2)-Shimura curve X, the period P over the Shimura subvariety  $S \times S^{\perp}$  (S for O(W) and  $S^{\perp}$  for O(W<sup> $\perp$ </sup>)) is given by:

$$\int_{S\times S^{\perp}} \int_{X} f(\tau)\theta(\phi)(\tau;h)d\mu dh \quad (d\mu = \eta^{-2}d\xi d\eta)$$
$$= \int_{X} f(\tau) \left( \int_{S^{\perp}} \theta(\phi^{\perp})(\tau;h^{\perp})dh^{\perp} \right) \cdot \left( \int_{S} \theta(\phi_{0})(\tau;h_{0})dh \right) d\mu.$$

Then invoke the Siegel–Weil formula to convert inner integrals into the Siegel-Weil Eisenstein series  $E(\phi)$  and  $E(\phi^{\perp})$ , reaching Rankin-Selberg integral

$$P = \int_X f(\tau) E(\phi^{\perp}) E(\phi_0) d\mu = L \text{-value.}$$

§1. Choice of V: For a Q-vector space V and a Q-algebra A, write  $V_A := V \otimes_{\mathbb{Q}} A$ . Let  $E := \mathbb{Q}[\sqrt{\Delta}]$  be a quadratic extension of Q with discriminant  $\Delta$ . Pick a quaternion algebra D over Q and put  $B := D \otimes_{\mathbb{Q}} E$ . We let  $1 \neq \sigma \in \text{Gal}(E/\mathbb{Q})$  act on B through the factor E. Recall

 $V = D_{\sigma} := \{ v \in B | v^{\sigma} = v^{\iota} \} \text{ for } v^{\iota} = \operatorname{Tr}_{B/E}(v) - v.$ 

The quadratic form is given by  $Q(v) = vv^{\sigma} = N(v) \in \mathbb{Q}$ . We have two cases of isomorphism classes of  $(D_{\mathbb{R}}, E_{\mathbb{R}})$ . Note  $E_{\mathbb{R}} = \mathbb{R} \times \mathbb{R}$ ; so, we have two cases Case I and Case H. The symbol "I" (resp. "H") indicate D is indefinite (resp. definite). The decomposition we take is

 $V = Z \oplus D_0$   $Z = \mathbb{Q}$  with quadratic form  $Q_Z(z) = z^2$ , and

 $D_0 := \{v_0 \in \sqrt{\Delta}D | \operatorname{Tr}_{D/\mathbb{Q}}(v) = 0\} \text{ with } Q_0(v) = vv^{\sigma} = N(v)$ 

Signature of  $D_0$  is (1,2) in Case I and (3,0) in Case H,  $O_{D_0}$  is almost  $D^{\times}$  and the same for  $O_{D_{\sigma}}$  and  $B^{\times}$ .

§2. Bruhat functions and majorant. On  $Z = \mathbb{Q}$ , for a Dirichlet character  $\psi$  modulo N, we regard  $\psi$  as a function supported on  $\widehat{\mathbb{Z}} \subset Z_{\mathbb{A}(\infty)} = \mathbb{A}^{(\infty)}$ . This  $\psi$  produces theta series  $\sum_{n \in \mathbb{Z}} \psi(n) n^j e(n^2 \tau)$  on  $\Gamma_0(4N^2)$  of character  $\psi\left(\frac{-1}{2}\right)$  and of weight  $j + \frac{1}{2}$ .

Take a maximal order R of D and take the characteristic function  $\phi_0$  of  $D_{0,\mathbb{A}} \cap \sqrt{\Delta} \widehat{R}$ . Here for any lattice L,  $\widehat{L} = L \otimes_{\mathbb{Z}} \widehat{\mathbb{Z}}$ . This  $\phi_0$  produces theta series on  $\Gamma_0(4\partial \Delta)$  of character  $\left(-\Delta\right)$ .

The theta series for  $D_{\sigma}$  of  $\psi \otimes \phi_0$  has level  $M = [4N^2, 4\partial \Delta]$ . We choose M so that C|M for the conductor C of F.

A positive definite symmetric matrix  $P \in M_n(\mathbb{R})$  (or the symmetric bilinear form on  $V_{\mathbb{R}}$  associated to P) is a positive majorant of a symmetric matrix S if  $PS^{-1} = SP^{-1}$  ( $\Leftrightarrow S^{-1}P = P^{-1}S$ ).

§3. Schwartz function  $\Psi$  on  $D_{\sigma,\mathbb{R}}$  in Case I. The recipe of Hecke–Siegel is to put  $\Psi(v) = H(v)\mathbf{e}(\xi N(v) + P(v)\eta\sqrt{-1})$  for  $\mathbf{e}(x) = \exp(2\pi\sqrt{-1}x)$  and a harmonic polynomial H, where  $P(v) = \frac{1}{2}p(v,v)$  with a positive majorant p of  $s(v,v') = \operatorname{Tr}_{B/E}(v^{\iota}v')$ . All positive majorants form the symmetric space  $\mathfrak{S}$  of  $O_{D_{\sigma}}$ .

We identify  $(D_{\sigma,\mathbb{R}}, N) = (M_2(\mathbb{R}), \det)$  by  $M_2(\mathbb{R}) \ni v \mapsto (v, v^t) \in D_{\sigma,\mathbb{R}} \subset D_{\mathbb{R}} \times D_{\mathbb{R}}$  and put for  $(z, w) \in \mathfrak{H} \times \mathfrak{H}$  on which  $B^{\times} \sim \operatorname{GO}_{D_{\sigma}}$ acts by  $\alpha(z, w) = (\alpha(z), \alpha^{\sigma}(w))$ . For  $(z, w) \in \mathfrak{H} \times \mathfrak{H}$ , a standard harmonic polynomial of  $v \in D_{\sigma}$  of degree k is given by  $[v; z, w]^k = s(v, p(z, w))^k$  for  $p(z, w) = (\frac{z}{1})(w, 1)J$ . For  $0 < k \in \mathbb{Z}$ ,  $\Psi(v; \tau, z, w) = \operatorname{Im}(\tau) \frac{[v; z, \overline{w}]^k}{(z-\overline{z})^k (w-\overline{w})^k} e(N(v)\overline{\tau} + i \frac{\operatorname{Im}(\tau)}{2|\operatorname{Im}(z)\operatorname{Im}(w)|} |[v; z, \overline{w}]|^2),$ for  $(\alpha, \beta) \in GL_2(\mathbb{R}) \times GL_2(\mathbb{R})$  (see §3.2.3),

 $\alpha p(z,w)\beta^{\iota} = p(\alpha(z),\beta(w))j(\alpha,z)j(\beta,w).$ 

This formula is due to Shimura. This function is not a tensor product of functions on  $Z_{\mathbb{R}}$  and  $D_{0,\mathbb{R}}$  which causes some difficulty later. For simplicity, we assume k = 2. See Section 3.2 for  $\Psi$ .

§4. Theta kernel. Let  $\phi$  be a Schwartz-Bruhat function on  $D_{\sigma,\mathbb{A}}$ . Let  $\mathsf{Mp}(\mathbb{A}) \twoheadrightarrow \mathsf{SL}_2(\mathbb{A})$  be the metaplectic cover constructed by Weil, and  $\phi \mapsto \mathbf{w}(g)\phi$  the Weil representation. Noting  $B^{\times} \twoheadrightarrow \mathsf{GO}_{D_{\sigma}}$  by  $v \mapsto h^{\iota}vh^{\sigma}$ , Siegel-Weil theta series  $\theta(g; h)$  is

 $\sum_{\alpha \in D_{\sigma}} (\mathbf{w}(g)\phi)(h^{\iota}\alpha h^{\sigma}) : \mathsf{SL}_{2}(\mathbb{Q}) \setminus \mathsf{Mp}(\mathbb{A}) \times B^{\times} \setminus B^{\times}_{\mathbb{A}} \to \mathbb{C}.$ 

Write  $\widehat{\Gamma} = \widehat{\Gamma}_{\phi} = \{ u \in B_{\mathbb{A}(\infty)}^{\times} | \theta(g, u^{\iota}hu^{\sigma}) = \theta(g, h) \}.$ 

In Case I, choose  $\phi = (\psi \otimes \phi_0) \Psi(v; \sqrt{-1}, \sqrt{-1}, \sqrt{-1})$  and for  $g_\tau = \eta^{-1/2} \begin{pmatrix} \eta & \xi \\ 0 & 1 \end{pmatrix}$   $(\tau = \xi + \eta \sqrt{-1} \in \mathfrak{H})$ , we specialize g to  $g_\tau$  and h to  $(g_z, g_w)$  for  $(\tau, z, w) \in \mathfrak{H} \times \mathfrak{H} \times \mathfrak{H}$ . Then

$$\theta(\tau;z,w) := \theta(g_{\tau};g_z,g_w) = \sum_{\alpha \in D_{\sigma}} (\psi \otimes \phi_0)(\alpha) \Psi(\alpha;\tau,z,w).$$

Set  $\theta^*(\phi)(f) := \int_{X_0(M)} f(\tau)\theta(\phi)(\tau; z, w)\eta^{k-2}d\xi d\eta$  (k = 2). Then  $\theta^*(\phi)(f)$  is a weight 2 quaternionic modular form on  $B^{\times}$  holomorphic in z and anti-holomorphic in w for  $f \in S_2^-(\Gamma_0(M), \psi^{-1}(\Delta))$ .

§5. Theta differential form. To compute the period on  $Sh_D = D^{\times}_+ \setminus (D^{\times}_{\mathbb{A}(\infty)} \times \mathfrak{H}) \subset Sh_B = B^{\times} \setminus (B^{\times}_{\mathbb{A}(\infty)} \times \mathfrak{Z}_B)$ , we convert  $\theta(\tau; z, w)$  into a sheaf valued differential 2-form. If n = k - 2 > 0, the sheaf comes from the  $B^{\times}$ -module

$$L_E(n; A) = \sum_{0 \le i,j \le n} A X^{n-j} Y^j X'^{n-i} Y'^i$$

with  $B^{\times}$ -action  $\gamma P(X, Y; X', Y') = P((X, Y)^t \gamma^{\iota}; (X', Y')^t \gamma^{\sigma \iota})$ . As we assumed k = 2 (i.e., n = 0), we have L(n; A) = A.

By putting  $\Theta = \theta(\phi)(\tau; z, w)dz \wedge d\overline{w}$  for n = k-2, we get  $\mathbb{C}$ -valued  $\Gamma_{\phi}$ -invariant differential form. The period we like to compute is

$$P = P_1(\theta^*(\phi)(f)) = \int_{Sh_D} \int_{X_0(M)} f(-\overline{\tau}) \Theta(\tau; z, z) d\xi d\eta.$$

We integrate over  $Sh_D$  by a measure  $d\mu$  given by  $y^{-2}dxdy$  over  $\mathfrak{H}$  and  $\int_{\widehat{\Gamma}} d\mu = 1$ .

§6. Siegel–Weil Eisenstein series; §4.4.2. Recall the explicit section  $\mathbf{r} : B \hookrightarrow Mp$  of the representation  $\mathbf{w}$  as follows:

 $\mathbf{r}(\operatorname{diag}[a, a^{-1}])\phi(v) = |a|_{\mathbb{A}}^{3/2}\phi(av), \quad \mathbf{r}\begin{pmatrix}1 & u\\0 & 1\end{pmatrix}\phi(v) = \mathbf{e}(uN(v))\phi(v).$ For the standard Borel subgroup  $B \subset SL_2$ , the function  $g \mapsto (\mathbf{r}(g)\phi)(0)$  is left  $B(\mathbb{Q})$  invariant. Siegel–Weil Eisenstein series is

$$E(\phi)(g;s) = \sum_{\gamma \in B(\mathbb{Q}) \setminus \mathsf{SL}_2(\mathbb{Q})} (\mathbf{w}(\gamma g)\phi)(0) |a(\gamma g)|_{\mathbb{A}}^s,$$

where  $g = \text{diag}[a(g), a(g)^{-1}] \begin{pmatrix} 1 & u \\ 0 & 1 \end{pmatrix} c$  for an element c in the maximal compact subgroup by Iwasawa decomposition.

The Siegel–Weil formula by Kudla-Rallis and Sweet is

 $E(\phi)(g; 0) = \int_{S} \theta(\phi)(g, h) d\omega$  for the Tamagawa measure  $d\omega$ .

The ratio  $\mathfrak{m} = \mathfrak{m}(\widehat{\Gamma}) = d\mu/d\omega$  is the mass of Siegel–Shimura, which is an arithmetic rational number times  $\zeta(2)/\pi$  in Case I and  $\zeta(2)/\pi^2$  in Case H. Later we describe the Tamagawa measure, the metaplectic group and Weil representation in more details. §7. Conclusion in Case I; §5.3. Decomposing  $v = a \oplus v_0$  in  $D_{\sigma} = Z \oplus D_0$ , we have for k = 2

$$[a+v_0;z,\overline{z}]^k = ([a;z,\overline{z}] + [v_0;z,\overline{z}])^k = \sum_{j=0}^k \binom{k}{j} (a(z-\overline{z}))^{k-j} [v_0;z,\overline{z}]^j.$$

Thus we have  $\phi = \sum_{j=0}^{k} {k \choose j} \phi_{k-j}^{Z} \phi_{j}^{D_{0}}$  with infinity part  $\Psi_{j}^{?}$  of  $\phi_{j}^{?}$  given by  $\Psi_{j}^{D_{0}} := (z - \overline{z})^{-j} [v_{0}; z, \overline{z}]^{j} \mathbf{e}(N(\mathfrak{x})\overline{\tau} + \frac{i \operatorname{Im}(\tau) |[v_{0}; z, \overline{z}]|^{2}}{2 \operatorname{Im}(z)^{2}}), \Psi_{j}^{Z} := a^{j} \mathbf{e}(a^{2}\tau)$ with  $(\mathbf{w}(b)\phi_{j}^{D_{0}})(0) = 0$  and  $E(\phi_{j}^{D_{0}})|_{B(\mathbb{A})} = 0$  unless j = 0, and we reach Rankin convolution of  $\theta(\phi_{k}^{Z}) = \sum_{n \in \mathbb{Z}} \psi(n) n^{k} \mathbf{e}(n^{2}z)$  and fover  $B(\mathbb{Q}) \setminus B(\mathbb{A}) / B(\widehat{\mathbb{Z}}) \cong [0, 1) \times \mathbb{R}^{\times}_{+}$ , which produces (see [Sh75])

$$\zeta(2)P = \mathfrak{m}2^{-2k} * (2\pi)^{-k} \Gamma(k) L^{(s)}(1, Ad(\rho_f) \otimes \left(\frac{\Delta}{-}\right))$$

with a simple constant \*. Here  $L^{(s)}$  means we remove Euler factors at p|C with either f|U(p) = 0 or  $\psi(p) = 0$ .

§8. Conclusion in Case H; §5.2. The choice of the Bruhat function  $\phi$  is the same as in Case I. As a  $\mathbb{C}$ -valued function, set

 $\Psi(\tau; v; \mathbf{x}) = \mathbf{e}(N(v)\tau).$ 

Again in exactly the same way, for

$$\theta^*(\phi)(f) := \int_{X_0(M)} \theta(\phi)(\tau; g) f(\tau) \eta^{k-2} d\xi d\eta \quad (k=2)$$

and  $P = \int_{S} \theta^{*}(\phi)(f) d\mu$ , we conclude for a simple constant c'

$$\zeta(2)P = 2\mathfrak{m} *' (2\pi)^{-k+1} \Gamma(k) L^{(s)}(1, Ad(\rho_f) \otimes \left(\frac{\Delta}{-}\right)).$$

Writing the point set  $S = \{x\}_{x \in Sh_R}$ ,  $\mathfrak{m}(\widehat{\Gamma}) = \sum_{x \in Sh_R} e_x^{-1} \doteq \zeta(2)$ for  $e_x = |\widehat{\Gamma} \cap \mathcal{O}_{D_0}(\mathbb{Q})|$  and  $P \doteq \sum_{x \in Sh_R} e_x^{-1} \theta^*(\phi)(f)(x)$ .

Thus the period formula is an adjoint analogue of the mass formula of Siegel–Shimura. The determination of  $\mathfrak{m}(\widehat{\Gamma})$  was finished by Shimura in 1999 for an arbitrary quadratic space over a totally real field (see §5.2.8 for the explicit formula for the mass).

§9. Schwartz–Bruhat functions; §3.1.3. For a  $\mathbb{Q}$ -vector space V, write  $V_p := V \otimes_{\mathbb{Z}} \mathbb{Z}_p$  (which is a vector space over a local field  $\mathbb{Q}_p$ ). A Bruhat function on  $V_p$  is a locally constant compactly supported function with values in  $\mathbb{C}$ . Write  $\mathcal{S}(V_p)$  for the space of Bruhat functions on  $V_p$ . For a real vector space  $V_{\infty}$ , we define  $\mathcal{S}(V_{\infty})$  to be the Schwartz space of functions on  $V_{\infty}$ . Thus  $\mathcal{S}(V_{\infty})$  is made of  $C^{\infty}$ -class functions with all derivatives rapidly decreasing as Euclidean norm of  $v \in V_{\infty}$  grows. In other words,  $\phi \in \mathcal{S}(V_{\infty})$  if and only if  $\phi$  is of  $C^{\infty}$ -class and for any polynomial P(v) and any m-th derivative  $\Phi$  of  $\phi$ ,  $|P(v)\Phi(v)|$  goes to 0 as  $|x| \to \infty$ . Writing  $V_{\mathbb{A}}$  for the adelization. We pick a lattice L of V and put  $\widehat{L} = \prod_p L_p \subset V_{\mathbb{A}(\infty)}$  with  $L_p = L \otimes_{\mathbb{Z}} \mathbb{Z}_p$ . A Schwartz-Bruhat function on  $V_{\mathbb{A}}$  is a finite linear combination of the function of the form  $\phi(x) = \prod_v \phi_v(x_v)$  with  $\phi_v \in \mathcal{S}(V_v)$  and  $\phi_p$  is the characteristic function of  $L_p$  for almost all p.

§10. Weil representation. Let  $v(u) = \begin{pmatrix} 1 & u \\ 0 & 1 \end{pmatrix}$ , diag $[a, b] = \begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix}$ and  $J = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ . Let  $\mathbf{e} : \mathbb{A}/\mathbb{Q} \to S^1$  be the additive character with  $\mathbf{e}(x_{\infty}) := e_{\infty}(2\pi\sqrt{-1}x_{\infty})$ . We put  $\mathbf{e}_{v} := \mathbf{e}|_{\mathbb{O}_{v}}$  and for a number field F, we write  $\mathbf{e}_F = \mathbf{e} \circ \operatorname{Tr}_{F/\mathbb{Q}}$  and  $\mathbf{e}_{F_v} = \mathbf{e}_v \circ \operatorname{Tr}_{F_v/\mathbb{Q}_v}$ . Let  $Q: V \rightarrow F$  be a non-degenerate quadratic form with symmetric bilinear form s(v,w) = Q(v+w) - Q(v) - Q(w) over a number field F. We have the following operator  $\mathbf{r}(g) \in Aut(\mathcal{S}(V_{7}))$  for  $m = \dim_F V$ , ? =  $v, \mathbb{A}$  with  $u \in F_v$  or  $F_{\mathbb{A}}$  and  $a \in F_v^{\times}$ :  $\mathbf{r}(v(u)) = \mathbf{e}_F(uQ(v))\phi(v), \ \mathbf{r}(\operatorname{diag}[a,a^{-1}]\phi(v) = |a|_{F_{\mathbb{A}}}^{m/2}\phi(av) \ \text{and}$  $\mathbf{r}(J)\phi(v) = \widehat{\phi}(-v) := \int_{V_2} \mathbf{e}_F(s(w,-v))\phi(w)dw$  (Fourier transform), where dw is normalized so that  $\widehat{\phi}(v) = \phi(-v)$ . If  $b, b' \in B(F_{\mathbb{A}})$  (upper triangular Borel subgroup), we extend r to  $\Omega = B(F_{\mathbb{A}})JB(F_{\mathbb{A}})$ by  $\mathbf{r}(bJb') := \mathbf{r}(b)\mathbf{r}(J)\mathbf{r}(b')$ . Then if  $g,h \in SL_2(F_7)$  either unipotent, diagonal or J,  $\mathbf{r}(gh) = \kappa(g,h)\mathbf{r}(g)\mathbf{r}(h)$  for a 2-cocycle  $\kappa$ on SL<sub>2</sub> with values in  $S^1$ . Write Mp( $F_7$ )  $\subset$  Aut( $\mathcal{S}(V_7)$ ) for the group generated by these operators. We have an extension (\*)  $1 \to S^1 \to \mathsf{Mp}(F_7) \xrightarrow{\pi_?} \mathsf{SL}_2(F_7) \to 1$  with  $\mathsf{Mp} \ni \mathbf{w}(g) \mapsto g \in \mathsf{SL}_2$ . Therefore, the group Mp acts on  $\mathcal{S}(V_7)$  by a representation w.

 $\S$ **11. Weil's theta series.** The extension (\*) is split in the following cases:

1.  $\dim_F V$  is even (the section is unique and if  $V = D_{\sigma,F_?}$  $b \mapsto \chi_V(a)\mathbf{r}(b)$  over  $B(F_{\mathbb{A}})$  if  $b = v(u) \operatorname{diag}[a, a^{-1}]$ );

- 2.  $b \mapsto \mathbf{r}(b)$  and also  $b \mapsto \chi_V(a)\mathbf{r}(b)$  as above over  $B(F_{\mathbb{A}})$ ;
- 3. Over  $\widehat{\Gamma}_0(4)$  (canonical);

4. Over  $SL_2(F)$  (canonical and coincides with  $b \mapsto \mathbf{r}(b)$  over B(F).

For the orthogonal group  $O_V$  for V and  $\phi \in S(V_A)$ , we define a function on  $Mp(F_A) \times O_V(F_A)$  by  $\theta(\phi)(g,h) = \sum_{\alpha \in V} \mathbf{w}(g)L(h)\phi(\alpha)$ , where  $(L(h)\phi)(v) = \phi(vh)$  (as usual,  $O(F_A)$  acts on  $V_{F_A}$  from the right). Weil showed that  $\theta(\phi)(g,h)$  is real analytic on  $Mp(F_\infty) \times O_V(F_\infty)$ , left invariant under  $SL_2(F) \times O_V(F)$  and right invariant under an open subgroup of  $Mp(F_{A(\infty)}) \times O_V(F_{A(\infty)})$ ; in short, an automorphic form on  $Mp \times O_V$ .

All the details are in Chapter 4.