

Determinacy Proofs for Long games

Itay Neeman

Department of Mathematics
University of California Los Angeles
Los Angeles, CA 90095-1555
ineeman@math.ucla.edu

1. Preliminaries:
 - (a) The games.
 - (b) Extenders, iteration trees.
 - (c) Auxiliary game representations.
 - (d) Example: Σ_2^1 determinacy.
2. Games of length $\omega \cdot \omega$ with Σ_2^1 payoff.
3. Continuously coded games with Σ_2^1 payoff.

Let $C \subset \mathbb{R}^{<\omega_1}$ be given.* Let $f: \mathbb{R} \rightarrow \mathbb{N}$, a partial function, be given. $G_{\text{cont-}f}(C)$ is played as follows:

I	$y_\alpha(0)$	$y_\alpha(2)$	
II		$y_\alpha(1)$	$y_\alpha(3)$...

In round α , I and II alternate playing natural numbers $y_\alpha(i)$, $i < \omega$, producing a real y_α .

If $f(y_\alpha)$ is not defined, the game ends. I wins iff $\langle y_0, y_1, \dots, y_\alpha \rangle \in C$.

Otherwise we set $n_\alpha = f(y_\alpha)$. If there exists $\xi < \alpha$ so that $n_\alpha = n_\xi$, the game ends. Again I wins iff $\langle y_0, y_1, \dots, y_\alpha \rangle \in C$.

Otherwise the game continues.

The game ends at a countable α ; the map $\xi \mapsto n_\xi$ embeds α into \mathbb{N} . This map is produced continuously in ξ . The game is said to have **continuously coded length**.

*Following standard abuse of notation, we use \mathbb{R} to denote \mathbb{N}^ω .

Let $C \subset \mathbb{R}^\omega = \mathbb{N}^{\omega \cdot \omega}$ be given. In $G_{\omega \cdot \omega}(C)$ the players play ω rounds as follows, producing $y_k \in \mathbb{R}$ for $k < \omega$.

I	$y_0(0)$	$y_1(0)$...
II	$y_0(1)$		$y_1(1)$...

I wins iff $\langle y_k \mid k < \omega \rangle$ belongs to C .

Let $C \subset \mathbb{R} = \mathbb{N}^\omega$ be given. In $G_\omega(C)$ the players play one round as follows, producing $y \in \mathbb{R}$.

I	$y(0)$	$y(2)$...
II	$y(1)$	$y(3)$...

I wins iff $y \in C$.

We intend to prove that $G_{\text{cont-}f}(C)$ are determined, for all continuous f and all Σ_2^1 payoff sets C .

As an illustrative case we will first prove that $G_{\omega \cdot \omega}(C)$ are determined, for all Σ_2^1 payoff sets C .

Before that, we will prove that $G_\omega(C)$ are determined for all Σ_2^1 sets $C \subset \mathbb{R}$.

Determinacy for games of length ω was proved by Martin and Steel.

Determinacy for games of fixed length $\omega \cdot \alpha$, α limit, was proved by Woodin.

Determinacy for games of continuously coded length was proved by Neeman.

An **extender** on κ is a directed system of measures on κ . If E is an extender on κ , we use $\text{dom}(E)$ to denote κ .

An extender E allows us to form an **ultrapower** of V , denoted $\text{Ult}(V, E)$, and an elementary **ultrapower embedding** $\pi: V \rightarrow \text{Ult}(V, E)$.

We use P, Q, M, N to denote models of ZFC.

We say that Q and Q^* **agree** to κ if $\mathcal{P}(\kappa) \cap Q^* = \mathcal{P}(\kappa) \cap Q$.

Suppose $Q \models "E \text{ is an extender on } \kappa"$. Suppose Q^* and Q agree to κ . Then E can be applied also to Q^* : We can form the **ultrapower** $\text{Ult}(Q^*, E)$, and an elementary **ultrapower embedding** $\sigma: Q^* \rightarrow \text{Ult}(Q^*, E)$.

$\text{Ult}(Q^*, E)$ needn't always be wellfounded. If it is wellfounded, we assume it's transitive.

An **iteration tree** \mathcal{T} of length ω consists of


- a tree order T on ω ,
- a sequence of models $\langle M_k \mid k < \omega \rangle$, and
- embeddings $j_{k,l}: M_k \rightarrow M_l$ for $k T l$.

Each model M_{l+1} for $l + 1 > 0$ is an ultrapower of a preceding model. More precisely: $M_{l+1} = \text{Ult}(M_k, E_l)$, where E_l an extender picked from M_l , and $k \leq l$ is the T predecessor of $l + 1$. $j_{k,l+1}$ is the ultrapower embedding.

$$\begin{array}{ccc}
 & M_{l+1} & \\
 & \uparrow & \\
 j_{k,l+1} & & E_l \in M_l \\
 & \uparrow & \\
 & M_k &
 \end{array}$$

(M_l and M_k must agree to $\text{dom}(E_l)$.)

An iteration tree on M is a tree with $M_0 = M$.




Our trees will generally have an **even branch**, M_0, M_2, M_4, \dots , giving rise to the direct limit M_{even} .

The tree structure on the odd models will usually be some permutation of $\omega^{<\omega}$. With each **odd branch** b we associate the direct limit M_b .

(In this example, $0 \ T \ 1$, $0 \ T \ 2$, $1 \ T \ 3$, $0 \ T \ 3$, etc.)

In the **iteration game*** on M , players “good” and “bad” collaborate to produce a sequence of iteration trees as follows:




“Bad” plays an iteration tree \mathcal{T}_ξ on M_ξ . “Good” plays a branch b_ξ through \mathcal{T}_ξ . We let $M_{\xi+1}$ be the direct limit model determined by b_ξ and proceed to the next round. For limit λ we let M_λ be the direct limit of M_ξ for preceding ξ . We start with $M_0 = M$.

If ever a model $(M_\xi, \xi < \omega_1)$ is reached which is illfounded, “bad” wins. Otherwise “good” wins.

*The definition given here is specialized to our context. The concept of iteration games is due to Martin–Steel.

We also consider iteration games where round ξ has the following form:



“Bad” plays an iteration tree \mathcal{T}_ξ on M_ξ . “Good” plays a branch b_ξ , giving rise to the direct limit, P_ξ .

Then “good” plays an extender E_ξ in P_ξ , with $\text{dom}(E_\xi)$ within the level of agreement between M_ξ and P_ξ . We set $M_{\xi+1} = \text{Ult}(M_\xi, E_\xi)$ and continue to the next round.

If ever a model (P_ξ or M_ξ , $\xi < \omega_1$) is reached which is illfounded, “bad” wins. Otherwise “good” wins.

We refer to this game too as an **iteration game**.

M is **iterable** if the good player has a winning strategy for each of the iteration games described above. We refer to such winning strategies as **iteration strategies**.

Countable elementary substructures of V are iterable in this sense (Martin–Steel).

Suppose $M \models \text{“}\delta \text{ is a Woodin cardinal”}$, and in V there are M -generics for $\text{col}(\omega, \delta)$. Let \dot{A} name a set of reals in $M^{\text{col}(\omega, \delta)}$.

Work with some $x \in \mathbb{R}$. We work to define an auxiliary game, $\mathcal{A}[x]$, of ω moves, taken from M . In this game I tries to witness that $x \in \dot{A}[h]$ for some generic h . II tries to witness the opposite.

The auxiliary game is played as follows:

I	...	l_n, \mathcal{X}_n, p_n	...
II	$\mathcal{F}_n, \mathcal{D}_n$...

In round n I plays

- $l = l_n$, a number $< n$, or $l_n = \text{“new”}$.
- \mathcal{X}_n , a set of names for reals of $M^{\text{col}(\omega, \delta)}$.
- p_n , a condition in $\text{col}(\omega, \delta)$.

II plays

- \mathcal{F}_n a function from \mathcal{X}_n into the ordinals.
- \mathcal{D}_n , a function from \mathcal{X}_n into $\{\text{dense sets in } \text{col}(\omega, \delta)\}$.

$$\mathcal{A}[x] : \begin{array}{c|ccc} \text{I} & \dots & l_n, \mathcal{X}_n, p_n & \dots \\ \hline \text{II} & & \mathcal{F}_n, \mathcal{D}_n & \dots \end{array}$$

If $l_n = \text{“new”}$ we make no requirements on I. Otherwise, we require $p_n < p_l$ and $\mathcal{X}_n \subset \mathcal{X}_l$. We further require that for every name $\dot{x} \in \mathcal{X}_n$:

1. p_n forces “ $\dot{x} \in \dot{A}$ ”.
2. p_n forces “ $\dot{x}(0) = \tilde{x}_0$ ”, ..., “ $\dot{x}(l) = \tilde{x}_l$ ”.
3. p_n belongs to $\mathcal{D}_l(\dot{x})$.

We make the following requirement on II:

4. For every name $\dot{x} \in \mathcal{X}_n$, $\mathcal{F}_n(\dot{x}) < \mathcal{F}_l(\dot{x})$.

If there is h so that $x \in \dot{A}[h]$, I can pick a name for x , play \mathcal{X}_n containing this name, and play $p_n \in h$. Condition 4 ensures defeat for II.

On the other hand, if there is an infinite run of $\mathcal{A}[x]$ where I covered all possible names and chains of conditions, condition 4 ensures that $x \notin \dot{A}[h]$ for all generic h .

Note 1. Rather than play the sets \mathcal{X}_n directly, I plays their *type*. I plays $\kappa_n < \delta$, and a set u_n of formulae with parameters in $M \parallel \kappa_n \cup \{\kappa_n, \delta, \dot{A}\}$.^{*} We take \mathcal{X}_n to be the set of names which satisfy all these formulae.

The fact that this still allows I enough control over her choice of \mathcal{X}_n has to do with our assumption that δ is a Woodin cardinal.

\mathcal{F}_n and \mathcal{D}_n are played similarly.

Observe that moves in $\mathcal{A}[x]$ are therefore elements of $M \parallel \delta$.

Note 2. The association $x \mapsto \mathcal{A}[x]$ is continuous: The rules governing the first $n + 1$ rounds of $\mathcal{A}[x]$ depend only on $x \upharpoonright n$.

We in fact defined an association $s \mapsto \mathcal{A}[s]$ ($s \in \omega^{<\omega}$, $\mathcal{A}[s]$ a game of $\text{lh}(s) + 1$ many rounds). This association belongs to M .

^{*}By $M \parallel \kappa_n$ we mean $V_{\kappa_n}^M$.

Recall that g is $\text{col}(\omega, \delta)$ -generic/ M . We alternate between thinking of g as a generic enumeration of δ , and as a generic enumeration of $M \parallel \delta$.

Let $\sigma_{\text{gen}}[x, g]$, a strategy for I in $\mathcal{A}[x]$ be defined as follows:


$\sigma_{\text{gen}}[x, g]$ plays in each round the *first* (with respect to the enumeration g) legal move.

Note. The association $x, g \mapsto \sigma_{\text{gen}}[x, g]$ is continuous.

Lemma 1. Suppose that there exists an infinite run of $\mathcal{A}[x]$, played according to $\sigma_{\text{gen}}[x, g]$. Then $x \notin \dot{A}[g]$. (This is only useful if $x \in M[g]$.)

Proof: In playing for I, $\sigma_{\text{gen}}[g, x]$ goes over all possible names and all possible generics. (This uses the genericity of the enumeration g .) So in fact $x \notin \dot{A}[h]$ for all generic h . \square

We wish to phrase a similar lemma with a strategy for II, which puts x in A . To do this we have to give II additional control. We let II “shift” the play board along an even branch of an iteration tree.



I l_0
 \mathcal{X}_0 \rightsquigarrow
 p_0

II \mathcal{F}_0
 \mathcal{D}_0

I l_1
 \mathcal{X}_1 \rightsquigarrow
 p_1

II \mathcal{F}_1
 \mathcal{D}_1

I l_2
 \mathcal{X}_2 \rightsquigarrow
 p_2

The game $\mathcal{A}^*[x]$ is played as follows:

I	...	l_n, \mathcal{X}_n, p_n	...
II	$E_{2n}, E_{2n+1}, \mathcal{F}_n, \mathcal{D}_n \quad \dots$		

At the start of round n we have a model M_{2n} , an embedding $j_{0,2n}: M \rightarrow M_{2n}$, and a position P_n of n rounds in $j_{0,2n}(\mathcal{A})[x]$.

I plays l_n, \mathcal{X}_n, p_n , a legal move in $j_{0,2n}(\mathcal{A})[x]$ following P_n .

II plays extenders E_{2n}, E_{2n+1} giving rise to models M_{2n+1}, M_{2n+2} , and to an embedding $j_{2n,2n+2}: M_{2n} \rightarrow M_{2n+2}$. (The T -predecessor of $2n + 1$ is $2l_n + 1$ if $l_n \neq$ "new" and $2n$ otherwise.)

We let $Q_n = j_{2n,2n+2}(P_n-, l_n, \mathcal{X}_n, p_n)$. (This is the "shifting" mentioned before.)

II plays $\mathcal{F}_n, \mathcal{D}_n$, a legal move in $j_{0,2n+2}(\mathcal{A})[x]$ following Q_n .

We let $P_{n+1} = Q_n-, \mathcal{F}_n, \mathcal{D}_n$ and proceed to the next round.

Definition. A **pivot** for x is a pair \mathcal{T}, \vec{a} so that

1. \mathcal{T} is an iteration tree on M , with an even branch.
2. \vec{a} is a run of $j_{\text{even}}(\mathcal{A})[x]$.
3. For every odd branch b of \mathcal{T} , there exists some h so that
 - (a) h is $\text{col}(\omega, j_b(\delta))$ -generic/ M_b ; and
 - (b) $x \in j_b(\dot{A})[h]$.

Any run of $\mathcal{A}^*[x]$ produces \mathcal{T}, \vec{a} which satisfy conditions 1 and 2.

Lemma 2. There exists $\sigma_{\text{piv}}[x, g]$, a strategy for II in $\mathcal{A}^*[x]$, so that every run according to $\sigma_{\text{piv}}[x, g]$ is a pivot.

The association $x, g \mapsto \sigma_{\text{piv}}[x, g]$ is continuous.

The proof of Lemma 2 draws heavily on the techniques of Martin–Steel’s “A proof of projective determinacy”. The assumption that δ is a Woodin cardinal is crucial.

To sum: Have continuous associations $x \mapsto \mathcal{A}[x]$; $x, g \mapsto \sigma_{\text{gen}}[x, g]$; $x \mapsto \mathcal{A}^*[x]$; and $x, g \mapsto \sigma_{\text{piv}}[x, g]$.

$\sigma_{\text{gen}}[x, g]$ is a strategy for I in $\mathcal{A}[x]$.

If \vec{a} is an infinite run of $\mathcal{A}[x]$ according to $\sigma_{\text{gen}}[x, g]$, then $x \notin \dot{A}[g]$.

$\sigma_{\text{piv}}[x, g]$ is a strategy for II in $\mathcal{A}^*[x]$.

If \mathcal{T} , \vec{a} is an infinite run of $\mathcal{A}^*[x]$ according to $\sigma_{\text{piv}}[x, g]$, then

for every odd branch b of \mathcal{T} , there exists some h so that

- h is $\text{col}(\omega, j_b(\delta))$ -generic/ M_b ; and
- $x \in j_b(\dot{A})[h]$.

Σ_2^1 determinacy:

Fix $A \subset \mathbb{R}$, a Σ_2^1 set (say the set of reals which satisfy a given Σ_2^1 statement ϕ).

Suppose there is an iterable class model M with a Woodin cardinal δ . Suppose that (in V) there is g which is $\text{col}(\omega, \delta)$ -generic/ M .

We intend to prove that (in V) $G_\omega(A)$ is determined.

Let $\dot{A} \in M$ name A . More precisely, \dot{A} names the set of reals of $M^{\text{col}(\omega, \delta)}$ which satisfy ϕ in $M^{\text{col}(\omega, \delta)}$.

We have $x \mapsto \mathcal{A}[x]$, $x, g \mapsto \sigma_{\text{gen}}[x, g]$, etc. as before.

Let G be the following game, defined and played inside M :

I	x_0	a_{0-I}	a_{1-I}	x_2	\dots
II		a_{0-II}	x_1	a_{1-II}	

I and II alternate playing natural numbers, producing together $x = \langle x_0, x_1, \dots \rangle \in \mathbb{R}$. In addition they play moves a_{0-I}, a_{0-II}, \dots in $\mathcal{A}[x]$.

II is the closed player; she wins if she can last all ω moves. Otherwise I wins.

G is a closed game, hence determined. A winning strategy exists in M .

Case 1: I wins G . Fix $\Sigma \in M$ a winning strategy for I (the open player).

We wish to show that I wins $G_\omega(A)$ in V . Let us play $G_\omega(A)$ against an imaginary opponent. We describe how to play, and win.

We construct a run $x \in \mathbb{R}$ of $G_\omega(A)$. At the same time we construct \mathcal{T}, \vec{a} , a run of $\mathcal{A}^*[x]$.

The participants in our construction are:

- The imaginary opponent: playing x_n for odd n .
- The strategy $\sigma_{\text{piv}}[g, x]$: playing for II in $\mathcal{A}^*[x]$.
- The strategy Σ and its shifts along the even branch of \mathcal{T} : playing x_n for even n and playing for I in $\mathcal{A}^*[x]$ (i.e. playing for I in shifts of $\mathcal{A}[x]$).

We obtain $x \in \mathbb{R}$ and \mathcal{T}, \vec{a} a run of $\mathcal{A}^*[x]$ according to $\sigma_{\text{piv}}[x, g]$.

We must check that x belongs to A .



Σ x_0

Σ l_0
 $\mathcal{X}_0 \rightsquigarrow$
 p_0

σ_{piv} \mathcal{F}_0
 \mathcal{D}_0

Oppnt x_1

$j_{0,2}(\Sigma)$ l_1
 $\mathcal{X}_1 \rightsquigarrow$
 p_1

σ_{piv} \mathcal{F}_1
 \mathcal{D}_1

$j_{0,4}(\Sigma)$ x_2

$j_{0,4}(\Sigma)$ l_2
 $\mathcal{X}_2 \rightsquigarrow$
 p_2

Note that x, \vec{a} is an infinite run of $j_{\text{even}}(G)$ according to $j_{\text{even}}(\Sigma)$.

Now Σ is a strategy for the open player in G . So there are no infinite runs according to Σ . But there is an infinite run according to $j_{\text{even}}(\Sigma)$. Thus M_{even} is **illfounded**.

M is iterable. So there exists some branch b of \mathcal{T} so that M_b is wellfounded. b must be an odd branch.

By Lemma 2, \mathcal{T}, \vec{a} is a pivot for x . Thus there is h so that

- h is $\text{col}(\omega, j_b(\delta))$ -generic/ M_b and
- $x \in j_b(\dot{A})[h]$.

This means that in $M_b[h]$, x satisfies the Σ_2^1 statement ϕ .

By absoluteness, x satisfies ϕ in V . (This uses the wellfoundedness of M_b .)

So $x \in A$ as required.

□(Case 1.)