

Math 3B: Lecture 4

Noah White

October 6, 2017

Last time

Last time, we spoke about

- Graphing using calculus

Last time

Last time, we spoke about

- Graphing using calculus
- Slanted asymptotes

Last time

Last time, we spoke about

- Graphing using calculus
- Slanted asymptotes
- Examples

Functions

A function is three pieces of information

Functions

A function is three pieces of information

- A domain, $D \subset \mathbb{R}$

Functions

A function is three pieces of information

- A domain, $D \subset \mathbb{R}$
- A range, $R \subset \mathbb{R}$, and

Functions

A function is three pieces of information

- A domain, $D \subset \mathbb{R}$
- A range, $R \subset \mathbb{R}$, and
- A rule $f : D \rightarrow R$ that assigns to every element of D an element of R .

Functions

A function is three pieces of information

- A domain, $D \subset \mathbb{R}$
- A range, $R \subset \mathbb{R}$, and
- A rule $f : D \rightarrow R$ that assigns to every element of D an element of R .

Functions

A function is three pieces of information

- A domain, $D \subset \mathbb{R}$
- A range, $R \subset \mathbb{R}$, and
- A rule $f : D \rightarrow R$ that assigns to every element of D an element of R .

Example

The functions

are all different functions!

Functions

A function is three pieces of information

- A domain, $D \subset \mathbb{R}$
- A range, $R \subset \mathbb{R}$, and
- A rule $f : D \rightarrow R$ that assigns to every element of D an element of R .

Example

The functions

- $f : \mathbb{R} \rightarrow \mathbb{R}; x \mapsto x^2$

are all different functions!

Functions

A function is three pieces of information

- A domain, $D \subset \mathbb{R}$
- A range, $R \subset \mathbb{R}$, and
- A rule $f : D \rightarrow R$ that assigns to every element of D an element of R .

Example

The functions

- $f : \mathbb{R} \rightarrow \mathbb{R}; x \mapsto x^2$
- $f : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}; x \mapsto x^2$

are all different functions!

Functions

A function is three pieces of information

- A domain, $D \subset \mathbb{R}$
- A range, $R \subset \mathbb{R}$, and
- A rule $f : D \rightarrow R$ that assigns to every element of D an element of R .

Example

The functions

- $f : \mathbb{R} \rightarrow \mathbb{R}; x \mapsto x^2$
- $f : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}; x \mapsto x^2$
- $f : \mathbb{R} \rightarrow \mathbb{R}_{\geq 0}; x \mapsto x^2$

are all different functions!

Global Maximums and minimums

Definition (Global maximum)

A function $f : D \rightarrow R$ has a global maximum at a if

$$f(x) \leq f(a) \quad \text{for all } x \in D$$

Global Maximums and minimums

Definition (Global maximum)

A function $f : D \rightarrow R$ has a global maximum at a if

$$f(x) \leq f(a) \quad \text{for all } x \in D$$


Definition (Global minimum)

A function $f : D \rightarrow R$ has a global minimum at a if

$$f(x) \geq f(a) \quad \text{for all } x \in D$$

Example of a global minimum

$f : \mathbb{R} \rightarrow \mathbb{R}; x \mapsto x^2$ has a min at $x = 0$



Example of a global maximum

$f : (-\infty, 0] \rightarrow \mathbb{R}; f(x) = x^3$ has a max at $x = 0$

