

This week you will practice writing differential equations modelling real world phenomena as well as understanding population models. You will also get practice solving separable differential equations.

Homework: The homework will be due on Friday 17 January, at 8am, the *start* of the lecture. It will consist of questions:

2 and 7.

*Numbers in parentheses indicate the question has been taken from the textbook:

S. J. Schreiber, *Calculus for the Life Sciences*, Wiley,

and refer to the section and question number in the textbook.

1. (6.1) Write a differential equation to model the situations described below. Do not try to solve.
 - (a) (6.1-1) The number of bacteria in a culture grows at a rate that is proportional to the number of bacteria present.
 - (b) (6.1-2) A sample of radium decays at a rate that is proportional to the amount of radium present in the sample.
 - (c) (6.1-5) According to Benjamin Gompertz (1779-1865) the growth rate of a population is proportional to the number of individuals present, where the factor of proportionality is an exponentially decreasing function of time.
 - (d) (6.1-7) The rate at which an epidemic spreads through a community of P susceptible people is proportional to the product of the number of people y who have caught the disease and the number $P - y$ who have not.
 - (e) (6.1-8) The rate at which people are implicated in a government scandal is proportional to the product of the number N of people already implicated and the number of people involved who have not yet been implicated.

2. (6.1) A population model is given by

$$\frac{dP}{dt} = P(100 - P).$$

- (a) (6.1-9) For what values is the population at equilibrium?
 - (b) (6.1-10) For what values is $\frac{dP}{dt} > 0$?
 - (c) (6.1-11) For what values is $\frac{dP}{dt} < 0$?
 - (d) (6.1-12) Describe how the fate of the population depends on the initial density.
3. (6.1) A population model is given by

$$\frac{dP}{dt} = P(P - 1)(100 - P).$$

- (a) (6.1-13) For what values is the population at equilibrium?
 - (b) (6.1-14) For what values is $\frac{dP}{dt} > 0$?
 - (c) (6.1-15) For what values is $\frac{dP}{dt} < 0$?
 - (d) (6.1-16) Describe how the fate of the population depends on the initial density.
4. (6.1) Radioactive decay: Certain types of atoms (e.g. carbon-14, xenon-133, lead-210, etc.) are inherently unstable. They exhibit random transitions to a different atom while emitting radiation in the process. Based on experimental evidence, Rutherford found in the early 20th century that the number, N , of atoms in a radioactive substance can be described by the equation

$$\frac{dN}{dt} = -\lambda N$$

where t is measured in years and $\lambda > 0$ is known as the *decay constant*. The decay constant is found experimentally by measuring the half life, τ of the radioactive substance (i.e. the time it takes for half of the substance to decay). Use this information in the following problems.

- (a) (6.1-18) Find a solution to the decay equation assuming that $N(0) = N_0$.
 - (b) (6.1-19) For xenon-133, the half-life is 5 days. Find λ . Assume t is measured in days.
 - (c) (6.1-20) For carbon-14 the half life is 5,568 years. Find the decay constant λ , assuming t is measured in years.
 - (d) (6.1-21) How old is a piece of human bone which contains just 60% of the amount of carbon-14 expected in a sample of bone from a living person, assuming the half life of carbon-14 is 5,568 years?
 - (e) (6.1-22) The Dead Sea Scrolls were written on parchment at about 100 B.C. What percentage of carbon-14 originally contained in the parchment remained when the scrolls were discovered in 1947?
5. (6.1-30) Hyperthyroidism is caused by a new growth of tumor-like cells that secrete thyroid hormones in excess to the normal hormones. If left untreated, a hyperthyroid individual can exhibit extreme weight loss, anorexia, muscle weakness, heart disease intolerance to stress, and eventually death. The most successful and least invasive treatment option is radioactive iodine-131 therapy.

This involves the injection of a small amount of radioactivity into the body. For the type of hyperthyroidism called Graves disease, it is usual for about 40 – 80% of the administered activity to concentrate in the thyroid gland. For functioning adenomas (hot nodules), the uptake is closer to 20 – 30%. Excess iodine-131 is excreted rapidly by the kidneys. The quantity of radioiodine used to treat hyperthyroidism is not enough to injure any tissue except the thyroid tissue, which slowly shrinks over a matter of weeks to months. Radioactive iodine is either swallowed in a capsule or sipped in solution through a straw. A typical dose is 5 – 15 millicuries. The half-life of iodine-131 is 8 days.

- (a) Suppose that it takes 48 hours for a shipment of iodine-131 to reach a hospital. How much of the initial amount shipped is left once it arrives at the hospital?
- (b) Suppose a patient is given a dosage of 10 millicuries of which 30% concentrates in the thyroid gland. How much is left one week later?
- (c) Suppose a patient is given a dosage of 10 millicuries of which 30% concentrates in the thyroid gland. How much is left 30 days later?