

EQUALITY CASES OF THE ALEXANDROV–FENCHEL INEQUALITY ARE NOT IN THE POLYNOMIAL HIERARCHY

SWEE HONG CHAN AND IGOR PAK

ABSTRACT. Describing the equality conditions of the *Alexandrov–Fenchel inequality* has been a major open problem for decades. We prove that in the case of convex polytopes, this description is not in the polynomial hierarchy unless the polynomial hierarchy collapses to a finite level. This is the first hardness result for the problem, and is a complexity counterpart of the recent result by Shenfeld and van Handel [SvH23], which gave a geometric characterization of the equality conditions. The proof involves Stanley’s *order polytopes* and employs poset theoretic technology.

1. INTRODUCTION

1.1. Foreword. Geometric inequalities play a central role in convex geometry, probability and analysis, with numerous combinatorial and algorithmic applications. The *Alexandrov–Fenchel (AF) inequality* lies close to the heart of convex geometry. It is one of the deepest and most general results in the area, generalizing a host of simpler geometric inequalities such as the *isoperimetric inequality* and the *Brunn–Minkowski inequality*, see §3.1.

The equality conditions for geometric inequalities are just as fundamental as the inequalities themselves, and are crucial for many applications, see §10.2. For simpler inequalities they tend to be straightforward and follow from the proof. As the inequalities become more complex, their proofs became more involved, and the equality cases become more numerous and cumbersome. This is especially true for the Alexandrov–Fenchel inequality, where the complete description of the equality cases remain open despite much effort and many proofs, see §3.2.

We use the language and ideas from computational complexity and tools from poset theory, to prove that the equality cases of the Alexandrov–Fenchel inequality cannot be explicitly described for convex polytopes in a certain formal sense. We give several applications to stability in geometric inequalities and to combinatorial interpretation of the defect of poset inequalities. We also raise multiple questions, both mathematical and philosophical, see Section 10.

1.2. Alexandrov–Fenchel inequality. Let $V(Q_1, \dots, Q_n)$ denote the *mixed volume* of convex bodies Q_1, \dots, Q_n in \mathbb{R}^n (see below). The *Alexandrov–Fenchel inequality* states that for convex bodies $K, L, Q_1, \dots, Q_{n-2}$ in \mathbb{R}^n , we have:

$$(AF) \quad V(K, L, Q_1, \dots, Q_{n-2})^2 \geq V(K, K, Q_1, \dots, Q_{n-2}) \cdot V(L, L, Q_1, \dots, Q_{n-2}).$$

Let polytope $K \subset \mathbb{R}^n$ be defined by a system of inequalities $A\mathbf{x} \leq \mathbf{b}$. We say that K is a *TU-polytope* if vector $\mathbf{b} \in \mathbb{Z}^n$, and matrix A is *totally unimodular*, i.e. all its minors have determinants in $\{0, \pm 1\}$. Note that all vertices of TU-polytopes are integral. Denote by EQUALITYAF the *equality verification problem of the Alexandrov–Fenchel inequality*, defined as the decision problem whether (AF) is an equality.

Theorem 1.1 (Main theorem). *Let $K, L, Q_1, \dots, Q_{n-2} \subset \mathbb{R}^n$ be TU-polytopes. Then the equality verification problem of the Alexandrov–Fenchel inequality (AF) is not in the polynomial hierarchy unless the polynomial hierarchy collapses to a finite level:*

$$\text{EQUALITYAF} \in \text{PH} \implies \text{PH} = \Sigma_m^P \quad \text{for some } m.$$

Informally, the theorem says that the equality cases of the Alexandrov–Fenchel inequality (AF) are *unlikely to have a description in the polynomial hierarchy*.¹ This is in sharp contrast with other geometric inequalities, including many special cases of (AF), when the equality cases have an explicit description, thus allowing an efficient verification (see §3.1).

Let us emphasize that constraining to TU-polytopes makes the theorem stronger rather than weaker. Indeed, one would hope that the equality verification problem is easy at least in the case when both vertices and facets are integral (cf. §10.3). In fact, we chose the *smallest* natural class of H-polytopes which contains all order polytopes (see below).

Let us quickly unpack the very strong claim of Theorem 1.1. In particular, the theorem implies that given the polytopes, the equality in (AF) cannot be decided in polynomial time: EQUALITYAF \notin P, nor even in probabilistic polynomial time: EQUALITYAF \notin BPP (unless PH collapses). Moreover, there can be no polynomial size certificate which verifies that (AF) is an equality: EQUALITYAF \notin NP, or a strict inequality: EQUALITYAF \notin coNP (ditto).

Our results can be viewed as a complexity theoretic counterpart of the *geometric description* of the Alexandrov–Fenchel inequality that was proved recently by Shenfeld and van Handel [SvH23]. In this context, Theorem 1.1 says that this geometric description is not computationally effective, and cannot be made so under standard complexity assumptions. From this point of view, the results in [SvH23] are optimal, at least for convex polytopes in the full generality (cf. §10.12).

Warning: Here we only give statements of the results without a context. Our hands are tied by the interdisciplinary nature of the paper with an extensive background in both convex geometry, poset theory and computational complexity. We postpone the definitions until Section 2, and the review until Section 3.

1.3. Stability. In particular, Theorem 1.1 prohibits certain *stability inequalities*. In the context of general inequalities, these results give quantitative measurements of how close are the objects of study (variables, surfaces, polytopes, lattice points, etc.) to the equality cases in some suitable sense, when the inequality is close to an equality, see e.g. [Fig13].

In the context of geometric inequalities, many sharp stability results appear in the form of *Bonnesen type inequality*, see [Oss79]. These are defined as the strengthening of a geometric inequality $f \geq g$ to $f - g \geq h$, such that $h \geq 0$, and $h = 0$ if and only if $f = g$.² They are named after the celebrated extension of the *isoperimetric inequality* by Bonnesen (see §3.3).

While there are numerous Bonnesen type inequalities of various strength for the Brunn–Minkowski inequalities and their relatives, the case of Alexandrov–Fenchel inequality (AF) remains unapproachable in full generality. Formally, define the *Alexandrov–Fenchel defect* as:

$$\delta(K, L, Q_1, \dots, Q_{n-2}) := V(K, L, Q_1, \dots, Q_{n-2})^2 - V(K, K, Q_1, \dots, Q_{n-2}) \cdot V(L, L, Q_1, \dots, Q_{n-2}).$$

One would want to find a bound of the form $\delta(\cdot) \geq \xi(\cdot)$, where ξ is a nonnegative computable function of the polytopes. The following result is an easy corollary from the proof of Theorem 1.1.

Corollary 1.2. *Suppose $\delta(K, L, Q_1, \dots, Q_{n-2}) \geq \xi(K, L, Q_1, \dots, Q_{n-2})$ is a Bonnesen type inequality, such that ξ is computable in polynomial time on all TU-polytopes. Then PH = NP.*

Informally, the corollary implies that for the stability of the AF inequality, one should either avoid polytopes altogether and require some regularity conditions for the convex bodies (as has been done in the past, see §3.3), or be content with functions ξ which are hard to compute (such inequalities can still be very useful, of course). See §10.10 for further implications.

To understand how the corollary follows from the proof of Theorem 1.1, the Bonnesen condition in this case states that $\xi(\cdot) = 0$ if and only if $\delta(\cdot) = 0$. Thus, the equality $\{\delta(\cdot) = 0\}$ can be decided in polynomial time on TU-polytopes, giving the assumption in the theorem.

¹The collapse in the theorem contradicts standard assumptions in computational complexity. A conjecture that the collapse does not happen is a strengthening of the $P \neq NP$ conjecture that remains out of reach, see §3.8.

²Following [Oss79], function h should also have a (not formally defined) “geometric description”.

1.4. Stanley inequality. We restrict ourselves to a subset of TU-polytopes given by the *order polytopes* (see §2.4). Famously, Stanley showed in [Sta81], that the Alexandrov–Fenchel inequality applied to certain such polytopes gives the *Stanley inequality*, that the numbers of certain linear extensions of finite posets form a log-concave sequence. This inequality is of independent interest in order theory (see §3.4), and is the starting point of our investigation.

Let $P = (X, \prec)$ be a poset with $|X| = n$ elements. Denote $[n] := \{1, \dots, n\}$. A *linear extension* of P is a bijection $f : X \rightarrow [n]$, such that $f(x) < f(y)$ for all $x \prec y$. Denote by $\mathcal{E}(P)$ the set of linear extensions of P , and let $e(P) := |\mathcal{E}(P)|$.

Let $x, z_1, \dots, z_k \in X$ and $a, c_1, \dots, c_k \in [n]$; we write $\mathbf{z} = (z_1, \dots, z_k)$ and $\mathbf{c} = (c_1, \dots, c_k)$, and we assume without loss of generality that $c_1 < \dots < c_k$. Let $\mathcal{E}_{\mathbf{z}\mathbf{c}}(P, x, a)$ be the set of linear extensions $f \in \mathcal{E}(P)$, such that $f(x) = a$ and $f(z_i) = c_i$ for all $1 \leq i \leq k$. Denote by $N_{\mathbf{z}\mathbf{c}}(P, x, a) := |\mathcal{E}_{\mathbf{z}\mathbf{c}}(P, x, a)|$ the number of such linear extensions. The *Stanley inequality* [Sta81] states that the sequence $\{N_{\mathbf{z}\mathbf{c}}(P, x, a), 1 \leq a \leq n\}$ is *log-concave*:

$$(Sta) \quad N_{\mathbf{z}\mathbf{c}}(P, x, a)^2 \geq N_{\mathbf{z}\mathbf{c}}(P, x, a+1) \cdot N_{\mathbf{z}\mathbf{c}}(P, x, a-1).$$

The problem of finding the equality conditions for (Sta) was first asked by Stanley in the original paper [Sta81, §3], see also [BT02, Question 6.3], [CPP23b, §9.9] and [MS24]. Formally, for every $k \geq 0$, denote by EQUALITYSTANLEY_k the *equality verification problem of the Stanley inequality with k fixed elements*, defined as the decision problem whether (Sta) is an equality. It was shown by Shenfeld and van Handel that $\text{EQUALITYSTANLEY}_0 \in \text{P}$, see [SvH23, Thm 15.3].

Theorem 1.3. *Let $k \geq 2$. Then the equality verification problem of the Stanley inequality (Sta) is not in the polynomial hierarchy unless the polynomial hierarchy collapses to a finite level:*

$$\text{EQUALITYSTANLEY}_k \in \text{PH} \implies \text{PH} = \Sigma_m^{\text{P}} \quad \text{for some } m.$$

In Section 5, we deduce Theorem 1.1 from Theorem 1.3. For the proof, any fixed k in (Sta) suffices, of course. In the opposite direction, we prove the following extension of the Shenfeld and van Handel’s result mentioned above:

Theorem 1.4. $\text{EQUALITYSTANLEY}_1 \in \text{P}$.

Together, Theorems 1.3 and 1.4 complete the analysis of equality cases of Stanley’s inequality.

1.5. Combinatorial interpretation. The problem of finding a *combinatorial interpretation* is fundamental in both enumerative and algebraic combinatorics, and was the original motivation of this investigation (see §3.7). Although very different in appearance and technical details, there are certain natural parallels with the stability problems discussed above.

Let $f \geq g$ be an inequality between two counting functions $f, g \in \#P$. We say that $(f - g)$ has a *combinatorial interpretation*, if $(f - g) \in \#P$. While many combinatorial inequalities have a combinatorial interpretation, for the Stanley inequality (Sta) this is an open problem. Formally, let

$$\Phi_{\mathbf{z}\mathbf{c}}(P, x, a) := N_{\mathbf{z}\mathbf{c}}(P, x, a)^2 - N_{\mathbf{z}\mathbf{c}}(P, x, a+1) \cdot N_{\mathbf{z}\mathbf{c}}(P, x, a-1)$$

denote the defect in (Sta). Let $\phi_k : (P, X^{k+1}, [n]^{k+1}) \rightarrow \mathbb{N}$ be the function computing $\Phi_{\mathbf{z}\mathbf{c}}(P, x, a)$.

Corollary 1.5. *For all $k \geq 2$, function ϕ_k does not have a combinatorial interpretation unless the polynomial hierarchy collapses to the second level:*

$$\phi_k \in \#P \implies \text{PH} = \Sigma_2^{\text{P}}.$$

To see some context behind this result, note that $N_{\mathbf{zc}}(P, x, a) \in \#P$ by definition, so $\phi_k \in \text{GapP}_{\geq 0}$, a class of nonnegative functions in $\text{GapP} := \#P - \#P$. We currently know very few functions which are in $\text{GapP}_{\geq 0}$ but not in $\#P$. The examples include

$$(\circledast) \quad (\#3\text{SAT}(F) - 1)^2, \quad (\#2\text{SAT}(F) - \#2\text{SAT}(F'))^2 \quad \text{and} \quad (e(P) - e(P'))^2,$$

where F, F' are CNF Boolean formulas and P, P' are posets [CP23a, IP22]. In other words, all three functions in (\circledast) *do not* have a combinatorial interpretation (unless PH collapses). The corollary provides the first *natural* example of a defect function that is $\text{GapP}_{\geq 0}$ but not in $\#P$.

The case $k = 0$, whether $\phi_0 \in \#P$, is especially interesting and remains a challenging open problem, see [CPP23b, §9.12] and [Pak22, Conj. 6.3]. The corollary suggests that Stanley's inequality (Sta) is unlikely to have a direct combinatorial proof, see §10.9.

To understand how the corollary follows from the proof of Theorem 1.3, note that $\phi_2 \in \#P$ implies that there is a polynomial certificate for the Stanley inequality being strict. In other words, we have $\text{EQUALITYSTANLEY}_2 \in \text{coNP}$, giving the assumption in the theorem.

Structure of the paper. We begin with definitions and notation in Section 2, followed by the lengthy background and literature review in Section 3 (see also §10.1). In the key Section 4, we give proofs of Theorems 1.1 and 1.3, followed by proofs of Corollaries 1.2 and 1.5. These results are reduced to several independent lemmas, which are proved one by one in Sections 5–8. We prove Theorem 1.4 in Section 9. This section is independent of the previous sections (except for notation in §6.1). We conclude with extensive final remarks and open problems in Section 10.

2. DEFINITIONS AND NOTATION

2.1. General notation. Let $[n] = \{1, \dots, n\}$, $\mathbb{N} = \{0, 1, 2, \dots\}$ and $\mathbb{Z}_{\geq 1} = \{1, 2, \dots\}$. For a subset $S \subseteq X$ and element $x \in X$, we write $S + x := S \cup \{x\}$ and $S - x := S \setminus \{x\}$. For a sequence $\mathbf{a} = (a_1, \dots, a_m)$, denote $|\mathbf{a}| := a_1 + \dots + a_m$. This sequence is *log-concave*, if $a_i^2 \geq a_{i-1}a_{i+1}$ for all $1 < i < m$.

2.2. Mixed volumes. Fix $n \geq 1$. For two sets $A, B \subset \mathbb{R}^n$ and constants $\alpha, \beta > 0$, denote by

$$\alpha A + \beta B := \{\alpha \mathbf{x} + \beta \mathbf{y} : \mathbf{x} \in A, \mathbf{y} \in B\}$$

the *Minkowski sum* of these sets. For a convex body $K \subset \mathbb{R}^n$ with affine dimension d , denote by $\text{Vol}_d(K)$ the volume of K . We drop the subscript when $d = n$.

One of the basic result in convex geometry is *Minkowski's theorem*, see e.g. [BZ88, §19.1], that the volume of convex bodies with affine dimension d behaves as a homogeneous polynomial of degree d with nonnegative coefficients:

Theorem 2.1 (Minkowski). *For all convex bodies $K_1, \dots, K_r \subset \mathbb{R}^n$ and $\lambda_1, \dots, \lambda_r > 0$, we have:*

$$(2.1) \quad \text{Vol}_d(\lambda_1 K_1 + \dots + \lambda_r K_r) = \sum_{1 \leq i_1, \dots, i_d \leq r} V(K_{i_1}, \dots, K_{i_d}) \lambda_{i_1} \cdots \lambda_{i_d},$$

where the functions $V(\cdot)$ are nonnegative and symmetric, and where d is the affine dimension of $\lambda_1 K_1 + \dots + \lambda_r K_r$ (which does not depend on the choice of $\lambda_1, \dots, \lambda_r$).

The coefficients $V(A_{i_1}, \dots, A_{i_d})$ are called *mixed volumes* of A_{i_1}, \dots, A_{i_d} . We refer to [HW20, Lei80, Schn14] for an accessible introduction to the subject.

2.3. Posets. For a poset $P = (X, \prec)$ and a subset $Y \subset X$, denote by $P_Y = (Y, \prec)$ a *subposet* of P . We use $(P - z)$ to denote a subposet P_{X-z} , where $z \in X$. Element $x \in X$ is *minimal* in P , if there exists no element $y \in X - x$ such that $y \prec x$. Define *maximal* elements similarly. Denote by $\min(P)$ and $\max(P)$ the set of minimal and maximal elements in P , respectively.

In a poset $P = (X, \prec)$, elements $x, y \in X$ are called *parallel* or *incomparable* if $x \not\prec y$ and $y \not\prec x$. We write $x \parallel y$ in this case. *Comparability graph* is a graph on X , with edges (x, y) where $x \prec y$. Element $x \in X$ is said to *cover* $y \in X$, if $y \prec x$ and there are no elements $z \in X$ such that $y \prec z \prec x$.

A *chain* is a subset $C \subset X$ of pairwise comparable elements. The *height* of poset $P = (X, \prec)$ is the maximum size of a chain. An *antichain* is a subset $A \subset X$ of pairwise incomparable elements. The *width* of poset $P = (X, \prec)$ is the size of the maximal antichain.

A *dual poset* is a poset $P^* = (X, \prec^*)$, where $x \prec^* y$ if and only if $y \prec x$. A *disjoint sum* $P + Q$ of posets $P = (X, \prec)$ and $Q = (Y, \prec')$ is a poset $(X \cup Y, \prec^\diamond)$, where the relation \prec^\diamond coincides with \prec and \prec' on X and Y , and $x \parallel y$ for all $x \in X, y \in Y$. A *linear sum* $P \oplus Q$ of posets $P = (X, \prec)$ and $Q = (Y, \prec')$ is a poset $(X \cup Y, \prec^\diamond)$, where the relation \prec^\diamond coincides with \prec and \prec' on X and Y , and $x \prec^\diamond y$ for all $x \in X, y \in Y$.

Posets constructed from one-element posets by recursively taking disjoint and linear sums are called *series-parallel*. Both *n-chain* C_n and *n-antichain* A_n are examples of series-parallel posets. *Forest* is a series-parallel poset formed by recursively taking disjoint sums (as before), and linear sums with one element: $C_1 \oplus P$. We refer to [Sta12, Ch. 3] for an accessible introduction, and to surveys [BW00, Tro95] for further definitions and standard results.

2.4. Poset polytopes. Let $P = (X, \prec)$ be a poset with $|X| = n$ elements. The *order polytope* $\mathcal{O}_P \subset \mathbb{R}^n$ is defined as

$$(2.2) \quad 0 \leq \alpha_x \leq 1 \quad \text{for all } x \in X, \quad \alpha_x \leq \alpha_y \quad \text{for all } x \prec y, \quad x, y \in X.$$

Similarly, the *chain polytope* (also known as the *stable set polytope*) $\mathcal{S}_P \subset \mathbb{R}^n$ is defined as

$$(2.3) \quad \beta_x \geq 0 \quad \text{for all } x \in X, \quad \beta_x + \beta_y + \dots \leq 1 \quad \text{for all } x \prec y \prec \dots, \quad x, y, \dots \in X.$$

In [Sta86], Stanley computed the volume of both polytopes:

$$(2.4) \quad \text{Vol}(\mathcal{O}_P) = \text{Vol}(\mathcal{S}_P) = \frac{e(P)}{n!}.$$

This connection is the key to many applications of geometry to poset theory and vice versa.

2.5. Terminology. For functions $f, g : X \rightarrow \mathbb{R}$, we write $f \geq g$, if $f(x) \geq g(x)$ for all $x \in X$. For an inequality $f \geq g$, the *defect* is a function $h := f - g$. The *equality cases* to describe the set of $x \in X$ such that $f(x) = g(x)$. Denote by $X_h := \{x \in X : h(x) = 0\} \subseteq X$ the subset of equality cases.

We use E_h to denote the *equality verification* of $f(x) = g(x)$, i.e. the decision problem

$$E_h := \{f(x) \stackrel{?}{=} g(x)\},$$

where $x \in X$ is an input. Since $E_h = \{x \in X_h\}$, this is a special case of the *inclusion problem*. We use V_h to denote the *verification* of $h(x) = a$, i.e. the decision problem

$$V_h := \{f(x) - g(x) \stackrel{?}{=} a\},$$

where $a \in \mathbb{R}$ and $x \in X$ are the input. Clearly, V_h is a more general problem than E_h .

For a subset $Y \subseteq X$, we use *description* for an equivalent condition for the inclusion problem $\{x \in Y\}$, where $x \in X$. We use *equality conditions* for a description of E_h . We say that equality cases of $f \geq g$ have a *description in the polynomial hierarchy* if $E_h \in \text{PH}$. In other words, there is a CNF Boolean formula $\Phi(y_1, y_2, y_3, \dots, x)$, such that

$$\forall x \in X : E_h \iff \exists y_1 \forall y_2 \exists y_3 \dots \Phi(y_1, y_2, y_3, \dots, x).$$

2.6. Complexity. We assume that the reader is familiar with basic notions and results in computational complexity and only recall a few definitions. We use standard complexity classes: P , FP , NP , $coNP$, $\#P$, Σ_m^P and PH . The notation $\{a =^? b\}$ is used to denote the decision problem whether $a = b$. We use the *oracle notation* R^S for two complexity classes $R, S \subseteq PH$, and the polynomial closure $\langle A \rangle$ for a problem $A \in PSPACE$. We will also use less common classes

$$\text{GapP} := \{f - g \mid f, g \in \#P\} \quad \text{and} \quad C=P := \{f(x) =^? g(y) \mid f, g \in \#P\}.$$

Note that $coNP \subseteq C=P$.

We also assume that the reader is familiar with standard decision and counting problems: $3SAT$, $\#3SAT$ and $PERMANENT$. Denote by $\#LE$ the problem of computing the number $e(P)$ of linear extensions. For a counting function $f \in \#P$, the *coincidence problem* is defined as:

$$C_f := \{f(x) =^? f(y)\}.$$

Note the difference with the equality verification problem E_{f-g} defined above. Clearly, we have both $E_{f-g} \in C=P$ and $C_f \in C=P$. Note also that $C_{\#3SAT}$ is both $C=P$ -complete and $coNP$ -hard.

The distinction between *binary* and *unary* presentation will also be important. We refer to [GJ78] and [GJ79, §4.2] for the corresponding notions of NP -completeness and *strong* NP -completeness. Unless stated otherwise, we use the word “*reduction*” to mean “polynomial Turing reduction”. We refer to [AB09, Gol08, Pap94] for definitions and standard results in computational complexity.

3. BACKGROUND AND HISTORICAL OVERVIEW

3.1. Geometric inequalities. The history of equality conditions of geometric inequalities goes back to antiquity, see e.g. [Blå05, Por33], when it was discovered that the *isoperimetric inequality* (Isop)

$$\ell(X)^2 \geq 4\pi a(X)$$

is an equality if and only if X is a circle. Here $\ell(X)$ is the perimeter and $a(X)$ is the area of a convex $X \subset \mathbb{R}^2$. This classical result led to numerous extensions and generalizations leading to the Alexandrov–Fenchel inequality (AF). We refer to [BZ88, Schn14] for a review of the literature.

Below we highlight only the most important developments to emphasize how the equality conditions become more involved as one moves in the direction of the AF inequality (see also §10.4 and §10.5). The celebrated *Brunn–Minkowski inequality* states that for all convex $K, L \subset \mathbb{R}^d$, we have:

$$(BM) \quad \text{Vol}(K + L)^{1/d} \geq \text{Vol}(K)^{1/d} + \text{Vol}(L)^{1/d},$$

see e.g. [Gar02] for a detailed survey. This inequality “plays an important role in almost all branches of mathematics” [Bar07]. Notably, both Brunn and Minkowski showed the equality in (BM) holds if and only if K is an expansion of L .

For the *mean width inequality*

$$(MWI) \quad s(K)^2 \geq 6\pi w(K) \text{Vol}(K),$$

for all convex $K \subset \mathbb{R}^3$, Minkowski conjectured (1903) the equality cases are the *cap bodies* (balls with attached tangent cones). Here $s(K)$ is the surface area and $w(K)$ is the *mean width* of K . Minkowski’s conjecture was proved by Bol (1943), see e.g. [BF34, BZ88].

The *Minkowski’s quadratic inequality* for three convex bodies $K, L, M \subset \mathbb{R}^3$, states:

$$(MQI) \quad V(K, L, M)^2 \geq V(K, K, M) \cdot V(L, L, M).$$

This is a special case of (AF) for $n = d = 3$. When $L = B_1$ is a unit ball and $K = M$, this gives (MWI). Favard [Fav33, p. 248] wrote that the equality conditions for (MQI) “*parait difficile à énoncer*” (“seem difficult to state”). There are even interesting families of convex polytopes that give equality cases (see e.g. [SvH23, Fig. 2.1]).

Shenfeld and van Handel [SvH22] gave a complete characterization of the equality cases of (MQI) as triples of convex bodies that are similarly truncated in a certain formal sense. Notably, for the full-dimensional H-polytopes in \mathbb{R}^3 , each with at most n facets, the equality conditions amount to checking $O(n)$ linear relations for distances between facet inequalities. This can be easily done in polynomial time.

3.2. Alexandrov–Fenchel inequality. For the AF inequality (AF), the equality conditions have long believed to be out of reach, as they would generalize those for (MWI) and (MQI). Alexandrov made a point of this in his original 1937 paper:

“Serious difficulties occur in determining the conditions for equality to hold in the general inequalities just derived” [Ale37, §4].

Half a century later, Burago and Zalgaller reviewed the literature and summarized:

“A conclusive study of all these situations when the equality sign holds has not been carried out, probably because they are too numerous” [BZ88, §20.5].

Schneider made a case for perseverance:

“As (AF) represents a classical inequality of fundamental importance and with many applications, the identification of the equality cases is a problem of intrinsic geometric interest. Without its solution, the Brunn–Minkowski theory of mixed volumes remains in an uncompleted state.” [Schn94, p. 426].

The AF inequality has a number of proofs using ideas from convex geometry, analysis and algebraic geometry, going back to two proofs by Alexandrov (Fenchel’s full proof never appeared). We refer to [BZ88, Schn14] for an overview of the older literature, especially [Schn14, p. 398] for historical remarks, and to [BL23, CP22, CKMS19, KK12, SvH19, Wang18] for some notable recent proofs. All these proofs use a limit argument at the end, which can create new equality cases that do not hold for generic convex bodies. This partially explains the difficulty of the problem (cf. §10.2 and [SvH22, Rem. 3.7]).

In [Ale37], Alexandrov gave a description of equality cases for combinatorially isomorphic polytopes. This is a large family of full-dimensional polytopes, for which every convex body is a limit. In particular, he showed that for the full-dimensional axis-parallel boxes $[\ell_1 \times \dots \times \ell_n]$, the equality in (AF) is equivalent to K and L being homothetic (cf. §10.6).

In the pioneering work [Schn85], Schneider published a conjectural description of the equality cases, corrected later by Ewald [Ewa88], see also [Schn14]. After many developments, this conjecture was confirmed for all smooth (full-dimensional) convex bodies Q_i [Schn90a], and for all (not necessarily full-dimensional) convex bodies $Q_1 = \dots = Q_{n-2}$, by Shenfeld and van Handel [SvH23]. Closer to the subject of this paper, in a remarkable development, the authors gave a geometric description of the equality cases for all convex polytopes. They explain:

“Far from being esoteric, it is precisely the case of convex bodies with empty interior (which is not covered by previous conjectures) that arises in combinatorial applications” [SvH23, §1.3].

The geometric description of the equality cases in [SvH23] is indirect, technically difficult to prove, and computationally hard in the degenerate cases.³ While we will not quote the full statement (Theorem 2.13 in [SvH23]), let us mention the need to find witnesses polytopes $M_i, N_i \subset \mathbb{R}^n$ which must satisfy certain conditions (Def. 2.10, *ibid.*) The second of these conditions is an equality of certain mixed volumes (Eq. (2.4), *ibid.*)

In [SvH23, §2.2.3], the authors write: “Condition (2.4) should be viewed merely as a normalization”.⁴ From the computational complexity point of view, asking for the equality of mixed

³It follows from our Theorem 1.1 that it has to be, see a discussion in §10.12.

⁴By that the authors of [SvH23] seem to mean that their description captured all the geometry in the problem, as opposed to the equality of mixed volumes which has no geometric content.

volumes (known to be hard to compute, see §3.8), lifts the problem outside of the polynomial hierarchy, to a hard coincidence problem (see §2.6). This coincidence problem eventually percolated into [MS24], see (3.3) below, which in turn led directly to this work.

3.3. Stability. *Bonnesen's inequality* is an extension of the isoperimetric inequality (Isop), which states that for every convex $X \subset \mathbb{R}^2$, we have:

$$(Bon) \quad \ell(X)^2 - 4\pi a(X) \geq 4\pi(R - r)^2,$$

where R is the smallest radius of the circumscribed circle, and r is the maximal radius of the inscribed circle.⁵ Moreover, Bonnesen proved [Bon29], that there is an *annulus* (thin shell) U between concentric circles of radii $R \geq r$, such that $\partial X \subseteq U$ and (Bon) holds. Note that the optimal such annulus can be computed in polynomial time, see [AAHS99].

Bonnesen's inequality (Bon) was an inspiration for many Bonnesen type inequalities [Oss78, Oss79, Gro90]. See also discrete versions in §10.4, and applications in computational geometry in [KS99]. There is now an extensive literature on stability inequalities in geometric and more general context, see e.g. [Fig13, Gro93].

There is an especially large literature on the stability of the Brunn–Minkowski inequality (BM). For major early advances by Diskant (1973), Groemer (1988) and others, see e.g. [Gro93] and references therein. We refer to [Fig14] for an overview of more recent results, including [FMP09, FMP10]. See also [EK14] for the thin shell type bounds, and [FJ17] for the stability of (BM) for *nonconvex sets*.

For the Alexandrov–Fenchel inequality (AF), there are very few stability results, all for the full dimensional convex bodies with various regularity conditions, see e.g. [Mar17, Schn90b].

3.4. Linear extensions. Linear extensions play a central role in enumerative combinatorics and order theory. They appear in connection with saturated chains in *distributive lattices*, *standard Young tableaux* and *P-partitions*, see e.g. [Sta12].

The world of inequalities for linear extensions has a number of remarkable results, some with highly nontrivial equality conditions. Notably, the *Björner–Wachs inequality* for $e(P)$ is an equality if and only if P is a forest [BW89, Thm 6.3], see also [CPP23b]. On the other hand, the celebrated *XYZ inequality* established by Shepp in [She82] (see also [AS16, §6.4]), has no nontrivial equality cases [Fis84].

An especially interesting example is the *Sidorenko inequality*

$$(3.1) \quad e(P) \cdot e(P^\circ) \geq n!$$

for posets P, P° on the same ground set with n elements, which have complementary comparability graphs [Sid91] (other proofs are given in [CPP23b, GG22]). Sidorenko also proved that the series-parallel posets are the only equality cases. This solves the equality verification problem of (3.1), since the recognition problem of series-parallel posets is in P, see [VTL82].

It was noticed in [BBS99], that the Sidorenko inequality follows from *Mahler's conjecture*, which states that for every convex centrally symmetric body $K \subset \mathbb{R}^n$, we have:

$$(3.2) \quad \text{Vol}(K) \cdot \text{Vol}(K^\circ) \geq \frac{4^n}{n!}.$$

To derive (3.1) from (3.2), take K to be the union of all axis reflections of the chain polytope \mathcal{S}_P defined in (2.3). Mahler's conjecture (3.2) is known for all axis symmetric convex bodies [StR81], but remains open in full generality [AASS20], in part due to the many equality cases [Tao08, §1.3].

⁵Note that when $X \subset \mathbb{R}$ is a nonzero interval, we have $r = 0$ and $\ell(X) = 4R$, so the inequality remains strict.

3.5. Stanley inequality. Stanley’s inequality (Sta) is of independent interest in order theory, having inspired a large literature especially in the last few years. The case $k = 0$ is especially interesting. The unimodality in this case was independently conjectured by Kislitsyn [Kis68] and Rivest, while the log-concavity was conjectured by Chung, Fishburn and Graham [CFG80], who established both conjectures for posets of width two. Stanley proved them in [Sta81] in full generality.⁶ The authors of [CFG80] called Rivest’s conjecture “tantalizing” and Stanley’s proof “very ingenious”.

The *Kahn–Saks inequality* is a generalization of the $k = 0$ case of (Sta), and is also proved from the AF inequality. This inequality was used to obtain the first positive result in the direction of the $\frac{1}{3} - \frac{2}{3}$ conjecture [KS84]. For posets of width two, both the $k = 0$ case of the Stanley inequality, and the Kahn–Saks inequality have natural q -analogues [CPP23a]. A generalization of Stanley’s inequality to *marked posets* was given in [LMS19].

For all $k \geq 0$, the *vanishing conditions* $\{N_{\mathbf{zc}}(P, x, a) =^? 0\}$ are in \mathbf{P} . This was shown by David and Jacqueline Daykin in [DD85, Thm 8.2], via explicit necessary and sufficient conditions. Recently, this result was rediscovered in [CPP23b, Thm 1.11] and [MS24, Thm 5.3]. Similarly, the *uniqueness conditions* $\{N_{\mathbf{zc}}(P, x, a) =^? 1\}$ are in \mathbf{P} by the result of Panova and the authors [CPP23b, Thm 7.5], where we gave explicit necessary and sufficient conditions. Both the vanishing and the uniqueness conditions give examples of equality cases of the Stanley inequality, which remained a “major challenge” in full generality [CPP23b, §9.10].

As we mentioned in the introduction, Shenfeld and van Handel resolved the $k = 0$ case of Stanley equality conditions by giving explicit necessary and sufficient conditions, which can be verified in polynomial time, see [SvH23]. Similar explicit necessary and sufficient conditions for the Kahn–Saks inequality were conjectured in [CPP23a, Conj. 8.7], and proved for posets of width two. Building on the technology in [SvH23], van Handel, Yan and Zeng gave the proof of this conjecture in [vHYZ23].

In [CP21], we gave a new proof of the $k = 0$ case of (Sta), using a *combinatorial atlas* technology. This is an inductive self-contained linear algebraic approach; see [CP22] for the introduction. We also gave a new proof of the Shenfeld and van Handel equality conditions, and generalized both results to *weighted linear extensions* (see §§1.16–18 in [CP21]).

In an important development, Ma and Shenfeld [MS24] advanced the technology of [SvH23], to give a clean albeit ineffective combinatorial description of the equality cases in full generality. In particular, they showed that (AF) is an equality if and only if

$$(3.3) \quad N_{\mathbf{zc}}(P, x, a - 1) = N_{\mathbf{zc}}(P, x, a) = N_{\mathbf{zc}}(P, x, a + 1).$$

They proceeded to give explicit necessary and sufficient conditions for these equalities in some cases (see §10.11). About the remaining cases that they called *critical* (see §9.2), they write: “It is an interesting problem to find [an explicit description] for critical posets” [MS24, Rem. 1.6]. Our Theorem 1.3 implies that such a description is unlikely, as it would imply a disproof of a major conjecture in computational complexity (see also §10.12).

3.6. Complexity aspects. There are two standard presentations of polytopes: *H-polytopes* described by the inequalities and *V-polytopes* described by the vertices. These two presentation types have very different nature in higher dimensions, see e.g. [DGH98]. We refer to [GK94, GK97] for an overview of standard complexity problems in geometry, and to [Schr86, §19], [Schr03, §5.16], for the background on totally unimodular matrices and TU-polytopes. Note also that testing whether matrix A is totally unimodular can be done in polynomial time, see [Sey80].

When the dimension n is bounded, H-polytopes and V-polytopes have the same complexity, so the volume and the mixed volumes are in \mathbf{FP} . Thus, the dimension n is unbounded throughout the paper. The volume of TU-polytopes is $\#\mathbf{P}$ -hard via reduction to KNAPSACK [DF88]. Note

⁶For $k \geq 1$, the inequality (Sta) is sometimes called the *generalized Stanley inequality*, see [CPP23b].

that for *rational* H-polytopes in \mathbb{R}^n , the volume denominators can be doubly exponential [Law91], thus not in PSPACE. This is why we constrain ourselves to TU-polytopes which is a subclass of H-polytopes that includes all order polytopes (see §5.1).

The mixed volume $V(Q_1, \dots, Q_n)$ coincides with the permanent when all Q_i are axis parallel boxes, see [vL82] and §10.6. Thus, computing the mixed volume is #P-hard even for the boxes, see [DGH98]. For rational H-polytopes, the vanishing problem $\{V(\cdot) \stackrel{?}{=} 0\}$ can be described combinatorially, and is thus in NP, see [DGH98, Est10]. It is equivalent to computing the rank of intersection of two geometric matroids (with a given realization), which is in P, see [Schr03, §41]. For TU-polytopes in \mathbb{R}^n , the uniqueness problem $\{V(\cdot) \stackrel{?}{=} \frac{1}{n!}\}$ is in NP by a result in [EG15].

The problem #LE is proved #P-complete by Brightwell and Winkler [BW91, Thm 1], and this holds even for posets of height two [DP20]. Linial noticed [Lin86], that this result and (2.4) together imply that the volume of H-polytopes is #P-hard even when the input is in unary. Linial also observed that the number of vertices of order polytopes is #P-complete (ibid.).

Now, fix $k \geq 0$, $x \in X$ and $\mathbf{z} \in X^k$. Clearly, we have:

$$e(P) = \sum_{a \in [n]} \sum_{\mathbf{c} \in [n]^k} N_{\mathbf{z}\mathbf{c}}(P, x, a),$$

where the summation has size $O(n^{k+1})$. Thus, computing $N_{\mathbf{z}\mathbf{c}}(P, x, a)$ is also #P-complete.

Finally, it was proved in [CP23a], that $C_{\#3SAT}$, $C_{\text{PERMANENT}}$ and $C_{\#LE}$ are not in PH, unless PH collapses to a finite level. The proof idea of Theorem 1.3 is inspired by these results.

3.7. Combinatorial interpretations. Finding a combinatorial interpretation is a standard problem throughout combinatorics, whenever a positivity phenomenon or an inequality emerges. Having a combinatorial interpretation allows one to deeper understand the underlying structures, give asymptotic and numerical estimates, as well as analyze certain algorithms. We refer to [Huh18, Sta89, Sta00] for an overview of inequalities in algebraic combinatorics and matroid theory, and to [Pak22] for a recent survey from the complexity point of view.

Recall that $\text{GapP} := \#P - \#P$ is a the class of difference of two #P functions, and let $\text{GapP}_{\geq 0}$ be a subclass of GapP of nonnegative functions. Thus, for every inequality $f \geq g$ of counting functions $f, g \in \#P$, we have $(f - g) \in \text{GapP}_{\geq 0}$. It was shown in [IP22, Prop. 2.3.1], that $\text{GapP}_{\geq 0} \neq \#P$, unless $\text{PH} = \Sigma_2^P$. The key example is

$$(\#3SAT(F) - \#3SAT(F'))^2,$$

see also the first function in (*). The other two functions in (*) were given in [CP23a]. A *natural* $\text{GapP}_{\geq 0}$ problem of computing S_n character squared: $[\chi^\lambda(\mu)]^2$, was proved not in #P (in unary), under the same assumptions [IPP22].

The idea that some natural combinatorial inequalities can have no combinatorial interpretations appeared in [Pak19]. A number of interesting examples were given in [IP22, §7], including the *Cauchy, Minkowski, Hadamard, Karamata* and *Ahlsvede–Daykin inequalities*, all proved not in #P under varying complexity assumptions.

Closer to the subject of this paper, Ikenmeyer and the second author showed that the AF defect $\delta(\cdot)$ is not in #P (unless $\text{PH} = \Sigma_2^P$), even for axis parallel rectangles in \mathbb{R}^2 whose edge length are given by #3SAT formulas [IP22, Thm 7.1.5]. This is a nonstandard model of computation. One can think of our Main Theorem 1.1 as a tradeoff: in exchange for needing a higher dimension, we now have unary input and the standard model of computation.

3.8. Complexity assumptions. The results in the paper use different complexity assumptions, and navigating between them can be confusing. Here is short list of standard implications:

$$\text{PH} \neq \Sigma_m^P \text{ for all } m \geq 2 \implies \text{PH} \neq \Sigma_2^P \implies \text{PH} \neq \text{NP} \implies \text{P} \neq \text{NP}.$$

In other words, the assumption in Theorems 1.1 and 1.3 is the strongest, while $P \neq NP$ is the weakest. Proving either of these would be a major breakthrough in theoretical computer science. Disproving either of these would bring revolutionary changes to the way the computational complexity understands the nature of computation. We refer to [Aar16, Wig19] for an extensive discussion, philosophy and implications in mathematics and beyond.

4. PROOF ROADMAP

The results in the paper follow from a series of largely independent polynomial reductions and several known results. In this section, we only state the reductions whose proofs will be given in the next few sections. We then deduce both theorems from these reductions.

4.1. Around Stanley equality. First, we show that Theorem 1.1 follows from Theorem 1.3. Recall the notation from the introduction. Let $P = (X, \prec)$ be a poset on $|X| = n$ elements. As before, let $x \in X$, $a \in [n]$, $\mathbf{z} \in X^k$, and $\mathbf{c} \in [n]^k$. Recall also

$$\text{EQUALITYAF} := \{V(K, L, Q_1, \dots, Q_{n-2})^2 \stackrel{?}{=} V(K, K, Q_1, \dots, Q_{n-2}) \cdot V(L, L, Q_1, \dots, Q_{n-2})\},$$

$$\text{EQUALITYSTANLEY}_k := \{N_{\mathbf{z}\mathbf{c}}(P, x, a)^2 \stackrel{?}{=} N_{\mathbf{z}\mathbf{c}}(P, x, a+1) \cdot N_{\mathbf{z}\mathbf{c}}(P, x, a-1)\}.$$

Proposition 4.1 (cf. [Sta81, §3]). *For all $k \geq 0$, EQUALITYSTANLEY_k reduces to EQUALITYAF .*

The proof of the proposition is given in Section 5, is very close to Stanley’s original proof of the inequality (Sta). The key difference is the observation that slices of order polytopes are TU-polytopes. Next, we need a simple technical result.

Lemma 4.2. *For all $k > \ell$, $\text{EQUALITYSTANLEY}_\ell$ reduces to EQUALITYSTANLEY_k .*

Proof. Let $P = (X, \prec)$ be a poset on n elements, and let $\mathbf{z} \in X^k$, $\mathbf{c} \in [n]^k$, $x \in X$, $a \in [n]$ be as in §1.4. Denote by $P' := P + A_{k-\ell}$ a poset obtained by adding $(k - \ell)$ independent elements $z'_1, \dots, z'_{k-\ell}$. Let $c'_i := n + i$, for all $1 \leq i \leq k - \ell$. For $\mathbf{z}' := (z_1, \dots, z_\ell, z'_1, \dots, z'_{k-\ell})$ and $\mathbf{c}' := (c_1, \dots, c_\ell, c'_1, \dots, c'_{k-\ell})$, we have:

$$N_{\mathbf{z}'\mathbf{c}'}(P', x, a) = N_{\mathbf{z}\mathbf{c}}(P, x, a).$$

Varying a , we conclude that EQUALITYSTANLEY_k is equivalent to $\text{EQUALITYSTANLEY}_\ell$ in this special case. This gives the desired reduction. \square

Next, we simplify the Stanley equality problem to the following *flatness problem*:

$$\text{FLATLE}_k := \{N_{\mathbf{z}\mathbf{c}}(P, x, a) \stackrel{?}{=} N_{\mathbf{z}\mathbf{c}}(P, x, a+1)\},$$

where $N_{\mathbf{z}\mathbf{c}}(P, x, a)$ are defined in §1.4. The idea is to ask whether a is in the flat part of the distribution of $f(x)$ (cf. Figure 15.1 in [SvH23]).

Lemma 4.3. *For all $k \geq 0$, FLATLE_k reduces to $\text{EQUALITYSTANLEY}_{k+2}$.*

We prove Lemma 4.3 in Section 6.

4.2. Relative numbers of linear extensions. Let $P = (X, \prec)$ be a poset on $|X| = n$ elements, and let $\min(P) \subseteq X$ be the set of minimal elements of P . For every $x \in \min(P)$, define the *relative number of linear extensions*:

$$(4.1) \quad \rho(P, x) := \frac{e(P)}{e(P - x)}.$$

In other words, $\rho(P, x) = \mathbb{P}[f(x) = 1]^{-1}$, where $f \in \mathcal{E}(P)$ is a uniform random linear extension of P . Denote by $\#RLE$ the problem of computing $\rho(P, x)$.

Lemma 4.4. *$\#RLE$ is polynomial time equivalent to $\#LE$.*

Proof. By definition, $\#RLE$ reduces to $\#LE$. In the opposite direction, let $P = (X, \prec)$ be a poset on $|X| = n$ elements. Fix a linear extension $g \in \mathcal{E}(P)$, and let $x_i := g^{-1}(i)$, $1 \leq i \leq n$. Denote by P_i a subposet of P restricted to x_i, \dots, x_n and observe that $x_i \in \min(P_i)$. We have:

$$e(P) = \frac{e(P_1)}{e(P_2)} \cdot \frac{e(P_2)}{e(P_3)} \cdots = \rho(P_1, x_1) \cdot \rho(P_2, x_2) \cdots,$$

which gives the desired reduction from $\#LE$ to $\#RLE$. \square

We relate RLE to flatness equality through the following series of reductions. Consider the following coincidence problem:

$$CRLE := \{ \rho(P, x) \stackrel{?}{=} \rho(Q, y) \},$$

where $P = (X, \prec)$, $Q = (Y, \prec')$ are posets, and $x \in \min(P)$, $y \in \min(Q)$.

Lemma 4.5 (see Theorem 7.1). *CRLE reduces to $FLATLE_0$.*

Next, consider the following decision problem:

$$QUADRLE := \{ \rho(P_1, x_1) \cdot \rho(P_2, x_2) \stackrel{?}{=} \rho(P_3, x_3) \cdot \rho(P_4, x_4) \},$$

where P_1, P_2, P_3, P_4 are finite posets and $x_i \in \min(P_i)$, for all $1 \leq i \leq 4$.

Lemma 4.6 (see Theorem 7.2). *QUADRLE reduces to CRLE.*

4.3. Verification lemma. Let $P = (X, \prec)$ be a poset on $|X| = n$ elements, and let $x \in \min(P)$. Consider

$$(4.2) \quad VERRLE := \left\{ \rho(P, x) \stackrel{?}{=} \frac{A}{B} \right\},$$

where A, B are coprime integers with $1 \leq B \leq A \leq n!$. We need the following:

Lemma 4.7 (Verification lemma). $NP^{(VERRLE)} \subseteq NP^{(QUADRLE)}$.

Note that the opposite direction “ \supseteq ” is also true and easy to prove. Indeed, suppose you have an oracle $VERRLE$. Guess the values $a_i := \rho(P_i, x_i) \in \mathbb{Q}$, verify that they are correct, and check that $a_1 \cdot a_2 = a_3 \cdot a_4$. This gives $QUADRLE$. We will only need the direction in the lemma which is highly nontrivial.

4.4. Putting everything together. We can now obtain all the results stated in the introduction, except for Theorem 1.4 which uses different tools and is postponed until Section 9.

Proof of Theorem 1.3. Recall that #LE is #P-complete [BW91] (see also §3.6). By Lemma 4.4, we conclude that #RLE is #P-hard. We then have:

$$(4.3) \quad \text{PH} \subseteq \text{P}^{\#P} \subseteq \text{P}^{\langle \#RLE \rangle} \subseteq \text{NP}^{\langle \text{VERRLE} \rangle}$$

where the first inclusion is Toda’s theorem [Toda91], the second inclusion is because #RLE is #P-hard, and the third inclusion is because one can simulate #RLE by first guessing and then verifying the answer.

Fix $k \geq 2$. Combining Lemmas 4.2, 4.3, 4.5 and 4.6, we conclude that QUADRLE reduces to EQUALITYSTANLEY_k. We have:

$$(4.4) \quad \text{NP}^{\langle \text{VERRLE} \rangle} \subseteq \text{NP}^{\langle \text{QUADRLE} \rangle} \subseteq \text{NP}^{\langle \text{EQUALITYSTANLEY}_k \rangle},$$

where the first inclusion is the Verification Lemma 4.7. Now, suppose EQUALITYSTANLEY_k ∈ PH. Then EQUALITYSTANLEY_k ∈ Σ_m^P for some m . Combining (4.3) and (4.4), this implies:

$$(4.5) \quad \text{PH} \subseteq \text{NP}^{\langle \text{EQUALITYSTANLEY}_k \rangle} \subseteq \text{NP}^{\Sigma_m^P} \subseteq \Sigma_{m+1}^P,$$

as desired. \square

As a byproduct of the proof, we get the same conclusion for the intermediate problems. This result is potentially of independent interest (cf. [CP23a]).

Corollary 4.8. *Problems VERRLE, QUADRLE, CRLE and FLATLE₀ are not in PH, unless PH = Σ_m^P for some m .*

Proof of Theorem 1.1. The result follows from Proposition 4.1 and Theorem 1.3. \square

Proof of Corollary 1.2. By the “Bonnesen type” assumption, we have

$$\{\xi(\cdot) \stackrel{?}{=} 0\} \iff \{\delta(\cdot) \stackrel{?}{=} 0\} = \text{EQUALITYAF}.$$

Since computing ξ is in FP, we have EQUALITYAF ∈ P. Then (4.5) for $k = 2$, and Proposition 4.1 give:

$$(4.6) \quad \text{PH} \subseteq \text{NP}^{\langle \text{EQUALITYSTANLEY}_2 \rangle} \subseteq \text{NP}^{\langle \text{EQUALITYAF} \rangle} \subseteq \text{NP}^P = \text{NP},$$

as desired. \square

Proof of Corollary 1.5. Suppose $\phi_k \in \#P$. By definition, we have:

$$\{\Phi_{\mathbf{zc}}(P, x, a) \neq^? 0\} \in \text{NP}.$$

In other words, we have EQUALITYSTANLEY_k ∈ coNP. Then (4.5) gives:

$$\text{PH} \subseteq \text{NP}^{\langle \text{EQUALITYSTANLEY}_k \rangle} \subseteq \text{NP}^{\text{coNP}} = \Sigma_2^P,$$

as desired. \square

5. AF EQUALITY FROM STANLEY EQUALITY

5.1. Slices of order polytopes. Let $P = (X, \prec)$ be a poset on $|X| = n$ elements. Recall the construction of order polytopes $\mathcal{O}_P \subseteq [0, 1]^n$ given in (2.2). Fix $z_1 \prec \dots \prec z_k$ and $1 \leq c_1 < \dots < c_k \leq n$. Denote $Z := \{z_1, \dots, z_k\}$ and let $Y := X \setminus Z$. For all $0 \leq i \leq k$, consider the following *slices* of the order polytopes:

$$S_i := \mathcal{O}_P \cap \{\alpha_x = 0 : x \prec z_i, x \in X\} \cap \{\alpha_x = 1 : x \succ z_{i+1}, x \in X\}.$$

Here the conditions $x \prec z_i$ and $x \succ z_{i+1}$ are vacuous when $i = 0$ and $i = k$, respectively. Note that $\dim S_i \leq n - k$ for all $0 \leq i \leq k$, since α_x is a constant on S_i for all $x \in Z$.⁷ The same argument implies that these slices are themselves order polytopes of subposets of P , a fact we do not need. Instead, we need the following simple result:

Lemma 5.1. *Slices S_i are TU-polytopes.*

Proof. Write K_i in the form $A \cdot (\alpha_y)_{y \in Y} \leq \mathbf{b}$. Observe that A has $\{-1, 0, 1\}$ entries, and so does \mathbf{b} . Every square submatrix B of A corresponds to taking a subposet with added rows of 0's, or with rows of 0's and a single ± 1 . By definition of \mathcal{O}_P , we can rearrange columns in B to make it upper triangular. Thus, $\det(B) \in \{-1, 0, 1\}$, as desired. \square

5.2. Proof of Proposition 4.1. Denote by $\mathcal{E}_{\mathbf{zc}}(P)$ the set of all linear extensions $f \in \mathcal{E}(P)$, such that $f(z_i) = c_i$ for all i , and let $N_{\mathbf{zc}}(P) := |\mathcal{E}_{\mathbf{zc}}(P)|$.

Let $S_0, \dots, S_k \subset \mathbb{R}^n$ be the slices defined above, and note that $\dim \langle S_0, \dots, S_k \rangle = n - k$. Stanley's original proof of (Sta) is based on the following key observation:

Lemma 5.2 ([Sta81, Thm 3.2]). *Let $z_1 \prec \dots \prec z_k$ and $1 \leq c_1 < \dots < c_k \leq n$. We have:*

$$(5.1) \quad V(\underbrace{S_0, \dots, S_0}_{c_1 - 1 \text{ times}}, \underbrace{S_1, \dots, S_1}_{c_2 - c_1 - 1 \text{ times}}, \dots, \underbrace{S_k, \dots, S_k}_{n - c_k \text{ times}}) = \frac{1}{(n - k)!} N_{\mathbf{zc}}(P).$$

Now let $z_i \leftarrow x$ and $c_i \leftarrow a$ for some i , such that $1 \leq c_1 < \dots < c_k \leq n$. By Lemma 5.2, the AF inequality (AF) becomes (Sta). By Lemma 5.1, slices $S_i \subset \mathbb{R}^n$ are TU-polytopes defined by $O(n^2)$ inequalities. This gives the desired reduction. \square

6. STANLEY EQUALITY FROM FLATNESS

6.1. Ma–Shenfeld poset notation. Recall the following terminology from [MS24]. For $s \in \{-1, 0, 1\}$ and $f \in \mathcal{E}_{\mathbf{zc}}(P, x, a + s)$, the *companions* in f are the elements in

$$\text{Com}(f) := \{f^{-1}(a - 1), f^{-1}(a), f^{-1}(a + 1)\} - x.$$

Note that $|\text{Com}(f)| = 2$ for all s as above. Let the *lower companion* $\text{lc}(f) \in \text{Com}(f)$ be the companion with smaller of the two values in f . Similarly, let the *upper companion* $\text{uc}(f) \in \text{Com}(f)$ be the companion with larger of the two values in f . Denote by $\mathcal{C}(x) \subset X$ the set of elements $y \in X$ comparable to x , i.e. $\mathcal{C}(x) := \{y \in X : x \prec y \text{ or } x \succ y\}$.

⁷In geometric language, slices S_i are sections of the order polytope \mathcal{O}_P with a k -dimensional affine subspace.

6.2. Proof of Lemma 4.3. Let $P = (X, \prec)$, and let $x, a, \mathbf{z} = (z_1, \dots, z_k)$ and $\mathbf{c} = (c_1, \dots, c_k)$ be an instance of FLATLE_k as above. To prove the reduction in the lemma, we construct a poset $Q = (Y, \prec)$ for which P is a subposet, and x, b, \mathbf{y} and \mathbf{x} , which give the desired instance $\text{EQUALITYSTANLEY}_{k+2}$.

Without loss of generality, we can assume that $\min(P) = \{z_0\}$ and $\max(P) = \{z_{k+1}\}$. In other words, assume that there are elements $z_0, z_{k+1} \in X$ such that $z_0 \prec y \prec z_{k+1}$ for all $y \in X$.

Let M_1, M_2, M_3 be given by

$$\begin{aligned} M_1 &:= |\{f \in \mathcal{E}_{\mathbf{z}\mathbf{c}}(P, x, a) : f^{-1}(a+1) \succ x\}|, \\ M_2 &:= |\{f \in \mathcal{E}_{\mathbf{z}\mathbf{c}}(P, x, a+1) : f^{-1}(a) \prec x\}|, \\ M_3 &:= |\{f \in \mathcal{E}_{\mathbf{z}\mathbf{c}}(P, x, a) : f^{-1}(a+1) \parallel x\}| \\ &= |\{f \in \mathcal{E}_{\mathbf{z}\mathbf{c}}(P, x, a+1) : f^{-1}(a) \parallel x\}|. \end{aligned}$$

Note that the two sets in the definition of M_3 are in bijection with each other via the map that swaps $f(a)$ with $f(a+1)$. It then follows from here that

$$N_{\mathbf{z}\mathbf{c}}(P, x, a) = M_1 + M_3 \quad \text{and} \quad N_{\mathbf{z}\mathbf{c}}(P, x, a+1) = M_2 + M_3.$$

This implies that

$$(6.1) \quad N_{\mathbf{z}\mathbf{c}}(P, x, a) = N_{\mathbf{z}\mathbf{c}}(P, x, a+1) \quad \Longleftrightarrow \quad M_1 = M_2.$$

Now, let $Q = (Y, \prec)$ be the poset $P + C_3$, i.e. $Y := X \cup \{u, v, w\}$ and with the additional relation $u \prec v \prec w$ and $\{u, v, w\}$ is incomparable to all elements in X . Let $\ell := \max\{i : c_i < a\}$ be the maximal index such that the corresponding element in \mathbf{z} is less than a . Let $b := a + 2$, and let

$$\begin{aligned} \mathbf{y} &:= (z_1, \dots, z_\ell, u, w, z_{\ell+1}, \dots, z_k) \in Y^{k+2}, \\ \mathbf{b} &:= (c_1, \dots, c_\ell, a, a+4, c_{\ell+1}+3, \dots, c_k+3) \in \mathbb{N}^{k+2}. \end{aligned}$$

In the notation above, for $s \in \{-1, 0, 1\}$ and $f \in \mathcal{E}_{\mathbf{y}\mathbf{b}}(Q, x, b+s)$, the companions in f are the elements in

$$\text{Com}(f) := \{f^{-1}(b-1), f^{-1}(b), f^{-1}(b+1)\} - x.$$

Let⁸

$$\begin{aligned} \mathcal{F}(b, \text{com}, \text{inc}) &:= \{f \in \mathcal{E}_{\mathbf{y}\mathbf{b}}(Q, x, b) : \text{lc}(f) \in \mathcal{C}(x), \text{uc}(f) \notin \mathcal{C}(x)\}, \\ \mathcal{F}(b, \text{inc}, \text{com}) &:= \{f \in \mathcal{E}_{\mathbf{y}\mathbf{b}}(Q, x, b) : \text{lc}(f) \notin \mathcal{C}(x), \text{uc}(f) \in \mathcal{C}(x)\}, \\ \mathcal{F}(b, \text{com}, \text{com}) &:= \{f \in \mathcal{E}_{\mathbf{y}\mathbf{b}}(Q, x, b) : \text{lc}(f) \in \mathcal{C}(x), \text{uc}(f) \in \mathcal{C}(x)\}, \\ \mathcal{F}(b, \text{inc}, \text{inc}) &:= \{f \in \mathcal{E}_{\mathbf{y}\mathbf{b}}(Q, x, b) : \text{lc}(f) \notin \mathcal{C}(x), \text{uc}(f) \notin \mathcal{C}(x)\}, \end{aligned}$$

and we write $F(b, \cdot, \cdot) := |\mathcal{F}(b, \cdot, \cdot)|$. Note that by construction it follows that, for all $f \in \mathcal{F}(b, \cdot, \cdot)$, we have

$$b-2 = f(u) < f(v) < f(w) = b+2,$$

so $f(v) \in \{b-1, b, b+1\}$, and thus v will always be a companion in f . Sets $\mathcal{F}(b+1, *, *)$ and $\mathcal{F}(b-1, *, *)$ are defined analogously.

⁸We warn the reader that from this point on our notation is substantially different from that in [MS24].

Claim 6.1. *We have:*

$$\begin{aligned}
F(b, \text{com}, \text{inc}) &= M_2, & F(b, \text{inc}, \text{com}) &= M_1, \\
F(b, \text{com}, \text{com}) &= 0, & F(b, \text{inc}, \text{inc}) &= 2M_3, \\
F(b+1, \text{com}, \text{inc}) &= M_2, & F(b+1, \text{inc}, \text{com}) &= M_2, \\
F(b+1, \text{com}, \text{com}) &= 0, & F(b+1, \text{inc}, \text{inc}) &= 2M_3, \\
F(b-1, \text{com}, \text{inc}) &= M_1, & F(b-1, \text{inc}, \text{com}) &= M_1, \\
F(b-1, \text{com}, \text{com}) &= 0, & F(b-1, \text{inc}, \text{inc}) &= 2M_3.
\end{aligned}$$

Proof. We only compute the values $F(b, *, *)$, as proof of the other cases is analogous. Denote by $\mathcal{E}_{\mathbf{zc}}(P)$ the set of all linear extensions $f \in \mathcal{E}(P)$, such that $f(z_i) = c_i$ for all i .

Let $\psi : \mathcal{E}_{\mathbf{yb}}(Q) \rightarrow \mathcal{E}_{\mathbf{zc}}(P)$ be the map given by $\psi(f) = g$, where

$$g(s) := \begin{cases} f(s) & \text{if } f(s) < f(u), \\ f(s) - 1 & \text{if } f(u) < f(s) < f(v), \\ f(s) - 2 & \text{if } f(v) < f(s) < f(w), \\ f(s) - 3 & \text{if } f(s) > f(w), \end{cases}$$

for all $s \in X$. It follows from the definition of $\text{lc}(f)$ and $\text{uc}(f)$, that

$$\mathcal{F}(b, \text{com}, \text{inc}) = \{f \in \mathcal{E}_{\mathbf{yb}}(Q, x, b) : f^{-1}(b-1) \prec x, f^{-1}(b+1) = v\},$$

It then follows that φ restricted to $\mathcal{F}(b, \text{com}, \text{inc})$ is a bijection onto

$$\{g \in \mathcal{E}_{\mathbf{zc}}(P, x, a+1) : g^{-1}(a+1) \prec x\},$$

which gives us $F(b, \text{com}, \text{inc}) = M_2$. Similar arguments gives $F(b, \text{inc}, \text{com}) = M_1$. Note that $F(b, \text{com}, \text{com}) = 0$, because v is always a companion in f but $v \parallel x$ by definition. Note also that

$$\begin{aligned}
\mathcal{F}(b, \text{inc}, \text{inc}) &= \{f \in \mathcal{E}_{\mathbf{yb}}(Q, x, b) : f^{-1}(b-1) \parallel x, f^{-1}(b+1) = v\} \\
&\cup \{f \in \mathcal{E}_{\mathbf{yb}}(Q, x, b) : f^{-1}(b+1) \parallel x, f^{-1}(b-1) = v\}.
\end{aligned}$$

It then follows that ψ restricted to $\mathcal{F}(b, \text{com}, \text{inc})$ is a bijection onto

$$\{g \in \mathcal{E}_{\mathbf{zc}}(P, x, a+1) : g^{-1}(a) \parallel x\} \cup \{g \in \mathcal{E}_{\mathbf{zc}}(P, x, a) : g^{-1}(a+1) \parallel x\},$$

which gives $F(b, \text{inc}, \text{inc}) = 2M_3$. This finishes proof of the claim. \square

By the claim, we have:

$$\begin{aligned}
N_{\mathbf{yb}}(Q, c, b) &= F(b, \text{com}, \text{inc}) + F(b, \text{inc}, \text{com}) + F(b, \text{com}, \text{com}) + F(b, \text{inc}, \text{inc}) \\
&= M_2 + M_1 + 2M_3.
\end{aligned}$$

Similarly, we have:

$$\begin{aligned}
N_{\mathbf{yb}}(Q, x, b+1) &= 2M_2 + 2M_3, \\
N_{\mathbf{yb}}(Q, x, b-1) &= 2M_1 + 2M_3.
\end{aligned}$$

We conclude:

$$\begin{aligned}
N_{\mathbf{yb}}(Q, x, b)^2 - N_{\mathbf{yb}}(Q, x, b+1) \cdot N_{\mathbf{yb}}(Q, x, b-1) \\
= (M_1 + M_2 + 2M_3)^2 - 4(M_1 + M_3)(M_2 + M_3) = (M_1 - M_2)^2.
\end{aligned}$$

This implies that

$$(6.2) \quad N_{\mathbf{yb}}(Q, x, b)^2 = N_{\mathbf{yb}}(Q, x, b+1) \cdot N_{\mathbf{yb}}(Q, x, b-1) \iff M_1 = M_2.$$

Lemma 4.3 now follows by combining (6.1) and (6.2). \square

7. FLATNESS FROM THE QUADRUPLE RELATIVE RATIO

Recall several key definitions from Section 4. Let $N(R, z, c)$ be the number of linear extensions $f \in \mathcal{E}(R)$ for which $f(z) = c$. Similarly, let

$$(7.1) \quad \text{FLATLE}_0 := \{ N(R, z, c) \stackrel{?}{=} N(R, z, c+1) \},$$

where $R = (Z, \prec^\circ)$ is a finite poset on $|Z| = \ell$ elements, $z \in Z$ and $1 \leq c \leq \ell$. Finally, let

$$(7.2) \quad \text{CRLE} := \{ \rho(P, x) \stackrel{?}{=} \rho(Q, y) \},$$

where $P = (X, \prec)$, $Q = (Y, \prec')$ are posets, and $x \in \min(P)$, $y \in \min(Q)$.

7.1. One poset from two. The following result give a quantitative version⁹ of Lemma 4.5.

Theorem 7.1. *CRLE reduces to FLATLE₀. More precisely, suppose we have a poset $P = (X, \prec)$ on $n = |X|$ elements, a poset $Q = (Y, \prec')$ on $m = |Y|$ elements, and $x \in \min(P)$, $y \in \min(Y)$. Then there exists a polynomial time construction of a poset $R = (Z, \prec^\circ)$ on $\ell := |Z| = m + n$ elements, $z \in Z$, and $c \in [\ell]$, such that (7.1) \Leftrightarrow (7.2).*

Proof. Let $P^* = (X, \prec^*)$ be the dual poset of P . Define $R = (Z, \prec^\circ)$ to be a poset on

$$Z := (X - x) \cup (Y - y) \cup \{w, z\},$$

where w, z are two new elements. Let the partial order \prec° coincide with \prec^* on $(X - x)$, and with \prec' on $(Y - y)$, with additional relations

$$(7.3) \quad p \prec^\circ z \prec^\circ q, \quad \text{for all } p \in X - x, q \in Y - y,$$

$$(7.4) \quad p \prec^\circ w \text{ if and only if } x \prec p, \quad \text{for all } p \in X - x,$$

$$(7.5) \quad w \prec^\circ q \text{ if and only if } y \prec' q, \quad \text{for all } q \in Y - y.$$

That is, we are taking the series sum $(P^* - x) \oplus \{z\} \oplus (Q - y)$, then adding an element w to emulate x in P for $f(w) < f(z)$, where $f \in \mathcal{E}(R)$, and emulate y in Q when $f(w) > f(z)$. It then follows from a direct calculation that

$$N(R, z, n+1) = e(P) \cdot e(Q - y).$$

Indeed, by (7.3), for every $f \in \mathcal{N}(R, z, n+1)$ we have:

$$\{f^{-1}(1), \dots, f^{-1}(n)\} = X - x + w, \quad \{f^{-1}(n+2), \dots, f^{-1}(m+n)\} = Y - y.$$

These two labelings define a linear extension of P^* after a substitution $w \leftarrow x$ given by (7.4), and a linear extension $Q - y$. By an analogous argument, we have:

$$N(R, z, n) = e(P^* - x) \cdot e(Q),$$

Set $c \leftarrow n$. Combining these two observations, we get

$$\frac{N(R, z, c+1)}{N(R, z, c)} = \frac{N(R, z, n+1)}{N(R, z, n)} = \frac{\rho(P, x)}{\rho(Q, y)},$$

which gives the desired reduction and proves the result. \square

⁹We do not actually need the precise bounds below, other than the fact that they are at most polynomial. However, these bounds help to clarify the construction.

7.2. Two posets from four. Now recall the decision problem

$$(7.6) \quad \text{QUADRLE} := \{ \rho(P_1, x_1) \cdot \rho(P_2, x_2) \stackrel{?}{=} \rho(P_3, x_3) \cdot \rho(P_4, x_4) \}.$$

The following result give a quantitative version of Lemma 4.6.

Theorem 7.2. *QUADRLE reduces to CRLE. More precisely, for every $P_i = (X_i, \prec_i)$ posets on $n_i = |X_i|$ elements, and $x_i \in \min(P_i)$, $1 \leq i \leq 4$, there exists a polynomial time construction of a poset $P = (X, \prec)$ on $n := |X| \leq n_1 + \max\{n_2, n_3\} + 1$ elements, of a poset $Q = (Y, \prec')$ on $m := |Y| \leq n_4 + \max\{n_2, n_3\} + 1$ elements, such that $(7.2) \Leftrightarrow (7.6)$.*

We now build toward the proof of this theorem.

Lemma 7.3. *Let $P = (X, \prec)$ and $Q = (Y, \prec')$ be posets with $m = |X|$ and $n = |Y|$ elements, respectively. Let $x \in \min(P)$ and $y \in \min(Q)$. Then there exists a poset $R = (Z, \prec^\circ)$ and $z \in \min(P)$, such that $|Z| = m + n + 1$ and*

$$\rho(R, z) = m + \left(1 + \frac{\rho(Q, y)}{\rho(P, x)}\right)^{-1}.$$

Proof. Let $P^* = (X, \prec^*)$ denotes the dual poset to P . Let $R := (Z, \prec^\circ)$ be given by

$$Z := (X - x) \cup (Y - y) \cup \{v, w, z\}.$$

where \prec° inherits the partial order \prec^* on $X - x$, the partial order \prec' on $Y - y$, and with additional relations:

$$\begin{aligned} p \prec^\circ v \prec^\circ q & \quad \forall p \in X - x, \quad q \in Y - y, \\ p \prec^\circ w & \Leftrightarrow p \prec^* x \quad \forall p \in X - x, \\ q \succ^\circ w & \Leftrightarrow q \succ' y \quad \forall q \in Y - y, \\ z \parallel_{\prec^\circ} p \quad \forall p \in X - x, \quad z \prec^\circ q \quad \forall q \in Y - y, \\ z \prec^\circ v, \quad z \parallel_{\prec^\circ} w. \end{aligned}$$

That is, we are taking the series sum $(P^* - x) \oplus \{v\} \oplus (Q - y)$, then adding an element w to emulate x in P for all $f(w) < f(v)$, emulate y in Q for all $f(w) > f(v)$, and finally adding z to track the value of $f(v)$. Here the linear extension $f \in \mathcal{E}(R)$ in each case. By construction, we have either $f(v) = m + 1$ or $f(v) = m + 2$.

Claim. We have:

$$(7.7) \quad e(R) = m e(P - x) e(Q) + (m + 1) e(P) e(Q - y).$$

Proof of claim. Let us show that the first term $m e(P - x) e(Q)$ is the number linear extensions $f \in \mathcal{E}(R)$ s.t. $f(v) = m + 1$. For such f we have:

$$\begin{aligned} \{f^{-1}(1), \dots, f^{-1}(m)\} &= X - x + z, \\ \{f^{-1}(m + 2), \dots, f^{-1}(m + n + 1)\} &= Y - y + w. \end{aligned}$$

Note that the restriction of f to $\{f^{-1}(1), \dots, f^{-1}(m)\}$ defines a linear extension of $(P^* - x + z)$. Additionally, note that the restriction of f to $\{f^{-1}(m + 2), \dots, f^{-1}(m + n + 1)\}$ defines a linear extension of Q . In total, we have $e(P^* - x + z) e(Q) = m e(P - x) e(Q)$ linear extensions f as above.

Similarly, let us show that the second term $(m + 1) e(P) e(Q - y)$ is the number of linear extensions $f \in \mathcal{E}(R)$ s.t. $f(v) = m + 2$. For such f we have:

$$\begin{aligned} \{f^{-1}(1), \dots, f^{-1}(m + 1)\} &= X - x + w + z, \\ \{f^{-1}(m + 3), \dots, f^{-1}(m + n + 1)\} &= Y - y. \end{aligned}$$

Note that the restriction of f to $\{f^{-1}(1), \dots, f^{-1}(m+1)\}$ defines a linear extension of $(P^* + z)$. Additionally, note that the restriction of f to $\{f^{-1}(m+3), \dots, f^{-1}(m+n+1)\}$ defines a linear extension of $(Q - y)$. In total, we have $e(P^* + z)e(Q - y) = (m+1)e(P)e(Q - y)$ linear extensions f as above. This completes the proof. \square

Following the argument in the claim we similarly have:

$$(7.8) \quad e(R - z) = e(P - x)e(Q) + e(P)e(Q - y).$$

Indeed, the term $e(P - x)e(Q)$ is the number of linear extensions $f \in \mathcal{E}(R)$ for which $f(v) = m$, and the term $e(P)e(Q - y)$ is the number of linear extensions $f \in \mathcal{E}(R)$ for which $f(v) = m + 1$. We omit the details.

Combing (7.7) and (7.8), we have:

$$\rho(R, z) = m + \frac{e(P)e(Q - y)}{e(P - x)e(Q) + e(P)e(Q - y)} = m + \left(1 + \frac{\rho(Q, y)}{\rho(P, x)}\right)^{-1},$$

as desired. \square

Lemma 7.4. *Let $P = (X, \prec)$ be a poset on $n = |X|$ elements, and let $x \in \min(P)$. Then there exists a poset $Q = (Y, \prec')$ and an element $y \in \min(Q)$, such that $|Y| = n + 1$ and*

$$\rho(Q, y) = 1 + \frac{1}{\rho(P, x)}.$$

Proof. Let $Y := X + z$, and let \prec' coincide with \prec on P , with added relations

$$z \prec' u \quad \text{for all } u \in X - x, \quad \text{and} \quad z \parallel x.$$

Note that $z \in \min(Q)$. Note also that

$$e(Q - z) = e(P) \quad \text{and} \quad e(Q) = e(P) + e(P - x),$$

since for every $f \in \mathcal{E}(Q)$ we either have $f(z) = 1$, or $f(z) = 2$ and thus $f(x) = 1$. We now take $y \leftarrow z$, and observe that

$$\rho(Q, y) = \frac{e(Q)}{e(Q - z)} = \frac{e(P) + e(P - x)}{e(P)} = 1 + \frac{1}{\rho(P, x)},$$

as desired. \square

Lemma 7.5. *Let $P = (X, \prec)$ be a poset on $n = |X|$ elements, and let $x \in \min(P)$. Then there exists a poset $Q = (Y, \prec')$, and $y \in \min(Q)$, such that $|Y| = n + 1$ and*

$$\rho(Q, y) = 1 + \rho(P, x).$$

Proof. Let Q be as in the proof of Lemma 7.4. Note that $x \in \min(Q)$, and that

$$e(Q - x) = e(P - x),$$

since z is the unique minimal element in $Q - x$. We now take $y \leftarrow x$, and observe that

$$\rho(Q, y) = \frac{e(Q)}{e(Q - x)} = \frac{e(P) + e(P - x)}{e(P - x)} = 1 + \rho(P, x),$$

as desired. \square

Proof of Theorem 7.2. By symmetry, we will without loss of generality assume that $n_2 \geq n_3$. By applying Lemma 7.3 followed by applying Lemma 7.5 for $n_2 - n_3$ many times, we get a poset $P = (X, \prec)$ and $x \in \min(P)$ such that

$$\rho(P, x) = (n_2 - n_3) + \left(n_3 + \left(1 + \frac{\rho(P_1, x_1)}{\rho(P_3, x_3)} \right)^{-1} \right) = n_2 + \left(1 + \frac{\rho(P_1, x_1)}{\rho(P_3, x_3)} \right)^{-1}.$$

Additionally, poset P has

$$|X| = n = (n_1 + n_3 + 1) + n_2 - n_3 = n_1 + \max\{n_2, n_3\} + 1 \text{ elements.}$$

On the other hand, by Lemma 7.3 we get a poset Q and $y \in \min(Q)$, s.t. such that

$$\rho(Q, y) = n_2 + \left(1 + \frac{\rho(P_4, x_4)}{\rho(P_2, x_2)} \right)^{-1}.$$

and with

$$m = n_2 + n_4 + 1 = n_4 + \max\{n_2, n_3\} + 1.$$

It now follows that

$$\rho(P, x) = \rho(Q, y) \iff \frac{\rho(P_1, x_1)}{\rho(P_3, x_3)} = \frac{\rho(P_4, x_4)}{\rho(P_2, x_2)},$$

as desired. \square

8. VERIFICATION LEMMA

The proof of the Verification Lemma 4.7 is different from other reductions which are given by parsimonious bijections. Before proceeding to the proof, we need several technical and seemingly unrelated results.

8.1. Continuous fractions. Given $a_0 \geq 0$, $a_1, \dots, a_s \in \mathbb{Z}_{\geq 1}$, where $s \geq 0$, the corresponding *continued fraction* is defined as follows:

$$[a_0; a_1, \dots, a_s] := a_0 + \frac{1}{a_1 + \frac{1}{\ddots + \frac{1}{a_s}}}$$

Numbers a_i are called *quotients*, see e.g. [HW08, §10.1]. We refer to [Knu98, §4.5.3] for a detailed asymptotic analysis of the quotients in connection with the Euclidean algorithm, and further references. The following technical result is key in the proof of the Verification Lemma.

Proposition 8.1 (cf. [KS21, §3]). *Let $a_0, \dots, a_s \in \mathbb{Z}_{\geq 1}$. Then there exists a poset $P = (X, \prec)$ of width two on $|X| = a_0 + \dots + a_s$ elements, and element $x \in \min(P)$, such that*

$$\rho(P, x) = [a_0; a_1, \dots, a_s].$$

Corollary 8.2. *Let $a_1, \dots, a_s \in \mathbb{Z}_{\geq 1}$. Then there exists a width two poset $P = (X, \prec)$ on $|X| = a_1 + \dots + a_s$ elements, and element $x \in \min(P)$, such that*

$$\frac{1}{\rho(P, x)} = [0; a_1, \dots, a_s].$$

Proof. This follows from $[a_1; a_2, \dots, a_s] = [0; a_1, \dots, a_s]^{-1}$. \square

Remark 8.3. Proposition 8.1 was proved implicitly in [KS21, §3]. Unfortunately, the notation and applications in that paper are very different from ours, so we chose to include a self-contained proof for completeness.

We now present the proof of Proposition 8.1, which uses the following corollary of Lemma 7.4 and Lemma 7.5.

Corollary 8.4. *Let $P = (X, \prec)$ be a width two poset on $n = |X|$ elements, let $x \in \min(P)$, and let $a \in \mathbb{Z}_{\geq 1}$. Then there exists a width two poset $Q = (Y, \prec')$ and $y \in \min(Q)$, such that $|Y| = n + a$ and*

$$\rho(Q, y) = a + \frac{1}{\rho(P, x)}.$$

Proof. Use Lemma 7.4 once, and Lemma 7.5 ($a - 1$) times. Also note that the operations used in Lemma 7.4 and Lemma 7.5 do not increase the width of the poset Q if the input poset P is not a chain. \square

Proof of Proposition 8.1. We use induction on s . For $s = 0$, let $P := C_{a_0-1} + \{x\}$ be a disjoint sum of two chains, and observe that $\rho(P, x) = a_0$.

Suppose the claim holds for $s - 1$, i.e. there exists a poset P_1 on $n = a_1 + \dots + a_s$ elements and $x_1 \in \min(P_1)$, such that $\rho(P_1, x_1) = [a_1; a_2, \dots, a_s]$, and with $|P_1| = a_1 + \dots + a_s$. By Corollary 8.4, there exists a poset Q on $a_0 + n$ elements, and $x \in \min(P)$, such that

$$\rho(P, x) = a_0 + \frac{1}{\rho(P_1, x_1)} = a_0 + \frac{1}{[a_1; a_2, \dots, a_s]} = [a_0; a_1, \dots, a_s].$$

This completes the proof. \square

8.2. Number theoretic estimates. For $A \in \mathbb{Z}_{\geq 1}$ and $m \in [A]$, consider the quotients in the continued fraction of m/A and their sum:

$$\frac{m}{A} = [0; a_1(m), \dots, a_s(m)] \quad \text{and} \quad S_A(m) := \sum_{i=1}^s a_i(m).$$

Note that every rational number can be represented by continued fractions in exactly two ways, and $S_A(m)$ are equal for both representations. Also note that

$$(8.1) \quad S_A(m) = S_{A'}(m'), \quad \text{where} \quad A' := \frac{A}{\gcd(A, m)} \quad \text{and} \quad m' := \frac{m}{\gcd(A, m)}$$

are normalized to be coprime integers. The following technical result will also be used in the proof of the Verification Lemma 4.7.

Proposition 8.5. *There exists a constant $C > 0$, such that for all coprime integers A, B which satisfy $C < B < A < 2B$, there exists an integer $m := m(A, B)$ such that $m < B$,*

$$S_A(m) \leq 2(\log A)^2 \quad \text{and} \quad S_B(m) \leq 2(\log B)^2.$$

We now build toward the proof of this result. We need the following technical result.

Lemma 8.6 (Yao–Knuth [YK75]). *We have:*

$$\frac{1}{n} \sum_{m \in [n]} S_n(m) = \frac{6}{\pi^2} (\log n)^2 + O((\log n)(\log \log n)^2) \quad \text{as } n \rightarrow \infty.$$

By the Markov inequality, it follows from Lemma 8.6 that

$$(8.2) \quad |\{m \in [n] : S_n(m) > 2(\log n)^2\}| \leq \frac{3}{\pi^2} n(1 + o(1)).$$

Proof of Proposition 8.5. Denote

$$\vartheta(A, B) := |\{m \in [B] : S_A(m) \leq 2(\log A)^2, S_B(m) \leq 2(\log B)^2\}|.$$

To prove the result, it sufficed to show that

$$\vartheta(A, B) = \Omega(B) \quad \text{as } C \rightarrow \infty.$$

Now, it follows from the inclusion-exclusion principle, that

$$\vartheta(A, B) \geq B - |\{m \in [B] : S_A(m) > 2(\log A)^2\}| - |\{m \in [B] : S_B(m) > 2(\log B)^2\}|.$$

On the other hand, we have:

$$|\{m \in [B] : S_A(m) > 2(\log A)^2\}| \leq |\{m \in [A] : S_A(m) > 2(\log A)^2\}| \stackrel{(8.2)}{\leq} \frac{3}{\pi^2} A(1 + o(1)),$$

and

$$|\{m \in [B] : S_B(m) > 2(\log B)^2\}| \stackrel{(8.2)}{\leq} \frac{3}{\pi^2} B(1 + o(1)).$$

Combining these inequalities, we get

$$\vartheta(A, B) \geq B - \frac{3}{\pi^2} (A + B)(1 + o(1)) \geq B(1 - \frac{9}{\pi^2})(1 - o(1)),$$

and the result follows since $(1 - \frac{9}{\pi^2}) > 0$. \square

Remark 8.7. The proof of Proposition 8.5 does not give a (deterministic) polynomial time algorithm to find the desired m , i.e. in $\text{poly}(\log A)$ time. There is, however, a relatively simple *probabilistic* polynomial time algorithm, cf. [CP23a, Rem. 5.31].

8.3. Bounds on relative numbers of linear extensions. The following simple bound is the final ingredient we need for the proof of the Verification Lemma.

Proposition 8.8 (see [CPP23c, EHS89]). *Let $P = (X, \prec)$ be a poset on $|X| = n$ elements, and let $x \in \min(X)$. Then $1 \leq \rho(P, x) \leq n$. Moreover, $\rho(P, x) = 1$ if and only if $\min(P) = \{x\}$, i.e. x is the unique minimal element.*

The lower bound holds for all $x \in X$, see e.g. [EHS89]. The upper bound is a special case of [CPP23c, Lem. 5.1]. We include a short proof for completeness.

Proof. The lower bound $e(P - x) \leq e(P)$ follows from the injection $\mathcal{E}(P - x) \rightarrow \mathcal{E}(P)$ that maps $f \in \mathcal{E}(P - x)$ into $g \in \mathcal{E}(P)$ by letting $g(x) \leftarrow 1$, $g(y) \leftarrow f(y) + 1$ for all $y \neq x$. For the second part, note that $e(P) - e(P - x)$ is the number of $f \in \mathcal{E}(P)$ such that $f(x) > 1$, so $e(P) - e(P - x) = 0$ implies $\min(P) = \{x\}$.

The upper bound $e(P) \leq ne(P - x)$ follows from the injection $\mathcal{E}(P) \rightarrow \mathcal{E}(P - x) \times [n]$ that maps $g \in \mathcal{E}(P)$ into a pair $(f, g(x))$ where $f \in \mathcal{E}(P - x)$ is defined as $f(y) \leftarrow g(y)$ if $g(y) < g(x)$, $f(y) \leftarrow g(y) - 1$ if $g(y) > g(x)$. \square

8.4. Proof of Verification Lemma 4.7. Recall the decision problem

$$\text{VERRLE} := \left\{ \rho(P, x) = \frac{A}{B} \right\},$$

where $P = (X, \prec)$ is a poset on $n = |X|$ elements, $x \in \min(P)$, and A, B are coprime integers with $B < A \leq n!$. We simulate VERRLE with an oracle for QUADRLE as follows.

By Proposition 8.8, we need only to consider the cases $1 < \frac{A}{B} \leq n$. Indeed, when $\rho(P, x) < 1$ or $\rho(P, x) > n!$, VERRLE does not hold. Additionally, when $\rho(P, x) = 1$, VERRLE holds if and only if P is a chain. Let $k := \lfloor \frac{A}{B} \rfloor$. As in the $s = 0$ part of the proof of Proposition 8.1, there exists a poset $P_3 = (X_3, \prec_3)$ with $|X_3| = k \leq n$, and an element $x_3 \in \min(P_3)$, such that $\rho(P_3, x_3) = k$.

Let A', B' be coprime integers such that

$$\frac{A}{B} = k \frac{A'}{B'}.$$

Then we have $B \leq B' < A' < 2B'$, $A' \leq A$ and thus $\log A' = O(n \log n)$. By Proposition 8.5, there is a positive integer $m \in [B']$, such that

$$S_{A'}(m) \leq 2(\log A')^2 \quad \text{and} \quad S_{B'}(m) \leq 2(\log B')^2.$$

At this point we *guess* such m . Since computing the quotients of m/A' can be done in polynomial time, we can verify in polynomial time that m satisfies the inequalities above.

By Corollary 8.2, we can construct posets $P_2 = (X_2, \prec_2)$, $P_4 = (X_4, \prec_4)$ with $x_2 \in \min(P_2)$, $x_4 \in \min(P_4)$, such that

$$\rho(P_2, x_2) = \frac{B'}{m} \quad \text{and} \quad \rho(P_4, x_4) = \frac{A'}{m}.$$

The corollary also gives us

$$|X_2| \leq S_{B'}(m) \leq 2(\log B')^2 = O(n^2(\log n)^2),$$

and we similarly have $|X_4| = O(n^2(\log n)^2)$. Since posets P_2, P_3 and P_4 have polynomial size, we can call QUADRLE to check

$$\{ \rho(P, x) \cdot \rho(P_2, x_2) \stackrel{?}{=} \rho(P_3, x_3) \cdot \rho(P_4, x_4) \}.$$

Observe that

$$\frac{\rho(P_3, x_3) \cdot \rho(P_4, x_4)}{\rho(P_2, x_2)} = \frac{m}{B'} \cdot k \cdot \frac{A'}{m} = \frac{A}{B}.$$

Thus, in this case QUADRLE is equivalent to VERRLE, as desired. \square

Remark 8.9. In our recent paper [CP24], we use ideas from the proof above to obtain further results for relative numbers of linear extensions. We also use stronger number theoretic estimates than those given by Lemma 8.6.

9. FIXING ONE ELEMENT

In this section we prove Theorem 1.4. The proof relies heavily on [MS24]. We also need the definition and basic properties of the *promotion* and *demotion* operations on linear extensions, see e.g. [Sta09] and [Sta12, §3.20].

9.1. Explicit equality conditions. For $k = 1$, the equality cases of Stanley's inequality (Sta) are tuples (P, x, z, a, c) , where $P = (X, \prec)$ is a poset on $n = |X|$ elements, $x, z \in X$, $a, c \in [n]$, and the following holds:

$$(9.1) \quad N_{zc}(P, x, a)^2 = N_{zc}(P, x, a+1) \cdot N_{zc}(P, x, a-1).$$

The subscripts here and throughout this section are no longer bold, to emphasize that $k = 1$. Recall also both the notation in §1.4, and the Ma–Shenfeld poset notation in §6.1.

Lemma 9.1. *Let $P = (X, \prec)$ be a poset on $n = |X|$ elements, let $x, z \in X$, $a, c \in [n]$. Then the equality (9.1) is equivalent to:*

$$(*) \quad \text{for every } f \in \mathcal{E}_{zc}(P, x, a+s), \ s \in \{0, \pm 1\}, \text{ we have } x \parallel \text{lc}(f) \text{ and } x \parallel \text{uc}(f).$$

We prove Lemma 9.1 later in this section.

Remark 9.2. For the case $k = 0$, the analogue of $(*)$ that companions of f are incomparable to x , was proved in [SvH23, Thm 15.3(c)]. However, $(*)$ fails for $k = 2$, as shown in the “hope shattered” Example 1.4 in [MS24]. Thus, Lemma 9.1 closes the gap between these two results. See §10.8 for potential complexity implications of this observation.

Note also that condition $(*)$ is in \mathbf{P} since can be equivalently described in terms of explicit conditions on the partial order (rather than in terms of linear extensions of the poset). This is proved in [SvH23, Thm 15.3(d)] for $k = 0$, and in [MS24, Eq. (1.6)] for $k = 1$.

Proof of Theorem 1.4. As before, let $P = (X, \prec)$ be a poset on $n = |X|$ elements, let $x, y, z \in X$ and $a, b, c \in [n]$. Denote by $N_{zc}(P, x, a, y, b)$ the number of linear extensions $f \in \mathcal{E}_{zc}(P, x, a)$ that additionally satisfy $f(y) = b$.

Now, condition $(*)$ in Lemma 9.1, can be rewritten as follows:

$$(9.2) \quad N_{zc}(P, x, a', y, b') = 0 \quad \text{for all } y \in \mathcal{C}(x) \quad \text{and} \quad a', b' \in \{a - 1, a, a + 1\}.$$

Indeed, each vanishing condition in (9.2) is checking whether there exists a companion of x in a linear extension that is comparable to x .

Recall that each vanishing condition in (9.2) is in \mathbf{P} , see references in §3.5. There are at most $6n$ instances to check, since for all $y \in X$ there are at most 6 choices of distinct a', b' in $\{a - 1, a, a + 1\}$. Therefore, EQUALITYSTANLEY₁ $\in \mathbf{P}$. \square

9.2. Ma–Shenfeld theory. We now present several ingredients needed to prove Lemma 9.1. We follow closely the Ma–Shenfeld paper [MS24], presenting several results from that paper.

In [MS24], Ma–Shenfeld defined the notions of *subcritical*, *critical*, and *supercritical* posets, which are directly analogous to the corresponding notions for polytopes given in [SvH23], cf. §3.2. As the precise definitions are rather technical, we will not state them here while still including key properties of those families that are needed to prove Lemma 9.1.

We start with the following hierarchical relationship between the three families:

$$\{\text{subcritical posets}\} \supseteq \{\text{critical posets}\} \supseteq \{\text{supercritical posets}\}.$$

A poset that is subcritical but not critical is called *sharp subcritical*, and a poset that is critical but not super critical is called *sharp critical*.

The equality conditions for (9.1) are directly determined by the classes to which the poset P belongs, as we explain below. We note that these families depend on the choices of P, x, a, z, c , which we omit from the notation to improve readability. Furthermore, without loss of generality we can assume that $z \notin \{a - 1, a, a + 1\}$, as otherwise one of the numbers in (9.1) are equal to 0, making the problem in \mathbf{P} (see above).

We now state two other properties of these families, which require the following definitions. Following [MS24], we add two elements z_0, z_{k+1} into the poset such that $z_0 \preceq y \preceq z_{k+1}$ for all $y \in X$, and we define $c_0 := 0$ and $c_{k+1} := n + 1$. A *splitting pair* is a pair of integers (r, s) in $\{0, \dots, k + 1\}$, such that $(r, s) \neq (0, k + 1)$.¹⁰

Lemma 9.3 ([MS24, Lemma 5.10]). *Let $P = (X, \prec)$ be a sharp subcritical poset. Then there exists a splitting pair (r, s) such that*

$$(9.3) \quad |\{u \in X : z_r \prec u \prec z_s\}| = c_s - c_r - 1.$$

We say that poset P is *split indecomposable* if, for every splitting pair (r, s) ,

$$|\{u \in X : z_r \prec u \prec z_s\}| \leq c_s - c_r - 2.$$

In particular, by Lemma 9.3 every sharp subcritical poset is not split indecomposable.

¹⁰In [MS24, Def 5.2], this pair is instead written as $(r + 1, s)$.

It was shown in [MS24], that we can without loss of generality assume that poset P is split indecomposable. Indeed, otherwise checking (9.1) can be reduced to checking the same problem for a smaller poset: either restricting to the set in (9.3), or removing this set from the poset, see [MS24, §6] for details. Thus we can without loss of generality assume that P is a critical poset.

Lemma 9.4 ([MS24, Lemma 5.11]). *Let P be a split indecomposable sharp critical poset. Then there exists a splitting pair (r, s) such that $c_r < a < c_s$ and*

$$(9.4) \quad |\{u \in X : z_r \prec u \prec z_s\}| = c_s - c_r - 2.$$

Remark 9.5. Lemmas 9.3 and 9.4 can be modified to imply that deciding whether poset P is subcritical, critical, or supercritical is in \mathbf{P} . We do not need this result for the proof of Lemma 9.1, so we omit these changes to stay close to the presentation in [MS24]. More generally, one can ask similar questions for \mathbf{H} -polytopes. While we believe that for \mathbf{TU} -polytopes these decision problems are still likely to be in \mathbf{P} , proving that would already be an interesting challenge beyond the scope of this paper.

Recall from §6.2 that $\mathcal{F}(a, \text{com}, \text{com})$ is the set of linear extensions in $\mathcal{E}_{zc}(P, x, a)$, such that both the lower and upper companions of x are incomparable to x . Next, $\mathcal{F}(a, \text{com}, \text{inc})$ is the set of linear extensions in $\mathcal{E}_{zc}(P, x, a)$, such that the lower companion is comparable to x , but the upper companion is incomparable to x . Similarly, $\mathcal{F}(a, \text{inc}, \text{com})$ is the set of linear extensions in $\mathcal{E}_{zc}(P, x, a)$, such that the lower companion is incomparable to x , but the upper companion is comparable to x . Let $\mathcal{F}(a - 1, \cdot, \cdot)$ and $\mathcal{F}(a + 1, \cdot, \cdot)$ be defined analogously. Finally, let $F(a + s, \cdot, \cdot) := |\mathcal{F}(a + s, \cdot, \cdot)|$ where $s \in \{0, \pm 1\}$, be the numbers of these linear extensions.

Lemma 9.6 ([MS24, Thm 1.5]). *Let P be a critical poset. Then (9.1) holds if and only if*

$$(9.5) \quad F(a - 1, \text{com}, \text{com}) = F(a, \text{com}, \text{com}) = F(a + 1, \text{com}, \text{com}) = 0 \quad \text{and}$$

$$(9.6) \quad \begin{aligned} F(a - 1, \text{com}, \text{inc}) &= F(a - 1, \text{inc}, \text{com}) = F(a, \text{com}, \text{inc}) \\ &= F(a, \text{inc}, \text{com}) = F(a + 1, \text{com}, \text{inc}) = F(a + 1, \text{inc}, \text{com}). \end{aligned}$$

Now note that $F(a - 1, \text{com}, \text{inc}) \leq F(a - 1, \text{inc}, \text{com})$, with the equality if and only if every upper companion of x is always incomparable to the lower companion of x . By an analogous arguments applied to $F(a, \cdot, \cdot)$ and $F(a + 1, \cdot, \cdot)$, we get the following corollary.

Corollary 9.7. *Let P be a critical poset. Suppose*

$$N_{zc}(P, x, a)^2 = N_{zc}(P, x, a + 1) \cdot N_{zc}(P, x, a - 1) \neq 0,$$

Then, for every linear extension $f \in \mathcal{E}(P)$ counted by (9.6), the upper companion is incomparable to the lower companion: $\text{uc}(f) \parallel \text{lc}(f)$.

Finally, we have equality conditions for supercritical posets.

Lemma 9.8 ([MS24, Thm 1.3]). *Let P be a supercritical poset. Then (9.1) holds if and only if equalities (9.5) and (9.6) hold, and additionally*

$$(9.7) \quad \text{all numbers in (9.6) are equal to 0.}$$

9.3. Proof of Lemma 9.1. Note that (9.5), (9.6) and (9.7) are equivalent to requiring that x is incomparable to both $\text{lc}(f)$ and $\text{uc}(f)$. Thus it suffices to show that, if P is a critical poset, then (9.7) holds.

Suppose to the contrary, that $P = (X, \prec)$ is a counterexample, and let $n := |X|$. Then P is a sharp critical poset. By taking the dual poset if necessary, we can assume, without loss of generality, that $c < a$. It then follows that the splitting pair (r, s) in Lemma 9.4 is $(1, 2)$. This means that $c_r = c$ and $c_s = n + 1$, so we have from (9.4) that

$$(9.8) \quad |\{u \in P : z \prec u\}| = n - c - 1.$$

Since (9.7) does not hold, there exists $f \in \mathcal{F}(a, \text{com}, \text{inc})$ and $h \in \mathcal{F}(a - 1, \text{com}, \text{inc})$. Let $y_1 := f^{-1}(a - 1)$ (i.e., the lower companion in f) and $y_2 := h^{-1}(a)$ (i.e., the lower companion in h). Note that we have $y_1 \prec x \prec y_2$. Let $m = f(y_2)$, and note that $m \geq a + 2$ by definition.

We claim: There exists a new linear extension $g \in \mathcal{E}(P)$ such that $g(y_2) = m - 1$, and such that $g \in \mathcal{F}(a, \text{com}, \text{inc})$ if $m > a + 2$, and $g \in \mathcal{F}(a, \text{com}, \text{com})$ if $m = a + 2$. Note that this suffices to prove the lemma, as by replacing f with g and decreasing m repeatedly, we get that $F(a, \text{com}, \text{com}) > 0$, which contradicts (9.5).

We now prove the claim. Since $h(y_2) = a < m = f(y_2)$, there exists $w \in X$ such that $f(w) < m$ and $w \parallel y_2$. Suppose w is such an element that maximizes $f(w)$. There are *three cases*:

First, suppose that $f(w) > a$. By the maximality assumption, every element ordered between w and y_2 according to f , is incomparable to w . Then we can promote w to be larger than y_2 . Note that the resulting linear extension $g \in \mathcal{E}(P)$ satisfies $g(y_2) = m - 1$, $g(y_1) = a - 1$ and $g(x) = a$, as desired.

Second, suppose that $c < f(w) < a$. By the maximality assumption, every element ordered between w and y_2 according to f , is incomparable to w . Then we can promote w to be larger than y_2 . The resulting linear extension $g' \in \mathcal{E}(P)$ satisfies $g'(y_2) = m - 1$. Note, however, that we have $g'(y_1) = a - 2$ and $g'(x) = a - 1$. In order to fix this, let $v := f^{-1}(a + 1)$. It follows from Corollary 9.7, that v is incomparable to y_1 and x . Thus we can demote v to be the smaller than y_1 . We obtain a new linear extension $g \in \mathcal{E}(P)$ that satisfies $g(y_1) = a - 1$ and $g(x) = a$, as desired.

Third, suppose that $f(w) < c$. Then, every element ordered between z and y_2 according to f , is less than y_2 . Note that there are $m - c - 1$ many such elements. On the other hand, it follows from (9.8), that there is exactly one element in $\{f^{-1}(c + 1), f^{-1}(c + 2), \dots, f^{-1}(n)\}$ that is incomparable to z . It then follows that there are at least $m - c - 2$ elements that are greater than z and less than y_2 , i.e.

$$(9.9) \quad |\{u \in X : z \prec u \prec y_2\}| \geq m - c - 2.$$

On the other hand, the existence of h implies that

$$(9.10) \quad |\{u \in X : z \prec u \prec y_2\}| \leq h(y_2) - c - 1 = a - c - 1 \leq m - c - 3,$$

a contradiction. This finishes the proof of the claim. \square

10. FINAL REMARKS

10.1. The basis of our work. Due to the multidisciplinary nature of this paper, we make a special effort to simplify the presentation. Namely, the proofs of our main results (Theorems 1.1 and 1.3), are largely self-contained in a sense that we only use standard results in combinatorics (Stanley's theorem in §5.2 and the Brightwell–Winkler's theorem in §3.6), computational complexity (Toda's theorem in §4.4), and number theory (Yao–Knuth's theorem in §8.2). In reality, the paper freely uses tools and ideas from several recent results worth acknowledging.

First, we heavily build on the recent paper by Shenfeld and van Handel [SvH23], and the followup by Ma and Shenfeld [MS24]. Without these results we would not know where to look for “bad posets” and “bad polytopes”. Additionally, the proof in §6.2 is a reworking and simplification of many technical results and ideas in [MS24].

Second, in §8.1 we use and largely rework the continued fraction approach by Kravitz and Sah [KS21]. There, the authors employ the *Stern–Brocot* and *Calkin–Wilf tree* notions, which we avoid in our presentation as we aim for different applications.

Third, in the heart of our proof of Theorem 1.3 in §4.4, we follow the complexity roadmap championed by Ikenmeyer, Panova and the second author in [IP22, IPP22]. Same for the heart of the proof of the Verification Lemma 4.7 in §8.4, which follows the approach in our companion paper [CP23a].

On the other hand, the proof of Theorem 1.4 given in Section 9, is the opposite of self-contained, as we rely heavily on both results and ideas in [MS24]. We also use properties the *promotion* and *demotion* operations on linear extensions, that were introduced by Schützenberger in the context of algebraic combinatorics, see [Schü72]¹¹. Panova and the authors employed this approach in a closely related setting in [CPP23a, CPP23b, CPP23c]. We emphasize once again that our proof of Theorem 1.4 is independent of the rest of the paper and is the only part that uses results in [MS24].

10.2. Equality cases. The reader unfamiliar with the subject may wonder whether equality conditions of known inequalities are worth an extensive investigation. Here is how Gardner addresses this question:

“If inequalities are silver currency in mathematics, those that come along with precise equality conditions are gold. Equality conditions are treasure boxes containing valuable information.” [Gar02, p. 360].

Closer to the subject of this paper, Shenfeld and van Handel explain the difficulty of finding equality conditions for (MQI) and (AF):

“In first instance, it may be expected that the characterization of the extremals of the Minkowski and Alexandrov–Fenchel inequalities should follow from a careful analysis of the proofs of these inequalities. It turns out, however, that none of the classical proofs provides information on the cases of equality: the proofs rely on strong regularity assumptions (such as smooth bodies or polytopes with restricted face directions) under which only trivial equality cases arise, and deduce the general result by approximation. The study of the nontrivial extremals requires one to work directly with general convex bodies, whose analysis gives rise to basic open questions in the foundation of convex geometry.” [SvH22, p. 962].

10.3. Polytopes. The family of TU-polytopes that we chose is very special in that these H-polytopes have integral vertices (but not a description in \mathbb{P} , as V-polytopes are defined to have). In [CP24+], we consider a family of *axis-parallel boxes* which have similar properties. Clearly, for general convex bodies there is no natural way to set up a computational problem that would not be immediately intractable (unless one moves to a more powerful computational model, see e.g. [BCSS98]).

10.4. Discrete isoperimetric inequality. For a discrete version of the isoperimetric inequality in the plane, one can consider convex polygons with given normals to edges. In this case, L’Huillier (1775) proved that the isoperimetric ratio is minimized for circumscribed polygons, see e.g. [Fej72, §I.4]. In the 1860s, Steiner and Lindelöf studied a natural generalization of this problem in \mathbb{R}^3 , but were unable to solve it in full generality.

At the turn of 20th century, Minkowski developed the *theory of mixed volumes*, motivated in part to resolve the Steiner–Lindelöf problem. He showed that among all polytopes with given normals, the isoperimetric ratio is minimized on circumscribed polytopes, see e.g. [Fej72, §V.7].

There are several Bonnesen type and stability versions of the discrete isoperimetric inequality, see e.g. [FRS85, IN15, Zhang98]. Let us single out a hexagon version used by Hales in his famous proof of the *honeycomb conjecture* [Hal01, Thm 4].

¹¹These operations were rediscovered in [DD85, Day84], where they are called *push up* and *push down*, respectively.

10.5. Brunn–Minkowski inequality. There are several proofs of the Brunn–Minkowski inequality (BM), but some of them do not imply the equality conditions, such as, e.g., the “brick-by-brick” inductive argument in [Mat02, §12.2]. Note also that Alexandrov’s proof of the *Minkowski uniqueness theorem* (of polytopes with given facet volumes and normals) relies on the equality conditions for the Brunn–Minkowski inequality, see [Ale50]. This is essential for Alexandrov’s “topological method”, and is the basis for the *variational principle* approach, see e.g. [Pak09].

10.6. Van der Waerden conjecture. The Alexandrov–Fenchel inequality (AF) came to prominence in combinatorics after Egorychev [Ego81] used it to prove the *van der Waerden conjecture*, that was proved earlier by Falikman [Fal81].¹² See [Knu81, vL82] for friendly expositions. This development set the stage for Stanley’s paper [Sta81]. The conjecture states that for every bistochastic $n \times n$ matrix A , we have

$$(\text{vdW}) \quad \text{per}(A) \geq \frac{n!}{n^n},$$

and the equality holds only if $A = (a_{ij})$ has uniform entries: $a_{ij} = \frac{1}{n}$ for all $1 \leq i, j \leq n$.

Note that Egorychev’s proof of the equality conditions for (vdW) used Alexandrov’s equality conditions (AF) for nondegenerate boxes, see §3.2 (cf. [Knu81, p. 735] and [vL82, §7]). In a followup paper [CP24+], we analyze the complexity of the Alexandrov–Fenchel equality condition for degenerate boxes. Note also that Knuth’s exposition in [Knu81] is essentially self-contained, while Gurvits’s proof of (vdW) completely avoids (AF), see [Gur08, LS10].

10.7. Matroid inequalities. Of the several log-concavity applications of the AF inequality given by Stanley in [Sta81] (see also [Sta86, §6]), one stands out as a special case of a Mason’s conjecture (Thm 2.9 in [Sta81]). The strongest of the three Mason’s conjectures states that the numbers $I(M, k) / \binom{n}{k}$ are log-concave, where $I(M, k)$ is the number of independent sets of size k in a matroid M on n elements. These Mason’s conjectures were recently proved in a long series of spectacular papers culminating with [AHK18, ALOV24, BH20], see also an overview in [Huh18, Kal22].

Curiously, the equality cases for these inequalities are rather trivial and can be verified in polynomial time [MNY21] (see also [CP21, §1.6]). Here we assume that the matroid is given in a concise presentation (such presentations include graphical, bicircular and representable matroids). Curiously, for the weighted extension of Mason’s third conjecture given in [CP21, Thm 1.6], the equality cases are more involved. It follows from [CP21, Thm 1.9], however, that this problem is in coNP. In other words, Theorem 1.3 shows that EQUALITYSTANLEY₂ is likely much more powerful.

Note that the defect $\psi(M, k) := I(M, k)^2 - I(M, k+1) \cdot I(M, k-1)$ is conjectured to be not in #P, see [Pak22, Conj. 5.3]. Clearly, the argument in the proof of Corollary 1.5 does not apply in this case. Thus, another approach is needed to prove this conjecture, just as another approach is needed to prove that $\phi_0 \notin \#P$ (see §1.4).

10.8. Complexity of equality cases. Recall that Theorem 1.1 does not imply that EQUALITYAF is NP-hard or coNP-hard, more traditional measures of computational hardness. This remains out of reach. Note, however, that EQUALITYSTANLEY_k is naturally in the class C=P, see §2.6.

Conjecture 10.1. EQUALITYSTANLEY_k is C=P-complete for large enough k .

If this holds for all $k \geq 2$, this would imply a remarkable dichotomy with $k \leq 1$ (see Theorem 1.4). To motivate the conjecture, recall from §3.6, that C=P-complete problem $C_{\#3\text{SAT}}$ is coNP-hard. See [CP23a] for more on the complexity of combinatorial coincidence problems.

Note that the proof of EQUALITYSTANLEY₂ \notin PH implies that EQUALITYAF \notin PH even when at most four polytopes are allowed to be distinct. It would be interesting to decide if this number can be reduced down to three. It is known that two distinct TU-polytopes are not enough. This follows from a combination of our argument that for supercritical cases (in the sense of [SvH23]), we have EQUALITYAF \in coNP, and an argument that for two polytopes the equality cases are supercritical.¹³

¹²According to Vladimir Gurvich’s *essay*, Egorychev was the referee of Falikman’s article which was submitted prior to Egorychev’s preprint.

¹³Ramon van Handel, personal communication, April 2023.

10.9. Injective proofs. In enumerative combinatorics, whenever one has an equality between the numbers counting certain combinatorial objects, one is tempted to find a *direct bijection* between the sides, see e.g. [Loe11, Pak05, Sta12]. Similarly, when presented an inequality $f \geq g$, one is tempted to find a *direct injection*, see e.g. [Pak19, Sta89]. In the context of linear extensions, such injections appear throughout the literature, see e.g. [Bre89, BT02, CPP23a, DD85, GG22, LP07].

Typically, a direct injection and its inverse are given by simple polynomial time algorithms, thus giving a combinatorial interpretation for the defect $(f - g)$. Therefore, if a combinatorial inequality is not in $\#P$, it is very unlikely that there is a proof by a direct injection. In particular, Corollary 1.5 implies that the Stanley inequality (Sta) most likely cannot be proved by a direct injection. This confirms an old speculation:

“It appears unlikely that Stanley’s Theorem for linear extensions quoted earlier can be proved using the kind of injection presented here.” [DDP84, §4].

Similarly, Corollary 1.5 suggests that the strategy in [CPP23b, §9.12] is unlikely to succeed, at least for $k \geq 2$.¹⁴

To fully appreciate how delicate is Corollary 1.5, compare it with a closely related problem. It is known that for all $k \geq 0$, the analogue of the Stanley inequality (Sta) holds for the number $\Omega(P, t)$ of *order preserving maps* $X \rightarrow [t]$, for all $t \in \mathbb{N}$. This was conjectured by Graham in [Gra82, p. 129] (see also [Gra83, p. 233]), motivated by the proof of the XYZ inequality [She82] (cf. §3.4). The result was proved in [DDP84, Thm 4] by a direct injection (see also [Day84, §4.2] for additional details of the proof). In other words, in contrast with ϕ_k , the defect of the analogue of (Sta) for order preserving maps has a combinatorial interpretation. Note that it is not known whether the defect of the XYZ inequality is in $\#P$, see [Pak22, Conj. 6.4].

10.10. Stability proofs. By analogy with the injective proofs, Corollary 1.2 suggests that certain proofs of the Alexandrov–Fenchel inequality are likely not possible. Here we are thinking of the mass transportation proof of characterization of the isoperimetric sets given in [FMP10, App.], following Gromov’s approach in [Gro86]. It would be interesting to make this idea precise.

10.11. Dichotomy of the equality cases. As we discuss in §9.2, it follows from the results in [MS24], that the equality verification of the Stanley inequality (Sta) can be decided in polynomial time for supercritical posets. In contrast, by Theorem 1.3, the problem is not in PH for critical posets.¹⁵ We believe that this dichotomy also holds for the equality cases of the Alexandrov–Fenchel inequality (AF) for classes of H-polytopes for which the scaled mixed volume is in $\#P$.

10.12. The meaning of it all. Finding the equality conditions of an inequality may seem like a straightforward unambiguous problem, but the case of the Alexandrov–Fenchel inequality shows that it is nothing of the kind. Even the words “equality conditions” are much too vague for our taste. What the problem asks is a *description of the equality cases*. But since many geometric and combinatorial inequalities have large families of equality cases, the word “description” becomes open-ended (cf. §2.5). How do you know when you are done? At what point are you satisfied with the solution and do not need further details?

These are difficult questions which took many decades to settle, and the answers depend heavily on the area. In the context of geometric inequalities discussed in §3.1, the meaning of “description” starts out simple enough. There is nothing ambiguous about discs as equality cases of the isoperimetric inequality in the plane (Isop), or pairs of homothetic convex bodies for the Brunn–Minkowski inequality (BM), or circumscribed polygons with given normals for the discrete isoperimetric inequality (see §10.4). Arguably, Bol’s equality cases of (MWI) are also unambiguous — in \mathbb{R}^3 , you literally know the cap bodies when you see them.

However, when it comes to Minkowski’s quadratic inequality (MQI), the exact meaning of “description” is no longer obvious. Shenfeld and van Handel write “The main results of this paper will provide a complete solution to this problem” [SvH22]. Indeed, their description of 3-dimensional triples of convex bodies cannot be easily improved upon, at least not in the case of convex polytopes (see §3.1). Some questions may still linger, but they are on the structure of the equality cases rather than on their recognition.¹⁶

¹⁴In [Gra83, p. 129], Graham asked if Stanley’s inequality can be proved using the AD and FKG inequalities. This seems unlikely, even though we do not know how to formalize this question.

¹⁵We further clarify this in our survey [CP23b, §10], written after this paper.

¹⁶For example, one can ask to characterize all possible triples of polytope graphs that arise as equality cases.

What Shenfeld and van Handel did, is finished off the geometric approach going back to Brunn, Minkowski, Favard, Fenchel, Alexandrov and others, further formalized by Schneider. “Maybe a published conjecture will stimulate further study of this question”, Schneider wrote in [Schn85]. This was prophetic, but that conjecture was not the whole story, as it turned out.

In [SvH23], the authors write again: “We completely settle the extremals of the Alexandrov–Fenchel inequality for convex polytopes.” Unfortunately, their description is extraordinarily complicated in higher dimensions, so the problem of *recognizing* the equality cases is no longer easy (see §3.2). And what good is a description if it cannot be used to recognize the equality cases?

In combinatorics, the issue of “description” has also been a major problem for decades, until it was fully resolved with the advent of computational complexity. For example, consider the following misleadingly simple description: “Let G be a planar cubic Hamiltonian graph.” Is that good enough? How can you tell if a given graph G is as you describe? We now know that the problem whether G is planar, cubic and Hamiltonian is NP-complete [GJT76]. But if you only need the “planar” condition, the problem is computationally easy, while the “cubic” condition is trivial. Consequently, “planar cubic Hamiltonian” should not be viewed as a “good” description, but if one must consider the whole class of such graphs, this description is (most likely) the best one can do.

Going over equality cases for various inequalities on the numbers of linear extensions, already gives an interesting picture. For the Björner–Wachs inequality (see §3.4), the recognition problem of forests is in P, of course. On the other hand, as we explain in §3.4, for the Sidorenko inequality (3.1), the recognition problem of series-parallel posets is in P for a more involved reason. On the opposite end of the spectrum, for the (rather artificial) inequality $(e(P) - e(Q))^2 \geq 0$, the equality verification is not in PH, unless PH collapses, see §3.7 and [CP23a, Thm 1.4].

In this language, for the $k = 0$ case of the Stanley inequality (Sta), the description of equality cases given in [SvH23] is trivially in P. Similarly, for the $k = 1$ case, the description of equality cases is also in P by Theorem 1.4. On the other hand, Theorem 1.3 shows that for $k \geq 2$, the description in [MS24] is (very likely) not in P. Under standard complexity assumptions, there is no description of the equality cases in P at all, or even in PH for that matter.

Now, the problem of *counting* the equality cases brings a host of new computational difficulties, making seemingly easy problems appear hard when formalized, see [Pak22]. Even for counting non-isomorphic forest posets on n elements, to show that this function is in #P one needs to define a *canonical labeling* to be able to distinguish the forests, to make sure each is counted exactly once, see e.g. [SW19].

In this language, Corollary 1.5 states that *there are no combinatorial objects* that can be counted to give the number of non-equality cases of the Stanley inequality, neither the non-equality cases themselves nor anything else. The same applies to the equality cases. Fundamentally, this is because you should not be able to efficiently tell if the instances you are observing are the ones you should be counting.

Back to the Alexandrov–Fenchel inequality (AF), the description of equality cases by Shenfeld and van Handel is a breakthrough in convex geometry, and gives a complete solution for a large family of (n -tuples of) convex polytopes (see §10.11). However, our Theorem 1.1 says that from the computational point of view, the equality cases are intractable in full generality. Colloquially, this says that *there is no good description* of the equality cases of the Alexandrov–Fenchel inequality, unless the world of computational complexity is not what we think it is. As negative as this may seem, this is what we call a complete solution indeed.

Acknowledgements. We are grateful to Karim Adiprasito, Sasha Barvinok, Károly Böröczky, Christian Ikenmeyer, Jeff Kahn, Joe O’Rourke, Aldo Pratelli, Matvey Soloviev and Richard Stanley for useful remarks on the subject. Special thanks to Yair Shenfeld and Ramon van Handel for many very helpful comments on the first draft of the paper, and to Greta Panova for the numerous helpful discussions.

An extended abstract of this paper is to appear in *Proc. STOC* (2024); we thank the Program Committee and the reviewers for helpful comments. These results were obtained when both authors were visiting the American Institute of Mathematics at their new location in Pasadena, CA. We are grateful to AIM for the hospitality. The first author was partially supported by the Simons Foundation. Both authors were partially supported by the NSF.

REFERENCES

- [Aar16] Scott Aaronson, $P \stackrel{?}{=} NP$, in *Open problems in mathematics*, Springer, Cham, 2016, 1–122; available at scottaaronson.com/papers/pnp.pdf
- [AHK18] Karim Adiprasito, June Huh and Eric Katz, Hodge theory for combinatorial geometries, *Annals of Math.* **188** (2018), 381–452.
- [AAHS99] Pankaj K. Agarwal, Boris Aronov, Sarel Har-Peled and Micha Sharir, Approximation and exact algorithms for minimum-width annuli and shells, in *Proc. 15th SoCG*, ACM, New York, 1999, 380–389.
- [Ale37] Alexander D. Alexandrov, To the theory of mixed volumes of convex bodies II. New inequalities between mixed volumes and their applications (in Russian), *Mat. Sb.* **2** (1937), 1205–1238; English translation in A. D. Alexandrov, *Selected works*, Part I, Ch. IV, CRC Press, 1996, 61–97.
- [Ale50] Alexander D. Alexandrov, *Convex polyhedra* (translated from the Russian 1950 ed.), Springer, Berlin, 2005, 539 pp.
- [AS16] Noga Alon and Joel H. Spencer, *The probabilistic method* (fourth ed.), John Wiley, Hoboken, NJ, 2016, 375 pp.
- [ALOV24] Nima Anari, Kuikui Liu, Shayan Oveis Gharan and Cynthia Vinzant, Log-concave polynomials III: Mason’s Ultra-log-concavity conjecture for independent sets of matroids, *Proc. AMS* (published electronically March 20, 2024), 13 pp.; [arXiv:1811.01600](https://arxiv.org/abs/1811.01600).
- [AB09] Sanjeev Arora and Boaz Barak, *Computational complexity. A modern approach*, Cambridge Univ. Press, Cambridge, 2009, 579 pp.
- [AASS20] Shiri Artstein-Avidan, Shai Sadozsky and Raman Sanyal, Geometric inequalities for anti-blocking bodies, *Commun. Contemp. Math.* **25** (2023), no. 3, Paper No. 2150113, 30 pp.
- [Bar07] Alexander Barvinok, Brunn–Minkowski inequalities for contingency tables and integer flows, *Adv. Math.* **211** (2007), 105–122.
- [BW89] Anders Björner and Michelle L. Wachs, q -hook length formulas for forests, *J. Combin. Theory, Ser. A* **52** (1989), 165–187.
- [Blå05] Viktor Blåsjö, The isoperimetric problem, *Amer. Math. Monthly* **112** (2005), 526–566.
- [BCSS98] Lenore Blum, Felipe Cucker, Michael Shub and Steve Smale, *Complexity and real computation*, Springer, New York, 1998, 453 pp.
- [BBS99] Béla Bollobás, Graham Brightwell and Alexander Sidorenko, Geometrical techniques for estimating numbers of linear extensions, *European J. Combin.* **20** (1999), 329–335.
- [Bon29] Tommy Bonnesen, *Les problèmes des isopérimètres et des isépiphanes* (in French), Gauthier-Villars, Paris, 1929, 175 pp.
- [BF34] Tommy Bonnesen and Werner Fenchel, *Theory of convex bodies* (translated from the German 1934 ed.), BCS, Moscow, ID, 1987, 172 pp.
- [BH20] Petter Brändén and June Huh, Lorentzian polynomials, *Annals of Math.* **192** (2020), 821–891.
- [BL23] Petter Brändén and Jonathan Leake, Lorentzian polynomials on cones, preprint (2023), 34 pp.; [arXiv:2304.13203](https://arxiv.org/abs/2304.13203).
- [Bre89] Francesco Brenti, Unimodal, log-concave and Pólya frequency sequences in combinatorics, *Mem. AMS* **81** (1989), no. 413, 106 pp.
- [BT02] Graham Brightwell and William T. Trotter, A combinatorial approach to correlation inequalities, *Discrete Math.* **257** (2002), 311–327.
- [BW00] Graham Brightwell and Douglas B. West, Partially ordered sets, Ch. 11 in *Handbook of discrete and combinatorial mathematics*, CRC Press, Boca Raton, FL, 2000, 717–752.
- [BW91] Graham Brightwell and Peter Winkler, Counting linear extensions, *Order* **8** (1991), 225–247.
- [BZ88] Yuri D. Burago and Victor A. Zalgaller, *Geometric inequalities*, Springer, Berlin, 1988, 331 pp.
- [CP21] Swee Hong Chan and Igor Pak, Log-concave poset inequalities, to appear in *Jour. AMR*; preprint (2021), 71 pp.; [arXiv:2110.10740](https://arxiv.org/abs/2110.10740).
- [CP22] Swee Hong Chan and Igor Pak, Introduction to the combinatorial atlas, *Expo. Math.* **40** (2022), 1014–1048.
- [CP23a] Swee Hong Chan and Igor Pak, Computational complexity of counting coincidences, to appear in *Theor. Comp. Sci.*; preprint (2023), 24 pp.; [arXiv:2308.10214](https://arxiv.org/abs/2308.10214).
- [CP23b] Swee Hong Chan and Igor Pak, Linear extensions of finite posets, preprint (2023), 55 pp.; [arXiv:2311.02743](https://arxiv.org/abs/2311.02743).
- [CP24] Swee Hong Chan and Igor Pak, Linear extensions and continued fractions, *European J. Combin.* **122** (2024), 104018, 14 pp.
- [CP24+] Swee Hong Chan and Igor Pak, Quadratic permanent equality is not in PH, in preparation.
- [CPP23a] Swee Hong Chan, Igor Pak and Greta Panova, Extensions of the Kahn–Saks inequality for posets of width two, *Combinatorial Theory* **3** (2023), no. 1, Paper No. 8, 34 pp.

- [CPP23b] Swee Hong Chan, Igor Pak and Greta Panova, Effective poset inequalities, *SIAM J. Discrete Math.* **37** (2023), 1842–1880.
- [CPP23c] Swee Hong Chan, Igor Pak and Greta Panova, On the cross-product conjecture for the number of linear extensions, to appear in *Canadian Jour. Math.*; preprint (2023), 24 pp.; [arXiv:2306.09240](https://arxiv.org/abs/2306.09240).
- [CFG80] Fan R. K. Chung, Peter C. Fishburn and Ronald L. Graham, On unimodality for linear extensions of partial orders, *SIAM J. Algebraic Discrete Methods* **1** (1980), 405–410.
- [CKMS19] Dario Cordero-Erausquin, Bo’az Klartag, Quentin Merigot and Filippo Santambrogio, One more proof of the Alexandrov–Fenchel inequality, *C.R. Math. Acad. Sci. Paris* **357** (2019), no. 8, 676–680.
- [DD85] David E. Daykin and Jacqueline W. Daykin, Order preserving maps and linear extensions of a finite poset, *SIAM J. Algebraic Discrete Methods* **6** (1985), 738–748.
- [DDP84] David E. Daykin, Jacqueline W. Daykin, and Michael S. Paterson, On log concavity for order-preserving maps of partial orders, *Discrete Math.* **50** (1984), 221–226.
- [Day84] Jacqueline W. Daykin, *Monotonic functions of finite posets*, Ph.D. thesis, University of Warwick, 1984, 174 pp.; available at wrap.warwick.ac.uk/131009
- [DP20] Samuel Dittmer and Igor Pak, Counting linear extensions of restricted posets, *Electronic J. Combin.* **27** (2020), Paper No. 4.48, 13 pp.
- [DF88] Martin Dyer and Alan M. Frieze, On the complexity of computing the volume of a polyhedron, *SIAM J. Comput.* **17** (1988), 967–974.
- [DGH98] Martin Dyer, Peter Gritzmann and Alexander Hufnagel, On the complexity of computing mixed volumes, *SIAM J. Comput.* **27** (1998), 356–400.
- [EHS89] Paul Edelman, Takayuki Hibi and Richard P. Stanley, A recurrence for linear extensions, *Order* **6** (1989), 15–18.
- [Ego81] Georgy P. Egorychev, The solution of van der Waerden’s problem for permanents, *Adv. in Math.* **42** (1981), 299–305.
- [EK14] Ronen Eldan and Bo’az Klartag, Dimensionality and the stability of the Brunn–Minkowski inequality, *Ann. Sc. Norm. Super. Pisa, Cl. Sci.* **13** (2014), 975–1007.
- [Est10] Alexander Esterov, Newton polyhedra of discriminants of projections, *Discrete Comput. Geom.* **44** (2010), 96–148.
- [EG15] Alexander Esterov and Gleb Gusev, Systems of equations with a single solution, *J. Symbolic Comput.* **68** (2015), 116–130.
- [Ewa88] Günter Ewald, On the equality case in Alexandrov–Fenchel’s inequality for convex bodies, *Geom. Dedicata* **28** (1988), 213–220.
- [Fal81] Dmitry I. Falikman, Proof of the van der Waerden conjecture regarding the permanent of a doubly stochastic matrix, *Math. Notes* **29** (1981), 475–479.
- [Fav33] Jean Favard, Sur les corps convexes (in French), *Journ. de Math.* **12** (1933), 219–282.
- [Fej72] László Fejes Tóth, *Lagerungen in der Ebene auf der Kugel und im Raum* (in German), Springer, Berlin, 1972, 238 pp.
- [Fig13] Alessio Figalli, Stability in geometric and functional inequalities, in *Proc. Eur. Congress of Mathematics*, EMS, Zürich, 2013, 585–599.
- [Fig14] Alessio Figalli, Quantitative stability results for the Brunn–Minkowski inequality, in *Proc. ICM Seoul*, Vol. III, Kyung Moon Sa, Seoul, 2014, 237–256.
- [FJ17] Alessio Figalli and David Jerison, Quantitative stability for the Brunn–Minkowski inequality, *Adv. Math.* **314** (2017), 1–47.
- [FMP09] Alessio Figalli, Francesco Maggi and Aldo Pratelli, A refined Brunn–Minkowski inequality for convex sets, *Ann. Inst. H. Poincaré C, Anal. Non Linéaire* **26** (2009), 2511–2519.
- [FMP10] Alessio Figalli, Francesco Maggi and Aldo Pratelli, A mass transportation approach to quantitative isoperimetric inequalities, *Invent. Math.* **182** (2010), 167–211.
- [Fis84] Peter C. Fishburn, A correlational inequality for linear extensions of a poset, *Order* **1** (1984), 127–137.
- [FRS85] J. Chris Fisher, Dieter Ruoff and James Shilleto, Perpendicular polygons, *Amer. Math. Monthly* **92** (1985), 23–37.
- [FKG71] Cees M. Fortuin, Pieter W. Kasteleyn, and Jean Ginibre, Correlation inequalities on some partially ordered sets, *Comm. Math. Phys.* **22** (1971), 89–103.
- [GG22] Christian Gaetz and Yibo Gao, The hull metric on Coxeter groups, *Comb. Theory* **2** (2022), no. 2, Paper No. 7, 15 pp.
- [Gar02] Richard J. Gardner, The Brunn–Minkowski inequality, *Bull. AMS* **39** (2002), 355–405.
- [GJ78] Michael R. Garey and David S. Johnson, “Strong” NP-completeness results: motivation, examples, and implications, *J. ACM* **25** (1978), 499–508.
- [GJ79] Michael R. Garey and David S. Johnson, *Computers and Intractability: A Guide to the Theory of NP-completeness*, Freeman, San Francisco, CA, 1979, 338 pp.

- [GJT76] Michael R. Garey, David S. Johnson and Robert E. Tarjan, The planar Hamiltonian circuit problem is NP-complete, *SIAM J. Comput.* **5** (1976), 704–714.
- [Gol08] Oded Goldreich, *Computational complexity. A conceptual perspective*, Cambridge Univ. Press, Cambridge, UK, 2008, 606 pp.
- [Gra82] Ronald L. Graham, Linear extensions of partial orders and the FKG inequality, in *Ordered sets*, Reidel, Boston, MA, 1982, 213–236.
- [Gra83] Ronald L. Graham, Applications of the FKG inequality and its relatives, in *Mathematical programming: the state of the art*, Springer, Berlin, 1983, 115–131.
- [GK94] Peter Gritzmann and Victor Klee, On the complexity of some basic problems in computational convexity. II. Volume and mixed volumes, in *Polytopes: abstract, convex and computational*, Kluwer, Dordrecht, 1994, 373–466.
- [GK97] Peter Gritzmann and Victor Klee, Computational convexity, in *Handbook of discrete and computational geometry*, CRC, Boca Raton, FL, 1997, 491–515.
- [Gro90] Helmut Groemer, Stability properties of geometric inequalities, *Amer. Math. Monthly* **97** (1990), 382–394.
- [Gro93] Helmut Groemer, Stability of geometric inequalities, in *Handbook of convex geometry*, Vol. A, North-Holland, Amsterdam, 1993, 125–150.
- [Gro86] Mikhael Gromov, Isoperimetric inequalities in Riemannian manifolds, *Lecture Notes in Math.* **1200**, Springer, Berlin, 1986, 114–129; available at tinyurl.com/2p93zyvz
- [Gur08] Leonid Gurvits, Van der Waerden/Schrijver–Valiant like conjectures and stable (aka hyperbolic) homogeneous polynomials: one theorem for all, *Electronic J. Combin.* **15** (2008), no. 1, RP 66, 26 pp.
- [Hal01] Thomas C. Hales, The honeycomb conjecture, *Discrete Comput. Geom.* **25** (2001), 1–22.
- [HW08] Godfrey H. Hardy and Edward M. Wright, *An introduction to the theory of numbers* (sixth ed., revised), Oxford Univ. Press, Oxford, 2008, 621 pp.
- [HW20] Daniel Hug and Wolfgang Weil, *Lectures on convex geometry*, Springer, Cham, 2020, 287 pp.
- [Huh18] June Huh, Combinatorial applications of the Hodge–Riemann relations, in *Proc. ICM Rio de Janeiro*, vol. IV, World Sci., Hackensack, NJ, 2018, 3093–3111.
- [IP22] Christian Ikenmeyer and Igor Pak, What is in $\#P$ and what is not?, preprint (2022), 82 pp.; extended abstract in *Proc. 63rd FOCS* (2022), 860–871; [arXiv:2204.13149](https://arxiv.org/abs/2204.13149).
- [IPP22] Christian Ikenmeyer, Igor Pak and Greta Panova, Positivity of the symmetric group characters is as hard as the polynomial time hierarchy, *Int. Math. Res. Not.* (2024), no. 10, 8442–8458.
- [IN15] Emanuel Andrei and Levon Nurbekyan, On the stability of the polygonal isoperimetric inequality, *Adv. Math.* **276** (2015), 62–86.
- [KS84] Jeff Kahn and Michael Saks, Balancing poset extensions, *Order* **1** (1984), 113–126.
- [Kal22] Gil Kalai, The work of June Huh, in *Proc. ICM* (2022, virtual), 16 pp.; available at tinyurl.com/dnnncve5
- [KK12] Kiumars Kaveh and Askold Khovanskii, Algebraic equations and convex bodies, in *Perspectives in analysis, geometry, and topology*, Birkhäuser, New York, 2012, 263–282.
- [Kis68] Sergey S. Kislitsyn, A finite partially ordered set and its corresponding set of permutations, *Math. Notes* **4** (1968), 798–801.
- [Knu81] Donald E. Knuth, A permanent inequality, *Amer. Math. Monthly* **88** (1981), 731–740; Erratum 798.
- [Knu98] Donald E. Knuth, *The art of computer programming. Vol. 2. Seminumerical algorithms* (third ed.), Addison-Wesley, Reading, MA, 1998, 762 pp.
- [KS21] Noah Kravitz and Ashwin Sah, Linear extension numbers of n -element posets, *Order* **38** (2021), 49–66.
- [KS99] Ravi Kumar and Dandapani Sivakumar, Roundness estimation via random sampling, in *Proc. 10th SODA*, ACM, New York, 1999, 603–612.
- [LP07] Thomas Lam and Pavlo Pylyavskyy, Cell transfer and monomial positivity, *J. Algebraic Combin.* **26** (2007), 209–224.
- [LS10] Monique Laurent and Alexander Schrijver, On Leonid Gurvits’s proof for permanents, *Amer. Math. Monthly* **117** (2010), 903–911.
- [Law91] Jim Lawrence, Polytope volume computation, *Math. Comp.* **57** (1991), 259–271.
- [Lei80] Kurt Leichtweiss, *Konvexe Mengen* (in German), DVW, Berlin, 1980, 330 pp.
- [Lin86] Nathan Linial, Hard enumeration problems in geometry and combinatorics, *SIAM J. Algebraic Discrete Methods* **7** (1986), 331–335.
- [LMS19] Ricky I. Liu, Karola Mészáros and Avery St. Dizier, Gelfand–Tsetlin polytopes: a story of flow and order polytopes, *SIAM J. Discrete Math.* **33** (2019), 2394–2415.
- [Loe11] Nicholas A. Loehr, *Bijective combinatorics*, CRC Press, Boca Raton, FL, 2011, 590 pp.
- [MS24] Zhao Yu Ma and Yair Shenfeld, The extremals of Stanley’s inequalities for partially ordered sets, *Adv. Math.* **436** (2024), Paper No. 109404, 72 pp.

- [Mar17] Yves Martinez-Maure, A stability estimate for the Aleksandrov–Fenchel inequality under regularity assumptions, *Monatsh. Math.* **182** (2017), 65–76.
- [Mat02] Jiří Matoušek, *Lectures on discrete geometry*, Springer, New York, 2002, 481 pp.
- [MNY21] Satoshi Murai, Takahiro Nagaoka and Akiko Yazawa, Strictness of the log-concavity of generating polynomials of matroids, *J. Combin. Theory, Ser. A* **181** (2021), Paper 105351, 22 pp.
- [Oss78] Robert Osserman, The isoperimetric inequality, *Bull. AMS* **84** (1978), 1182–1238.
- [Oss79] Robert Osserman, Bonnesen–style isoperimetric inequalities, *Amer. Math. Monthly* **86** (1979), 1–29.
- [Pak05] Igor Pak, Partition bijections, a survey, *Ramanujan J.* **12** (2006), 5–75.
- [Pak09] Igor Pak, *Lectures on discrete and polyhedral geometry*, monograph draft (2009), 440 pp.; available at math.ucla.edu/~pak/book.htm
- [Pak19] Igor Pak, Combinatorial inequalities, *Notices AMS* **66** (2019), 1109–1112; an expanded version of the paper is available at tinyurl.com/py8sv5v6
- [Pak22] Igor Pak, What is a combinatorial interpretation?, in *Open Problems in Algebraic Combinatorics*, AMS, Providence, RI, to appear, 58 pp.; [arXiv:2209.06142](https://arxiv.org/abs/2209.06142).
- [Pap94] Christos H. Papadimitriou, *Computational Complexity*, Addison-Wesley, Reading, MA, 1994, 523 pp.
- [Por33] Thomas I. Porter, A history of the classical isoperimetric problem, in *Contributions to the Calculus of Variations*, Univ. Chicago Press, 1933, 475–523; available at tinyurl.com/526frnkf
- [StR81] Jean Saint-Raymond, Sur le volume des corps convexes symétriques (in French), in *Initiation Seminar on Analysis*, Publ. Math. Univ. Pierre et Marie Curie, Paris, 1981, 25 pp.
- [Schn85] Rolf Schneider, On the Aleksandrov–Fenchel inequality, in *Discrete geometry and convexity*, New York Acad. Sci., New York, 1985, 132–141.
- [Schn90a] Rolf Schneider, On the Aleksandrov–Fenchel inequality for convex bodies. I., *Results Math.* **17** (1990), 287–295.
- [Schn90b] Rolf Schneider, A stability estimate for the Aleksandrov–Fenchel inequality, with an application to mean curvature, *Manuscripta Math.* **69** (1990), 291–300.
- [Schn94] Rolf Schneider, Equality in the Aleksandrov–Fenchel inequality — present state and new results, in *Intuitive geometry*, North-Holland, Amsterdam, 1994, 425–438; available at tinyurl.com/5bwrvu9n
- [Schn14] Rolf Schneider, *Convex bodies: the Brunn–Minkowski theory* (second ed.), Cambridge Univ. Press, Cambridge, UK, 2014, 736 pp.
- [Schr86] Alexander Schrijver, *Theory of linear and integer programming*, John Wiley, Chichester, 1986, 471 pp.
- [Schr03] Alexander Schrijver, *Combinatorial optimization. Polyhedra and efficiency*, vols. A–C, Springer, Berlin, 2003, 1881 pp.
- [Schü72] Marcel-Paul Schützenberger, Promotion des morphismes d’ensembles ordonnés (in French), *Discrete Math.* **2** (1972), 73–94.
- [SW19] Pascal Schweitzer and Daniel Wiebking, A unifying method for the design of algorithms canonizing combinatorial objects, in *Proc. 51st STOC* (2019), 1247–1258.
- [Sey80] Paul D. Seymour, Decomposition of regular matroids, *J. Combin. Theory, Ser. B* **28** (1980), 305–359.
- [SvH19] Yair Shenfeld and Ramon van Handel, Mixed volumes and the Bochner method, *Proc. AMS* **147** (2019), 5385–5402.
- [SvH22] Yair Shenfeld and Ramon van Handel, The extremals of Minkowski’s quadratic inequality, *Duke Math. J.* **171** (2022), 957–1027.
- [SvH23] Yair Shenfeld and Ramon van Handel, The extremals of the Alexandrov–Fenchel inequality for convex polytopes, *Acta Math.* **231** (2023), 89–204.
- [She82] Lawrence A. Shepp, The XYZ conjecture and the FKG inequality, *Ann. Probab.* **10** (1982), 824–827.
- [Sid91] Alexander Sidorenko, Inequalities for the number of linear extensions, *Order* **8** (1991), 331–340.
- [Sta81] Richard P. Stanley, Two combinatorial applications of the Aleksandrov–Fenchel inequalities, *J. Combin. Theory, Ser. A* **31** (1981), 56–65.
- [Sta86] Richard P. Stanley, Two poset polytopes, *Discrete Comput. Geom.* **1** (1986), no. 1, 9–23.
- [Sta89] Richard P. Stanley, Log-concave and unimodal sequences in algebra, combinatorics, and geometry, in *Graph theory and its applications*, New York Acad. Sci., New York, 1989, 500–535.
- [Sta00] Richard P. Stanley, Positivity problems and conjectures in algebraic combinatorics, in *Mathematics: frontiers and perspectives*, AMS, Providence, RI, 2000, 295–319.
- [Sta09] Richard P. Stanley, Promotion and evacuation, *Electronic J. Combin.* **16** (2009), no. 2, RP 9, 24 pp.
- [Sta12] Richard P. Stanley, *Enumerative Combinatorics*, vol. 1 (second ed.) and vol. 2, Cambridge Univ. Press, 2012 and 1999, 626 pp. and 581 pp.
- [Tao08] Terence Tao, *Structure and Randomness. Pages from year one of a mathematical blog*, AMS, Providence, RI, 2008, 298 pp.; section §1.3 is available at wp.me/p3qzP-h
- [Toda91] Seinosuke Toda, PP is as hard as the polynomial-time hierarchy, *SIAM J. Comput.* **20** (1991), 865–877.

- [Tro95] William T. Trotter, Partially ordered sets, in *Handbook of combinatorics*, vol. 1, Elsevier, Amsterdam, 1995, 433–480.
- [VTL82] Jacobo Valdes, Robert E. Tarjan and Eugene L. Lawler, The recognition of series parallel digraphs, *SIAM J. Comput.* **11** (1982), 298–313.
- [vHYZ23] Ramon van Handel, Alan Yan and Ximeng Zeng, The extremals of the Kahn–Saks inequality, preprint (2023), 29 pp.; [arXiv:2309.13434](#).
- [vL82] Jacobus H. van Lint, The van der Waerden conjecture: two proofs in one year, *Math. Intelligencer* **4** (1982), no. 2, 72–77.
- [Wang18] Xu Wang, A remark on the Alexandrov–Fenchel inequality, *J. Funct. Anal.* **274** (2018), 2061–2088.
- [Wig19] Avi Wigderson, *Mathematics and computation*, Princeton Univ. Press, Princeton, NJ, 2019, 418 pp.; available at math.ias.edu/avi/book
- [YK75] Andrew C. Yao and Donald E. Knuth, Analysis of the subtractive algorithm for greatest common divisors, *Proc. Nat. Acad. Sci. USA* **72** (1975), 4720–4722.
- [Zhang98] Xin-Min Zhang, Schur-convex functions and isoperimetric inequalities, *Proc. AMS* **126** (1998), 461–470.

(Swee Hong Chan) DEPARTMENT OF MATHEMATICS, RUTGERS UNIVERSITY, PISCATWAY, NJ 08854.
Email address: sweehong.chan@rutgers.edu

(Igor Pak) DEPARTMENT OF MATHEMATICS, UCLA, LOS ANGELES, CA 90095.
Email address: pak@math.ucla.edu