VANISHING OF SCHUBERT COEFFICIENTS IS IN AM ∩ coAM ASSUMING THE GRH

IGOR PAK* AND COLLEEN ROBICHAUX*

ABSTRACT. The *Schubert vanishing problem* is a central decision problem in algebraic combinatorics and Schubert calculus, with applications to representation theory and enumerative algebraic geometry. The problem has been studied for over 50 years in different settings, with much progress given in the last two decades.

We prove that the Schubert vanishing problem is in AM assuming the *Generalized Riemann Hypothesis* (GRH). This complements our earlier result in [PR24b], that the problem is in coAM assuming the GRH. In particular, this implies that the Schubert vanishing problem is unlikely to be coNP-hard, as we previously conjectured in [PR24b].

The proof is of independent interest as we formalize and expand the notion of a *lifted formulation* partly inspired by algebraic computations of Schubert problems, and *extended formulations* of linear programs. We use the result by Mahajan–Vinay [MV97] to show that the determinant has a lifted formulation of polynomial size. We combine this with Purbhoo's algebraic criterion [Pur06] to derive the result.

FOREWORD

Despite appearances, the results of the paper do not require much of the background to state, see below. There is, however, a great deal of combinatorial and algebraic background needed to understand and appreciate the motivations. This occupies the rest of Section 1.

The heart of the proof is a combinatorial result that we call Determinant Lemma 2.2, which states that as a polynomial in commuting variables, the determinant has a lifted formulation of a polynomial size. Here the lifted formulations are an algebraic analogue of extended formulations for convex polyhedra, while the result is a statement in algebraic complexity theory. We present both the background and a short proof of the Determinant Lemma in Section 2.

Then, in Section 3 we present the proof of the main result. In Appendix A, we present four different definitions of Schubert polynomials, elucidating their different properties. Finally, in Appendix B we include various quotes on the history of the problem and its significance.

1. Schubert vanishing problem

1.1. Main result. Schubert polynomials $\mathfrak{S}_w \in \mathbb{N}[x_1, x_2, \ldots]$ indexed by permutations $w \in S_n$, are celebrated generalizations of Schur polynomials. Schubert polynomials satisfy partial symmetries, but are not symmetric in general. They were introduced by Lascoux and Schützenberger [LS82, LS85] to represent cohomology classes of Schubert varieties in the complete flag variety, building on the earlier works by Demazure [Dem74] and Bernstein–Gelfand–Gelfand [BGG73]. We refer to [Las95] for a historical introduction.

Schubert polynomials have been extensively studied from algebraic, combinatorial, representation theoretic, and computational points of view. Fundamentally, they represent a combinatorial approach to problems in enumerative algebraic geometry, answering questions of the form: *How* many lines in the space intersect four given lines in general position?¹ See [Mac91, Man01] for

April 5, 2025.

^{*}Department of Mathematics, UCLA, Los Angeles, CA 90095, USA. Email: {pak,robichaux}@math.ucla.edu.

¹The answer is 2 in this case, see e.g. [KL72]. Making rigorous sense of the natural generalization of this problem was the goal of *Hilbert's fifteenth problem* (1900). Resolving it required a major effort, resulted in several (equivalent) formal definitions, see e.g. [Kle76].

classic introductory surveys, [Knu16, Knu22] for overviews of recent results, [AF24, KM05] for geometric aspects, and [Pak24, §10] for an overview of computational complexity aspects.

Schubert coefficients are defined as structure constants:

$$\mathfrak{S}_u \cdot \mathfrak{S}_v = \sum_{w \in S_\infty} c_{u,v}^w \mathfrak{S}_w$$

for $u, v \in S_n$. It is known that $c_{u,v}^w \in \mathbb{N}$ for all $u, v, w \in S_\infty$, as they have a geometric meaning which generalize the number of intersection points of lines, see e.g. [AF24, Ful97]. Schubert polynomials and Schubert coefficients also emerge in representation theory [BS00, RS95], category theory [KP04], matroid theory [AB07], and the pole placement problem in linear systems theory [Byr89, EG02].

Since Schubert polynomials are Schur polynomials for *Grassmannian permutations* (permutations with one descent), Schubert coefficients generalize the *Littlewood–Richardson coefficients*. The *Schubert vanishing problem* is a decision problem

SCHUBERT VANISHING :=
$$\{c_{u,v}^w = {}^? 0\}$$
.

This is an extremely well studied problem, both for its own sake, and as a stepping stone towards understanding the nature of Schubert coefficients. The main result of this paper is the following:

Theorem 1.1 (Main theorem). SCHUBERTVANISHING $\in AM \cap coAM$ assuming the GRH.

Here the GRH stands for the Generalized Riemann Hypothesis, that all nontrivial zeros of Lfunctions $L(s, \chi_k)$ have real part $\frac{1}{2}$. The main theorem comes as a surprising improvement over the main result in [PR24b], which states that SCHUBERTVANISHING \in coAM assuming the GRH. See an extensive discussion of the prior work later in this section.

1.2. Schubert vanishing. It turns out, the Schubert vanishing problem can be stated in an elementary language in the style of classical geometry, without the use of Schubert polynomials. Since we will not need to use them, several formal definitions of Schubert polynomials are included in Appendix A in case the reader is curious.

Let $Z = \mathbb{C}^n$ be a fixed vector space with a basis $\{e_1, \ldots, e_n\}$. A complete flag F_{\bullet} is a sequence of subspaces $\{\mathbf{0}\} = F_0 \subset F_1 \subset F_2 \subset \ldots \subset F_n = Z$, where dim $(F_i) = i$ for all $0 \leq i \leq n$. Let \mathcal{F}_n denote the set of complete flags in Z. A coordinate flag $E_{\bullet} \in \mathcal{F}_n$ is a complete flag $\{\mathbf{0}\} = E_0 \subset E_1 \subset E_2 \subset \ldots \subset E_n = Z$, where $E_i = \mathbb{C}\langle e_1, \ldots, e_i \rangle$. A permutation flag $E_{\bullet}^w \in \mathcal{F}_n$ is a coordinate flag corresponding to the basis $\{e_{w(1)}, \ldots, e_{w(n)}\}$.

 \neg SCHUBERTVANISHING (= SCHUBERTNONVANISHING)

Input: $n \times n$ integral matrices $(a_{ij}), (b_{ij}), (c_{ij})$ **Decide:** $\forall U_{\bullet}, V_{\bullet} \in \mathcal{F}_n$ s.t. $U_i \cap V_{n-i} = \{\mathbf{0}\}$ for all $1 \leq i < n$, $\exists W_{\bullet} \in \mathcal{F}_n$ s.t. $\dim(W_i \cap E_j) = a_{ij}, \dim(W_i \cap U_j) \geq b_{ij}$ and $\dim(W_i \cap V_j) \geq c_{ij}$, for all $1 \leq i, j \leq n$

The naming choices for these complete flags is not accidental. It is easy to see that for every complete flag W_{\bullet} , the dimension matrix $\dim(W_i \cap E_j) = a_{ij}$ coincides with a dimension matrix for some permutation flag E_{\bullet}^w for some $w \in S_n$, so we have $a_{ij} = |\{w(1), \ldots, w(j)\} \cap \{1, \ldots, i\}|$. Similarly, taking $b_{ij} = |\{u(1), \ldots, u(j)\} \cap \{1, \ldots, i\}|$ and $c_{ij} = |\{v(1), \ldots, v(j)\} \cap \{1, \ldots, i\}|$ provides a translation between two equivalent formulations of the Schubert vanishing problem.

1.3. **Prior work: general results.** Much of the work on the problem has been a healthy collaboration and occasional competition of combinatorial and algebraic tools. Below we give a somewhat ahistorical overview, leaving the special cases until the end.

As stated in §1.2 above, the Schubert vanishing problem is not a priori decidable since there are uncountably many pairs of complete flags U_{\bullet}, V_{\bullet} to be checked. In fact, over \mathbb{R} the existence of W_{\bullet} can depend on U_{\bullet}, V_{\bullet} even if these two complete flags are generic; over large finite fields a major result by Vakil that this does not happen [Vak06].

Over \mathbb{C} , the problem simplifies significantly. We can always assume that U_{\bullet} is a permutation flag, while V_{\bullet} is a *generic flag*, but making the notion of "generic" quantitative is highly nontrivial and not well-understood in explicit terms. Heuristically, this phenomenon is a variation on the *polynomial identity testing* (PIT), where the problem is in BPP over large finite fields, while over \mathbb{R} the problem is believed to be not in PH.²

By extending combinatorial tools of Lascoux and Schützenberger, it was shown in [BB93, BJS93, FS94] that the Schubert-Kostka numbers $K_{w,\alpha} := [x_1^{\alpha_1} x_2^{\alpha_2} \cdots] \mathfrak{S}_w$ are nonnegative integers, and moreover that they are in #P as a counting function. This shows that SCHUBERTVANISHING \in PSPACE. The effective Möbius inversion argument in [PR24a, Thm 1.4] easily implies that computing Schubert coefficients is in GapP = #P - #P, and that SCHUBERTVANISHING $\in C_{=}P$. This was also observed earlier by Morales as a consequence of the Postnikov-Stanley formula [PS09, §17], see [Pak24, Prop. 10.2] for the explanation. Until recently, it was believed that SCHUBERTVANISHING \notin PH, and potentially even $C_{=}P$ -complete, see a discussion in [PR24a, §2.2].

In a different direction, a direct description of an algebraic system was given by Billey and Vakil in [BV08, Thm 5.4], which has exactly $c_{u,v}^w$ solutions for generic values of certain variables. They also describe the system of conditions for these variables being generic under the assumption that the set of solutions is 0-dimensional [BV08, Cor. 5.5]. The authors do not give a complexity analysis for this system; see [PR24b, §8.1] for further details and a complexity discussion.

In [HS17], Hein and Sottile introduced an algebraic system similar in flavor that they called a *lifted square formulation*, giving a practical algorithm for computing Schubert coefficients $c_{u,v}^w$. Their system had additional variables compared to the Billey–Vakil system, and allowed polynomial equations to have smaller (polynomial) size. This property was critical in the analysis given in [PR24a] (see below). We note that systems in other *numerical Schubert calculus* papers [HHS16, L+21] do not have polynomial size and are based on different principles.

Another approach was given by Purbhoo [Pur06] (see also [Bel06, §2]). He introduced a fundamentally different algebraic system which gives necessary and sufficient condition for vanishing of Schubert coefficients. Just like the Billey–Vakil and Hein–Sottile system, some variables were required to be generic to ensure that the set of solutions is 0-dimensional, a difficult condition to analyze computationally. An important feature of this approach is the dual nature of the system, as it gives an algebraic certificate for vanishing rather than non-vanishing.

In [PR24b, Thm 1.4], we proved that SCHUBERTVANISHING \in coAM assuming the GRH. We modified the Hein–Sottile system to prove the inclusion \neg SCHUBERTVANISHING \in HNP, the parametric version of the *Hilbert Nullstellensatz*. Then we used a recent result in [A+24], that HNP \in AM assuming the GRH, which is an extension of Koiran's celebrated result for the Hilbert Nullstellensatz [Koi96].

Our Main Theorem 1.1 that SCHUBERTVANISHING \in AM, is a complementary result proved by using a superficially similar inclusion SCHUBERTVANISHING \in HNP. The proof is based on Purbhoo's algebraic system. Curiously, we also used Purbhoo's algebraic system to show that SCHUBERTVANISHING \in coNP_C \cap P_R, a complexity result in the *Blum–Shub–Smale model of computation* over general fields [PR24b, Appendix B].

1.4. **Prior work: special cases.** There is a large number of sufficient conditions for vanishing of Schubert coefficients scattered across the literature. These were made both in an attempt to

²Over \mathbb{R} , PIT is equivalent to the existential theory of the reals ($\exists \mathbb{R}$), see e.g. [Sch10].

better understand the problem from a combinatorial point of view, and as a partial result towards its eventual resolution. Immediate from the combinatorial and geometric interpretations of the Schubert coefficients, we have the *dimension condition* that if $inv(u) + inv(v) \neq inv(w)$ then $c_{u,v}^w = 0$, where inv(w) denotes the number of inversions of w.

Further, we have the following conditions, which can be verified in polynomial time:

- the number of descents condition of Lascoux and Schützenberger [LS82],
- strong Bruhat order condition, see e.g. [SY22, §5.1] combined with [Man01, Prop. 2.1.11],
- Knutson's descent cycling condition [Knu01] (see also [PW24, Cor. 4.15]),
- permutation array condition by Billey and Vakil [BV08, Thm 5.1] (see also [AB07, Prop. 9.7]),
- St. Dizier and Yong's condition on certain filling of Rothe diagrams [SY22, Thm A], and
- Hardt and Wallach's condition on empty rows in Rothe diagrams [HW24, Cor. 5.12].

For Grassmannian permutations, Schubert coefficients are the *Littlewood–Richardson* (LR) coefficients, see e.g. [Mac91, Man01]. In this special case the vanishing problem is in P as a corollary of the Knutson–Tao saturation theorem [DM06, MNS12]. This is one of several important special cases where Schubert coefficients $c_{u,v}^w$ have a known combinatorial interpretation. In such cases, the combinatorial interpretation can be interpreted as NP (sufficient) conditions for non-vanishing. Notable examples include:

- Purbhoo's root game conditions [Pur04, Pur06], and
- Knutson and Zinn-Justin's several tiling conditions [KZ17, KZ23].

We refer to [SY22, §5] and [PR24b, §1.6] for technical details, comparisons, and further background on all these conditions.

1.5. **Implications.** Let us emphasize several implications of the main result.

1.5.1. New type of problem. From the computational complexity point of view, having a new natural problem in $AM \cap coAM$ is quite curious since this problem is apparently different from known problems in this class. Indeed, other problems in $AM \cap coAM$ include various "equivalence problems": graph isomorphism [GMW91] (see also [BHZ87]), code equivalence [PR97] (see also [BW24]), ring isomorphism [KS06], permutation group isomorphism [BCGQ11], and tensor isomorphism [GQ23].

In fact, the problems listed above are in lower complexity classes. For example, famously, GRAPH ISOMORPHISM is in NP \cap coAM, see e.g. [KST93], and is in the perfect/statistical zero knowledge classes PZK \subseteq SZK \subseteq AM \cap coAM [Vad99]. We refer to [BBM11] for a rare example of a problem that is in AM \cap coAM, but not necessarily in SZK.

1.5.2. Positive rule via derandomization. A major open problem in algebraic combinatorics is whether Schubert coefficients have a combinatorial interpretation [Sta00, Problem 11], see also §B.1 and §B.2. In the language of computational complexity this is asking whether this counting problem is in #P, see a detailed discussion in [Pak24, §10] (cf. also [Ass23]). This would imply that SCHUBERTVANISHING is in coNP. In the combinatorial language, this is saying that positivity of Schubert coefficients problem $\{c_{u,v}^w > ^? 0\}$ has a positive rule [PR24c].

The special cases mentioned in §1.4 suggest that both Schubert vanishing and Schubert non-vanishing might have a positive rule, i.e., that SCHUBERTVANISHING \in NP \cap coNP. Until recently this conclusion would seem fantastical and out of reach. Now, it was pointed out in [PR24b, PR24c], that a derandomization result by Miltersen–Vinodchandran [MV05] (extending [KvM02]), implies that AM = NP assuming some languages in NE \cap coNE require nondeterministic exponential size circuits.

For our purposes, a weaker derandomization assumption would also suffice: it was shown by Gutfreund, Shaltiel and Ta-Shma [GST03], that if EXP requires exponential time even for AM protocols (we call this GST assumption), then $AM \cap coAM = NP \cap coNP$. In a combinatorial language, this GST assumption combined with the GRH imply that there exists a positive rule for both vanishing and positivity of Schubert coefficients. While the latter is quite natural from a

combinatorial interpretation of Schubert coefficients point of view, the former is quite surprising, see below.

1.5.3. Computational hardness of Schubert vanishing. It has been known for a while that the vanishing of Schubert coefficients is computationally hard, see e.g. an extensive discussion in [BV08, §5.2] and §B.3. Here are two versions of the problem available in the literature.

Question 1.2 (Adve, Robichaux and Yong [ARY19, Question 4.3]). Is SCHUBERTVANISHING NPhard?

Conjecture 1.3 (Pak and Robichaux [PR24b, Conj. 1.6]). SCHUBERT VANISHING is coNP-hard.

The following result resolved both the question and the conjecture under standard assumptions:

Corollary 1.4. SCHUBERTVANISHING is not NP-hard, assuming the GRH and $PH \neq \Sigma_2^p$. Similarly, SCHUBERTVANISHING is not coNP-hard, assuming the GRH and $PH \neq \Pi_2^p$.

The corollary follows immediately from the Main Theorem 1.1 and a result of Boppana, Håstad and Zachos [BHZ87, Thm 2.3]. The corollary implies that the vanishing of Schubert coefficients is quite different from the vanishing of *Kronecker coefficients*, which is known to be coNP-hard even for partitions given in unary [IMW17]. See also [Pan23, §5.2] for further details and references.

1.6. Notation. We use $\mathbb{N} = \{0, 1, 2, ...\}$ and $[n] = \{1, ..., n\}$. We use $e_1, ..., e_n$ to denote the standard basis in \mathbb{C}^n , and **0** to denote zero vector. We use bold symbols such as $\boldsymbol{x}, \boldsymbol{y}, \boldsymbol{\alpha}, \boldsymbol{\beta}$ to denote sets and vectors of variables, and bars such as $\vec{\boldsymbol{x}}$ and $\vec{\boldsymbol{y}}$, to denote complex vectors. We also use \vec{f} to denote a sequence of polynomials (f_1, \ldots, f_m) .

In computational complexity, we use only standard notation and complexity classes. We refer to [AB09, Gol08, Pap94] for the definitions and extensive background, and to [Aar16] for the extensive introduction.

2. LIFTED AND COMPACT FORMULATIONS

The following section is completely independent of the rest of the paper and discusses Hilbert's Nullstellensatz and the technology of lifted formulations.

2.1. Hilbert's Nullstellensatz. Let $\mathbf{K} = \mathbb{C}[x_1, \dots, x_k]$ for some s > 0. We use $\mathbf{x} = (x_1, \dots, x_k)$. Consider a system

(2.1)
$$f_1(\boldsymbol{x}) = \ldots = f_m(\boldsymbol{x}) = 0 \text{ where } f_i \in \mathbf{K}$$

and denote by $\boldsymbol{S}(\overrightarrow{f}) \subseteq \mathbb{C}^k$ the set of solutions, where $\overrightarrow{f} = (f_1, \ldots, f_m)$.

Hilbert's weak Nullstellensatz is a fundamental result in algebra, which states that a polynomial system has no solutions over \mathbb{C} if and only if there exist $(g_1, \ldots, g_m) \in \mathbf{K}^m$, such that

$$\sum_{i=1}^m f_i g_i = 1.$$

Now let $f_1, \ldots, f_m \in \mathbb{Z}[x_1, \ldots, x_k]$. The decision problem HN (*Hilbert's Nullstellensatz*), asks if the polynomial system (2.1) has a solution over \mathbb{C}^3 . Here and everywhere below, the *size* of the polynomial system (2.1) is defined as

$$\phi(\overrightarrow{f}) := \sum_{i=1}^{m} \deg(f_i) + \sum_{i=1}^{m} s(f_i),$$

where s(g) denotes the sum of bit-lengths of coefficients in the polynomial g. Famously, Koiran showed that HN is in the polynomial hierarchy:

³By the Nullstellensatz, this is equivalent to asking if there is a solution over $\overline{\mathbb{Q}}$.

Theorem 2.1 ([Koi96, Thm 2]). HN is in AM assuming the GRH.

For the proof, Koiran's needs the existence of primes in certain intervals and with modular conditions, thus the GRH assumption. We refer to [A+24] for detailed overview of the problem and references to the earlier work.

2.2. Lifted formulations. Let $\mathbf{L} = \mathbb{C}[x_1, \ldots, x_k, y_1, \ldots, y_\ell]$ for some $k, \ell > 0$. We say that a system

(2.2)
$$g_1(\boldsymbol{x}, \boldsymbol{y}) = \ldots = g_\ell(\boldsymbol{x}, \boldsymbol{y}) = 0 \text{ where } g_i \in \mathbf{L}$$

is a *lifted formulation* of the system (2.1), if

(2.3)
$$\forall \boldsymbol{x} \in \boldsymbol{S}(\overrightarrow{f}) \exists \boldsymbol{y} \in \mathbb{C}^{\ell} : (\boldsymbol{x}, \boldsymbol{y}) \in \boldsymbol{S}(\overrightarrow{g}).$$

In other words, a natural projection $\iota : \mathbb{C}^{k+\ell} \to \mathbb{C}^k$ maps solutions of (2.1) into solutions of (2.2): $\iota(\mathbf{S}(\overrightarrow{g})) = \mathbf{S}(\overrightarrow{f})$. Clearly, HN for the system (2.1) is equivalent to HN for the system (2.2).

It may seem unintuitive that the size of a system of polynomials can become smaller for a lifted formulation, but this is not uncommon in symmetric situations. For example, polynomial $x^{2^r} = z$ has exponential size (in r), while its lifted formulation

$$y_1 = x^2, y_2 = y_1^2, \dots, y_r = y_{r-1}^2, y_r = z$$

has linear size.

The idea of lifted formulations is completely standard in numerical analysis and especially numerical algebraic geometry, see e.g. [HS17, L+21] for some recent work in the context of Schubert polynomials. Our own motivation comes from the literature on extended formulations of polyhedra, see below.

Lifted formulation (2.2) is called *compact* if $\phi(\vec{g}) = \text{poly}(k)$. A major obstacle in [PR24b], see Remark A.2, was that the determinant polynomial equation has exponential size. The following lemma shows that the determinant has a compact lifted formulation.

Lemma 2.2 (Determinant lemma). Let $X = (x_{ij})$ be an $n \times n$ matrix of variables. Then the determinant equation

(Det) $\det X = z$

has a lifted formulation of size $O(n^3)$.

This lemma is key for our proof of Theorem 1.1 as we use it essentially as a black box. We expect it to have further geometric and representation theoretic applications in the future.

2.3. **Prior work: extended formulations.** We modeled the notions above after extended formulations, the celebrated area of combinatorial optimization.

For a convex polyhedron $P \subset \mathbb{R}^k$ defined by inequalities, an *extended formulation* is a polyhedron $Q \subset \mathbb{R}^\ell$ which projects onto P. When Q is defined by poly(k) inequalities, such extended formulation is called *compact*. In such cases, one efficiently solve *linear programming* problems on P, by solving them on Q and projecting solutions onto P, even if P has exponentially many facets.

A prototypical example is the permutohedron $P_n \subset \mathbb{R}^n$ defined as a convex hull of points $(\sigma(1), \ldots, \sigma(n))$, where $\sigma \in S_n$. This polytope is (n-1)-dimensional, has n! vertices and (2^n-2) defining inequalities. The permutohedron has a compact extended formulation as a projection of the *Birkhoff polytope* $B_n \subset \mathbb{R}^{n^2}$, see e.g. [Pas14]. This is a $(n-1)^2$ -dimensional polytope of bistochastic matrices, with n! vertices given by 0/1 matrices, and with n^2 defining inequalities.

The idea of using extended formulations was formalized by Yannakakis in a remarkable paper [Yan91], where he showed that some combinatorial polytopes have compact extended formulations. More importantly, Yannakakis shows that matching and TSP polytopes, *do not* have compact extended formulations, under certain "symmetry" constraints. One can view the latter result as an obstacle to solving TSP, a benchmark NP-hard problem, using a polynomial size linear program.

 $\overline{7}$

These results inspired a long series of papers. Notably, the notion of extended formulations was generalized to *positive semidefinite lifts* in [GPT13]. The symmetry constraints were eventually removed in major advances [F+15, Rot17]. We refer to [CCZ13, CCZ14, KWY11] for the introduction to the area, background in combinatorial optimization and further references.

Our proof of the Determinant Lemma 2.2 is partially motivated by a beautiful construction of an extended formulation of size $O(n \log n)$ given by Goemans [Goe15], which is optimal and improves the $O(n^2)$ size of the Birkhoff polytope construction. His construction employs the structure and efficiency of the AKS sorting [AKS83], to simulate the sorting with linear inequalities. In effect, Goemans's construction is completely oblivious to the exact details the AKS sorting, which in turn uses explicit construction of expanders also as a black box.

2.4. **Prior work: complexity of the determinant.** Computing the permanent vs. determinant is a classical problem going back to Valiant [Val79b], leading to $VP = {}^{?}VNP$ problem. In turn, this problem led to foundations of the *Geometric Complexity Theory* (GCT), see e.g. [Aar16, BI18]. Of course, much of the effort is on lower bounds, a subject tangential to this paper.

There are several different ways one can restrict the computational model, e.g. [LR17] gives an exponential lower bound for the permanent under symmetry constraints. Similarly, the *Algebraic Complexity Theory* (ACT) restricts algebraic computations to straight line programs, and is the closest to our need. We refer to [BCS97] for the careful treatment of ACT and the background, and to [CKL24] for a recent treatment of the DETERMINANT in the context of (more general) algebraic branching problems.

We note that every straight-line program can be realized as a lifted formulation, but the smallest size lifted formulation can in principle be much smaller. This is not unusual; see [IL17] by Ikenmeyer and Landsberg, which compares complexity of computing the determinant and the permanent for different computational models. We note that commutativity of variables is crucial in this setting, since computing determinant over noncommutative rings requires exponential time [Nis91].

In the standard model, the differences between permanent vs. determinant become stark. Famously, the permanent of integer matrices is #P-complete [Val79a], while the determinant is GapLcomplete [Toda91, Vin91]. Another natural complete problem in GapL is the DIRECTEDPATHD-IFFERENCE, which inputs an acyclic digraph with a source s, two sinks t_+, t_- , and outputs the difference in the number of paths $s \to t_+$ and $s \to t_-$. In a beautiful paper [MV97], Mahajan and Vinay gave a parsimonious reduction of DIRECTEDPATHDIFFERENCE from the DETERMINANT, which we use in our proof of Lemma 2.2.

2.5. Proof of the Determinant Lemma 2.2. Recall the construction in [MV97]; see also [IL17, §3] for a concise presentation and examples. The authors construct an explicit directed acyclic graph Γ_n with the following properties:

- vertices of Γ_n have layers $\{0, \ldots, n\}$, and directed edges connect layers ℓ to $(\ell + 1)$,
- Γ_n has $O(n^3)$ vertices and $O(n^4)$ edges,
- \circ all edges in Γ_n have weights x_{ij} ,
- Γ_n has a unique source s at layer 0 and two sinks t_+, t_- at layer n,

• the sum of weighted paths $s \to t_+$ minus the sum of weighted paths $s \to t_-$ is det(X).

Here the weight of a path is a product of weights of its edges. In [MV97], the authors use this construction to show that det(X) can be computed in polynomial time using a straight line program (in contrast with the *Gaussian algorithm* which requires a circuit). We use the same graph construction to construct a compact lifted formulation as follows.

For a vertex v in Γ_n , denote by y_v the corresponding variables. Start the lifted formulation with an equation $y_s = 1$. For every vertex $v \neq s$, add an equation

$$y_v = \sum_{(w,v)\in\Gamma_n} y_w \cdot \operatorname{weight}(w,v),$$

where the summation is over all directed edges (w, v) in the graph Γ_n . Finally, add an equation

$$z = y_{t_+} - y_{t_-}$$
.

By construction, each y_v counts the sum of weighted paths $s \to v$. Thus, we obtain a lifted formulation for (Det) of size $O(n^3)$.

Remark 2.3. Note that just like in the Goemans's construction in [Goe15], the specifics of Γ_n are irrelevant, only the explicit nature and polynomial size are important. As we mentioned above, any straight-line computation of a polynomial $f(\mathbf{x})$ can be simulated by a lifted formulation, but not vice versa. It would be interesting to see if the $O(n^3)$ bound in the lemma can be improved. Given that we have $(n^2 + 1)$ variables, can one find a better lower bound for the smallest lifted formulation of (Det)?

3. Proof of the main theorem

3.1. Parametric Hilbert's Nullstellensatz. For the proof of the Main Theorem 1.1, we need the following strengthening of Theorem 2.1 to finite algebraic extensions. Let

 $f_1,\ldots,f_m \in \mathbb{Z}(y_1,\ldots,y_k)[x_1,\ldots,x_s].$

The decision problem HNP (*Parametric Hilbert's Nullstellensatz*) asks if the polynomial system (2.1) has a solution over $\overline{\mathbb{C}(y_1, \ldots, y_k)}$. In a remarkable recent work, Ait El Manssour, Balaji, Nosan, Shirmohammadi, and Worrell extended Theorem 2.1 to HNP :

Theorem 3.1 ([A+24, Thm 1]). HNP is in AM assuming the GRH.

We prove Theorem 1.1 as an application of Theorem 3.1. More precisely, recall that in [PR24b] the authors proved that \neg SCHUBERTVANISHING reduces to HNP. This and Theorem 3.1 immediately imply that SCHUBERTVANISHING \in coAM assuming the GRH. In this paper we prove the following counterpart:

Lemma 3.2 (Main lemma). SCHUBERTVANISHING reduces to HNP.

The lemma, combined with Theorem 3.1, immediately implies Main Theorem 1.1.

3.2. **Purbhoo's criterion.** Let $G = GL_n(\mathbb{C})$ be the general linear group. This is a matrix group lying in an ambient vector space $V \simeq \mathbb{C}^{n^2}$. Let B denote the Borel subgroup, i.e. the group of upper triangular matrices. Let N denote the subgroup of unipotent matrices, i.e. the group of upper triangular matrices with 1's on the diagonal. We have:

$$\mathsf{N} \subset \mathsf{B} \subset \mathsf{G} \subset V.$$

Let \mathfrak{n} denote the Lie algebra of N, i.e. the set of strictly upper triangular matrices (with 0's on the diagonal). We think of \mathfrak{n} as a subspace of V.

For a permutation $w \in S_n$, define $R_w := \mathfrak{n} \cap (w \cdot B_-)$, where $B_- = B^T$ is the subgroup of lower triangular matrices. One can think of $R_w = (r_{ij})$ as strictly upper triangular matrices with $r_{ij} = 0$ for all i < j and w(i) < w(j). For example, we have $R_e = \{\mathbf{0}\}$ for an identity permutation e = (1, 2, ..., n), and $R_{w_o} = \mathfrak{n}$ for the long permutation $w_o = (n, n - 1, ..., 1)$.

Lemma 3.3 (*Purbhoo's criterion* [Pur06, Corollary 2.6]). For generic $\rho, \omega, \tau \in \mathbb{N}$, we have:

$$c_{u,v}^{w} \neq 0 \quad \Longleftrightarrow \quad \rho R_{u} \rho^{-1} + \omega R_{v} \omega^{-1} + \tau R_{w \circ w} \tau^{-1} = \mathfrak{n}$$

Here the sum is the usual sum of vector subspaces of V.

3.3. **Proof of Main Lemma 3.2.** Considering the converse of Lemma 3.3, we consider the equation

(3.1)
$$c_{u,v}^{w} = 0 \quad \Longleftrightarrow \quad \rho R_{u} \rho^{-1} + \omega R_{v} \omega^{-1} + \tau R_{w \circ w} \tau^{-1} \subsetneq \mathfrak{n}.$$

By the dimension condition we assume inv(u) + inv(v) = inv(w). Then define

$$S_u := \{ x_{ij} \mathbf{e}_{ij} : i < j, u(i) > u(j) \},\$$

$$S_v := \{ y_{ij} \mathbf{e}_{ij} : i < j, v(i) > v(j) \}, \text{ and }\$$

$$S_{w \circ w} := \{ z_{ij} \mathbf{e}_{ij} : i < j, (w \circ w)(i) > (w \circ w)(j) \}$$

Here \mathbf{e}_{ij} is the $n \times n$ matrix with a 1 in position (i, j) and 0's elsewhere. Let $\mathbf{x}, \mathbf{y}, \mathbf{z}$ denote sets of those variables x_{ij}, y_{ij}, z_{ij} appearing therein. Then these sets form bases of R_u, R_v , and $R_{w \circ w}$, respectively. Note that for $\pi \in S_n$, dim $(R_\pi) = \operatorname{inv}(\pi)$. Thus since $\operatorname{inv}(u) + \operatorname{inv}(v) = \operatorname{inv}(w)$, dim $(R_u) + \dim(R_v) + \dim(R_{w \circ w}) = \binom{n}{2}$.

To construct generic $\rho \in N$, let $\rho = (\rho_{ij})$, where

$$\rho_{ij} = \begin{cases} \alpha_{ij} & \text{if } i < j, \\ 1 & \text{if } i = j, \\ 0 & \text{otherwise} \end{cases}$$

Here these α_{ij} are formal parameters. Similarly build generic $\omega, \tau \in \mathbb{N}$ in terms of parameters β_{ij}, γ_{ij} , respectively. Then define the sets of these parameters $\boldsymbol{\alpha} = \{\alpha_{ij}\}, \boldsymbol{\beta} = \{\beta_{ij}\}, \text{ and } \boldsymbol{\gamma} = \{\gamma_{ij}\},$ where the indices range over $1 \leq i < j \leq n$.

Again, we form $\tilde{\rho} = (\tilde{\rho}_{ij})$, where

$$\widetilde{\rho}_{ij} = \begin{cases} a_{ij} & \text{if } i < j, \\ 1 & \text{if } i = j, \\ 0 & \text{otherwise.} \end{cases}$$

Note that here we treat α_{ij} as variables. Similarly build generic $\tilde{\omega}, \tilde{\tau}$ in terms of variables b_{ij}, c_{ij} , respectively. Then define the sets of variables $\boldsymbol{a} = \{a_{ij}\}, \boldsymbol{b} = \{b_{ij}\}, \text{ and } \boldsymbol{c} = \{c_{ij}\}, \text{ where the indices range over } 1 \leq i < j \leq n$.

Then we obtain bases for $\rho R_u \tilde{\rho}$, $\omega R_v \tilde{\omega}$, and $\tau R_{w_o w} \tilde{\tau}$, respectively:

$$T_u := \rho S_u \widetilde{\rho},$$

$$T_v := \omega S_v \widetilde{\omega}, \text{ and }$$

$$T_{w_v w} := \tau S_{w_v w} \widetilde{\tau}.$$

Note $\dim(R_u) + \dim(R_v) + \dim(R_{w \circ w}) = \#T_u + \#T_v + \#T_{w \circ w} = \binom{n}{2} = \dim(\mathfrak{n}).$

By ignoring all entries weakly below the main diagonal in each matrix, we view each element in $T := T_u \cup T_v \cup T_{w \circ w}$ as a $\binom{n}{2}$ -vector. Let M be the $\binom{n}{2} \times \binom{n}{2}$ matrix formed by the vectors in T. Then the right-hand side of Equation (3.1) holds if and only if det(M) = 0, when we set $\tilde{\rho} = \rho^{-1}$, $\tilde{\omega} = \omega^{-1}$, and $\tilde{\tau} = \tau^{-1}$.

Let $\mathcal{S}(u, v, w_{\circ}w)$ be the system formed by the constraints:

$$\left\{ \begin{array}{l} \rho \cdot \widetilde{\rho} \,=\, \mathsf{Id}_n\,,\\ \omega \cdot \widetilde{\omega} \,=\, \mathsf{Id}_n\,,\\ \tau \cdot \widetilde{\tau} \,=\, \mathsf{Id}_n\,,\\ \det(M) \,=\, 0\,, \end{array} \right.$$

where the last constraint is replaced by its lifted formulation using the Determinant Lemma 2.2. Here $\mathcal{S}(u, v, w_{\circ}w)$ uses variables $\mathbf{a} \cup \mathbf{b} \cup \mathbf{c} \cup \mathbf{x} \cup \mathbf{y} \cup \mathbf{z}$ and parameters $\mathbf{\alpha} \cup \mathbf{\beta} \cup \mathbf{\gamma}$.

Note that matrix entries in M have size $O(n^2)$, as they are monomials of degree at most 2 whose nonzero coefficients are in $\boldsymbol{\alpha} \cup \boldsymbol{\beta} \cup \boldsymbol{\gamma}$. By the Determinant Lemma 2.2, the last equation and thus the whole system $\mathcal{S}(u, v, w_{\circ}w)$ has size $O(n^{12})$. Now, the right-hand side of Equation (3.1) holds if and only if $\mathcal{S}(u, v, w_{\circ}w)$ is satisfiable over $\mathbb{C}(\alpha, \beta, \gamma)$. Since $\alpha \cup \beta \cup \gamma$ are algebraically independent, $\mathcal{S}(u, v, w_{\circ}w)$ has a solution over $\mathbb{C}(\alpha, \beta, \gamma)$ if and only if $\mathcal{S}(u, v, w_{\circ}w)$ has a solution over \mathbb{C} for a generic choice of evaluations $\overrightarrow{\alpha}, \overrightarrow{\beta}, \overrightarrow{\gamma}$ of α , β, γ . Thus by Lemma 3.3, the result follows.

3.4. Further applications. As noted in §1.4 (see also §A.4), the Schubert structure constants $c_{u,v}^w$ are also the structure constants arising from multiplying Schubert classes in the cohomology ring of the complete flag variety [LS82]. More generally, we may consider structure constants arising from multiplying Schubert classes in $H^*(G/B)$, for a complex reductive Lie group $G \in \{GL_n, SO_{2n+1}, Sp_{2n}, SO_{2n}\}$. Here $B \subset G$ the Borel subgroup, the subgroup of upper triangular matrices in G.

These generalized flag varieties G/B may be referred to the type A, B, C, D flag varieties, respectively. Let $c_{u,v}^w(Y)$ denote the corresponding type Y structure constants, where $Y \in \{A, B, C, D\}$.⁴ Here u, v, w are elements in the Weyl group \mathcal{W} in type Y. See [PR24b, §4.2] for a brief overview, or [AF24] for a detailed exposition. So far, we had focused on the type A structure constants, i.e. the GL_n case.

Continuing this generalization, we may employ the type Y Schubert polynomials \mathfrak{S}_w^Y of Billey– Haiman [BH95] to compute these structure coefficients:

$$\mathfrak{S}_u^Y \cdot \mathfrak{S}_v^Y = \sum_{w \in \mathcal{W}} c_{u,v}^w(Y) \mathfrak{S}_w^Y.$$

Thus we consider the type Y Schubert vanishing problem:

SchubertVanishing
$$(Y) := \{c_{u,v}^w(Y) = {}^? 0\}.$$

The main result of this paper extends to all classical type:

Theorem 3.4 (Schubert vanishing for all types). SCHUBERT VANISHING(Y) \in AM \cap coAM assuming the GRH, for all $Y \in \{A, B, C, D\}$.

Proof. By [PR24b, Theorem 2.4], SCHUBERTVANISHING(Y) \in coAM assuming the GRH for $Y \in \{A, B, C\}$. Further, as noted in [PR24b, Remark A.2], the analogous result in type D was prevented by a lingering determinantal equation det(ω) = 1. The Determinant Lemma 2.2 resolves this issue, proving that SCHUBERTVANISHING(D) \in coAM.⁵

Further, the generality of [Pur06, Corollary 2.6] gives a vanishing criterion in types A, B, C, D. For brevity, we suppress the details of the translation to types B, C, D and instead review the general framework.

Take the unipotent subgroup $\mathbb{N} \subset \mathbb{B} \subset \mathbb{G}$ of type Y. Let \mathfrak{n} denote the Lie algebra of \mathbb{N} . Then for $w \in \mathcal{W}$, define $R_w := \mathfrak{n} \cap (w\mathbb{B}_-w^{-1})$, where \mathbb{B}_- are the lower triangular matrices in \mathbb{G} . Then for generic elements $\rho, \omega, \tau \in \mathbb{N}$:

$$c_{u,v}^w = 0 \quad \Longleftrightarrow \quad \rho R_u \rho^{-1} + \omega R_v \omega^{-1} + \tau R_{w \circ w} \tau^{-1} \subsetneq \mathfrak{n}.$$

The argument used for Main Lemma 3.2 works verbatim to translate the right-hand side into a system of polynomial equations. The only adjustment is that we may impose additional equations to ensure $\rho, \omega, \tau \in \mathsf{G}$, as specified in the relevant sections of [PR24b]. Using the Determinant Lemma 2.2, the resulting systems have polynomial size. This shows SCHUBERTVANISHING(Y) reduces to HNP for $Y \in \{A, B, C, D\}$. Thus the result follows using Theorem 3.1.

⁴For non-classical types E_6 , E_7 , E_8 , F_4 and G_2 , there is only a finite number of Schubert coefficients, so the problem is uninteresting from the computational complexity point of view.

⁵In the Appendix C of a revised version of [PR24b], written jointly with David Speyer, the authors circumvent this issue in a different way and also prove that SCHUBERTVANISHING(D) \in coAM.

Acknowledgements. We are grateful to Kevin Purbhoo for sharing his insights which partially motivated this paper. We thank Sara Billey, Nickolas Hein, Minyoung Jeon, Allen Knutson, Leonardo Mihalcea, Greta Panova, Oliver Pechenik, Maurice Rojas, Mahsa Shirmohammadi, Alex Smith, Frank Sottile, David Speyer, Avi Wigderson, James Worrell, Weihong Xu, Alex Yong and Paul Zinn-Justin for interesting discussions and helpful comments. The first author was partially supported by the NSF grant CCF-2302173. The second author was partially supported by the NSF grant DMS-2302279.

References

- [Aar16] Scott Aaronson, $P \stackrel{?}{=} NP$, in Open problems in mathematics, Springer, Cham, 2016, 1–122.
- [ARY19] Anshul Adve, Colleen Robichaux and Alexander Yong, Vanishing of Littlewood–Richardson polynomials is in P, Comput. Complexity 28 (2019), 241–257.
- [A+24] Rida Ait El Manssour, Nikhil Balaji, Klara Nosan, Mahsa Shirmohammadi and James Worrell, A parametric version of the Hilbert Nullstellensatz, in *Proc. 8th SOSA* (2025), 444–451.
- [AKS83] Miklós Ajtai, János Komlós and Endre Szemerédi, Sorting in $c \log n$ parallel steps, Combinatorica **3** (1983), 1–19; extended abstract in Proc. 15th STOC (1983), 1–9.
- [AF24] David Anderson and William Fulton, *Equivariant cohomology in algebraic geometry*, Cambridge Univ. Press, Cambridge, UK, 2024, 446 pp.
- [AB07] Federico Ardila and Sara Billey, Flag arrangements and triangulations of products of simplices, Adv. Math. 214 (2007), 495–524.
- [AB09] Sanjeev Arora and Boaz Barak, *Computational complexity. A modern approach*, Cambridge Univ. Press, Cambridge, 2009, 579 pp.
- [Ass23] Sami Assaf, Schubert structure constants via Kohnert combinatorics, NSF DMS-2246785 Award (June 14, 2023); abstract is available at tinyurl.com/4zs2md43
- [BCGQ11] László Babai, Paolo Codenotti, Joshua A. Grochow and Youming Qiao, Code equivalence and group isomorphism, in Proc. 22nd SODA (2011), 1395–1408.
- [Bel06] Prakash Belkale, Geometric proofs of Horn and saturation conjectures, J. Algebraic Geom. 15 (2006), 133–173.
- [BW24] Huck Bennett and Kaung Myat Htay Win, Relating code equivalence to other isomorphism problems, Designs, Codes and Cryptography (published online Dec. 16, 2024), 23 pp.
- [BS00] Arkady Berenstein and Reyer Sjamaar, Coadjoint orbits, moment polytopes, and the Hilbert–Mumford criterion, J. Amer. Math. Soc. 13 (2000), 433–466.
- [BB93] Nantel Bergeron and Sara Billey, RC-graphs and Schubert polynomials, *Exp. Math.* 2 (1993), 257–269.
- [BGG73] I. N. Bernstein, I. M. Gelfand, and S. I. Gelfand, Schubert cells, and the cohomology of the spaces G/P, Uspehi Mat. Nauk 28 (1973), 3–26.
- [BBM11] Nayantara Bhatnagar, Andrej Bogdanov and Elchanan Mossel, The computational complexity of estimating MCMC convergence time, in *Proc. 14th APPROX* (2011), 424–435.
- [BH95] Sara Billey and Mark Haiman, Schubert polynomials for the classical groups, *Jour. AMS* 8 (1995), 443–482.
- [BGP25+] Sara Billey, Yibo Gao and Brendan Pawlowski, Introduction to the cohomology of the flag variety, monograph in preparation, 2025, 161 pp.
- [BJS93] Sara Billey, William Jockusch and Richard P. Stanley, Some combinatorial properties of Schubert polynomials, J. Algebraic Combin. 2 (1993), 345–374.
- [BV08] Sara Billey and Ravi Vakil, Intersections of Schubert varieties and other permutation array schemes, in *Algorithms in algebraic geometry*, Springer, New York, 2008, 21–54.
- [BI18] Markus Bläser and Christian Ikenmeyer, Introduction to geometric complexity theory, preprint (2018), 148 pp.; to appear in *Theory of Computing Graduate Surveys*; available at tinyurl.com/mpan48e6
- [BHZ87] Ravi B. Boppana, Johan Håstad and Stathis Zachos, Does coNP have short interactive proofs?, Inform. Process. Lett. 25 (1987), 127–132.
- [Bor53] Armand Borel, Sur la cohomologie des espaces fibrés principaux et des espaces homogènes de groupes de Lie compacts, Ann. of Math. (2) 57 (1953), 115–207.
- [BCS97] Peter Bürgisser, Michael Clausen and M. Amin Shokrollahi, Algebraic complexity theory, Springer, Berlin, 1997, 618 pp.
- [Byr89] Christopher I. Byrnes, Pole assignment by output fedback, in Three decades of mathematical system theory, Springer, Berlin, 1989, 31–78.
- [CKL24] Abhranil Chatterjee, Mrinal Kumar and Ben Lee Volk, Determinants vs. algebraic branching programs, in Proc. 15th ITCS (2024), paper 27, 13 pp.

- [CCZ13] Michele Conforti, Gérard Cornuéjols and Giacomo Zambelli, Extended formulations in combinatorial optimization, Ann. Oper. Res. 204 (2013), 97–143.
- [CCZ14] Michele Conforti, Gérard Cornuéjols and Giacomo Zambelli, *Integer programming*, Springer, Cham, 2014, 456 pp.
- [DM06] Jesús A. De Loera and Tyrrell B. McAllister, On the computation of Clebsch–Gordan coefficients and the dilation effect, *Experiment. Math.* **15** (2006), 7–19.
- [Dem74] Michel Demazure, Désingularisation des variétés de Schubert généralisées, Ann. Sci. École Norm. Sup. 7 (1974), 53–88.
- [EG02] Alexandre Eremenko and Andrei Gabrielov, Pole placement by static output feedback for generic linear systems, *SIAM J. Control Optim.* **41** (2002), 303–312.
- [F+15] Samuel Fiorini, Serge Massar, Sebastian Pokutta, Hans Raj Tiwary and Ronald de Wolf, Exponential lower bounds for polytopes in combinatorial optimization, J. ACM 62 (2015), no. 2, Art. 17, 23 pp.; extended abstract in Proc. 44th STOC (2012), 95–106.
- [FS94] Sergey Fomin and Richard P. Stanley, Schubert polynomials and the nil-Coxeter algebra, Adv. Math. 103 (1994), 196–207.
- [Ful97] William Fulton, Young tableaux, Cambridge Univ. Press, Cambridge, UK, 1997, 260 pp.
- [Goe15] Michel X. Goemans, Smallest compact formulation for the permutahedron, *Math. Program.* **153** (2015), no. 1, Ser. B, 5–11.
- [Gol08] Oded Goldreich, Computational complexity. A conceptual perspective, Cambridge Univ. Press, Cambridge, UK, 2008, 606 pp.
- [GMW91] Oded Goldreich, Silvio Micali and Avi Wigderson, Proofs that yield nothing but their validity, or All languages in NP have zero-knowledge proof systems, J. Assoc. Comput. Mach. 38 (1991), 691–729; extended abstract in Proc. 27th FOCS (1986), 174–187.
- [GPT13] João Gouveia, Pablo A. Parrilo and Rekha R. Thomas, Lifts of convex sets and cone factorizations, Math. Oper. Res. 38 (2013), 248–264.
- [GQ23] Joshua Grochow and Youming Qiao, On the complexity of isomorphism problems for tensors, groups, and polynomials I: Tensor isomorphism-completeness, SIAM J. Comput. 52 (2023), 568–617; extended abstract in Proc. 12th ICTS (2021), Art. No. 31, 19 pp.; arXiv:1907.00309.
- [GST03] Dan Gutfreund, Ronen Shaltiel and Amnon Ta-Shma, Uniform hardness versus randomness tradeoffs for Arthur–Merlin games, *Comput. Complexity* **12** (2003), 85–130.
- [HW24] Andrew Hardt and David Wallach, When do Schubert polynomial products stabilize?, preprint (2024), 32 pp.; arXiv:2412.06976.
- [HHS16] Jonathan D. Hauenstein, Nickolas Hein and Frank Sottile, A primal-dual formulation for certifiable computations in Schubert calculus, Found. Comput. Math. 16 (2016), 941–963.
- [HS17] Nickolas Hein and Frank Sottile, A lifted square formulation for certifiable Schubert calculus, J. Symbolic Comput. 79 (2017), 594–608.
- [IL17] Christian Ikenmeyer and Joseph M. Landsberg, On the complexity of the permanent in various computational models, J. Pure Appl. Algebra 221 (2017), 2911–2927.
- [IMW17] Christian Ikenmeyer, Ketan D. Mulmuley and Michael Walter, On vanishing of Kronecker coefficients, Comp. Complexity 26 (2017), 949–992.
- [IP22] Christian Ikenmeyer and Igor Pak, What is in #P and what is not?, preprint (2022), 82 pp.; extended abstract in Proc. 63rd FOCS (2022), 860-871; arXiv:2204.13149.
- [KWY11] Volker Kaibel, Laurence A. Wolsey and Mihalis Yannakakis, Extended formulations, Optima 85 (April, 2011), 2–10; available at https://www.mathopt.org/Optima-Issues/optima85.pdf
- [KS06] Neeraj Kayal and Nitin Saxena, Complexity of ring morphism problems, *Comput. Complexity* **15** (2006), 342–390.
- [Kle74] Steven L. Kleiman, The transversality of a general translate, *Compositio Math.* 28 (1974), 287–297.
- [Kle76] Steven L. Kleiman, Problem 15: Rigorous foundation of Schubert's enumerative calculus, in Mathematical developments arising from Hilbert problems, AMS, Providence, RI, 1976, 445–482.
- [KL72] Steven L. Kleiman and Dan Laksov, Schubert calculus, Amer. Math. Monthly 79 (1972), 1061–1082.
- [KvM02] Adam R. Klivans and Dieter van Melkebeek, Graph nonisomorphism has subexponential size proofs unless the polynomial-time hierarchy collapses, SIAM J. Comput. 31 (2002), 1501–1526; extended abstract in Proc. 31st STOC (1999), 659–667.
- [Knu01] Allen Knutson, Descent-cycling in Schubert calculus, Experiment. Math. 10 (2001), 345–353.
- [Knu16] Allen Knutson, Schubert calculus and puzzles, in Adv. Stud. Pure Math. 71, Math. Soc. Japan, Tokyo, 2016, 185–209.
- [Knu22] Allen Knutson, Schubert calculus and quiver varieties, in *Proc. ICM* (2022, virtual), Vol. VI, EMS Press, 4582–4605.
- [KM05] Allen Knutson and Ezra Miller, Gröbner geometry of Schubert polynomials, Annals of Math. 161 (2005), 1245–1318.

12

- [KZ17] Allen Knutson and Paul Zinn-Justin, Schubert puzzles and integrability I: invariant trilinear forms, preprint (2017), 51 pp.; arXiv:1706.10019.
- [KZ23] Allen Knutson and Paul Zinn-Justin, Schubert puzzles and integrability III: separated descents, preprint (2023), 42 pp.; arXiv:2306.13855.
- [KST93] Johannes Köbler, Uwe Schöning and Jacobo Torán, The graph isomorphism problem: its structural complexity, Birkhäuser, Boston, MA, 1993, 160 pp.
- [Koi96] Pascal Koiran, Hilbert's Nullstellensatz is in the polynomial hierarchy, J. Complexity 12 (1996), 273–286; see also DIMACS Tech. Report 96-27 (1996), 18 pp.
- [KP04] Witold Kraśkiewicz and Piotr Pragacz, Schubert functors and Schubert polynomials, European J. Combin. 25 (2004), 1327–1344.
- [LR17] Joseph M. Landsberg and Nicolas Ressayre, Permanent v. determinant: an exponential lower bound assuming symmetry and a potential path towards Valiant's conjecture, *Differential Geom. Appl.* 55 (2017), 146–166; extended abstract in *Proc. 7th ITCS* (2016), 29–35.
- [Las95] Alain Lascoux, Polynômes de Schubert: une approche historique (in French), Discrete Math. 139 (1995), 303–317.
- [LS82] Alain Lascoux and Marcel-Paul Schützenberger, Polynômes de Schubert (in French), C. R. Acad. Sci. Paris Sér I, Math. 294 (1982), 447–450.
- [LS85] Alain Lascoux and Marcel-Paul Schützenberger, Schubert polynomials and the Littlewood–Richardson rule, *Lett. Math. Phys.* **10** (1985), 111–124.
- [Len10] Cristian Lenart, Growth diagrams for the Schubert multiplication, J. Combin. Theory, Ser. A —textbf117 (2010), 842–856.
- [L+21] Anton Leykin, Abraham Martín del Campo, Frank Sottile, Ravi Vakil and Jan Verschelde, Numerical Schubert calculus via the Littlewood–Richardson homotopy algorithm, Math. Comp. 90 (2021), 1407– 1433.
- [Mac91] Ian G. Macdonald, Notes on Schubert polynomials, Publ. LaCIM, UQAM, Montreal, 1991, 116 pp.; available at tinyurl.com/382f7an7
- [MV97] Meena Mahajan and Vishwanathan Vinay, Determinant: combinatorics, algorithms, and complexity, Chicago J. Theoret. Comput. Sci. **1997** (1997), Article 5, 26 pp.
- [Man01] Laurent Manivel, Symmetric functions, Schubert polynomials and degeneracy loci, SMF/AMS, Providence, RI, 2001, 167 pp.
- [MV05] Peter B. Miltersen and N. Variyam Vinodchandran, Derandomizing Arthur–Merlin games using hitting sets, *Comput. Complexity* 14 (2005), 256–279; extended abstract in *Proc. 4th FOCS* (1999), 71–80.
- [MNS12] Ketan D. Mulmuley, Hariharan Narayanan and Milind Sohoni, Geometric complexity theory III. On deciding nonvanishing of a Littlewood–Richardson coefficient, J. Algebraic Combin. 36 (2012), 103–110.
- [Nis91] Noam Nisan, Lower bounds for non-commutative computation, in *Proc. 23rd STOC* (1991), 410–418. [Pak24] Igor Pak, What is a combinatorial interpretation?, in *Open Problems in Algebraic Combinatorics*, AMS
- [Pak24] Igor Pak, What is a combinatorial interpretation?, in Open Problems in Algebraic Combinatorics, AMS, Providence, RI, 2024, 191–260.
- [PR24a] Igor Pak and Colleen Robichaux, Signed combinatorial interpretations in algebraic combinatorics, preprint (2024), 24 pp.; to appear in Algebraic Combinatorics; extended abstract to appear in Proc. 37th FPSAC (2025); arXiv:2406.13902.
- [PR24b] Igor Pak and Colleen Robichaux, Vanishing of Schubert coefficients, preprint (2024), 30 pp.; arXiv: 2412.02064; extended abstract to appear in Proc. 57th STOC (2025).
- [PR24c] Igor Pak and Colleen Robichaux, Positivity of Schubert coefficients, preprint, 7 pp.; arXiv:2412.18984.
- [Pan23] Greta Panova, Computational complexity in algebraic combinatorics, to appear in Current Developments in Mathematics, International Press, Boston, MA, 30 pp.; arXiv:2306.17511.
- [Pap94] Christos H. Papadimitriou, Computational Complexity, Addison-Wesley, Reading, MA, 1994, 523 pp.
- [Pas14] Kanstantsin Pashkovich, Tight lower bounds on the sizes of symmetric extensions of permutahedra and similar results, Math. Oper. Res. 39 (2014), 1330–1339.
- [PW24] Oliver Pechenik and Anna Weigandt, An inverse Grassmannian Littlewood–Richardson rule and extensions, Forum Math. Sigma 12 (2024), Paper No. e114, 24 pp.
- [PR97] Erez Petrank and Ron M. Roth, Is code equivalence easy to decide?, IEEE Trans. Inform. Theory 43 (1997), 1602–1604.
- [PS09] Alexander Postnikov and Richard P. Stanley, Chains in the Bruhat order, J. Algebraic Combin. 29 (2009), 133–174.
- [Pur04] Kevin Purbhoo, Vanishing and non-vanishing criteria for branching Schubert calculus, Ph.D. thesis, UC Berkeley, 2004, 96 pp.; available at tinyurl.com/4bbueuwn
- [Pur06] Kevin Purbhoo, Vanishing and nonvanishing criteria in Schubert calculus, Int. Math. Res. Notices 2006 (2006), Art. 24590, 38 pp.
- [RS95] Victor Reiner and Mark Shimozono, Key polynomials and a flagged Littlewood-Richardson rule, J. Combin. Theory, Ser. A 70 (1995), 107–143.

- [Rot17] Thomas Rothvoss, The matching polytope has exponential extension complexity, J. ACM 64 (2017), no. 6, Art. 41, 19 pp.; extended abstract in Proc. 46th STOC (2014), 263–272.
- [Sch10] Marcus Schaefer, Complexity of some geometric and topological problems, in Lecture Notes Comput. Sci. 5849, Springer, Berlin, 2010, 334–344.
- [SY22] Avery St. Dizier and Alexander Yong, Generalized permutahedra and Schubert calculus, Arnold Math. J. 8 (2022), 517–533.
- [Sta00] Richard P. Stanley, Positivity problems and conjectures in algebraic combinatorics, in *Mathematics: frontiers and perspectives*, AMS, Providence, RI, 2000, 295–319.
- [Toda91] Seinosuke Toda, Counting problems computationally equivalent to computing the determinant, *Tech. Report CSIM* **91-07**, 21 pp.
- [Vad99] Salil Vadhan, A study of statistical zero-knowledge proofs, Ph.D. thesis, MIT, Cambridge, MA, 1999, 190 pp.
- [Vak06] Ravi Vakil, Schubert induction, Annals of Math. 164 (2006), 489–512.
- [Val79a] Leslie G. Valiant, The complexity of enumeration and reliability problems, SIAM J. Comput. 8 (1979), 410–421.
- [Val79b] Leslie G. Valiant, Completeness classes in algebra, in *Proc. 11th STOC* (1979), 249–261.
- [Vin91] Vishwanathan Vinay, Counting auxiliary pushdown automata and semi-unbounded arithmetic circuits, in Proc. 6th SCT (1991), 270–284.
- [Yan91] Mihalis Yannakakis, Expressing combinatorial optimization problems by linear programs, J. Comput. System Sci. 43 (1991), 441–466; extended abstract in Proc. 20th STOC (1988), 223–228.

Appendix A. Schubert polynomials

Although we do not use Schubert polynomials to state or prove the main result, the concept they represent is fundamental to fully understand the meaning of Main Theorem 1.1. We thus include several different definitions for reader's convenience. We refer to [Man01, Knu16] for more on these definitions and further background.

A.1. Divided differences. The following is the original definition due to Lascoux and Schützenberger [LS82]. For a permutation $w_{\circ} = (n, n - 1, ..., 2, 1)$, let

$$\mathfrak{S}_{w_\circ} := x_1^{n-1} x_2^{n-2} \cdots x_{n-1}.$$

A permutation $w \in S_n$ is said to have a *descent* at *i*, if w(i) > w(i+1). Denote by Des(w) the *set* of descents of w, and by $\text{des}(\sigma) := |\text{Des}(\sigma)|$ the number of descents. Define the divided difference operator

$$\partial_i F := rac{F - s_i F}{x_i - x_{i+1}}$$

where the transposition $s_i := (i, i + 1)$ acts on $F \in \mathbb{C}[x_1, \ldots, x_n]$ by transposing the variables. For all $i \in \text{Des}(w)$, let

$$\mathfrak{S}_{ws_i} := \partial_i \mathfrak{S}_w,$$

and define all Schubert polynomials recursively. It follows that $\mathfrak{S}_w \in \mathbb{Z}[\mathbf{x}]$ are homogeneous polynomials of degree $\operatorname{inv}(w)$. Here $\operatorname{inv}(w) := \{(i, j) : i < j, w(i) > w(j)\}$ is the number of inversions in w.

To show that Schubert polynomials are well defined, one needs to check that

$$\partial_i \partial_j = \partial_j \partial_i$$
 for all $|i - j| \ge 2$, and $\partial_i \partial_{i+1} \partial_i = \partial_{i+1} \partial_i \partial_{i+1}$,

which follow from a straightforward computation.

This definition is elementary, easy to use, and can be generalized in various directions. The disadvantage of this definition is a nonobvious combinatorial nature of the coefficients. One can only conclude that $[\mathbf{x}^{\alpha}]\mathfrak{S}_{w} \in \mathbb{Z}$, but not that $[\mathbf{x}^{\alpha}]\mathfrak{S}_{w} \in \mathbb{N}$.

A.2. Working forward, not backward. Using a standard embedding S_n into S_{n+1} by adding a fixed point (n + 1), one can define a limit object S_{∞} to be the set of bijections $\sigma : \mathbb{N} \to \mathbb{N}$, where $\sigma(i) = i$ for all but finitely many *i*. We can now define Schubert polynomials forward, starting with $\mathfrak{S}(1) := 1$. Use the following rules:

$$\mathfrak{S}_{ws_i} := \partial_i \mathfrak{S}_w$$
 if $i \in \operatorname{Des}(w)$ and $\mathfrak{S}_{ws_i} := 0$ if $i \notin \operatorname{Des}(w)$.

In this setting, it is easy to see that Schubert polynomials are uniquely defined. The existence becomes a substantive result, but other things become more apparent, e.g. the symmetry properties of the construction.

Note that when $\partial_i F = 0$, the polynomial F is symmetric in variables x_i and x_{i+1} . Thus one can think of Schubert polynomials as partially symmetric. In particular, it is easy to see that for the *Grassmannian permutations*, defined as permutations w with des(w) = 1, we have \mathfrak{S}_w are symmetric polynomials. For example, when $u_{n,k} = (1, \ldots, n-k, n, n-k+1, \ldots, n-1)$, we have:

$$\mathfrak{S}_{u_{n,k}}(x_1, \dots, x_n) = e_k(x_1, \dots, x_n) = \sum_{1 \le i_1 < \dots < i_k \le n} x_{i_1} \cdots x_{i_k}$$

is the elementary symmetric polynomial.



FIGURE A.1. Graphs in RC(1432) and the corresponding Schubert polynomial $\mathfrak{S}_{1432} = x_1 x_2 x_3 + x_1^2 x_3 + x_1 x_2^2 + x_2^2 x_3 + x_1^2 x_2$ with monomials in this order.

A.3. **Pipe dreams.** For a permutation $w \in S_n$, denote by RC(w) the set of *RC-graphs* (also called *pipe dreams*), defined as tilings of a staircase shape with *crosses* and *elbows* as in the figure below, such that:

(i) curves start in row k on the left and end in column w(k) on top, for all $1 \le k \le n$, and (ii) no two curves intersect twice.

It follows from these conditions that every $H \in \mathrm{RC}(w)$ has exactly $\mathrm{inv}(w)$ crosses.

The Schubert polynomial $\mathfrak{S}_w \in \mathbb{N}[x_1, x_2, \ldots]$ is defined as

(A.1)
$$\mathfrak{S}_w(\boldsymbol{x}) := \sum_{H \in \mathrm{RC}(w)} \boldsymbol{x}^H \quad \text{where} \quad \boldsymbol{x}^H := \prod_{(i,j) : H(i,j) = \boxplus} x_i.$$

In other words, \boldsymbol{x}^{H} is the product of x_{i} 's over all crosses $(i, j) \in H$, see Figure A.1. As mentioned above, note that Schubert polynomials stabilize when fixed points are added at the end, e.g. $\mathfrak{S}_{1432} = \mathfrak{S}_{14325}$.

In this setting the combinatorial nature of Schubert polynomials is more transparent. Notably, one can show that for Grassmannian permutations $w \in S_{\infty}$, Schubert polynomial \mathfrak{S}_w coincides with the Schur function corresponding to the partition given by the *Rothe diagram*

$$\mathbf{R}(w) := \left\{ (w(j), i) : i < j, w(i) > w(j) \right\} \subset \mathbb{N}^2.$$

It follows from this definition that the coefficients $[\mathbf{x}^{\alpha}]\mathfrak{S}_w$ are nonnegative integers, and moreover that they are in $\#\mathsf{P}$ as a counting function.

A.4. Geometric definition. Let $G = GL_n(\mathbb{C})$ be the general linear group. Take $B \subset G$ to be the Borel subgroup of upper triangular matrices in G. Similarly define $B_- \subset G$ the opposite Borel subgroup of lower triangular matrices in G.

The complete flag variety is defined as $\mathcal{F}_n := \mathsf{G}/\mathsf{B}$. Under the left action of B_- , the variety \mathcal{F}_n has finitely many orbits X_w° , indexed by permutations $w \in S_n$. These are called *Schubert cells*.

The Schubert varieties X_w are the Zariski closures of the Schubert cells X_w° . The Schubert classes $\{\sigma_w\}_{w \in \mathcal{W}}$ are the Poincaré duals of Schubert varieties. These form a \mathbb{Z} -linear basis of the cohomology ring $H^*(\mathcal{F}_n)$. Borel's ring isomorphism [Bor53] maps

$$\Phi: H^*(\mathcal{F}_n) \longrightarrow \mathbb{Z}[x_1, x_2, \dots, x_n] / \langle e_i(x_1, x_2, \dots, x_n) : i \in [n] \rangle$$

where e_i are elementary symmetric polynomials. Schubert polynomials are polynomial representatives of Schubert classes: $\Phi(\sigma_w) = \mathfrak{S}_w$.

In this setting, the Schubert coefficients $c_{u,v}^w$ are defined as structure constants:

$$\sigma_u \smile \sigma_v \,=\, \sum_{w \in S_n} \, c^w_{u,v} \, \sigma_w \,.$$

By the *Kleiman transversality* [Kle74], coefficients $c_{u,v}^w$ count the number of points in the intersection of generically translated Schubert varieties:

(A.2)
$$c_{u,v}^w = \# \{ X_u(F_{\bullet}) \cap X_v(G_{\bullet}) \cap X_{w_{\circ}w}(E_{\bullet}) \},$$

where F_{\bullet} , G_{\bullet} and E_{\bullet} are generic flags. This definition implies the S_3 -symmetries of Schubert coefficients:

$$c_{u,v}^{w_{\circ}w} \,=\, c_{v,u}^{w_{\circ}w} \,=\, c_{u,w}^{w_{\circ}v} \,=\, c_{w,u}^{w_{\circ}v} \,=\, c_{v,w}^{w_{\circ}u} \,=\, c_{w,v}^{w_{\circ}u} \,.$$

Appendix B. Quotes and historical remarks

B.1. Formulation of the problem. The problem of finding a combinatorial interpretation of Schubert coefficients goes back to Lascoux and Schützenberger in 1980s, and was restated by numerous authors. As the area evolved, so did the language and the formulation of the problem. For example, in his celebrated survey, Stanley states the problem as follows:

"Find a combinatorial interpretation of the 'Schubert intersection coefficients' $c_{u,v}^w$, thereby combinatorially reproving that they are nonnegative." [Sta00, Problem 11]

In the introduction to his monograph, Manivel singles out the problem as the main mystery in the area:

"We note that Schubert polynomials are far from having revealed all of their secrets. We know almost nothing, for example, about their multiplication, and about a rule of Littlewood–Richardson type which must govern them." [Man01, p. 3]

Lenart motivates the problem by the geometry, and as an effort to avoid the geometry altogether (cf. [Ass23]). He also singles out the vanishing problem as a motivation:

"A famous open problem in algebraic combinatorics, known as the Schubert problem [...] is to find a combinatorial description of the Schubert structure constants (and, in particular, a proof of their nonnegativity which bypasses geometry). The importance of this problem stems from the geometric significance of the Schubert structure constants, and from the fact that a combinatorial interpretation for these coefficients would facilitate a deeper study of their properties (such as their symmetries, vanishing, etc.). The Schubert problem proved to be a very hard problem, resisting many attempts to be solved." [Len10]

Despite many remarkable developments, these sentiments continue to hold as underscored by Knutson, who used a starkly different language:

"We cannot emphasize strongly enough that the name of the game is to give manifestly nonnegative formulæ for the [Schubert coefficients]." [Knu22, §1.4]^a

^aOriginal emphasis.

Knutson then emphasizes the vanishing problem as the first motivation:⁶

"For applications (including real-world engineering applications) it is more important to know that some structure constant c is positive, than it is to know its actual value. This is much more easily studied with a noncancelative formula." (ibid.)

B.2. Substance of the problem. There is a great deal of uncertainty in algebraic combinatorics as to what exactly constitutes a "combinatorial interpretation". This is best illustrated by the following formulation of the [main problem] in the most recent monograph:

"Find an interpretation for the Schubert structure constants in terms of counting some sort of combinatorial objects such as paths in Bruhat order, Mondrian tableaux, labeled diagrams, permutation arrays, or n-dimensional chess games." [BGP25+, Open Problem 3.112]

⁶Two other motivations are computational efficiency and "possibility for categorification".

These combinatorial objects are all in #P, making the problem harder and more narrow than it already is (or simpler, since without any complexity assumptions *any number* is the number of *some* paths in Bruhat order). We maintain that #P as the only known robust notion of a "combinatorial interpretation", and refer the reader to [IP22, Pak24] for the explanation behind this reasoning. We only mention in passing that combinatorial objects in the problem above come from well-known attempts to resolve the problem. Curiously, the authors hedge themselves:

"Note, the Schubert structure constants already count the number of points in a certain type of generic 0-dimensional intersection [...] Perhaps one could call this a combinatorial interpretation, since they do count something! However, it is very difficult to test if flags are truly in generic position, even though presumably almost anything you could choose would suffice." [BGP25+, Remark 3.113]

This remark goes straight to the core of the issue and underscores the need for the formal approach. Fundamentally, this paper is an attempt to understand the computational hardness of counting these intersections.

B.3. Complexity of the problem. Prior to [PR24b, PR24c], the effort to analyze the hardness of the Schubert vanishing problem was largely unsuccessful:

"It is well known that solving Schubert problems are 'hard'. To our knowledge, no complete analysis of the algorithmic complexity is known." [BV08, p. 41]

In this quote, Billey and Vakil are fully cognizant that counting 0-dimensional intersections can be the basis of the algorithm, as they describe in the paper. After employing a mixture of theoretical analysis and experimental evidence, they conclude:

"Of course this allows one in theory to solve all Schubert problems, but the number and complexity of the equations conditions grows quickly to make this prohibitive for large n." [BV08, p. 24]

Nothing in this paper suggests that computing Schubert coefficients can be made efficient; we are nowhere close to practical applications. Note, however, our lifted formulation approach is different from that by Billey–Vakil's experimental effort:

"It is well known in that solving more equations with fewer variables is not necessarily an improvement. More experiments are required to characterize the 'best' method of computing Schubert problems. We are limited in experimenting with this solution technique to what a symbolic programming language like Maple can do in a reasonable period of time. The examples in the next section will illustrate how this technique is useful in keeping both the number of variables and the complexity of the rank equations to a minimum.' [BV08, p. 43]

In contrast, we are happy to increase the number of variables to ensure that resulting polynomials have a poly-size support, at which point Theorem 3.1 can be applied.