arXiv:2202.03995v2 [math.CO] 2 Dec 2022

CASTELNUOVO-MUMFORD REGULARITY OF LADDER DETERMINANTAL VARIETIES AND PATCHES OF GRASSMANNIAN SCHUBERT VARIETIES

JENNA RAJCHGOT, COLLEEN ROBICHAUX, AND ANNA WEIGANDT

ABSTRACT. We give degree formulas for Grothendieck polynomials indexed by vexillary permutations and 1432-avoiding permutations via tableau combinatorics. These formulas generalize a formula for degrees of symmetric Grothendieck polynomials which appeared in previous joint work of the authors with Y. Ren and A. St. Dizier.

We apply our formulas to compute Castelnuovo-Mumford regularity of classes of generalized determinantal ideals. In particular, we give combinatorial formulas for the regularities of all one-sided mixed ladder determinantal ideals. We also derive formulas for the regularities of certain Kazhdan-Lusztig ideals, including those coming from open patches of Schubert varieties in Grassmannians. This provides a correction to a conjecture of Kummini-Lakshmibai-Sastry-Seshadri (2015).

1. INTRODUCTION

Castelnuovo-Mumford regularity is a fundamental measure of the complexity of a graded module. In this paper, we use Schubert calculus techniques to provide explicit, easy-tocompute, combinatorial formulas for the Castelnuovo-Mumford regularity of classes of generalized determinantal ideals. The classes we treat include one-sided mixed ladder determinantal ideals and ideals defining patches of Schubert varieties in Grassmannians. These two classes of ideals are connected by work of N. Gonciulea and C. Miller [13].

Let k be a field. Let $S = k[x_1, \ldots, x_n]$ be a polynomial ring with the standard grading, deg $(x_i) = 1$, and let $I \subseteq S$ be a homogeneous ideal. When S/I is Cohen-Macaulay, as is the case throughout this paper, the regularity of S/I is known to satisfy

(1.1)
$$\operatorname{reg}(S/I) = \operatorname{deg} K(S/I;t) - \operatorname{ht}_S(I),$$

where K(S/I;t) is the *K*-polynomial of S/I and $ht_S(I)$ is the height of I in S. Using this fact, the authors, in joint work with Y. Ren and A. St. Dizier [32], gave a combinatorial formula which computes the regularity of coordinate rings of *Grassmannian matrix Schubert varieties*. The key technical ingredient was a formula of C. Lenart [26] regarding symmetric *Grothendieck polynomials*. In the present paper, we extend our work from [32].

Date: December 1, 2022.

Jenna Rajchgot was partially supported by NSERC Grant RGPIN-2017-05732.

Colleen Robichaux was supported by the NSF GRFP under Grant No. DGE 1746047 and NSF RTG Grant No. DMS 1937241.

Anna Weigandt was partially supported by Bill Fulton's Oscar Zariski Distinguished Professor Chair funds.

1.1. Summary of results. We give a combinatorial formula for degrees of Grothendieck polynomials indexed by 1432-avoiding permutations (see Theorem 1.3) and vexillary (2143-avoiding) permutations (see Theorem 1.7). These formulas in turn, allow us to compute the regularity of the corresponding matrix Schubert varieties. In particular, our formula in the vexillary setting provides a formula for Castelnuovo-Mumford regularity of one-sided mixed ladder determinantal varieties (see Section 7 for details). Our formulas naturally generalize the Grassmannian formula of [32] (see Section 2.5 for details).

Theorem 1.1. Given $w \in S_n$ so that w is 1432-avoiding,

$$\operatorname{reg}(S/I_w) = \sum_{k=1}^n \rho_d(\sigma_k(w))$$

Defined in Section 2, these $\sigma_k(w)$ are certain subsets of the *Rothe diagram* D(w) of w, and $\rho_d(\sigma_k(w))$ denotes the size of the largest diagonal in $\sigma_k(w)$.

Example 1.2. Let w = 1462375. In the images below, the elements of $\sigma_k(w)$ are shaded, with a maximal diagonal path in $\sigma_k(w)$ marked with \times for $k \in [5]$. For k = 6, 7, we have $\sigma_k(w) = \emptyset$, and so we omit the figures.



Theorem 1.1 computes $reg(S/I_w) = 3 + 1 + 0 + 1 + 1 = 6$.

Theorem 1.1 is a direct consequence of the following:

Theorem 1.3. If $w \in S_n$ is 1432-avoiding, then

$$\deg(\mathfrak{G}_w) = \#D(w) + \sum_{k=1}^n \rho_d(\sigma_k(w)).$$

For $w \in S_n$, #D(w) is the Coxeter length of w. See Section 2 for the definitions of Grothendieck polynomials \mathfrak{G}_w and Rothe diagrams D(w). The proof of Theorem 1.3 appears in Section 4.

Example 1.4. Returning to w as in Example 1.2, Theorem 1.3 with Theorem 1.1 give that

$$\deg(\mathfrak{G}_w) = \#D(w) + \operatorname{reg}(S/I_w) = 6 + (3+1+0+1+1) = 12.$$

 \diamond

We have similar diagrammatic regularity and degree formulas in the vexillary setting.

Theorem 1.5. Given $v \in S_n$ so that v is vexillary,

$$\operatorname{reg}(S/I_v) = \sum_{k=1}^n \rho_a(\tau_k(v)).$$

Here, the $\tau_k(v)$ are certain subsets of the Young diagram $\lambda(v)$ associated to v and $\rho_a(\tau_k(v))$ denotes the size of the largest antidiagonal in $\tau_k(v)$. See Section 2 for details.

Example 1.6. Let v = 169247358. In the diagrams below, the elements of $\tau_i(v)$ are shaded for i = 1, 2, 3 in $\lambda(v)$. In particular, a maximal size antidiagonal path contained in $\tau_i(v)$ has been marked with \times 's.



Applying Theorem 1.5, we have $reg(S/I_v) = 4 + 3 + 1 = 8$.

Theorem 1.5 is a direct consequence of the following:

Theorem 1.7. Suppose $v \in S_n$ is vexillary. Then

$$\deg(\mathfrak{G}_v) = \#D(v) + \sum_{i=1}^n \rho_a(\tau_k(v)).$$

The proof of Theorem 1.7 appears in Section 4.

Example 1.8. Returning to v as in Example 1.6, Theorem 1.7 with Theorem 1.5 give that

$$\deg(\mathfrak{G}_v) = \#D(v) + \operatorname{reg}(S/I_v) = 13 + (4+3+1) = 21.$$

We also provide formulas for the regularity of certain homogeneous Kazhdan-Lusztig ideals $J_{v,w}$. When v is a 321-avoiding permutation, we provide a formula in terms of *pipe dreams* (see Proposition 6.4). When v and w are both Grassmannian, we provide an easily computable formula by computing the degree of the corresponding K-polynomial in terms of a vexillary Grothendieck polynomial.

1.2. Connections to the literature. Concurrently with this work the third author, with O. Pechenik and D. Speyer [31], derived a combinatorial formula for the regularity of matrix Schubert varieties indexed by arbitrary permutations. In contrast with our diagrammatic combinatorics, the formula in [31] is phrased in terms of a new statistic on permutations. In work released around the same time as the present paper, E. Hafner [17] obtained a new proof of the vexillary case of [31] in terms of bumpless pipe dreams. Her results illustrate the connection from the formula in [31] to our vexillary formula through bumpless pipe dreams. A. Yong also has recent work related to the present paper, where he studies regularities of tangent cones of Schubert varieties [37].

For certain mixed ladder determinantal ideals, regularity formulas can be deduced through *a*-invariant formulas of S. Ghorpade and C. Krattenthaler [11]. The ladder determinantal ideals they consider have certain restrictions on their rank conditions. Consequently, their one-sided ideals are special cases of the ideals that we consider. See Section 7 for further discussion.

Our formula for regularity of patches of Grassmannian Schubert varieties (Theorem 6.5) provides a correction to a conjecture of M. Kummini, V. Lakshmibai, P. Sastry, and C. S. Seshadri (see [23, Conjecture 7.5]). This correction was conjectured in our previous paper (see [32, Conjecture 5.6]).

 \diamond

4

1.3. Outline of the paper. In Section 2, we introduce the necessary combinatorial background. In Section 3, we give tableau interpretations of our Grothendieck degree formulas. We provide proofs of our degree formulas for vexillary and 1432-avoiding permutations in Section 4. Section 5 describes the connection between Grothendieck polynomials and regularity and proves our main theorems. Section 6 applies these regularity formulas to correct the conjecture of [23]. Section 7 further applies our main theorems to give combinatorial formulas for the regularity of one-sided ladder determinantal ideals.

Acknowledgements. We would like to thank Philippe Nadeau and Alexander Yong for helpful comments and conversations. We would also like to thank Elisa Gorla for helpful communications about the literature. Finally, we would like to thank the anonymous referee for their helpful comments.

2. Combinatorial degree formulas

2.1. Grothendieck polynomials. We start by defining *Grothendieck polynomials*, introduced by A. Lascoux and M. P. Schützenberger [25] in their study of the K-theory of the complete flag variety. Let S_n denote the **symmetric group** on n letters, i.e., the set of bijections from $[n] := \{1, 2, ..., n\}$ to itself. We write permutations in one-line notation unless otherwise specified and define $w_i := w(i)$ for $i \in [n]$. The symmetric group S_n acts on $\mathbb{Z}[x_1, \ldots, x_n]$ by $w \cdot f(x_1, \ldots, x_n) = f(x_{w_1}, x_{w_2}, \ldots, x_{w_n})$. Let $s_i \in S_n$ be the simple transposition $(i \ i + 1)$, written here in cycle notation. For $f \in \mathbb{Z}[x_1, x_2, \ldots, x_n]$, define

$$\partial_i f = \frac{f - s_i f}{x_i - x_{i+1}}, \text{ and } \pi_i f = \partial_i (1 - x_{i+1}) f.$$

We recursively define Grothendieck polynomials as follows. Let $w_0 = n n - 1 \dots 1$ be the **longest permutation** in S_n . Define

$$\mathfrak{G}_{w_0}(\mathbf{x}) = \mathfrak{G}_{w_0}(x_1, x_2, \dots, x_n) = x_1^{n-1} x_2^{n-2} \cdots x_{n-1}$$

For $w \neq w_0$ there exists some $i \in [n-1]$ such that $w_i > w_{i+1}$. Then we define $\mathfrak{G}_{ws_i}(\mathbf{x}) = \pi_i(\mathfrak{G}_w(\mathbf{x}))$. Since the π_i satisfy the same braid and commutation relations as the simple transpositions, $\mathfrak{G}_w(\mathbf{x})$ is well defined.

Write $x \oplus y := x + y - xy$. We define the **double Grothendieck polynomials** using the same recurrence, starting from the initial condition

$$\mathfrak{G}_{w_0}(\mathbf{x};\mathbf{y}) = \prod_{1 < i+j \leq n} (x_i \oplus y_j).$$

Here, the ∂_i 's only act on the x_i 's, leaving the y_j 's fixed.

2.2. **Permutations.** First we recall some background on the symmetric group with [27] as a reference. The **permutation matrix** of w, which we also denote by w, is the 0, 1-matrix with 1's at (i, w_i) for all $i \in [n]$ and 0's elsewhere. To each permutation we associate a **rank function** defined by

$$r_w(i,j) = \#\{(k,w_k) : k \le i, w_k \le j\}$$

The **Rothe diagram** of $w \in S_n$ is the subset

$$D(w) = \{(i, j) \in [n] \times [n] : w_i > j \text{ and } w_j^{-1} > i\}.$$

Visually, D(w) is the set of cells remaining in the $n \times n$ grid after plotting the points (i, w_i) for each $i \in [n]$ and striking out any cells which appear weakly below or weakly to the right of these points, as shown in Example 2.1.

Let $\ell(w) := \#D(w)$ denote the **Coxeter length** of w. The **code** of w is the tuple $code(w) = (c_1, \ldots, c_n)$ where c_i records the number of cells in the *i*th row of D(w). Let

$$L(code(w)) := max\{i \in [n] : c_i > 0\}.$$

The essential set of w is the subset of D(w)

$$\mathsf{Ess}(w) = \{(i, j) \in D(w) : (i + 1, j), (i, j + 1) \notin D(w)\}$$

The dominant component Dom(w) is the connected component of D(w) containing (1,1).

Example 2.1. For $w = 72416835 \in S_8$, D(w) is the following:



Here, we have $\mathsf{Ess}(w) = \{(1,6), (3,1), (3,3), (6,3), (6,5)\}, \mathsf{code}(w) = (6,1,2,0,2,2,0,0), and \mathsf{Dom}(w) = \{(1,i) : i \in [6]\} \cup \{(2,1), (3,1)\}.$

A subset $D \subseteq [n] \times [n]$ is a **diagonal path** if

 $D = \{(i_1, j_1), \dots, (i_k, j_k) : i_1 < i_2 < \dots < i_k \text{ and } j_1 < j_2 < \dots < j_k\}.$

Given $S \subseteq [n] \times [n]$ write $\rho_d(S)$ for the size of the largest diagonal path in contained S.

A permutation $w \in S_n$ is 1432-avoiding if there does not exist a 1432 *pattern*, i.e., indices h < i < j < k such that w has the pattern $w_h < w_k < w_j < w_i$. For example, w = 23715846 is not 1432-avoiding; we underlined the positions of a 1432 pattern. For w 1432-avoiding, let

$$\sigma_k(w) = \{ (i, j) \in D(w) : i > k \text{ and } j > w_k \},\$$

i.e., $\sigma_k(w)$ is the set of cells in D(w) which are strictly southeast of (k, w_k) . Example 1.2 gives an example of diagonal paths in $\sigma_k(w)$.

A partition $\lambda = (\lambda_1, ..., \lambda_k)$ is a weakly decreasing sequence of non negative integers. We write $|\lambda| = \lambda_1 + \cdots + \lambda_k$. The **Young diagram** of a partition λ is the set $\{(i, j) \in \mathbb{Z}_{>0} \times \mathbb{Z}_{>0} : 1 \leq j \leq \lambda_i\}$. We often conflate Young diagrams with their partitions. Given partitions λ and μ , we write $\lambda \subseteq \mu$ to mean that the Young diagram of λ is contained in the Young diagram of μ .

Given $w \in S_n$, let $\mu(w)$ be the partition whose Young diagram is

$$\bigcup_{(i,j)\in D(w)} [1,i] \times [1,j],$$

i.e., $\mu(w)$ is the smallest partition whose Young diagram contains D(w).

A permutation $v \in S_n$ is **vexillary** if it does not contain a 2143 *pattern*, i.e., indices i < j < k < l such that $v_j < v_i < v_l < v_k$. For example, v = 72581364 is not vexillary





FIGURE 1. Let v = 169247358. Pictured on the left is the filling of the cells $(i, j) \in D(v)$ with $r_v(i, j)$. On the right is $\lambda(v)$ filled with F_v .

since the underlined indices form a 2143 pattern. A vexillary permutation v has shape $\lambda(v)$, where $\lambda(v)$ is $\mathsf{code}(v) = (c_1, \ldots, c_n)$ sorted into decreasing order. A vexillary permutation v has flag

$$\phi(v) = (\phi_1 \leqslant \phi_2 \leqslant \cdots \leqslant \phi_m)$$
, where

 $\phi_i = \max\{j : (j,k) \in \mu(v) \text{ lies in the same diagonal as } (i,\lambda_i(v))\}.$

Note that we can think of $\lambda(v)$ as the partition with the property that each of the diagonals of its Young diagram has the same number of cells as the corresponding diagonal of D(v). Observe that $\phi(v)$ tells us how the positions of these boxes changed between D(v) to $\lambda(v)$.

Fill the diagonals of $\lambda(v)$ with $r_v(i, j)$ for the corresponding cells $(i, j) \in D(v)$, so that the entries are (weakly) increasing along diagonals. Write F_v for this filling (see Figure 1 for an example). Let

$$\tau_k(v) = \{ (i,j) \in \lambda(v) : F_v(i,j) \ge k \}.$$

A subset $A \subseteq [n] \times [n]$ is an **antidiagonal path** if

 $A = \{(i_1, j_1), \dots, (i_k, j_k) : i_1 < i_2 < \dots < i_k \text{ and } j_1 > j_2 > \dots > j_k\}.$

Given $S \subseteq [n] \times [n]$ write $\rho_a(S)$ for the largest antidiagonal path in S. See Example 1.6 for an example of antidiagonal paths in $\tau_k(v)$.

A permutation $g \in S_n$ is **Grassmannian** if it has a unique descent, i.e. a unique $k \in [n-1]$ such that $g_k > g_{k+1}$.

2.3. Pipe complexes. Let $a = (a_1, \ldots, a_k)$ be a word on the alphabet [n-1]. We say a is a reduced word for w if $w = s_{a_1} \cdots s_{a_k}$ and $\ell(w) = k$.

Define an algebra over \mathbb{Z} with generators $\{e_w : w \in S_n\}$ and multiplication given by

$$e_w e_{s_i} = \begin{cases} e_{ws_i} & \text{if } \ell(ws_i) > \ell(w) \\ e_w & \text{if } \ell(ws_i) < \ell(w) \end{cases}$$

The **Demazure product** $\delta(a)$ of a word $a = (a_1, \ldots, a_k)$ is defined by computing

$$e_{s_{a_1}}\cdots e_{s_{a_k}}=e_{\delta(a)}.$$

Label cells in D(v) along rows so that the first cell in row *i* is labeled *i*, the next i + 1, and so on. Given $P \subseteq D(v)$ let a_P be the word obtained by reading the labels of the elements in P within rows from right to left, starting at the top row and working downwards. Let

 $Pipes(v, w) = \{P \subseteq D(v) : a_P \text{ is a reduced word for } w\}.$

Likewise, let

$$\overline{\mathsf{Pipes}}(v,w) = \{P \subseteq D(v) : \delta(a_P) = w\}$$

For any $P \subseteq [n] \times [n]$, we assign it the **t**-weight

$$\operatorname{wt}_{\mathbf{t}}(P) = \prod_{(i,j)\in P} t_{ij}.$$

Pictorially, we represent $P \subseteq D(v)$ by marking $(i, j) \in D(v)$ with a + whenever $(i, j) \in P$. We define the **unspecialized Grothendieck polynomial** to be

(2.1)
$$\mathfrak{G}_{v,w}(\mathbf{t}) = \sum_{P \in \overline{\mathsf{Pipes}}(v,w)} (-1)^{\#P-\ell(w)} \mathtt{wt}_{\mathbf{t}}(P).$$

Note that by setting $v = w_0$, we can recover double Grothendieck polynomials by specializing the variables in $\mathfrak{G}_{w_0,w}(\mathbf{t})$:

 $\mathfrak{G}_w(\mathbf{x};\mathbf{y}) = \mathfrak{G}_{w_0,w}(x_1 \oplus y_1, x_2 \oplus y_2, \dots, x_n \oplus y_n).$

2.4. Excited Young Diagrams. Fix partitions $\lambda \subseteq \mu$. Let

$$D_{top}(\mu, \lambda) = \{(i, j) : i \in [k] \text{ and } j \in [\lambda_i]\}.$$

We call $D \subseteq \mu$ a **diagram**, represented graphically by marking these cells in D with +'s. An **excited move** is a mutation of a local 2×2 subsquare of the form

$$(2.2) \qquad \qquad + \qquad \mapsto \qquad + \cdot$$

Here, the mutated subsquare must be entirely contained within μ .

We write $\mathsf{ExcitedYD}(\mu, \lambda)$ for the set of $D \subseteq \mu$ which can be obtained by a sequence of excited moves starting from $D_{top}(\mu, \lambda)$. Such diagrams are called **excited Young diagrams**. There is a unique element of $\mathsf{ExcitedYD}(\mu, \lambda)$ to which no excited moves may be applied, see e.g., [34, Lemma 7.4]. Call this $D_{bot}(\mu, \lambda)$.

We also consider K-theoretic excited moves of the form

$$(2.3) \qquad \qquad + \qquad \mapsto \qquad + \qquad + \qquad ,$$

again, where all cells pictured are contained in μ . Write $\mathsf{KExcitedYD}(\mu, \lambda)$ for the set of diagrams which can be obtained from $D_{\mathsf{top}}(\mu, \lambda)$ by a sequence of excited and K-theoretic excited moves. We weight $D \in \mathsf{KExcitedYD}(\mu, \lambda)$ by

$$\operatorname{wt}(D) = \prod_{(i,j)\in D} (x_i \oplus y_j).$$

Proposition 2.2. If $v \in S_n$ is vexillary, then

$$\mathfrak{G}_{v}(\mathbf{x};\mathbf{y}) = \sum_{D \in \mathsf{KExcitedYD}(\mu(v),\lambda(v))} (-1)^{\#D - |\lambda(v)|} \operatorname{wt}(D).$$

Proof. This follows by noting the flagged set-valued tableaux (and diagonal pipe dreams) of [22] can be identified with $\mathsf{KExcitedYD}(\mu(v), \lambda(v))$. See e.g., [16] and [34] for further details.

Lemma 2.3. Fix partitions $\lambda \subseteq \mu$. There exists a unique permutation $v \in S_{\infty}$ so that $D(v) = D_{bot}(\mu, \lambda)$. In particular, v is vexillary.

Proof. That $D_{bot}(\mu, \lambda) \in \mathsf{ExcitedYD}(\mu, \lambda)$ is the diagram of a vexillary permutation follows from [34, Proposition 7.6].

Example 2.4. Let $\lambda = (5, 4, 2, 1, 0)$ and $\mu = (6, 6, 4, 4, 4)$. Then v = 5713624 is the unique vexillary permutation so that $D(v) = D_{bot}(\mu, \lambda)$.

Above are $D_{top}(\mu, \lambda)$, $D_{bot}(\mu, \lambda)$, and D(v), respectively.

Theorem 2.5. Fix Grassmannian permutations g and u with descent at position k so that $\lambda(g) \subseteq \lambda(u)$. Let v be the vexillary permutation such that $D(v) = D_{bot}(\lambda(u), \lambda(g))$. Then

$$\deg(\mathfrak{G}_{u,g}(\mathbf{t})) = \deg(\mathfrak{G}_v(\mathbf{x})).$$

Proof. Write $\operatorname{code}(u) = (c_1, \ldots, c_n)$. Since u is Grassmannian with descent at position k, $\lambda(u) = (c_k, c_{k-1}, \ldots, c_1)$ (see [27, Section 2.2]). In particular, this means we can identify each cell in D(u) with cells in $D_{\lambda(u)} = \{(i, j) : 1 \leq j \leq c_i\}$ by left justifying cells in D(u) within rows.

Under this identification, we map each element of $\overline{\mathsf{Pipes}}(u,g)$ to a subset of $D_{\lambda(u)}$. Call this set of diagrams L. It is immediate that $L \subseteq \overline{\mathsf{Pipes}}(w_0,g)$. In particular, this implies elements of L are connected by (flipped) K-theoretic moves, i.e., replacements of the form:

$$+$$
 \mapsto $+$

and

$$+$$
 \mapsto $+$

By flipping the first k rows vertically, we see that elements of L are in bijection with elements of KExcitedYD($\lambda(u), \lambda(g)$). Thus, we have a (degree preserving) bijection between elements of $\overline{\mathsf{Pipes}}(u, g)$ and KExcitedYD($\lambda(u), \lambda(g)$).

Then by Equation (2.1) and Proposition 2.2, we conclude $\deg(\mathfrak{G}_{u,g}(\mathbf{t})) = \deg(\mathfrak{G}_v(\mathbf{x}))$. \Box

8

+	++	+]	++	++	++	+		+]•
			-					-				+
				+						ЦШ	l 🕇	+
				+		+				•		

 \diamond

Example 2.6. Let g = 1247356 and u = 1457236. An element of Pipes(u, g) and its corresponding K-theoretic excited Young diagram are pictured below.



2.5. Connections to the Grassmannian degree formula. In previous work with Ren and St. Dizier [32], the authors presented a formula to compute the degree of symmetric Grothendieck polynomials. If $u \in S_n$ is Grassmannian with descent k, then the symmetric Grothendieck polynomial is $\mathfrak{G}_{u(\lambda)}(x_1, \ldots, x_k) := \mathfrak{G}_u(x_1, \ldots, x_n)$. Since Grassmannian permutations are both 1432-avoiding and vexillary, our new degree formulas are two different generalizations of this formula. We illustrate these connections here.

Write $\delta^{(k)} = (k, k - 1, ..., 1)$. Let $\mathsf{sv}(\lambda) = \max\{k : \delta^{(k)} \subseteq \lambda\}$. Given a partition $\lambda = (\lambda_1, \ldots, \lambda_k)$, let $\mathsf{trunc}^{(i)}(\lambda)$ be the partition obtained by removing the first *i* columns of the Young diagram of λ . Then:

Theorem 2.7 ([32]). If $\lambda = (\lambda_1, \ldots, \lambda_k)$, then

$$\deg(\mathfrak{G}_{\lambda}(x_1,\ldots,x_k)) = |\lambda| + \sum_{i=1}^k \mathsf{sv}(\mathsf{trunc}^{(\lambda_i)}(\lambda)).$$

Theorem 2.7 can be recovered using Theorem 1.3 or Theorem 1.7. We illustrate this in the example below.

Example 2.8. Let $\lambda = (3, 2, 2, 0)$ and k = 4. The Grassmannian permutation associated to the pair (λ, k) is w = 1457236. The first line below computes the formula in Theorem 2.7 where the *i*th Young diagram has $\mathsf{trunc}^{(\lambda_i)}(\lambda)$ shaded, with $\delta^{(k)}$ marked with \times 's for $k = \mathsf{sv}(\mathsf{trunc}^{(\lambda_i)}(\lambda))$.



Below, we demonstrate the rule given in Theorem 1.3. Here, we have $\sigma_k(w)$ shaded, with the longest diagonal marked with \times 's.



 \diamond

Now, we use the formula from Theorem 1.7. In each Young diagram, we have shaded $\tau_k(w)$, with the longest antidiagonals marked with \times 's.



Thus we see all three formulas compute $\deg(\mathfrak{G}_{\lambda}(x_1,\ldots,x_k)) = |\lambda| + 3 + 1 + 1 = 12.$

3. TABLEAU FORMULAS FOR GROTHENDIECK POLYNOMIALS

Since their introduction, Grothendieck polynomials have been studied with a number of combinatorial formulas ([9, 26, 4]). For our degree formulas, we will focus on two tableau formulas in the special cases of 1432-avoiding permutations and vexillary permutations. Furthermore, in each of these cases, we construct a tableau whose weight contributes to the top degree terms of the corresponding Grothendieck polynomial.

3.1. Set-valued Rothe tableaux. A set-valued Rothe tableau T of shape D(w) is a filling of D(w) with nonempty subsets of $\mathbb{Z}_{>0}$ such that for boxes $a, b \in D(w)$:

- if a lies north of b in the same column, then $\max T(a) < \min T(b)$, and
- if a lies west of b in the same row, then $\min T(a) \ge \max T(b)$,

where T(a) denotes the set of entries of T in box a. Let $\mathsf{SVT}(D(w))$ be the collection of such tableaux. We say a tableau $T \in \mathsf{SVT}(D(w))$ is flagged by $\phi = (\phi_1, \phi_2, \ldots, \phi_n)$ if for each box b in row i of D(w), max $T(b) \leq \phi_i$ for all i. For a 1432-avoiding $w \in S_n$, let

 $\mathsf{FSVD}(w) = \{T \in \mathsf{SVT}(D(w)) : T \text{ is flagged by } (1, 2, \dots, n)\}.$

Example 3.1. Below is some $T \in \mathsf{FSVD}(w)$ for w = 1462375.



 \diamond

Theorem 3.2. [8, Theorem 1.1] For $w \in S_n$ 1432-avoiding, \mathfrak{G}_w has the following expansion:

(3.1)
$$\mathfrak{G}_w(\mathbf{x}, \mathbf{y}) = \sum_{T \in \mathsf{FSVD}(w)} (-1)^{\#T - \#D(w)} \prod_{e \in T} x_{\operatorname{val}(e)} \oplus y_{\lambda_{r(e)} + \phi_{r(e)} - c(e) - \operatorname{val}(e) + 1}.$$

where the product is over entries e in T whose value is val(e) and c(e), r(e) are the column and row indices of e. For $T \in \mathsf{FSVD}(w)$, let #T denote the number of entries in T. We say $T \in \mathsf{FSVD}(w)$ is **maximal** if $T' \in \mathsf{FSVD}(w)$ implies $\#T' \leq \#T$. Now we give a construction of $T_w \in \mathsf{FSVD}(w)$ for a given 1432-avoiding w. Theorem 1.3 proves T_w is maximal. Let $\mathsf{md}(D)$ denote the northmost then we stmost maximal diagonal path of $D \subseteq [n]^2$. For $\mathsf{md}(\sigma_k(w)) \neq \emptyset$, let

 $\mathsf{NE}(\mathsf{md}(\sigma_k(w))) = \{(i,j) \in D(w) - \mathsf{md}(\sigma_k(w)) : (i,j) \text{ lies northeast of } \mathsf{md}(\sigma_k(w))\}.$

Take $T_0 \in \mathsf{SVT}(D(w))$ such that $T_0(i, j) = i$ for $i \in [L(\mathsf{code}(w))]$. For $k \in [L(\mathsf{code}(w)) - 1]$, let $T_k \in \mathsf{SVT}(D(w))$ such that for $(i, j) \in D(w)$:

$$T_{k}(i,j) := \begin{cases} T_{k-1}(i,j) \cup \{\min T_{k-1}(i,j) - 1\} & \text{if } (i,j) \in \mathsf{md}(\sigma_{k}(w)), \\ T_{k-1}(i,j) - 1 & \text{if } (i,j) \in \mathsf{NE}(\mathsf{md}(\sigma_{k}(w))), \\ T_{k-1}(i,j) & \text{otherwise}, \end{cases}$$

where T(i, j) - 1 is entrywise subtraction. Let $T_w := T_{L(\mathsf{code}(w))}$.

Example 3.3. Below we construct T_w for w = 1462375.



Lemma 3.4. Suppose w in S_n is 1432-avoiding. Then $T_w \in \mathsf{FSVD}(w)$.

Proof. We proceed by showing $T_k \in \mathsf{FSVD}(w)$ for $k \in [L(\mathsf{code}(w))]$ by induction on k. By construction, $T_0 \in \mathsf{FSVD}(w)$. Suppose $T_{k-1} \in \mathsf{FSVD}(w)$ for some $k \in [L(\mathsf{code}(w))]$. If $\mathsf{md}(\sigma_k(w)) = \emptyset$, the result follows the inductive assumption since $T_k = T_{k-1}$.

Otherwise, since $T_{k-1} \in \mathsf{FSVD}(w)$ by construction of T_k ,

$$\max T_k(i,j) \leqslant \max T_{k-1}(i,j) \leqslant i.$$

Similarly since T_{k-1} is decreasing along rows, T_k is clearly decreasing along rows. By definition of T_k , any (i, j) can be decremented no more than i - 1 times, so no entry can be decremented to 0. Thus it remains to show T_k increases down columns. Consider some $(i, j) \in \operatorname{md}(\sigma_k(w))$. Let

$$i' = \max\{x < i : (x, j) \in D(w)\}.$$

 \diamond



FIGURE 2. Let v be as in Figure 2.2. Then $\lambda(v) = (6, 4, 2, 1)$ and $\mu(v) = (8, 8, 8, 5, 5, 5)$. Pictured on the left is $\lambda(v) \subset \mu(v)$ with the diagonals used to compute $\phi(v) = (3, 3, 6, 6)$ drawn in red. To the right is an element of $\mathsf{FSVT}(v)$.

Since T_{k-1} is increasing down columns, it suffices to show that $\max T_k(i', j) < \min T_k(i, j)$. If $(i', j) \in \mathsf{NE}(\mathsf{md}(\sigma_k(w)))$ or does not exist, the result follows by the construction of T_k .

Otherwise, by the definition of $\operatorname{md}(\sigma_k(w))$, it follows that i = i' + h for some h > 1. Then for $s \in [i']$, by the definitions of T_s and $\operatorname{md}(\sigma_s(w))$,

(3.2)
$$\max T_s(i',j) + h \leq \min T_s(i,j).$$

Thus if $k \leq i'$, we are done. If k > i', it follows that $\max T_k(i', j) = \max T_{i'}(i', j)$ and $\min T_k(i, j) \geq \min T_{i'}(i, j) - h + 1$, so by Equation (3.2),

$$\max T_k(i', j) \le \min T_k(i, j) - 1.$$

Thus $T_k \in \mathsf{FSVD}(w)$.

3.2. Set-valued Young tableaux. A set-valued tableau *T* of shape $\lambda = (\lambda_1, \lambda_2, ..., \lambda_n)$ is a filling of λ with nonempty subsets of $\mathbb{Z}_{>0}$ such that for boxes $(a, b) \in \lambda$:

- if a lies north of b, then $\max T(a) < \min T(b)$, and
- if a lies west of b, then $\max T(a) \leq \min T(b)$,

where T(a) denotes the set of entries of T in box a. Let $\mathsf{SVT}(\lambda)$ be the collection of such tableaux. We say a tableau $T \in \mathsf{SVT}(\lambda)$ is flagged by $\phi = (\phi_1, \phi_2, \ldots, \phi_n)$ if for each box b in row i of λ , we have $\max T(b) \leq \phi_i$. For a vexillary permutation v, let

$$\mathsf{FSVT}(v) = \{T \in \mathsf{SVT}(\lambda(v)) : T \text{ is flagged by } \phi(v)\}.$$

An example of some $T \in \mathsf{FSVT}(169247358)$ is given in Figure 2. We note that many different choices of flagging can result in the same underlying set of tableaux. See [28, Remark 3.10] for further commentary.

Theorem 3.5. [22, Theorem 5.8] If $v \in S_n$ is vexillary, the double Grothendieck polynomial $\mathfrak{G}_v(\mathbf{x}; \mathbf{y})$ has the following expansion:

(3.3)
$$\mathfrak{G}_{v}(\mathbf{x};\mathbf{y}) = \sum_{T \in \mathsf{FSVT}(v)} (-1)^{\#T-|\lambda|} \prod_{e \in T} x_{\operatorname{val}(e)} \oplus y_{\operatorname{val}(e)+c(e)-r(e)},$$

where the product is over entries in T whose value is val(e) and c(e), r(e) are the column and row indices of e.

$$\square$$

For $T \in \mathsf{FSVT}(v)$ let #T denote the number of entries in T. We say $T \in \mathsf{FSVT}(v)$ is **maximal** if $\#T = \max\{\#U : U \in \mathsf{FSVT}(v)\}$. Now we give a construction of $U_v \in \mathsf{SVT}(v)$ for a given vexillary v. Theorem 1.7 proves U_v is maximal.

Let $\operatorname{ma}(\lambda)$ denote the northmost then we stmost maximal antidiagonal path of $\bigcup_i \mu_i \subseteq \lambda$. For $\operatorname{ma}(\tau_k(v)) \neq \emptyset$, let

$$\mathsf{SE}(\mathsf{ma}(\tau_k(v))) = \{(i,j) \in \lambda - \mathsf{ma}(\tau_k(v)) : (i,j) \text{ lies southeast of } \mathsf{ma}(\tau_k(v))\}$$

Take $U_0 \in \mathsf{SVT}(\lambda)$ such that $U_0(i, j) = i$ for $i \in [\ell(\lambda)]$. For $k \in [\ell(\lambda) - 1]$, let $U_k \in \mathsf{SVT}(\lambda)$ such that for $(i, j) \in \lambda$:

$$U_{k}(i,j) := \begin{cases} U_{k-1}(i,j) \cup \{\max U_{k-1}(i,j)+1\} & \text{if } (i,j) \in \operatorname{ma}(\tau_{k}(v)), \\ U_{k-1}(i,j)+1 & \text{if } (i,j) \in \operatorname{SE}(\operatorname{ma}(\tau_{k}(v))), \\ U_{k-1}(i,j) & \text{otherwise}, \end{cases}$$

where U(i, j) + 1 is entrywise addition. Let $U_v := U_{\ell(\lambda)}$.

Remark 3.6. By a similar argument to Lemma 3.4, it follows that $U_v \in \mathsf{FSVT}(v)$. By Theorem 1.7, it follows that U_v is maximal.

Example 3.7. Let v = 169247358. From Figure 2.2, we saw $\lambda(v) = (6, 4, 2, 1)$. Furthermore, $\phi(v) = (3, 3, 6, 6)$. Below is the construction of U_v from U_0 .



4. PROOFS OF DEGREE FORMULAS

In this section, we prove our Grothendieck degree formulas for 1432-avoiding permutations and vexillary permutations to deduce our main theorems.

4.1. **Proof of Theorem 1.3.** Recall, md(D) is the northmost then we that diagonal path of $D \subseteq [n]^2$ and

$$\sigma_k(w) = \{(i, j) \in D(w) : i > k, j > w(k)\}.$$

For brevity, define $f_d(w) = \#D(w) + \sum_{k=1}^n \rho_d(\sigma_k(w))$. We start by recalling a lemma from [8].

Lemma 4.1 ([8, Lemma 2.4]). Let $w \neq w_0$ be a 1432-avoiding permutation. If r is the first ascent of w, then ws_r is also 1432-avoiding.

Proposition 4.2. If w is 1432-avoiding, there exists $T \in \mathsf{FSVD}(w)$ such that

$$\#T = f_d(w).$$

In particular, $\deg(\mathfrak{G}_w) \ge f_d(w)$.

Proof. This follows by Lemma 3.4 since $\#T_w = f_d(w)$ by construction.

Lemma 4.3. Let $w \neq w_0$ be a 1432-avoiding permutation and suppose r is the first ascent of w. If there is a maximal diagonal path in $\sigma_r(w)$ which has no cells in row r+1, then there exists a maximal set-valued Rothe tableau for w such that the entries in row r+1 restricted to $\sigma_r(w)$ are all strictly less than r+1.

Proof. Suppose w is such that there is a maximal diagonal path in $\sigma_r(w)$ which has no cells in row r + 1. Consider maximal $T \in \mathsf{FSVD}(w)$ such that T has boxes containing r + 1 in row r + 1 restricted to $\sigma_r(w)$. We will construct $T' \in \mathsf{FSVD}(w)$ such that #T' = #T and such that the entries in row r + 1 restricted to $\sigma_r(w)$ are all strictly less than r + 1.

Let \mathbf{b}_1 denote the box containing the eastmost occurrence of r + 1 in row r + 1 in T. For $1 < i \leq \# \mathrm{md}(\sigma_r(w))$, we define $\mathbf{b}_i \in \sigma_r(w)$ as the box containing the northmost, then eastmost occurrence of r + i in T, in the region strictly east of \mathbf{b}_{i-1} . Thus $\{\mathbf{b}_i\}_{i \in [\sigma_r(w)]}$ forms a diagonal path.

Let \mathbf{c}_i denote the northmost box of $\sigma_r(w)$ lying directly south of \mathbf{b}_i for each $i \in [\# \mathrm{md}(\sigma_r(w))]$. By the assumption that there is a maximal diagonal path in $\sigma_r(w)$ which has no cells in row r + 1 and the definition of \mathbf{b}_i , $\{\mathbf{c}_i\}_{i \in [k]}$ exists for some $1 \leq k \leq \# \mathrm{md}(\sigma_r(w))$. Let P be constructed as follows:

 $P = {\mathbf{b}_1} \cup {\mathbf{b}_i : \mathbf{b}_{i-1} \in P \text{ and } \mathbf{c}_{i-1} \text{ lies in the same row as } \mathbf{b}_i}.$

Let $P' = {c_i : b_i \in P}$. By maximality of T, it follows that ${r + i, r + i - 1} \subseteq T(b_i)$ for each $i \in [\#P]$. Take T' such that

$$T'(x,y) := \begin{cases} T(\mathbf{b}_i) \setminus \{r+i\} & \text{if } (x,y) = \mathbf{b}_i, \\ T(\mathbf{c}_i) \cup \{r+i\} & \text{if } (x,y) = \mathbf{c}_i, \\ T(x,y) - 1 & \text{if } (x,y) \text{ lies directly between } \mathbf{c}_i \text{ and } \mathbf{b}_{i+1}, \\ T(x,y) - 1 & \text{if } (x,y) = (r+1,y) \in \sigma_r(w), \text{ lying west of } \mathbf{b}_1, \\ T(x,y) & \text{otherwise.} \end{cases}$$

It is straightforward to check $T' \in \mathsf{FSVD}(w)$. Since #T = #T', T' is of the desired form. \Box

Proposition 4.4. Suppose $w \in S_n$ is 1432-avoiding. Let r denote the position of the first ascent of w and $\{c_m < \cdots < c_0\} = \{w_r \leq i \leq w_{r+1} : (r+1,i) \in D(w)\}$. Then

$$D(w \cdot s_r) = (D(w) - \{(r+1, c_i) : 0 \le i \le m\}) \cup \{(r, c_i) : 0 \le i \le m\} \cup \{(r, w_r)\}.$$

Proof. This follows by the definition of D(w), since r is the first ascent of w.

Lemma 4.5. Let $w \neq w_0$ be a 1432-avoiding permutation, and suppose r is the first ascent of w. If there is a maximal diagonal path in $\sigma_r(w)$ which has no cells in row r + 1, then

$$f_d(w) + 1 = f_d(ws_r).$$

Otherwise,

$$f_d(w) = f_d(ws_r).$$

Proof. By Proposition 4.4, $\#D(ws_r) = \#D(w) + 1$. Further, since r was the first ascent of $w, (r, w_r) \in \mathsf{Dom}(ws_r)$. Further we see

$$\rho_d(\sigma_k(w)) = \rho_d(\sigma_k(ws_r))$$
 for $k \neq r, r+1$

by Proposition 4.4. By definition of r,

$$\rho_d(\sigma_{r+1}(w)) = \rho_d(\sigma_r(ws_r)).$$

Finally, by Proposition 4.4, $\sigma_{r+1}(ws_r) = \sigma_r(w) - \bigcup_{i=0}^{\ell} (r, c_i)$. Thus $\rho_d(\sigma_r(w)) = \rho_d(\sigma_{r+1}(ws_r))$ if there is a maximal diagonal path in $\sigma_r(w)$ which has no cells in row r+1. Otherwise, $\rho_d(\sigma_r(w)) = \rho_d(\sigma_{r+1}(ws_r)) + 1$, so the result follows by the definition of f_d .

Proof of Theorem 1.3. We proceed by induction on $\ell(w_0) - \ell(w)$. In the base case, $w = w_0$ and the formula is immediate since $\deg(w_0) = \ell(w_0) = \#D(w_0) = f_d(w_0)$.

Now pick $w \in S_n$ so that $w \neq w_0$. Assume the formula holds for all $w' \in S_n$ so that $\ell(w') > \ell(w)$. Let r be the first ascent of w. Let R denote the set of boxes in $\sigma_r(w)$ lying in row r+1. By Proposition 4.4, one obtains $D(ws_r)$ from D(w) by shifting all cells in R up one row and then placing a new cell in position (r, w_r) .

Consider $T \in \mathsf{FSVD}(w)$. We will construct $T' \in \mathsf{FSVD}(ws_r)$ from T by the following:

$$T'(x-1,y) := \begin{cases} r & \text{if } (x-1,y) = (r,w_r), \\ T(x,y) - \{r+1\} \cup \{r\} & \text{if } (x,y) \in R, \ r+1 \in T(x,y), \text{ and } r \notin T(x,y), \\ T(x,y) - \{r+1\} & \text{if } (x,y) \in R, \ r+1 \in T(x,y), \text{ and } r \in T(x,y), \\ T(x,y) & \text{if } (x,y) \in R, \ r+1 \notin T(x,y), \\ T(x-1,y) & \text{otherwise.} \end{cases}$$

Thus $T' \in \mathsf{FSVD}(ws_r)$ and $\#T' \ge \#T$, giving $\deg(\mathfrak{G}_w) \le \deg(\mathfrak{G}_{ws_r})$. We have two cases to check.

Case 1: Suppose all maximal diagonal paths in $\sigma_r(w)$ have a cell in row r+1. We have

$$f_d(w) \leq \deg(\mathfrak{G}_w) \qquad \text{(by Proposition 4.2)}$$
$$\leq \deg(\mathfrak{G}_{ws_r})$$
$$= f_d(ws_r) \qquad \text{(by inductive hypothesis).}$$

By Lemma 4.5, $f_d(w) = f_d(ws_r)$. Thus, $f_d(w) = \deg(\mathfrak{G}_w)$.

Case 2: Suppose there exists a maximal diagonal path in $\sigma_r(w)$ which has no cells in row r+1. By Lemma 4.3, there exists a maximal tableau T for w so that boxes in R have entries less than r + 1. Using the above construction for $T' \in \mathsf{FSVD}(ws_r)$, it follows that #T' = #T + 1. As a consequence, $\deg(\mathfrak{G}_w) < \deg(\mathfrak{G}_{ws_r})$.

Thus,

$$f_d(w) \leq \deg(\mathfrak{G}_w) \qquad \text{(by Proposition 4.2)}$$
$$< \deg(\mathfrak{G}_{ws_r})$$
$$= f_d(ws_r) \qquad \text{(by inductive hypothesis)}$$
$$= f_d(w) + 1 \qquad \text{(by Lemma 4.5).}$$

Thus $f_d(w) = \deg(\mathfrak{G}_w)$.

4.2. Proof of Theorem 1.7. If v is vexillary, we associate to v the following statistic:

(4.1)
$$f_a(v) = \#D(v) + \sum_{i=1}^n \rho_a(\tau_i(v))$$

Note that by definition, $\#D(v) = |\lambda(v)|$. The goal of this section is to prove Theorem 1.7, i.e., to show if v is vexillary, then $\deg(\mathfrak{G}_v) = f_a(v)$. We start with some lemmas.

We follow [22] for combinatorial background. The **maximal corner** (r, s) of w is the position of the right most cell in the last row of D(w). Let $t_{i,j}$ denote the transposition (i j). Define $w_P := wt_{r,w^{-1}(s)}$. Then w_P is the unique permutation such that

(4.2)
$$D(w_P) = D(w) - \{(r, s)\}.$$

Grothendieck polynomials satisfy a recurrence known as **transition**. Recall $t_{i,j}$ denotes the transposition (i, j). Let $w_P := wt_{r,w^{-1}(s)}$. Let $i_1 < i_2 < \cdots < i_k$ be the list of those indices i < r for which $\ell(w_P) + 1 = \ell(w_P t_{i,r})$.

Theorem 4.6 ([24]). Given $w \in S_n$, with maximal corner (r, s) and $t_{i_i,r}$'s as above,

$$\mathfrak{G}_w = \mathfrak{G}_{w_P} + (x_r - 1)(\mathfrak{G}_{w_P} \star (1 - t_{i_1,r})(1 - t_{i_2,r}) \cdots (1 - t_{i_k,r})),$$

where $\mathfrak{G}_v \star u := \mathfrak{G}_{vu}$.

When v is vexillary, there is at most one index i < r for which $\ell(v_P) + 1 = \ell(v_P t_{i,r})$. When such an index exists, we define $v_C = v_P t_{i,r}$. In this case, Theorem 4.6 specializes to

(4.3)

$$\mathfrak{G}_{v} = \mathfrak{G}_{v_{P}} + (x_{r} - 1)(\mathfrak{G}_{v_{P}} \star (1 - t_{i,r}))$$

$$= \mathfrak{G}_{v_{P}} + (x_{r} - 1)(\mathfrak{G}_{v_{P}} - \mathfrak{G}_{v_{C}})$$

$$= x_{r}\mathfrak{G}_{v_{P}} + (1 - x_{r})\mathfrak{G}_{v_{C}}.$$

If no such index exists, then necessarily $(r, s) \in \mathsf{Dom}(v)$ and we have

(4.4)
$$\mathfrak{G}_{v} = \mathfrak{G}_{v_{P}} + (x_{r} - 1)(\mathfrak{G}_{v_{P}})$$
$$= x_{r}\mathfrak{G}_{v_{P}}.$$

Lemma 4.7. Fix any permutation w and suppose the maximal corner $(r, s) \in \text{Dom}(w)$. Then $\deg(\mathfrak{G}_w) = \deg(\mathfrak{G}_{w_P}) + 1$.

Proof. This is an immediate consequence of Equation (4.4) since multiplying any nonzero polynomial by x_r increases the degree by 1.

Given a permutation w, the cell $(r, s) \in D(w)$ is called **accessible** if

- (1) $(r, s) \notin \mathsf{Dom}(w)$ and
- (2) there are no other cells which occur weakly southeast of (r, s) in D(w).

The maximal corner is an accessible box if and only if there exists i < r such that $\ell(w_P) + 1 = \ell(w_P t_{i,r})$. For vexillary permutations, there can be at most one such i, so we define $v_C = v_P t_{i,r}$ in this case. We may construct v_C graphically as follows. Consider the cells in D(v) which sit weakly northwest of the accessible box in its connected component. Move each of these diagonally one step in the northwest direction. This new diagram is the $D(v_C)$.

Lemma 4.8. Fix v vexillary, where the maximal corner (r, s) is an accessible box. Then $\deg(\mathfrak{G}_v) = \max\{\deg(\mathfrak{G}_{v_P}), \deg(\mathfrak{G}_{v_C})\} + 1.$

Proof. The monomials of Grothendieck polynomials alternate in sign based on degree. As such, Equation (4.3) is cancellation free. Therefore, the top degree monomials in \mathfrak{G}_v must come from $x_r \mathfrak{G}_{v_P}$ or $x_r \mathfrak{G}_{v_C}$.

Lemma 4.9. For v vexillary, if the maximal corner $(r, s) \in \text{Dom}(v)$, then $f_a(v) = f_a(v_P) + 1$.

Proof. By Equation (4.2), $\lambda(v_P)$ is obtained by removing the corresponding (boundary) cell from $\lambda(v)$. The label of this cell in F_v is zero since $(r, s) \in \mathsf{Dom}(v)$. At all other positions, F_v matches F_{v_P} . As such, $\tau_i(v) = \tau_i(v_P)$ for all i > 0. Therefore,

$$d(v) = |\lambda(v)| + \sum_{i=1}^{n} \rho_a(\tau_i(v))$$

= $|\lambda(v_P)| + 1 + \sum_{i=1}^{n} \rho_a(\tau_i(v_P))$
= $f_a(v_P) + 1.$

Lemma 4.10. Fix v vexillary and suppose the maximal corner (r, s) is an accessible box.

- (1) $f_a(v) \ge f_a(v_C) + 1$.
- (2) If (r, s) is the only cell in its row within its connected component in D(v), then $f_a(v) = f_a(v_C) + 1$.
- (3) $f_a(v) \ge f_a(v_P) + 1.$
- (4) If (r, s) is not the only cell in its row within its connected component in D(v), then $f_a(v) = f_a(v_P) + 1$.

Proof. Throughout, let (a, b) denote the position of the box in $\lambda(v)$ which corresponds to (r, s). Write $k = F_v(a, b)$. By assumption since $(r, s) \notin \mathsf{Dom}(v), k \ge 1$.

(1) To get F_{v_C} from F_v , take all labels weakly northwest of (a, b) with label k and decrease the value of these labels by 1. As such, $\tau_i(v) \supseteq \tau_i(v_C)$ for all i. Furthermore, since (r, s)has label $k, \tau_k(v) \supseteq \tau_k(v_C)$. In particular, $\tau_k(v_C)$ is obtained from $\tau_k(v)$ by removing a rectangular strip. Since this strip contains (r, s), removing this rectangle removes the last row of $\tau_k(v)$ entirely (and anything north of this row) by the definition of (r, s). Therefore, any antidiagonal path in $\tau_k(v_C)$ can be completed to a larger antidiagonal path in $\tau_k(v)$ by adding a box row r. As such, $\rho_a(\tau_k(v)) > \rho_a(\tau_k(v_C))$ and so

$$f_a(v) = |\lambda(v)| + \sum_{i=1}^n \rho_a(\tau_i(v))$$

> $|\lambda(v_C)| + \sum_{i=1}^n \rho_a(\tau_i(v_C))$
= $f_a(v_C).$

Since these are all integers, $f_a(v) \ge f_a(v_C) + 1$.

(2) Since there is a single box in the same row as (a, b) in $\tau_k(v)$ and this box is not in $\tau_k(v_C)$ (nor any boxes in its same column) we claim $\rho_a(\tau_k(v)) = \rho_a(\tau_k(v_C)) + 1$. For all other *i*, we have $\tau_i(v) = \tau_i(v_C)$ and so $\rho_a(\tau_i(v)) = \rho_a(\tau_i(v_C))$. Therefore, $f_a(v) = f_a(v_C) + 1$. (3) Using Equation (4.2), $F_{v_P}(i, j) = F_v(i, j)$ for all $(i, j) \in \lambda(v_P)$. As such,

(4.5)
$$\tau_i(v_P) = \begin{cases} \tau_i(v) & \text{if } i < k \\ \tau_i(v) - \{(a,b)\} & \text{otherwise.} \end{cases}$$

In particular, $\tau_i(v) \supseteq \tau_i(v_P)$ for all *i*. Therefore,

$$\sum_{i=1}^n \rho_a(\tau_i(v)) \ge \sum_{i=1}^n \rho_a(\tau_i(v_P)).$$

Then

$$f_a(v) = |\lambda(v)| + \sum_{i=1}^n \rho_a(\tau_i(v))$$

$$\geq |\lambda(v_P)| + 1 + \sum_{i=1}^n \rho_a(\tau_i(v_P))$$

$$= f_a(v_P) + 1.$$

(4) By assumption, $(r, s - 1) \in D(v)$. As such, if $(a, b) \in \tau_i(v)$ then $(a, b - 1) \in \tau_i(v)$ as well. Fix an antidiagonal path of cells in $\tau_i(v)$. If it does not use (a, b), then it is also an antidiagonal path of cells in $\tau_i(v_P)$. If it does use (a, b), then we can construct a new antidiagonal path of cells of the same size by replacing (a, b) with (a, b - 1). By (4.5), we see that this new antidiagonal path is also in $\tau_i(v_P)$. As such, $\rho_a(\tau_i(v)) = \rho_a(\tau_i(v))$. Then we conclude $f_a(v) = f_a(v_P) + 1$.

Proof. Fix v vexillary. The statement is trivial for the identity, so assume $\ell(v) \ge 1$. We will proceed by induction on the position of the maximal corner (r, s) (ordering cells of the grid lexicographically). In the base case, v = 21, we confirm $\deg(\mathfrak{G}_v) = 1 = f_a(v)$.

Assume the formula holds for any vexillary v' whose maximal corner occurs before (r, s), i.e., $\deg(\mathfrak{G}_{v'}) = f_a(v')$.

Case 1: $(r, s) \in \mathsf{Dom}(v)$. By Equation (4.2), the maximal corner of v_P occurs before (r, s). Furthermore, v_P is vexillary. As such,

$$deg(\mathfrak{G}_v) = deg(\mathfrak{G}_{v_P}) + 1 \qquad (by \text{ Lemma 4.7})$$
$$= f_a(v_P) + 1 \qquad (by \text{ induction hypothesis})$$
$$= f_a(v) \qquad (by \text{ Lemma 4.9}).$$

Case 2: $(r, s) \notin Dom(v)$ (i.e., it is an accessible box).

Both v_P and v_C are vexillary and their maximal corners (when defined) occur before (r, s). We know by Lemma 4.8 and the induction hypothesis that

(4.6)
$$\deg(\mathfrak{G}_{v}) = \max\{\deg(\mathfrak{G}_{v_{P}}), \deg(\mathfrak{G}_{v_{C}})\} + 1 = \max\{f_{a}(v_{P}), f_{a}(v_{C})\} + 1.$$

In particular, $1 + f_a(v_P) \leq \deg(\mathfrak{G}_v)$ and $1 + f_a(v_C) \leq \deg(\mathfrak{G}_v)$. Applying Lemma 4.10 to (4.6), we see that $\deg(\mathfrak{G}_v) \leq f_a(v)$. By parts (2) and (4) of Lemma 4.10, since (r, s) is an accessible box, $1 + f_a(v_P) = f_a(v)$ or $1 + f_a(v_C) = f_a(v)$. Then $f_a(v) = \deg(\mathfrak{G}_v)$.

5. Castelnuovo-Mumford regularity of Schubert determinantal ideals

We begin this section by recalling the connection between the Castelnuovo-Mumford regularity in the Cohen-Macaulay setting and the degree of a K-polynomial (Subsection 5.1). We then provide some background on Schubert determinantal ideals, explain how to express Castelnuovo-Mumford regularity of Schubert determinantal ideals in terms of degrees of Grothendieck polynomials, and prove Theorems 1.1 and 1.5 (Subsection 5.2).

5.1. Castelnuovo-Mumford regularity and connections to K-polynomials. Let $S = \mathbb{k}[x_1, \ldots, x_n]$ be a polynomial ring over the field \mathbb{k} , and assume that S is positively \mathbb{Z}^d -graded so that $\deg_0 S = \mathbb{k}$. Let M be a finitely generated graded S-module. The multigraded Hilbert series of M is a formal power series in indeterminates t_1, \ldots, t_d :

$$H(M; \mathbf{t}) = \sum_{\mathbf{a} \in \mathbb{Z}^d} \dim_{\mathbb{K}}(M_{\mathbf{a}}) \mathbf{t}^{\mathbf{a}} = \frac{K(M; \mathbf{t})}{\prod_{i=1}^n (1 - \mathbf{t}^{\mathbf{a}_i})}, \ \deg(x_i) = \mathbf{a}_i.$$

The numerator $K(M; \mathbf{t}) \in \mathbb{k}[\mathbf{t}^{\pm 1}]$ is called the **K-polynomial** of M. When S has the standard grading, that is $\deg(x_i) = 1$, the K-polynomial is a Laurent polynomial in a single indeterminate t.

For the rest of this subsection, assume that S has the standard grading, and let $I \subseteq S$ be a homogeneous ideal. There is a minimal free resolution

$$0 \to \bigoplus_{j} S(-j)^{\beta_{l,j}(S/I)} \to \bigoplus_{j} S(-j)^{\beta_{l-1,j}(S/I)} \to \dots \to \bigoplus_{j} S(-j)^{\beta_{0,j}(S/I)} \to S/I \to 0$$

where $l \leq n$ and S(-j) is the free S-module obtained by shifting the degrees of S by j. The **Castelnuovo-Mumford regularity** of S/I, denoted reg(S/I), is defined as

 $\operatorname{reg}(S/I) := \max\{j - i : \beta_{i,j}(S/I) \neq 0\}.$

When S/I is Cohen-Macaulay, we have that

(5.1)
$$\operatorname{reg}(S/I) = \operatorname{deg} K(S/I;t) - \operatorname{ht}_S I,$$

where $ht_S I$ denotes the height of the ideal I. See, for example, [2, Lemma 2.5] for justification of this formula. In this paper, we use Equation (5.1) to compute Castelnuovo-Mumford regularity of coordinate rings of certain matrix Schubert varieties and certain standardgraded Kazhdan-Lusztig varieties.

5.2. Regularity of Schubert determinantal ideals and proofs of Theorems 1.1 and 1.5. We begin by recalling basic facts about Schubert determinantal ideals. Fix an $n \times n$ permutation matrix w. Let $X = (x_{ij})$ be an $n \times n$ matrix of distinct indeterminates, and let $X_{[p],[q]}$ denote the matrix formed by intersecting the first p rows of X and the first q columns of X. Let $\Bbbk[\mathbf{x}] := \Bbbk[x_{ij} : 1 \leq i, j \leq n]$. The Schubert determinantal ideal $I_w \subseteq \Bbbk[\mathbf{x}]$ is the ideal

 $I_w = \langle \text{minors of size } r_w(i,j) + 1 \text{ in } X_{[i],[j]} : (i,j) \in \mathsf{Ess}(w) \rangle.$

By [10], I_w is a prime ideal, and $\mathbb{k}[\mathbf{x}]/I_w$ is Cohen-Macaulay. Recall that $\mathbb{k}[\mathbf{x}]/I_w$ is the coordinate ring of the matrix Schubert variety $\overline{B_-wB_+} \subseteq \operatorname{Mat}_{\mathbb{k}}(n,n)$ where $B_- \leq \operatorname{GL}_n(\mathbb{k})$ is the Borel subgroup of invertible lower triangular matrices, $B_+ \leq \operatorname{GL}_n(\mathbb{k})$ is the Borel subgroup of invertible upper triangular matrices, and $\operatorname{Mat}_{\mathbb{k}}(n,n)$ is the affine space of $n \times n$ matrices with entries in \mathbb{k} . Schubert determinantal ideals are homogeneous with respect to the standard grading of $\mathbb{k}[\mathbf{x}]$.

Proof of Theorems 1.1 and 1.5. We first recall how to express the regularity of $\mathbb{k}[\mathbf{x}]/I_w$ in terms of the degree of a Grothendieck polynomial. This was originally discussed in [32]. By [10], we have $h_{\mathbb{k}[\mathbf{x}]}I_w = \#D(w)$. It then follows by (5.1) that

$$\operatorname{reg}(\mathbb{k}[\mathbf{x}]/I_w) = \operatorname{deg} K(\mathbb{k}[\mathbf{x}]/I_w) - \#D(w).$$

By [3, Theorem 2.1] (see also [21, Theorem A]), $K(S/I_w; t) = \mathfrak{G}_w(1-t, \ldots, 1-t)$. Furthermore,

$$\deg \mathfrak{G}_w(1-t,\ldots,1-t) = \deg \mathfrak{G}_w(x_1,\ldots,x_n)$$

since the the coefficients in the homogeneous components $\mathfrak{G}_w(x_1,\ldots,x_n)$ all have the same sign (see, for example, [21]). Thus,

(5.2)
$$\operatorname{reg}(\Bbbk[\mathbf{x}]/I_w) = \operatorname{deg} \mathfrak{G}_w(x_1, \dots, x_n) - \#D(w)$$

Theorems 1.1 and 1.5 are now immediate from Theorems 1.3 and 1.7.

6. Regularity of homogeneous Kazhdan-Lusztig ideals

In this section, we recall the basics of Kazhdan-Lusztig ideals $J_{v,w}$ (Section 6.1) and provide preliminary combinatorial formulas for regularity of Kazhdan-Lusztig ideals $J_{v,w}$ when v is a 321-avoiding permutation (Section 6.2). We then provide an easily-computable combinatorial formula for the regularity of open patches of Schubert varieties in Grassmannians (Section 6.3). This proves a (generalization of a) conjecture from [32] giving a correction to a conjecture of [23].

6.1. Kazhdan-Lusztig ideals. We next recall Kazhdan-Lusztig ideals, which were introduced by A. Woo and A. Yong in [35] to study singularities of Schubert varieties. Given a permutation matrix $v \in S_n$, consider the matrix $M^{(v)}$ which has 1's at locations (i, v_i) , indeterminate z_{ij} in location $(i, j) \in D(v)$, and 0's elsewhere. Let $\mathbb{k}[\mathbf{z}^v] := \mathbb{k}[z_{ij} : (i, j) \in D(v)]$. Given $w \in S_n$, define the Kazhdan-Lusztig ideal $J_{v,w} \subseteq \mathbb{k}[\mathbf{z}^v]$ to be

$$J_{v,w} = \langle \text{minors of size } r_w(i,j) + 1 \text{ in } M^{(v)}_{[i],[j]} : (i,j) \in \mathsf{Ess}(w) \rangle$$

which is not the unit ideal precisely when $w \leq v$ in Bruhat order. The Kazhdan-Lusztig ideal $J_{v,w}$ is the prime defining ideal of the intersection of the Schubert variety $B_{-}\backslash\overline{B_{-}wB_{+}} \subseteq B_{-}\backslash\operatorname{GL}_{n}(\Bbbk)$ with the opposite Schubert cell $B_{-}\backslash B_{-}vB_{-}$ (see [35, Corollary 3.3] and the preceding discussion). Furthermore, $\Bbbk[\mathbf{z}^{v}]/J_{v,w}$ is Cohen-Macaulay. This follows by [19, Lemma A.4] together with the Cohen-Macaulayness of Schubert varieties [33]. See [35, Section 3.2] for further discussion.

Kahzdan-Lusztig ideals are not always homogeneous with respect to the standard grading on $\mathbb{k}[\mathbf{z}^{(v)}]$. However, when v is 321-avoiding, and hence when v is a Grassmannian permutation, $J_{v,w}$ is homogeneous with respect to the standard grading, see e.g., [20, Footnote on

pg. 25]. Some further partial results on the problem of when Kazhdan-Lusztig ideals are homogeneous with respect to the standard grading [35, Problem 5.5] can be found in the recent preprint [30].

6.2. Preliminaries on regularity of Kazhdan-Lusztig ideals $J_{v,w}$ where v is 321avoiding. We next describe a formula for the regularity of $\mathbb{k}[\mathbf{z}^v]/J_{v,w}$ where $J_{v,w}$ is a standard-graded Kazhdan-Lusztig ideal. This formula will be in terms of $\mathfrak{G}_w(\mathbf{x}; \mathbf{y})$, a double Grothendieck polynomial. Let $G_w(\mathbf{x}; \mathbf{y})$ denote the double Grothendieck polynomials in [21], so that $G_w(\mathbf{x}; \mathbf{y}) = \mathfrak{G}_w(\mathbf{1} - \mathbf{x}; \mathbf{1} - \frac{\mathbf{1}}{\mathbf{y}})$. We also let $G_{v,w}(\mathbf{t}) = \mathfrak{G}_{v,w}(1 - \mathbf{t})$.

The torus T^n acts on the opposite Schubert cell $B_{-}\backslash B_{-}vB_{-}$ by right multiplication. This induces a grading on $\mathbb{k}[\mathbf{z}^v]$ where variable z_{ij} in the matrix $M^{(v)}$ has degree $e_{v(i)} - e_j$, where $e_i \in \mathbb{Z}^n$ denotes the *i*th standard basis vector. By [36, Theorem 4.5], the *K*-polynomial of $\mathbb{k}[\mathbf{z}^v]/J_{v,w}$ for this \mathbb{Z}^n -grading is given by

(6.1)
$$K(\mathbb{k}[\mathbf{z}^{v}]/J_{v,w};\mathbf{t}) = G_w(t_{v(1)},\ldots,t_{v(n)};t_1,\ldots,t_n) = G_{v,w}(t_{ij} \mapsto t_{v(i)}/t_j).$$

Note that the conventions in [36] differ from ours.

In the case where v is 321-avoiding, there is a coarsening of the grading $f : \mathbb{Z}^n \to \mathbb{Z}$ which gives each $z_{ij} \in \mathbb{K}[z_{ij}]$ degree 1. Specifically, take $f(e_i) = 1$ if there exists k > i such that $v^{-1}(k) < v^{-1}(i)$ and $f(e_i) = 0$ otherwise (see e.g., the footnote on page 25 of [20]). Then the K-polynomial of $\mathbb{K}[\mathbf{z}^v]/J_{v,w}$, with respect to the standard grading, is

(6.2)
$$K(\mathbb{k}[\mathbf{z}^{v}]/J_{v,w};t) = G_w(t^{f(e_{v(1)})}, \dots, t^{f(e_{v(n)})};t^{-f(e_1)}, \dots, t^{-f(e_n)}) = G_{v,w}(t_{ij} \mapsto t^{f(e_{v(i)})+f(e_j)}).$$

Example 6.1. Let v = 34512 and w = 21435. Using Equation (2.1), we may compute $\mathfrak{G}_{v,w}(\mathbf{t}) = t_{11}t_{31} + t_{11}t_{22} - t_{11}t_{22}t_{31}$. For the \mathbb{Z}^n -grading, the substitution provided in Equation (6.1) yields

$$K(\mathbb{k}[\mathbf{z}^{v}]/J_{v,w};\mathbf{t}) = (1 - \frac{t_3}{t_1})(1 - \frac{t_5}{t_1}) + (1 - \frac{t_3}{t_1})(1 - \frac{t_4}{t_2}) - (1 - \frac{t_3}{t_1})(1 - \frac{t_5}{t_1})(1 - \frac{t_4}{t_2}).$$

Using Theorem 3.2, we may compute

$$\mathfrak{G}_w(\mathbf{x};\mathbf{y}) = \frac{x_1}{y_1}\frac{x_3}{y_1} + \frac{x_1}{y_1}\frac{x_2}{y_2} + \frac{x_1}{y_1}\frac{x_1}{y_3} - \frac{x_1}{y_1}\frac{x_1}{y_3}\frac{x_2}{y_2} - \frac{x_1}{y_1}\frac{x_1}{y_3}\frac{x_3}{y_1} - \frac{x_1}{y_1}\frac{x_2}{y_2}\frac{x_3}{y_1} + \frac{x_1}{y_1}\frac{x_1}{y_3}\frac{x_2}{y_2}\frac{x_3}{y_1}$$

Combining this with Equation (6.2) yields

$$K(\mathbb{k}[\mathbf{z}^{v}]/J_{v,w};\mathbf{t}) = G_w(1,1,1,t,t;t^{-1},t^{-1},1,1,1) = 2(1-t)^2 - (1-t)^3$$

under the \mathbb{Z} -grading.

Lemma 6.2. Let $v \in S_n$ such that v is 321-avoiding. If $(i, j) \in D(v)$, then $f(e_j) = 1$ and $f(e_{v_i}) = 0$.

Proof. Since $(i, j) \in D(v)$, there is $k = v_i$ such that $v_i > j$ and $i < v_j^{-1}$, thus $f(e_j) = 1$. If $f(e_{v_i}) = 1$, then there is $k > v_i$ such that $v_k^{-1} < i$. This would then imply that there is a 321-pattern in v. In particular, we would have $v_k^{-1} < i < v_j^{-1}$, with $j < v_i < k$. As v is 321-avoiding, we conclude that $f(e_{v_i}) = 0$.

 \diamond

Lemma 6.3. Let $v, w \in S_n$ such that v is 321-avoiding and $w \leq v$. Then

(6.3)
$$K(\mathbb{k}[\mathbf{z}^{v}]/J_{v,w};t) = G_{v,w}(t_{ij} \mapsto t) = \sum_{P \in \overline{\mathsf{Pipes}}(v,w)} (-1)^{\#P-\ell(w)} (1-t)^{\#P}$$

Proof. The coarsening of the grading $f : \mathbb{Z}^n \to \mathbb{Z}$ combined with Lemma 6.2 ensures that $t^{f(e_{v(i)})+f(e_j)} = t$ for $(i,j) \in D(v)$. Thus, the result follows by Equations (2.1) and (6.2) together with the fact that $G_{v,w}(\mathbf{t}) = \mathfrak{G}_{v,w}(1-\mathbf{t})$.

We will use the following to prove the main result of this section (Theorem 6.5).

Proposition 6.4. Let $v, w \in S_n$ such that v is 321-avoiding and $w \leq v$. Then,

(6.4)
$$\deg K(\Bbbk[\mathbf{z}^{v}]/J_{v,w};t) = \deg \mathfrak{G}_{v,w}(\mathbf{t}).$$

Furthermore, the Castelnuovo-Mumford regularity of $\mathbb{k}[\mathbf{z}^v]/J_{v,w}$ is given by

(6.5)
$$\operatorname{reg}(\mathbb{k}[\mathbf{z}^{v}]/J_{v,w}) = \operatorname{deg} \mathfrak{G}_{v,w}(\mathbf{t}) - \#D(w) = \max\{\#P \mid P \in \overline{\mathsf{Pipes}}(v,w)\} - \#D(w).$$

Proof. Equation (6.4) is immediate from Lemma 6.3. Equation (6.5) follows from Equations (6.4), (5.1) and the fact that $\operatorname{ht}_{\Bbbk[\mathbf{z}^v]} J_{v,w} = \#D(w)$.

6.3. Castelnuovo-Mumford regularity of patches of Grassmannian Schubert varieties. In [32], we gave a counterexample to a conjecture of Kummini-Lakshmibai-Sastry-Seshadri from [23] on the Castelnuovo-Mumford regularity of coordinate rings of standard open patches of certain Schubert varieties in Grassmannians. We then gave a conjecture of a correct formula [32, Conjecture 5.6]. In this short subsection, we prove a generalization of this conjecture.

Identify the Grassmannian $\operatorname{Gr}(k, n)$ with $P \setminus \operatorname{GL}_n(\Bbbk)$ where $P \subseteq \operatorname{GL}_n(\Bbbk)$ is the parabolic subgroup of block lower triangular matrices with block sizes k and n-k down the diagonal. Let u and g be a pair of Grassmannian permutations with descent at k. The Kazhdan-Lusztig ideal $J_{u,g}$ is the prime defining ideal of the intersection of the Schubert variety $P \setminus \overline{PgB_+} \subseteq P \setminus \operatorname{GL}_n(\Bbbk)$ with the open set $P \setminus PuB_- \subseteq P \setminus \operatorname{GL}_n(\Bbbk)$. The following theorem gives the regularity of the coordinate rings of these open sets of Grassmannian Schubert varieties.

Theorem 6.5. Fix Grassmannian permutations g and u with descent at position k so that $\lambda(g) \subseteq \lambda(u)$. Let v be the vexillary permutation such that $D(v) = D_{bot}(\lambda(u), \lambda(g))$. Then,

$$\operatorname{reg}(\mathbb{k}[\mathbf{z}^{u}]/J_{u,g}) = \operatorname{deg}(\mathfrak{G}_{v}(\mathbf{x})) - |\lambda(g)| = \sum_{i=1}^{n} \rho_{a}(\tau_{i}(v)).$$

Proof. The first equality follows due to Equation (6.5), Theorem 2.5, and the fact that $\operatorname{ht}_{\Bbbk[\mathbf{z}^u]}J_{u,g} = |\lambda(g)|$. The second equality is then immediate by Theorem 1.7 and the fact that $|\lambda(g)| = \#D(v)$ by construction of v.

We note that [32, Conjecture 5.6] concerned the special case of the above theorem where $u = (n - k + 1) (n - k + 2) \dots n + 2 \dots (n - k)$, written in one line notation.

REGULARITY OF LADDER DETERMINANTAL IDEALS

7. Regularity of ladder determinantal ideals

Our next goal is to provide a formula for the Castelnuovo-Mumford regularity of any one-sided ladder determinantal ideal. Ladder determinantal ideals are generalized determinantal ideals which were introduced by S. S. Abhyankar [1] to study singularities of Schubert varieties. There has since been substantial interest in their properties. For example, see [29, 18, 5, 6, 7, 12, 13, 22, 14, 15, 11] and references therein. The work of Ghorpade and Krattenthaler [11] on *a*-invariants of certain ladder determinantal ideals is most closely related to our results. This is discussed in more detail at the end of Section 7.1.

7.1. **One-sided ladder determinantal ideals.** A ladder L is a Young diagram (in English notation) filled with distinct indeterminates. Observe that a ladder is determined by a collection of southeast corners $L^{SE} = \{(a_i, b_i)\}_{i \in [s]}$ ordered northeast to southwest. Label the northwest corner of L to be (0, 0). Take (a_{s+1}, b_{s+1}) to be the southwestmost corner of the ladder and take (a_1, b_1) be the northeastmost corner of the ladder.

Let \mathcal{P} denote the lattice path from (a_{s+1}, b_{s+1}) to (a_1, b_1) which travels along the boundary of the ladder, so that cells weakly northwest of the \mathcal{P} are in L and boxes weakly southeast of \mathcal{P} are not in L. Let $P = \{(c_j, d_j)\}_{j \in [s']}$ denote a collection of distinguished points along \mathcal{P} . To each $(c_j, d_j) \in P$, assign a value $r_j \in \mathbb{Z}_{>0}$. Let $L_{I,J}$ denote the subset of L with row indices in I and column indices in J for $I, J \subseteq [n]$.

Let $\Bbbk[L]$ denote the polynomial ring generated by these indeterminate entries. Define the one-sided mixed ladder determinantal ideal $I_{L,\mathbf{r}}$:

$$I_{L,\mathbf{r}} = \langle \text{minors of size } r_j \text{ in } L_{[c_i],[d_i]} : j \in [s'] \rangle \subseteq \Bbbk[L].$$

Letting I_j denote the ideal of $r_j \times r_j$ minors of $L_{[c_j],[d_j]}$, one observes that

$$I_{L,\mathbf{r}} = \sum_{j \in [s']} I_j.$$

Following [21], we assume

(7.1)
$$0 < c_1 - r_1 < c_2 - r_2 < \dots < c_{s'} - r_{s'}$$
 and $0 < d_1 - r_1 < d_2 - r_2 < \dots < d_{s'} - r_{s'}$

so that $I_j \subseteq I_k$ for any $j \neq k, j, k \in [s']$. As outlined in [10, Proposition 9.6], L can be identified with a vexillary matrix Schubert variety \overline{X}_v where $\mathsf{Ess}(v)$ are the boxes indexed by P and the ranks satisfy $r_v(c_j, d_j) = r_j - 1$.

Example 7.1. To the left is a ladder L. Then $L^{SE} = \{(5,3), (3,5)\}$ with marked points and corresponding ranks given in red. To the right is the associated permutation v.



Then

$$I_{L,\mathbf{r}} = \langle 3 - \text{minors of } L_{[5],[3]}, 2 - \text{minors of } L_{[3],[3]}, 3 - \text{minors of } L_{[3],[5]} \rangle$$

= $\langle \det(L_{[3],\{3,4,5\}}), 2 - \text{minors of } L_{[3],[3]}, \det(L_{\{3,4,5\},[3]}) \rangle.$

For certain one-sided mixed ladder determinantal ideals, regularity formulas can be deduced through *a*-invariant formulas of Ghorpade-Krattenthaler [11]. Their formulas give results in the case in which $(r_1, r_2, \ldots, r_{s'}) = (1, 2, \ldots, t, t - 1, \ldots, 1)$ for some $t \in \mathbb{Z}_{>0}$, where Equation (7.1) is not imposed. Thus, for example, *L* as in Example 7.1 is not in the class of ladders considered in [11]. We note that an algorithm for *a*-invariant formulas is given in [11] for two-sided mixed ladder determinantal ideals with the same restriction on ranks.

7.2. One-sided ladder determinantal ideals via Grassmannian Kazhdan-Lusztig ideals. We now recall that each one-sided ladder determinantal ideal is a Kazhdan-Lusztig ideal $\mathcal{N}_{u,g}$ where u and g are Grassmannian permutations. This was first shown by Gonciulea-Miller [13, Theorem 4.7.3]; we include it here for completeness.

Take a ladder L with $L^{SE} = \{(a_i, b_i)\}_{i \in [s]}$, and marked points $P = \{(c_j, d_j)\}_{j \in [s']}$ assigning ranks r_j . Define $u \in S_{x+y}$ as the concatenation of partial permutations u_i , where for $i \in [s]$

(7.2)
$$u_{i} = \mathsf{id}_{a_{i}-a_{i+1}} + b_{i} + a_{0} - a_{i}, \text{ and} u_{s+1} = [x+y] \setminus \bigcup_{i \in [s]} u_{i}.$$

Set $(c_0, d_0) := (a_0, b_0)$ with $r_0 = 1$ and $(c_{s'+1}, d_{s'+1}) := (a_{s+1}, b_{s+1})$ with $r_{s'+1} = 1$. Define $g \in S_{x+y}$ as the concatenation of partial permutations g_i , where for $i \in [s'+1]$,

(7.3)
$$g_{i} = \mathsf{id}_{k_{i}-k_{i-1}} + k_{i-1} + h_{i-1}, \text{ and} \\ g_{s'+2} = [x+y] \setminus \bigcup_{i \in [s'+1]} g_{i}.$$

where $k_i = c_0 - c_i + r_i - 1$ and $h_i = d_i - r_i$.

Note that Equation (7.1) and the assumption that each indeterminate appears in at least one minor ensure that $L(\operatorname{code}(u)) = L(\operatorname{code}(g))$ and $u_j \ge g_j$ for each $j \in [x + y]$. Then by [13, Theorem 4.7.3] we have the following:

Proposition 7.2. Given a one-sided ladder determinantal ideal $I_{L,r}$ and u, g as above, $J_{u,g}$ and $I_{L,r}$ share the same generators.

Example 7.3. For L as in Example 7.1, below are D(u) and D(g) for the u, g as defined in Equations (7.2) and (7.3).





As a consequence to Proposition 7.2, the K-polynomial of each one-sided ladder determinantal ideal can be expressed both as a single Grothendieck polynomial and as a specialized double Grothendieck polynomial. Combining this with [10], we have:

Corollary 7.4. Given a one-sided ladder L with marked points $P = \{(c_j, d_j)\}_{j \in [s']}$ assigning ranks r_j ,

$$\operatorname{reg}(S/I_L) = \operatorname{reg}(S/J_{u,g}) = \sum_{k=1}^n \rho_a(\tau_k(v)),$$

where u, g are as defined in Equations (7.2) and (7.3). Here v is the vexillary permutation such that $\mathsf{Ess}(v)$ are the boxes indexed by P and $r_v(c_j, d_j) = r_j - 1$.

References

- S. S. Abhyankar. Enumerative combinatorics of Young tableaux, volume 115 of Monographs and Textbooks in Pure and Applied Mathematics. Marcel Dekker, Inc., New York, 1988.
- [2] B. Benedetti and M. Varbaro. On the dual graphs of Cohen-Macaulay algebras. Int. Math. Res. Not. IMRN, 2015(17):8085-8115, 2015.
- [3] A. S. Buch. Grothendieck classes of quiver varieties. Duke Math. J., 115(1):75–103, 2002.
- [4] Anders Skovsted Buch. A Littlewood-Richardson rule for the K-theory of Grassmannians. Acta Math., 189(1):37–78, 2002.
- [5] A. Conca. Ladder determinantal rings. J. Pure Appl. Algebra, 98(2):119–134, 1995.
- [6] A. Conca. Gorenstein ladder determinantal rings. J. London Math. Soc. (2), 54(3):453–474, 1996.
- [7] A. Conca and J. Herzog. Ladder determinantal rings have rational singularities. Adv. Math., 132(1):120– 147, 1997.
- [8] Neil J. Y. Fan and Peter L. Guo. Set-valued Rothe tableaux and Grothendieck polynomials. Adv. in Appl. Math., 128:Paper No. 102203, 28, 2021.
- S. Fomin and A. N. Kirillov. Grothendieck polynomials and the Yang-Baxter equation. In Formal power series and algebraic combinatorics/Séries formelles et combinatorie algébrique, pages 183–189. DIMACS, Piscataway, NJ, 1994.
- [10] W. Fulton. Flags, Schubert polynomials, degeneracy loci, and determinantal formulas. Duke Math. J., 65(3):381–420, 1992.
- [11] S. R. Ghorpade and C. Krattenthaler. Computation of the *a*-invariant of ladder determinantal rings. J. Algebra Appl., 14(9):1540014, 24, 2015.
- [12] N. Gonciulea and V. Lakshmibai. Singular loci of ladder determinantal varieties and Schubert varieties. J. Algebra, 229(2):463–497, 2000.
- [13] N. Gonciulea and C. Miller. Mixed ladder determinantal varieties. J. Algebra, 231(1):104–137, 2000.
- [14] E. Gorla. Mixed ladder determinantal varieties from two-sided ladders. J. Pure Appl. Algebra, 211(2):433–444, 2007.
- [15] E. Gorla, J. C. Migliore, and U. Nagel. Gröbner bases via linkage. J. Algebra, 384:110–134, 2013.
- [16] W. Graham and V. Kreiman. Excited Young diagrams, equivariant K-theory, and Schubert varieties. Trans. Amer. Math. Soc., 367(9):6597–6645, 2015.
- [17] E. Hafner. Vexillary Grothendieck polynomials via bumpless pipe dreams, 2022. arXiv:2201.12432.
- [18] J. Herzog and N. V. Trung. Gröbner bases and multiplicity of determinantal and Pfaffian ideals. Adv. Math., 96(1):1–37, 1992.
- [19] D. Kazhdan and G. Lusztig. Representations of Coxeter groups and Hecke algebras. Invent. Math., 53(2):165–184, 1979.
- [20] A. Knutson. Frobenius splitting, point-counting, and degeneration, 2009. arXiv:0911.4941.
- [21] A. Knutson and E. Miller. Gröbner geometry of Schubert polynomials. Ann. of Math., 161(3):1245–1318, 2005.

- [22] A. Knutson, E. Miller, and A. Yong. Gröbner geometry of vertex decompositions and of flagged tableaux. J. Reine Angew. Math., 630:1–31, 2009.
- [23] M. Kummini, V. Lakshmibai, P. Sastry, and C. S. Seshadri. Free resolutions of some Schubert singularities. *Pacific J. Math.*, 279(1-2):299–328, 2015.
- [24] A. Lascoux. Transition on Grothendieck polynomials. In Physics and combinatorics, 2000 (Nagoya), pages 164–179. World Sci. Publ., River Edge, NJ, 2001.
- [25] A. Lascoux and M. P. Schützenberger. Structure de Hopf de l'anneau de cohomologie et de l'anneau de Grothendieck d'une variété de drapeaux. C. R. Acad. Sci. Paris Sér. I Math., 295(11):629–633, 1982.
- [26] C. Lenart. Combinatorial aspects of the K-theory of Grassmanians. Ann. Comb., 4(1):67–82, 2000.
- [27] L. Manivel. Symmetric functions, Schubert polynomials and degeneracy loci. SMF/AMS Texts and Monographs. American Mathematical Society, Providence, 2001. Translated from the 1998 French original by John R. Swallow.
- [28] A. H. Morales, I. Pak, and G. Panova. Hook formulas for skew shapes I. q-analogues and bijections. J. Combin. Theory Ser. A, 154:350–405, 2018.
- [29] H. Narasimhan. The irreducibility of ladder determinantal varieties. J. Algebra, 102(1):162–185, 1986.
- [30] E. Neye. A Gröbner basis for Schubert patch ideals, 2021. arXiv:2111.13778.
- [31] O. Pechenik, D. Speyer, and A. Weigandt. Castelnuovo-Mumford regularity of matrix Schubert varieties, 2021. arXiv:2111.10681.
- [32] J. Rajchgot, Y. Ren, C. Robichaux, A. St. Dizier, and A. Weigandt. Degrees of symmetric Grothendieck polynomials and Castelnuovo-Mumford regularity. Proc. Amer. Math. Soc., 149(4):1405–1416, 2021.
- [33] A. Ramanathan. Schubert varieties are arithmetically Cohen-Macaulay. Invent. Math., 80(2):283–294, 1985.
- [34] A. Weigandt. Bumpless pipe dreams and alternating sign matrices. J. Combin. Theory Ser. A, 182:Paper No. 105470, 52, 2021.
- [35] A. Woo and A. Yong. Governing singularities of Schubert varieties. J. Algebra, 320(2):495–520, 2008.
- [36] A. Woo and A. Yong. A Gröbner basis for Kazhdan-Lusztig ideals. Amer. J. Math., 134(4):1089–1137, 2012.
- [37] A. Yong. Castelnuovo-Mumford regularity and Schubert geometry, 2022. arXiv:2202.06362.

(JR) DEPT. OF MATHEMATICS AND STATISTICS, MCMASTER UNIVERSITY, HAMILTON, ON L8S 4K1, CANADA Email address: rajchgoj@mcmaster.ca

(CR) DEPT. OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, LOS ANGELES LOS ANGELES, CA 90095, USA

 $Email \ address: \ {\tt robichaux@ucla.edu}$

(AW) DEPT. OF MATHEMATICS, MASSACHUSETTS INSTITUTE OF TECHNOLOGY, CAMBRIDGE, MA 02142, USA

Email address: weigandt@mit.edu