

EQUIVARIANT COHOMOLOGY, SCHUBERT CALCULUS, AND EDGE LABELED TABLEAUX

COLLEEN ROBICHAUX, HARSHIT YADAV, AND ALEXANDER YONG

ABSTRACT. This chapter concerns edge labeled Young tableaux, introduced by H. Thomas and the third author. It is used to model equivariant Schubert calculus of Grassmannians. We survey results, problems, conjectures, together with their influences from combinatorics, algebraic and symplectic geometry, linear algebra, and computational complexity. We report on a new shifted analogue of edge labeled tableaux. Conjecturally, this gives a Littlewood–Richardson rule for the structure constants of the D. Anderson–W. Fulton ring, which is related to the equivariant cohomology of isotropic Grassmannians.

To William Fulton on his eightieth birthday, for inspiring generations.

1. INTRODUCTION

1.1. Purpose. Singular cohomology is a functor between the categories

$\{\text{topological spaces, continuous maps}\} \rightarrow \{\text{graded algebras, homomorphisms}\}.$

The cohomology functor links the geometry of Grassmannians to symmetric functions and Young tableaux. However, this does not take into account the large torus action on the Grassmannian. A similar functor for topological spaces with continuous group actions is equivariant cohomology.

What are equivariant analogues for these centerpieces of algebraic combinatorics?

We posit a comprehensive answer, with applications, and future perspectives.

1.2. Schubert calculus. Let $X = \text{Gr}_k(\mathbb{C}^n)$ be the Grassmannian of k -dimensional planes in \mathbb{C}^n . The group GL_n of invertible $n \times n$ matrices acts transitively on X by change of basis. Let $B_- \subset \text{GL}_n$ be an opposite Borel subgroup of lower triangular matrices. B_- acts on X with finitely many orbits X_λ° where λ is a partition (identified with its Young diagram, in English notation) that is contained in the $k \times (n - k)$ rectangle Λ . These *Schubert cells* satisfy $X_\lambda^\circ \cong \mathbb{C}^{k(n-k)-|\lambda|}$ where $|\lambda| = \sum_i \lambda_i$. Their closures, the *Schubert varieties*, satisfy

$$X_\lambda := \overline{X_\lambda^\circ} = \coprod_{\mu \supseteq \lambda} X_\mu^\circ.$$

Let ν^\vee be the 180° -rotation of $k \times (n - k) \setminus \nu$. Suppose $|\lambda| + |\mu| + |\nu^\vee| = k(n - k) = \dim X$. By Kleiman transversality [29], there is a dense open $\mathcal{O} \subset \text{GL}_n \times \text{GL}_n \times \text{GL}_n$ such that

$$c_{\lambda, \mu}^\nu := \#\{g_1 \cdot X_\lambda \cap g_2 \cdot X_\mu \cap g_3 \cdot X_{\nu^\vee}\} \in \mathbb{Z}_{\geq 0}$$

is independent of $(g_1, g_2, g_3) \in \mathcal{O}$. Each $c_{\lambda, \mu}^\nu$ is called a *Littlewood–Richardson coefficient*. Modern Schubert calculus is concerned with these coefficients, as well as their generalizations/analogues (from varying the space X or cohomology theory).

Date: June 11, 2020.

Let $\sigma_\lambda \in H^{2|\lambda|}(X)$ be the Poincaré dual to X_λ . These *Schubert classes* form a \mathbb{Z} -linear basis of $H^*(X)$ and

$$\sigma_\lambda \smile \sigma_\mu = \sum_{\nu \subseteq \Lambda} c_{\lambda, \mu}^\nu \sigma_\nu.$$

The *Schur function* s_λ is the generating series $s_\lambda = \sum_T x^T$ for semistandard Young tableaux of shape λ , *i.e.*, row weakly increasing and column strictly increasing fillings of λ with elements of \mathbb{N} . The weight of T is $x^T := \prod_i x_i^{\#i \in T}$. For example, if $\lambda = (2, 1)$, the semistandard tableaux are

$$\begin{array}{|c|c|} \hline 1 & 1 \\ \hline 2 & \\ \hline \end{array} \quad \begin{array}{|c|c|} \hline 1 & 2 \\ \hline 2 & \\ \hline \end{array} \quad \begin{array}{|c|c|} \hline 1 & 3 \\ \hline 2 & \\ \hline \end{array} \quad \begin{array}{|c|c|} \hline 1 & 1 \\ \hline 3 & \\ \hline \end{array} \quad \begin{array}{|c|c|} \hline 1 & 2 \\ \hline 3 & \\ \hline \end{array} \quad \begin{array}{|c|c|} \hline 1 & 3 \\ \hline 3 & \\ \hline \end{array} \quad \begin{array}{|c|c|} \hline 2 & 2 \\ \hline 3 & \\ \hline \end{array} \quad \begin{array}{|c|c|} \hline 2 & 3 \\ \hline 3 & \\ \hline \end{array} \quad \dots$$

Hence $s_{(2,1)} = x_1^2 x_2 + x_1 x_2^2 + x_1 x_2 x_3 + x_1^2 x_3 + x_1 x_2 x_3 + x_1 x_3^2 + x_2^2 x_3 + x_2 x_3^2 + \dots$. Schur functions form a \mathbb{Z} -linear basis of Sym , the ring of symmetric functions in infinitely many variables.

The map $\sigma_\lambda \mapsto s_\lambda$ induces a ring isomorphism

$$H^*(X) \cong \text{Sym}/I$$

where I is the ideal $\langle s_\lambda : \lambda \not\subseteq \Lambda \rangle$. Therefore in Sym ,

$$(1) \quad s_\lambda \cdot s_\mu = \sum_{\nu} c_{\lambda, \mu}^\nu s_\nu.$$

To compute Schubert calculus of X , it suffices to determine (1) by working with Schur *polynomials* in only finitely many variables x_1, \dots, x_k . Better yet, the *jeu de taquin* theory of Young tableaux, introduced by M.-P. Schützenberger [58] gives a combinatorial rule for computing $c_{\lambda, \mu}^\nu$; this is summarized in Section 3. What we have discussed thus far constitutes textbook material on Schubert calculus; see, *e.g.*, W. Fulton's [18].

1.3. Overview. This chapter describes an equivariant analogue of M.-P. Schützenberger's theory, due to H. Thomas and the third author; in short, one replaces Young tableaux with *edge labeled tableaux*. Now, we hasten to offer an *apologia*: such tableaux are not the only combinatorial model to compute equivariant Schubert calculus. For example, one has work of A. Molev-B. Sagan [43] and the *puzzles* of A. Knutson-T. Tao [33]. The latter has had important recent followup, see, *e.g.*, A. Knutson-P. Zinn-Justin's [36] and the references therein. One also has the tableaux of V. Kreiman [39] or A. Molev [42].

That said, we wish to argue how the edge labeled model is a handy and flexible viewpoint. It has been applied to obtain equivariant analogues of a number of theorems (delineated in Section 3 and 5). Another application, due to O. Pechenik and the third author [49], is to Schubert calculus for the equivariant K -theory of X . Translating the combinatorics into puzzle language allowed for a proof (of a correction) of a conjecture of A. Knutson-R. Vakil about the same structure constants [48]. However, as we wish to restrict to equivariant cohomology proper, this direction is not part of our discussion.

There is an important frontier to cross, that is, the still unsolved problem of finding a combinatorial rule for equivariant Schubert calculus of maximal orthogonal and Lagrangian Grassmannians. The non-equivariant story is explained in Section 7. We explain the problem in Section 8 together with some recent developments of C. Monical [44] and of the authors [55]. The latter work shows that the combinatorial problems concerning the two spaces are essentially equivalent.

This brings us to the principal new announcement of this work (Section 9): the notion of *shifted edge labeled tableaux*. We define an analogue of *jeu de taquin* and use this to conjecturally define an associative ring (which we prove to also be commutative). The introduction of this ring is stimulated by recent work of D. Anderson-W. Fulton (see Section 10) who define a ring connected to the equivariant cohomology of Lagrangian Grassmannians. Conjecturally, the two rings are isomorphic. This provides our strongest evidence to date of the applicability of the edge labeled approach to the aforementioned open problem; we know of no similar results using other combinatorial models.

As this work is partially expository and partly an announcement, we limited the number of complete proofs in order to keep the focus on the high-level research objectives. Where possible, we have sketched arguments (with references) and indicated those results which may be taken as an exercise for the interested reader. These exercises are warmups for the conjectures and open problems contained herein.

2. EQUIVARIANT COHOMOLOGY OF GRASSMANNIANS

2.1. Generalities. We recall some general notions about equivariant cohomology. References that we consulted are L. Tu's synopsis [69], J. Tymoczko's exposition [70] and A. Knutson-T. Tao's [33, Section 2].

Let \mathcal{M} be a topological space with the continuous action of a topological group G . If G acts freely on \mathcal{M} , then in fact the equivariant cohomology ring $H_G^*(\mathcal{M})$ is $H^*(\mathcal{M}/G)$, see [70, Proposition 2.1]. However, in general the action is not free, and \mathcal{M}/G might be, e.g., non-Hausdorff. Borel's *mixing space construction* introduces a contractible space EG on which G acts freely. Thus G 's diagonal action on $EG \times \mathcal{M}$ is free and

$$H_G^*(\mathcal{M}) := H^*(EG \times \mathcal{M}/G).$$

The space EG is the total space of the *universal principal G -bundle* $\pi : EG \rightarrow BG$ where $BG = EG/G$ is the classifying space of G . Here, universality means that if $\rho : P \rightarrow \mathcal{M}$ is any G -bundle, there exists a unique map $f : \mathcal{M} \rightarrow BG$ (up to homotopy) such that $P \cong f^*(EG)$. By functoriality, the constant map $c : \mathcal{M} \rightarrow \{pt\}$ induces a homomorphism $c^* : H_G^*(pt) \rightarrow H_G^*(\mathcal{M})$. Hence $H_G^*(\mathcal{M})$ is a module over $H_G^*(pt)$ by $\beta \cdot \kappa := c^*(\beta)\kappa$ for $\beta \in H_G^*(pt)$ and $\kappa \in H_G^*(\mathcal{M})$. While ordinary (singular) cohomology of a point is \mathbb{Z} , $H_G^*(pt)$ is *big*. For instance, if $G = T$ is an n torus $(S^1)^n$, then $H_T^*(pt) = \mathbb{Z}[t_1, \dots, t_n]$.

If we presume G is a algebraic group acting on a smooth algebraic variety M , these notions have versions in the algebraic category; see, e.g., D. Anderson's [2].

2.2. The Grassmannian. Concretely, if $\lambda = (\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_k \geq 0)$ then

$$(2) \quad X_\lambda = \{V \in X \mid \dim(V \cap F^{n-k+i-\lambda_i}) \geq i, 1 \leq i \leq k\},$$

where $F^d = \text{span}(e_n, e_{n-1}, \dots, e_{n-d+1})$ and e_i is the i -th standard basis vector; see [19, Section 9.4] for details.

Let $T \subset GL_n$ be the torus of invertible diagonal matrices. Then from (2), X_λ is T -stable. Therefore, X_λ admits an equivariant Schubert class ξ_λ in the T -equivariant cohomology ring $H_T^*(X)$. By what we have recounted in Section 2.1, $H_T^*(X)$ is a module over

$$(3) \quad H_T^*(pt) := \mathbb{Z}[t_1, t_2, \dots, t_n].$$

The equivariant Schubert classes form a $H_{\top}^*(pt)$ -module basis of $H_{\top}^*(X)$. Therefore,

$$(4) \quad \xi_{\lambda} \cdot \xi_{\mu} = \sum_{\nu \subseteq \Lambda} C_{\lambda, \mu}^{\nu} \xi_{\nu},$$

where $C_{\lambda, \mu}^{\nu} \in H_{\top}^*(pt)$. For more details about equivariant cohomology specific to flag varieties we point the reader to [2] and S. Kumar's textbook [37, Chapter XI].

Let $\beta_i := t_i - t_{i+1}$. D. Peterson conjectured, and W. Graham [22] proved

Theorem 2.1 (Equivariant positivity [22]). $C_{\lambda, \mu}^{\nu} \in \mathbb{Z}_{\geq 0}[\beta_1, \dots, \beta_{n-1}]$.¹

In fact, $\deg C_{\lambda, \mu}^{\nu} = |\lambda| + |\mu| - |\nu|$ and $C_{\lambda, \mu}^{\nu} = 0$ unless $|\lambda| + |\mu| \geq |\nu|$. In the case $|\lambda| + |\mu| = |\nu|$, $C_{\lambda, \mu}^{\nu} = c_{\lambda, \mu}^{\nu}$ is the Littlewood–Richardson coefficient of Section 1.

Fix a grid with n rows and $m \geq n + \lambda_1 - 1$ columns. The *initial diagram* places λ in the northwest corner of this grid. For example, if $\lambda = (3, 2, 0, 0)$, the initial diagram for λ is the first of the three below.

$$\begin{bmatrix} + & + & + & \cdot & \cdot & \cdot \\ + & + & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \end{bmatrix} \quad \begin{bmatrix} + & + & \cdot & \cdot & \cdot & \cdot \\ + & \cdot & \cdot & + & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & + & \cdot & \cdot \end{bmatrix} \quad \begin{bmatrix} + & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & + & \cdot & \cdot & \cdot \\ \cdot & + & \cdot & \cdot & + & \cdot \\ \cdot & \cdot & \cdot & + & \cdot & \cdot \end{bmatrix}$$

A *local move* is a change of any 2×2 subsquare of the form

$$\begin{array}{cc} + & \cdot \\ \cdot & \cdot \end{array} \mapsto \begin{array}{cc} \cdot & \cdot \\ \cdot & + \end{array}$$

A *plus diagram* is any configuration of $+$'s in the grid resulting from some number of local moves starting from the initial diagram for λ . We have given two more examples of plus diagrams for $\lambda = (3, 2, 0, 0)$.

Let $\text{Plus}(\lambda)$ denote the set of plus diagrams for λ . If $P \in \text{Plus}(\lambda)$, let

$$\text{wt}_x(P) = x_1^{\alpha_1} x_2^{\alpha_2} \cdots x_n^{\alpha_n}.$$

Here, α_i is the number of $+$'s in the i th row of P . For instance, if P is the rightmost diagram shown above, $\text{wt}_x(P) = x_1 x_2 x_3^2 x_4$. A more refined statistic is

$$\text{wt}_{x,y}(P) = \prod_{(i,j)} x_i - y_j.$$

The product is over those (i, j) with a $+$ in row i and column j of P . For the same P ,

$$\text{wt}_{x,y}(P) = (x_1 - y_1)(x_2 - y_3)(x_3 - y_2)(x_3 - y_5)(x_4 - y_4).$$

Let $X = \{x_1, x_2, \dots, x_n\}$ and $Y = \{y_1, y_2, \dots, y_{n+\lambda_1-1}\}$ be two collections of indeterminates. The *factorial Schur function* is

$$s_{\lambda}(X; Y) = \sum_{P \in \text{Plus}(\lambda)} \text{wt}_{x,y}(P).$$

¹Actually, W. Graham proved that for any generalized flag variety H/B , the equivariant Schubert structure constant is expressible as a nonnegative integer polynomial in the simple roots of the (complex, semisimple) Lie group H .

This description arises in, *e.g.*, [31]. Moreover, it is an exercise to show

$$s_\lambda(X) = \sum_{P \in \text{Plus}(\lambda)} \text{wt}_x(P) = s_\lambda(X; 0, 0, \dots).$$

The factorial Schur polynomials form a $\mathbb{Z}[Y]$ -linear basis of $\text{Sym} \otimes_{\mathbb{Q}} \mathbb{Z}[Y]$. In addition,

$$(5) \quad s_\lambda(X; Y) s_\mu(X; Y) = \sum_{\nu} C_{\lambda, \mu}^{\nu} s_{\nu}(X; Y).$$

For example, one checks that

$$s_{(1,0)}(x_1, x_2; Y)^2 = s_{(2,0)}(x_1, x_2; Y) + s_{(1,1)}(x_1, x_2; Y) + (y_3 - y_2) s_{(1,0)}(x_1, x_2; Y).$$

In view of (3), the definition of $C_{\lambda, \mu}^{\nu}$ in terms of (5) gives a definition of $H_{\mathbb{T}}^*(X)$ that suffices for our combinatorial ends.

2.3. Equivariant restriction. We may further assume G is an algebraic n -torus T which acts on \mathcal{M} with finitely many isolated fixed points \mathcal{M}^T . A feature of equivariant cohomology is that the inclusion \mathcal{M}^T into \mathcal{M} induces an *injection*

$$H_{\mathbb{T}}^*(\mathcal{M}) \hookrightarrow H_{\mathbb{T}}^*(\mathcal{M}^T) \cong \bigoplus_{\mathcal{M}^T} H_{\mathbb{T}}^*(pt) \cong \bigoplus_{\mathcal{M}^T} \mathbb{Z}[t_1, \dots, t_n].$$

For each T -invariant cycle Y in \mathcal{M} , one has an equivariant cohomology class $[Y]_{\mathbb{T}} \in H_{\mathbb{T}}^*(\mathcal{M})$. This injection says that this class is a $\#\{\mathcal{M}^T\}$ -tuple of polynomials $[Y]|_x$ where $x \in \mathcal{M}^T$. Each polynomial $[Y]|_x$ is an *equivariant restriction*. Under certain assumptions on \mathcal{M} , which cover all generalized flag manifolds (such as Grassmannians), one has a divisibility condition on the restrictions. This alludes to the general and influential package of ideas contained in Goresky–Kottwitz–MacPherson (“GKM”) theory; we refer to [21] as well as J. Tymoczko’s survey [70].

One has more precise results (predating [21]) for any generalized flag variety. Work of B. Kostant–S. Kumar [38] combined with a formula of H. Anderson–J. Jantzen–W. Soergel [4] describes $[Y]|_x$ where Y is a Schubert variety and x is one of the T -fixed points. For restriction formulas specific to the Grassmannian $M = \text{Gr}_k(\mathbb{C}^n)$, see the formula of T. Ikeda–H. Naruse [28, Section 3] in terms of *excited Young diagrams*.² From either formula, one sees immediately that

$$(6) \quad \xi_\lambda|_\mu = 0 \text{ unless } \lambda \subseteq \mu,$$

and

$$(7) \quad \xi_\mu|_\mu \neq 0.$$

The central difference between this picture of equivariant cohomology of Grassmannians and the Borel-type presentation (due to A. Arabia [5]) is that multiplication can be done *without relations* and computed by pointwise multiplication of the restriction polynomials. In particular, each instance of (4) gives rise to $\binom{n}{k}$ many polynomial identities. For example, combining (4) and (6) gives

$$\xi_\lambda|_\mu \cdot \xi_\mu|_\mu = C_{\lambda, \mu}^{\mu} \xi_\mu|_\mu.$$

²One can give another formula in terms of certain specializations of the factorial Schur polynomial; see, *e.g.*, [28, Theorem 5.4] and the associated references.

By (7), this implies

$$(8) \quad \xi_\lambda|_\mu = C_{\lambda,\mu}^\mu,$$

which is a fact first noted (for generalized flag varieties) by A. Arabia [5]. It follows that:

$$(9) \quad \xi_\lambda \cdot \xi_{(1)} = \xi_\lambda|_{(1)} \xi_\lambda + \sum_{\lambda^+} \xi_{\lambda^+},$$

where λ^+ is obtained by adding a box to λ ; see, *e.g.*, [33, Proposition 2]. Alternatively, it is an exercise to derive it from a vast generalization due to C. Lenart-A. Postnikov's [40, Corollary 1.2].

Since $H_T^*(X)$ is an associative ring, one has

$$(\xi_\lambda \cdot \xi_\mu) \cdot \xi_{(1)} = \xi_\lambda \cdot (\xi_\mu \cdot \xi_{(1)}),$$

which when expanded using (4) and (9) gives a recurrence that uniquely determines the structure coefficients; we call this the *associativity recurrence*. Since we will not explicitly need it in this chapter we leave it as an exercise (see [66, Lemma 3.3] and the references therein).³

3. YOUNG TABLEAUX AND JEU DE TAQUIN

There are several combinatorial rules for the Littlewood–Richardson coefficient $c_{\lambda,\mu}^\nu$. The one whose theme will pervade this chapter is the *jeu de taquin* rule, which, moreover is the first *proved* rule for the coefficients [58].

Let ν/λ be a skew shape. A *standard tableau* T of shape ν/λ is a bijective filling of ν/λ with $1, 2, \dots, |\nu/\lambda|$ such that the rows and columns are increasing. Let $\text{SYT}(\nu/\lambda)$ be the set of all such tableaux. An *inner corner* c of λ/μ is a maximally southeast box of μ . For $T \in \text{SYT}(\lambda/\mu)$, a *jeu de taquin slide* $\text{jdt}_c(T)$ is obtained as follows. Initially place \bullet in c , and apply one of the following *slides*, according to how T looks near c :

$$(J1) \quad \begin{array}{|c|c|} \hline \bullet & a \\ \hline b & \\ \hline \end{array} \mapsto \begin{array}{|c|c|} \hline b & a \\ \hline \bullet & \\ \hline \end{array} \text{ (if } b < a, \text{ or } a \text{ does not exist)}$$

$$(J2) \quad \begin{array}{|c|c|} \hline \bullet & a \\ \hline b & \\ \hline \end{array} \mapsto \begin{array}{|c|c|} \hline a & \bullet \\ \hline b & \\ \hline \end{array} \text{ (if } a < b, \text{ or } b \text{ does not exist)}$$

Repeat application of (J1) or (J2) on the new box c' where \bullet arrives at. End when \bullet arrives at a box d of λ that has no labels south or east of it. Then $\text{jdt}_c(T)$ is obtained by erasing \bullet .


A *rectification* of $T \in \text{SYT}(\lambda/\mu)$ is defined iteratively. Pick an inner corner c_0 of λ/μ and compute $T_1 := \text{jdt}_{c_0}(T) \in \text{SYT}(\lambda^{(1)}/\mu^{(1)})$. Let c_1 be an inner corner of $\lambda^{(1)}/\mu^{(1)}$ and compute $T_2 := \text{jdt}_{c_1}(T_1) \in \text{SYT}(\lambda^{(2)}/\mu^{(2)})$. Repeat $|\mu|$ times, arriving at a standard tableau of straight (*i.e.*, partition) shape. Let $\text{Rect}_{\{c_i\}}(T)$ be the result.

Theorem 3.1 (First fundamental theorem of jeu de taquin). *$\text{Rect}_{\{c_i\}}(T)$ is independent of the choice of sequence of successive inner corners $\{c_i\}$.*

Theorem 3.1 permits one to speak of *the* rectification $\text{Rect}(T)$.

³We give an analogue (22) of the associativity recurrence in our proof of Theorem 8.2.

Example 3.2. For instance, here are two different rectification orders for a tableau T .




Theorem 3.3 (Second fundamental theorem of jeu de taquin). *The cardinality*

$$(10) \quad \#\{T \in \text{SYT}(\nu/\lambda) : \text{Rect}(T) = U\}$$

is independent of the choice of $U \in \text{SYT}(\mu)$.

Example 3.4. Below are the tableaux $T \in \text{SYT}((3, 2, 1)/(2, 1))$ such that $\text{Rect}(T) = U \in \text{SYT}((2, 1))$. Of the tableaux below, T_1, T_2 rectify to U_1 and T_3, T_4 rectify to U_2 .



For proofs of Theorems 3.1 and 3.3 we recommend the self-contained argument found in M. Haiman's [24], which is based on his theory of *dual equivalence*.

Theorem 3.5 ([58], Jeu de taquin computes the Littlewood–Richardson coefficient). *Fix $U \in \text{SYT}(\mu)$. Then $c'_{\lambda, \mu}$ equals the number (10).*

Example 3.6. Continuing Example 3.4, fix $U := U_1$. Then $c_{(2,1), (2,1)}^{(3,2,1)} = 2 = \#\{T_1, T_2\}$. \square

It is convenient to fix a choice of tableau U in Theorem 3.5. Namely, let $U = S_\mu$ be the superstandard tableau of shape μ , which is obtained by filling the boxes of μ in English reading order with $1, 2, 3, \dots$. This is the choice made in Example 3.6. A larger instance is

$$S_{(5,3,1)} = \begin{array}{|c|c|c|c|c|} \hline 1 & 2 & 3 & 4 & 5 \\ \hline 6 & 7 & 8 & & \\ \hline 9 & & & & \\ \hline \end{array}.$$

There are a number of ways to prove a Littlewood–Richardson rule such as Theorem 3.5. We describe the two that we will refer to in this chapter:

- “*Bijjective argument*”: In terms of (1), the most direct is to establish a bijection between pairs (A, B) of semistandard tableau of shape λ and μ respectively and pairs (C, D) where $C \in \text{SYT}(\nu/\lambda)$ such that $\text{Rect}(C) = S_\mu$ and D is a semistandard tableau of shape ν . This can be achieved using the *Robinson–Schensted correspondence*.
- “*Associativity argument*”: This was used by A. Knutson–T. Tao–C. Woodward [35] and A. Buch–A. Kresch–H. Tamvakis [12]. Define a putative ring $(R, +, \star)$ with additive basis

$\{[\lambda] : \lambda \subseteq \Lambda\}$ and product

$$[\lambda] \star [\mu] := \sum_{\nu \subseteq \Lambda} \bar{c}_{\lambda, \mu}^{\nu} [\nu],$$

where $\bar{c}_{\lambda, \mu}^{\nu}$ is a collection of nonnegative integers. Assume one can prove \star is commutative and associative and moreover agrees with Pieri's rule, i.e., $\bar{c}_{\lambda, (p)}^{\nu} = 0$ unless ν/λ is a horizontal strip of size p , and equals 1 otherwise. Then it follows that $c_{\lambda, \mu}^{\nu} = \bar{c}_{\lambda, \mu}^{\nu}$. One can apply this to prove Theorem 3.5. The proofs of commutativity and associativity use Theorem 3.1. Typically, the hard step is the proof of associativity, which explains the nomenclature for this proof technique. \square

Another formulation of the Littlewood–Richardson rule is in terms of semistandard Young tableaux of shape ν/λ and content μ . These are fillings T of ν/λ with μ_i many i 's, and such that the rows are weakly increasing and columns are strictly increasing. The *row reading word* is obtained by reading the entries of T along rows, from right to left and from top to bottom. Such a word $(w_1, w_2, \dots, w_{|\nu/\lambda|})$ is *ballot* if for every fixed $i, k \geq 1$,

$$\#\{j \leq k : w_j = i\} \geq \#\{j \leq k : w_j = i + 1\}.$$

A tableau is ballot if its reading word is ballot.

Theorem 3.7 (Ballot version of the Littlewood–Richardson rule). $c_{\lambda, \mu}^{\nu}$ equals the number of semistandard tableaux of shape ν/λ and content μ that are ballot.

Example 3.8. Suppose $\lambda = (3, 2, 1), \mu = (3, 2, 1), \nu = (4, 4, 3, 1)$. Below are the 3 semistandard tableaux of shape ν/λ and content μ that are ballot.

			1
		1	2
	1	2	
3			

			1
		1	2
	1	3	
2			

			1
		1	2
	2	3	
1			

Proof sketch for Theorem 3.7: Given a semistandard tableau T , one creates a standard tableau T' of the same shape by replacing all μ_1 many 1's by $1, 2, 3, \dots, \mu_1$ from left to right and then replacing the (original) μ_2 many 2's by $\mu_1 + 1, \mu_1 + 2, \dots, \mu_1 + \mu_2$ etc. This process is called *standardization*. Standardizing the tableaux in Example 3.8 respectively gives:

			3
		2	5
	1	4	
6			

			3
		2	5
	1	6	
4			

			3
		2	5
	4	6	
1			

We claim standardization induces a bijection between the rules of Theorem 3.7 and Theorem 3.5. More precisely, if T is furthermore ballot, then T' satisfies $\text{rect}(T') = S_{\mu}$. We leave it as an exercise to establish this, e.g., by induction on $|\lambda|$. \square

There is a polytopal description of the Littlewood–Richardson rule derivable from Theorem 3.7. We first learned this from a preprint version of [46]. Suppose T is a semistandard tableau of shape ν/λ . Set

$$r_k^i = r_k^i(T) = \#\{k' \text{'s in the } i\text{th row of } T\}.$$

Let μ denote the content of T and $\ell(\mu)$ be the number of nonzero parts of μ . By convention, let $r_{\ell(\mu)+1}^i = 0, r_k^{\ell(\mu)+1} = 0$.

Now consider the following linear inequalities, constructed to describe the tableaux from Theorem 3.7 that are counted by $c_{\lambda,\mu}^\nu$.

- (A) Non-negativity: $r_k^i \geq 0, \forall i, k$.
- (B) Shape constraints: $\lambda_i + \sum_k r_k^i = \nu_i, \forall i$.
- (C) Content constraints: $\sum_i r_k^i = \mu_k, \forall k$.
- (D) Tableau constraints: $\lambda_{i+1} + \sum_{j \leq k} r_j^{i+1} \leq \lambda_i + \sum_{j' < k} r_{j'}^i, \forall i$.
- (E) Ballot constraints: $\sum_{i' < i} r_k^{i'} \geq r_{k+1}^i + \sum_{i' < i} r_{k+1}^{i'}, \forall i, k$.

Define a polytope

$$\mathcal{P}_{\lambda,\mu}^\nu = \{(r_k^i) : \text{(A)–(E)}\} \subseteq \mathbb{R}^{\ell(\nu) \cdot \ell(\mu)}.$$

The following is a straightforward exercise, once one assumes Theorem 3.7:

Theorem 3.9 (Polytopal Littlewood–Richardson rule). $c_{\lambda,\mu}^\nu$ counts the number of integer lattice points in $\mathcal{P}_{\lambda,\mu}^\nu$.

Example 3.10. Using λ, μ, ν as in Example 3.8, $\mathcal{P}_{\lambda,\mu}^\nu$ has 3 integer lattice points $R(T_j) = (r(T_j)_{ik}) = (r_k^i(T_j))$ below. Each $r_k^i(T_j) = \#\{k' \text{'s in the } i\text{th row of } T_j\}$, where the T_j are as in Example 3.8.

$$R(T_1) = \begin{pmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ 1 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad R(T_2) = \begin{pmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} \quad R(T_3) = \begin{pmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ 0 & 1 & 1 \\ 1 & 0 & 0 \end{pmatrix}$$

Another conversation concerns $\text{LR}_r = \{(\lambda, \mu, \nu) : \ell(\lambda), \ell(\mu), \ell(\nu) \leq r : c_{\lambda,\mu}^\nu \neq 0\}$.

Corollary 3.11. LR_r is a semigroup, i.e., if $(\lambda, \mu, \nu), (\alpha, \beta, \gamma) \in \text{LR}_r$, then $(\lambda + \alpha, \mu + \beta, \nu + \gamma) \in \text{LR}_r$.

Proof. Since $(\lambda, \mu, \nu), (\alpha, \beta, \gamma) \in \text{LR}_r$, by Theorem 3.9 there exists lattice points $(r_k^i) \in \mathcal{P}_{\lambda,\mu}^\nu$ and $(\bar{r}_k^i) \in \mathcal{P}_{\alpha,\beta}^\gamma$. By examination of the inequalities (A)–(E), clearly $(r_k^i + \bar{r}_k^i)$ is a lattice point in $\mathcal{P}_{\lambda+\alpha,\mu+\beta}^{\nu+\gamma}$, and we are done by another application of Theorem 3.9. \square

This *Littlewood–Richardson semigroup* LR_r is discussed in A. Zelevinsky’s article [74]. That work concerns the Horn and saturation conjectures (we will discuss these in Section 5). The point that a polytopal rule for $c_{\lambda,\mu}^\nu$ implies the semigroup property already appears in *ibid*. It is also true that LR_r is *finitely generated*. This is proved in A. Elashvili’s [15], who credits the argument to M. Brion–F. Knop from “August–September, 1989”. This argument (which applies more generally to tensor product multiplicities of any reductive Lie group) is not combinatorial. For another demonstration, see the proof of Proposition 6.8. Despite subsequent advances in understanding Littlewood–Richardson coefficients, the following remains open:

Problem 3.12 (cf. Problems A and C of [74]). Explicitly give a finite (minimal) list of generators of LR_r .

In connection to the work of Section 5, a closely related problem has been solved by P. Belkale [9]. His paper determines the extremal rays of the rational polyhedral cone defined by the points of LR_r .

4. EDGE LABELED TABLEAUX AND JEU DE TAQUIN

The history of the combinatorics of $C_{\lambda,\mu}^\nu$ is interesting in its own right. The first combinatorial rule for $C_{\lambda,\mu}^\nu$ is due to A. Molev-B. Sagan [43], who solved an even more general problem. Under the obvious specialization, this rule is not positive in the sense of Theorem 2.1. The first such rule, in terms of *puzzles*, was found by A. Knutson and T. Tao [33]. Subsequently, visibly equivalent tableaux rules were independently found by V. Kreiman [39] and A. Molev [42]. Later, P. Zinn-Justin [75] studied the puzzle rule of [33] based on the quantum integrability of the tiling model that underlies puzzles. Coming full circle, a point made in [75, Section 6.5] is that the rule of [43] *does* provide a positive rule after all, under the “curious identity” of [75, Section 6.4].

This work takes a different view. It is about the *edge labeled tableaux* introduced by H. Thomas and the third author [66]. A *horizontal edge* of ν/λ is an east-west line segment which either lies along the lower boundary of λ or ν , or which separates two boxes of ν/λ . An *equivariant filling* of ν/λ is an assignment of elements of $[N] := \{1, 2, \dots, N\}$ to the boxes of ν/λ or to a horizontal edge of ν/λ . While every box contains a single label, each horizontal edge holds an element of $2^{[N]}$. An equivariant filling is *standard* if every label in $[N]$ appears exactly once and moreover any box label is:

- strictly smaller than any label in its southern edge and the label in the box immediately below it;
- strictly larger than any label in its northern edge and the label in the box immediately above it; and
- weakly smaller than the label in the box immediately to its right.

(No condition is placed on the labels of adjacent edges.) Let $\text{EqSYT}(\nu/\lambda, \ell)$ be the set of equivariant standard tableaux with entries from $[N]$.

Example 4.1. Let $\nu/\lambda = (4, 3, 2)/(3, 2, 1) \subseteq \Lambda = 3 \times 4$ and

$$T = \begin{array}{|c|c|c|c|} \hline & & & 5 \\ \hline & & 1, 2 & \\ & & 4 & \\ \hline & & & \\ \hline & 6 & & \\ & 3 & & \\ \hline \end{array}$$

Then $T \in \text{EqSYT}(\nu/\lambda, 6)$. □

As with ordinary tableaux, a rectification of $T \in \text{EqSYT}(\nu/\lambda, N)$ is defined iteratively. Given an inner corner c and $T \in \text{EqSYT}(\nu/\lambda, N)$, if none of the following possibilities applies, terminate the slide. Otherwise use the unique applicable case (in the left tableaux listed to the below, c contains the \bullet):

$$(J1) \begin{array}{|c|c|} \hline \bullet & a \\ \hline b & \\ \hline \end{array} \mapsto \begin{array}{|c|c|} \hline b & a \\ \hline \bullet & \\ \hline \end{array} \quad (\text{if } b < a, \text{ or } a \text{ does not exist})$$

$$(J2) \begin{array}{|c|c|} \hline \bullet & a \\ \hline b & \\ \hline \end{array} \mapsto \begin{array}{|c|c|} \hline a & \bullet \\ \hline b & \\ \hline \end{array} \quad (\text{if } a < b, \text{ or } b \text{ does not exist})$$

$$(J3) \begin{array}{|c|c|} \hline \bullet & a \\ \hline S & \\ \hline \end{array} \mapsto \begin{array}{|c|c|} \hline a & \bullet \\ \hline S & \\ \hline \end{array} \quad (\text{if } a < \min(S))$$

$$(J4) \begin{array}{|c|c|} \hline \bullet & a \\ \hline S & \\ \hline \end{array} \mapsto \begin{array}{|c|c|} \hline s & a \\ \hline S & \\ \hline \end{array} \quad (\text{if } s := \min(S) < a \text{ and } S' := S \setminus \{s\})$$

This *equivariant jeu de taquin slide* into c is denoted by $\text{Ejdt}_c(T)$. Clearly, $\text{Ejdt}_c(T)$ is also a standard equivariant filling.

The rectification of T , denoted $\text{Erect}(T)$, is the result of successively using Ejdt_c by choosing c that is eastmost among all choices of inner corners at each stage.

Example 4.2. Continuing Example 4.1, we use “•” to indicate the boxes that are moved into during $\text{Erect}(T)$. The rectification of the third column is as follows:

$$(11) \quad \begin{array}{|c|c|c|c|} \hline & & \bullet & 5 \\ \hline & & 1, 2 & \\ \hline & & 4 & \\ \hline & 6 & & \\ \hline \end{array} \mapsto \begin{array}{|c|c|c|c|} \hline & & 1 & 5 \\ \hline & & 2 & \\ \hline & & 4 & \\ \hline & 6 & & \\ \hline \end{array}$$

The rectification of the second column given by:

$$(12) \quad \begin{array}{|c|c|c|c|} \hline & & 1 & 5 \\ \hline & & 2 & \\ \hline & \bullet & 4 & \\ \hline & 6 & & \\ \hline \end{array} \mapsto \begin{array}{|c|c|c|c|} \hline & \bullet & 1 & 5 \\ \hline & 4 & 2 & \\ \hline & 6 & & \\ \hline \end{array} \mapsto \begin{array}{|c|c|c|c|} \hline & 1 & \bullet & 5 \\ \hline & 4 & 2 & \\ \hline & 6 & & \\ \hline \end{array} \mapsto \begin{array}{|c|c|c|c|} \hline & 1 & 2 & 5 \\ \hline & 4 & & \\ \hline & 6 & & \\ \hline \end{array}$$

and finally the rectification of the first column given by:

$$(13) \quad \begin{array}{|c|c|c|c|} \hline & 1 & 2 & 5 \\ \hline & 4 & & \\ \hline \bullet & 6 & & \\ \hline \end{array} \mapsto \begin{array}{|c|c|c|c|} \hline & 1 & 2 & 5 \\ \hline \bullet & 4 & & \\ \hline 3 & 6 & & \\ \hline \end{array} \mapsto \begin{array}{|c|c|c|c|} \hline & 1 & 2 & 5 \\ \hline 3 & 4 & & \\ \hline \bullet & 6 & & \\ \hline \end{array} \mapsto \begin{array}{|c|c|c|c|} \hline \bullet & 1 & 2 & 5 \\ \hline 3 & 4 & & \\ \hline 6 & & & \\ \hline \end{array} \mapsto \dots \mapsto \begin{array}{|c|c|c|c|} \hline 1 & 2 & 5 & \\ \hline 3 & 4 & & \\ \hline 6 & & & \\ \hline \end{array}$$

Here the “ $\mapsto \dots \mapsto$ ” refers to slides moving the • right in the first row. \square

We now define $\text{Ejdtwt}(T) \in \mathbb{Z}[t_1, \dots, t_n]$ for a standard tableau T . Each box x in $\Lambda = k \times (n - k)$ has a (Manhattan) *distance* from the lower-left box: suppose x has matrix coordinates (i, j) , then

$$\text{dist}(x) := k + j - i.$$

Next, assign $x \in \Lambda$ the weight $\beta(x) = t_{\text{dist}(x)} - t_{\text{dist}(x)+1}$.

If after rectification of a column, the label l still remains an edge label, $\text{Ejdtfactor}(l)$ is declared to be zero. Otherwise, say an edge label l passes through a box x if it occupies x during the equivariant rectification of the column of T in which l begins. Let the boxes passed be x_1, x_2, \dots, x_s . Also, when the rectification of a column is complete, suppose the filled boxes strictly to the right of x_s are y_1, \dots, y_t . Set

$$\text{Ejdtfactor}(l) = (\beta(x_1) + \beta(x_2) + \dots + \beta(x_s)) + (\beta(y_1) + \beta(y_2) + \dots + \beta(y_t)).$$

Notice that since the boxes $x_1, \dots, x_s, y_1, \dots, y_t$ form a hook inside ν , $\text{Ejdtfactor}(i) = t_e - t_f$ with $e < f$. Now define

$$\text{Ejdtwt}(T) := \prod_l \text{Ejdtfactor}(l),$$

where the product is over all edge labels l of T .

Theorem 4.3 (Edge labeled jeu de taquin rule [66]).

$$C_{\lambda, \mu}^\nu = \sum_T \text{Ejdtwt}(T),$$

where the sum is over all $T \in \text{EqSYT}(\nu/\lambda, |\mu|)$ such that $\text{Erect}(T) = S_\mu$.

Since each $\text{Ejdtfactor}(l)$ is a positive sum of the indeterminates $\beta_i = t_i - t_{i+1}$, Theorem 4.3 expresses $C'_{\lambda,\mu}$ as a polynomial with positive coefficients in the β_i 's, in agreement with Theorem 2.1.

Example 4.4. Below each $x \in \Lambda = 2 \times 3$ is filled with $\beta(x)$:

$t_2 - t_3$	$t_3 - t_4$	$t_4 - t_5$
$t_1 - t_2$	$t_2 - t_3$	$t_3 - t_4$

T_1, T_2 below are those $T \in \text{EqSYT}((3, 3)/(2, 2), 3)$ such that $\text{Erect}(T) = S_{(2,1)}$ with nonzero weight:

 $T_1 =$

		2
1		3

 $T_2 =$

		2
	1	3

 $T_3 =$

		$\frac{1}{2}$
		$\frac{3}{3}$

While T_3 also satisfies $\text{Erect}(T) = S_{(2,1)}$, it has weight zero since there is an edge label when one has completed rectifying the third column.

Theorem 4.3 asserts

$$\begin{aligned} C_{(2,2),(2,1)}^{(3,3)} &= \text{Ejdtwt}(T_1) + \text{Ejdtwt}(T_2) \\ &= [((t_1 - t_2) + (t_2 - t_3)) + (t_3 - t_4)] + [((t_2 - t_3) + (t_3 - t_4)) + (t_4 - t_5)] \\ &= (t_2 - t_5) + (t_1 - t_4). \end{aligned}$$

In order to prove Theorem 4.3, one wishes to adapt the general strategy indicated at the end of Section 2.3. If we define $\overline{C}'_{\lambda,\mu} := \sum_T \text{Ejdtwt}(T)$ (as in Theorem 4.3) and show this collection of numbers satisfies the associativity recurrence, one shows $C'_{\lambda,\mu} = \overline{C}'_{\lambda,\mu}$, as desired.⁴

The rule of Theorem 4.3 appears to be too “rigid” to carry out this strategy. Instead, in [66], a more “flexible version” in terms of semistandard edge labeled tableaux is introduced, together with a corresponding collection of jeu de taquin slides. While we do not wish to revisit the rather technical list of slide rules here, one of the consequences is a ballot rule for $C'_{\lambda,\mu}$, generalizing Theorem 3.7, which we explain now.

An equivariant tableau is *semistandard* if the box labels weakly increase along rows (left to right), and all labels strictly increase down columns. A single edge may be labeled by a set of integers, without repeats; the smallest of them must be strictly greater than the label of the box above, and the largest must be strictly less than the label of the box below.

Example 4.5. Below is an equivariant semistandard Young tableau on $(4, 2, 2)/(2, 1)$.

		$\frac{1}{2,3}$	$\frac{1}{2}$
	1		
2	3		

The content of this tableau is $(3, 3, 2)$. □

⁴It would be quite interesting to give a “bijective argument” (in the sense of Section 3) using the combinatorial description of the factorial Schur polynomials.

Let $\text{EqSSYT}(\nu/\lambda)$ be the set of all equivariant semistandard Young tableaux of shape ν/λ . A tableau $T \in \text{EqSSYT}(\nu/\lambda)$ is *ballot* if, for every column c and every label ℓ ,

$$(\# \ell\text{'s weakly right of column } c) \geq (\#(\ell + 1)\text{'s weakly right of column } c).$$

Given a tableau $T \in \text{EqSSYT}(\nu/\lambda)$, a (box or edge) label ℓ is *too high* if it appears weakly above the upper edge of a box in row ℓ . In the above example, all edge labels are too high. (When there are no edge labels, the semistandard and ballot conditions imply no box label is too high, but in general the three conditions are independent.)

Suppose an edge label ℓ lies on the bottom edge of a box x in row i . Let $\rho_\ell(x)$ be the number of times ℓ appears as a (box or edge) label strictly to the right of x . We define

$$(14) \quad \text{Eballotfactor}(\ell, x) = t_{\text{dist}(x)} - t_{\text{dist}(x)+i-\ell+1+\rho_\ell(x)}.$$

When the edge label is not too high, this is always of the form $t_p - t_q$, for $p < q$. (In particular, it is nonzero.) Finally, define

$$(15) \quad \text{Eballotwt}(T) = \prod \text{Eballotfactor}(\ell, x),$$

the product being over all edge labels ℓ .

Theorem 4.6 (Edge labeled ballot rule [66, Theorem 3.1]). $C_{\lambda, \mu}^\nu = \sum_T \text{Eballotwt}(T)$, where the sum is over all $T \in \text{EqSSYT}(\nu/\lambda)$ of content μ that are ballot and have no label which is too high.

Example 4.7. In Example 4.4, we saw $C_{(2,2),(2,1)}^{(3,3)} = (t_2 - t_5) + (t_1 - t_4)$. Below has $x \in \Lambda$ is filled with $\text{dist}(x)$.

2	3	4
1	2	3

We see T_1, T_2 below are those $T \in \text{EqSSYT}((3,3)/(2,2))$ of content $(2,1)$ that are ballot and have no label which is too high.

$$T_1 = \begin{array}{|c|c|c|} \hline & & 1 \\ \hline & & 2 \\ \hline 1 & & \\ \hline \end{array} \quad T_2 = \begin{array}{|c|c|c|} \hline & & 1 \\ \hline & & 2 \\ \hline & 1 & \\ \hline \end{array}$$

Thus we see as Theorem 4.3 states, since for both cases $p_\ell(x) = p_1(x) = 1$,

$$\begin{aligned} C_{(2,2),(2,1)}^{(3,3)} &= \text{Eballotwt}(T_1) + \text{Eballotwt}(T_2) \\ &= \text{Eballotfactor}(1, (2,1)) + \text{Eballotfactor}(1, (2,2)) \\ &= (t_1 - t_{1+2-1+1+1}) + (t_2 - t_{2+2-1+1+1}) \\ &= (t_1 - t_4) + (t_2 - t_5), \end{aligned}$$

so this rule agrees.

5. NONVANISHING OF LITTLEWOOD–RICHARDSON COEFFICIENTS, SATURATION AND HORN INEQUALITIES

For which triples of partitions (λ, μ, ν) does $c_{\lambda, \mu}^\nu \neq 0$?

The importance of this question comes from a striking equivalence to a 19th century question about linear algebra. This equivalence was first suggested as a question by R. C. Thompson in the 1970s; see R. Bhatia's survey [10, pg. 308] (another survey on this topic is W. Fulton's [20]). Suppose A, B, C are three $r \times r$ Hermitian matrices and $\lambda, \mu, \nu \in \mathbb{R}^r$ are the respective lists of eigenvalues (written in decreasing order). The *eigenvalue problem for Hermitian matrices* asks

Which eigenvalues (λ, μ, ν) can occur if $A + B = C$?

After work of H. Weyl, K. Fan, V. B. Lidskii-H. Weilandt and others, A. Horn recursively defined a list of inequalities on triples $(\lambda, \mu, \nu) \in \mathbb{R}^{3r}$ (see Theorem 5.2(2) below). He conjectured that these give a complete solution to the eigenvalue problem [25]. That these inequalities (or equivalent ones) are necessary has been proved by several authors, including B. Totaro [68] and A. Klyachko [30]. A. Klyachko also established that his list of inequalities is sufficient, giving the first solution to the eigenvalue problem.

Also, A. Klyachko showed that his inequalities give an asymptotic solution to the problem of which Littlewood–Richardson coefficients $c_{\lambda, \mu}^{\nu}$ are nonzero. That is, suppose λ, μ, ν are partitions with at most r parts. A. Klyachko proved that if $c_{\lambda, \mu}^{\nu} \neq 0$, then $(\lambda, \mu, \nu) \in \mathbb{Z}_{\geq 0}^{3r}$ satisfies his inequalities. Conversely, he showed that if $(\lambda, \mu, \nu) \in \mathbb{Z}_{\geq 0}^{3r}$ satisfy his inequalities then $c_{N\lambda, N\mu}^{N\nu} \neq 0$ for some $N \in \mathbb{N}$. (Here, $N\lambda$ is the partition with each part of λ stretched by a factor of N .) Subsequently, A. Knutson-T. Tao [32] sharpened the last statement, and established:

Theorem 5.1 (Saturation theorem). *$c_{\lambda, \mu}^{\nu} \neq 0$ if and only if $c_{N\lambda, N\mu}^{N\nu} \neq 0$ for any $N \in \mathbb{N}$.*

Combined with [30], it follows that A. Klyachko's solution agrees with A. Horn's conjectured solution.⁵ Let $[r] := \{1, 2, \dots, r\}$. For any

$$I = \{i_1 < i_2 < \dots < i_d\} \subseteq [r]$$

define the partition

$$\tau(I) := (i_d - d \geq \dots \geq i_2 - 2 \geq i_1 - 1).$$

This bijects subsets of $[r]$ of cardinality d with partitions whose Young diagrams are contained in a $d \times (r - d)$ rectangle. The following combines the main results of [30, 32]:

Theorem 5.2. ([30], [32]) *Let λ, μ, ν be partitions with at most r parts such that*

$$(16) \quad |\lambda| + |\mu| = |\nu|.$$

The following are equivalent:

- (1) $c_{\lambda, \mu}^{\nu} \neq 0$.
- (2) *For every $d < r$, and every triple of subsets $I, J, K \subseteq [r]$ of cardinality d such that $c_{\tau(I), \tau(J)}^{\tau(K)} \neq 0$, we have*

$$(17) \quad \sum_{i \in I} \lambda_i + \sum_{j \in J} \mu_j \geq \sum_{k \in K} \nu_k.$$

⁵The proof in a preprint version of [32] used the polytopal Littlewood–Richardson rule of Bernstein–Zelevinsky; we refer to the survey of A. Buch [11]. The published proof is formulated in terms of the *Honeycomb model*. It is an easy exercise to prove the “ \Rightarrow ” of the equivalence using either of the Littlewood–Richardson rules found in Section 3. A proof of the converse using such rules would be surprising. Another solution, due to H. Derksen-J. Weyman [14] was given in the setting of *semi-invariants of quivers*.

(3) There exist $r \times r$ Hermitian matrices A, B, C with eigenvalues λ, μ, ν such that $A+B = C$.

Remark 5.3. The logic of the proof of Theorem 5.2 in [32] is to show Theorem 5.1. In fact, Theorem 5.1 also follows from the equivalence (1) \iff (2) of Theorem 5.2. This is since the Horn inequalities from (2) are homogeneous. This point seems to have been first noted in P. Belkale’s [8] which moreover gives a geometric proof of Theorem 5.2. \square

Remark 5.4. The equivalence (1) \iff (2) immediately implies the semigroup property of LR_r (Corollary 3.11). \square

Remark 5.5. P. Belkale’s doctoral thesis [6] (published in [7]) shows that a much smaller list of inequalities than those in Theorem 5.2(2) suffice. Namely, replace the condition “ $c_{\tau(\lambda), \tau(\mu)}^{\tau(\nu)} \neq 0$ ” with “ $c_{\tau(\lambda), \tau(\mu)}^{\tau(\nu)} \neq 1$ ”. A. Knutson-T. Tao-C. Woodward [34] showed that the inequalities in this shorter list are minimal, *i.e.*, none can be dispensed with. \square

Remark 5.6. Theorem 5.2 gives a different proof that LR_r is finitely generated as a semigroup. We will give the argument in the proof of Proposition 6.8, below.

While Theorem 5.2 characterizes nonvanishing of $c_{\lambda, \mu}^{\nu}$, the inequalities are recursive and non-transparent to work with. The Littlewood–Richardson rules of Section 3 require one to search for a valid tableau in a possibly large search space. K. Purbhoo [51] (see also [52]) developed a general and intriguing *root game*, which in the case of Grassmannians can be “won” if and only if $c_{\lambda, \mu}^{\nu} \neq 0$.

Theorem 5.1 permits a determination of the *formal* computational complexity of the nonvanishing decision problem “ $c_{\lambda, \mu}^{\nu} \neq 0$ ” (in the bit length of the input (λ, μ, ν) , where one assumes arithmetic operations take constant time). This was resolved independently by T. McAllister-J. De Loera [13] and K. D. Mulmuley-H. Narayanan-M. Sohoni, [46], by a neat argument that combines Theorem 5.1 with celebrated developments in linear programming:

Theorem 5.7. *The decision problem of determining if $c_{\lambda, \mu}^{\nu} \neq 0$ is in the class P of polynomial problems.*

Proof. By Theorem 3.9, $c_{\lambda, \mu}^{\nu} \neq 0$ if and only if the polytope $\mathcal{P}_{\lambda, \mu}^{\nu}$ has a lattice point. Clearly, if $\mathcal{P}_{\lambda, \mu}^{\nu} \neq \emptyset$, it has a rational vertex. In this case, a dilation $N\mathcal{P}_{\lambda, \mu}^{\nu}$ contains a lattice point. One checks from the definitions that $N\mathcal{P}_{\lambda, \mu}^{\nu} = \mathcal{P}_{N\lambda, N\mu}^{N\nu}$ which means $c_{N\lambda, N\mu}^{N\nu} \neq 0$. Thus, by Theorem 5.1,

$$c_{\lambda, \mu}^{\nu} \neq 0 \iff c_{N\lambda, N\mu}^{N\nu} \neq 0 \iff \mathcal{P}_{\lambda, \mu}^{\nu} \neq \emptyset.$$

To determine if $\mathcal{P}_{\lambda, \mu}^{\nu} \neq \emptyset$, one needs to decide feasibility of any linear programming problem involving $\mathcal{P}_{\lambda, \mu}^{\nu}$. One appeals to ellipsoid/interior point methods for polynomiality.⁶ Actually, our inequalities are of the form $Ax \leq b$ where the vector b is integral and the entries of A are from $\{-1, 0, 1\}$. Hence our polytope is *combinatorial* and so one achieves a strongly polynomial time complexity using É. Tardos’ algorithm; see [64, 23]. \square

In contrast, H. Narayanan [47] proved that counting $c_{\lambda, \mu}^{\nu}$ is a #P-complete problem in L. Valiant’s complexity theory of counting problems [71]. In particular, this means that no polynomial time algorithm for computing $c_{\lambda, \mu}^{\nu}$ can exist unless $P = NP$ (it is widely

⁶The Klee–Minty cube shows that the practically efficient simplex method has exponential worst-case complexity.

expected that $P \neq NP$). It is curious that the counting problem is (presumably) hard, whereas the nonzeroness version is polynomial time. This already occurs for the original #P-complete problem from [71], i.e., to compute the permanent of an $n \times n$ matrix $M = (m_{ij})$ where $m_{ij} \in \{0, 1\}$. Now, determining if $\text{per}(M) > 0$ is equivalent to deciding the existence of matching in a bipartite graph that has incidence matrix M ; the algorithm of J. Edmonds-R. Karp provides the polynomial-time algorithm.

6. EQUIVARIANT NONVANISHING, SATURATION, AND FRIEDLAND'S INEQUALITIES

We now turn to the equivariant analogues of results from the previous section.

For which triples of partitions (λ, μ, ν) does $C_{\lambda, \mu}^\nu \neq 0$?

In [3] it was shown that this Schubert calculus question is also essentially equivalent to an eigenvalue problem. Recall that a Hermitian matrix M *majorizes* another Hermitian matrix M' if $M - M'$ is positive semidefinite (its eigenvalues are all nonnegative). In this case, we write $M \geq M'$. S. Friedland [17] studied the following question:

Which eigenvalues (λ, μ, ν) can occur if $A + B \geq C$?

His solution, given as linear inequalities, includes Klyachko's inequalities, a trace inequality and some extra inequalities. Later, W. Fulton [19] proved the extra inequalities are unnecessary (where his inequalities are those of Theorem 6.2(2) below), leading to a natural extension of the equivalence (2) \iff (3) of Theorem 5.2.

One would like an extension of the equivalence with (1) of Theorem 5.2 as well. Now, D. Anderson, E. Richmond and the third author [3] proved:

Theorem 6.1 (Equivariant saturation). $C_{\lambda, \mu}^\nu \neq 0$ if and only if $C_{N\lambda, N\mu}^{N\nu} \neq 0$ for any $N \in \mathbb{N}$.⁷

Actually, Theorem 6.1 is proved by establishing the equivalence (1) \iff (2) below.

Theorem 6.2 ([3], [17], [19]). *Let λ, μ, ν be partitions with at most r parts such that*

$$(18) \quad |\lambda| + |\mu| \geq |\nu| \text{ and } \max\{\lambda_i, \mu_i\} \leq \nu_i \text{ for all } i \leq r.$$

The following are equivalent:

- (1) $C_{\lambda, \mu}^\nu \neq 0$.
- (2) *For every $d < r$, and every triple of subsets $I, J, K \subseteq [r]$ of cardinality d such that $c_{\tau(I), \tau(J)}^{\tau(K)} \neq 0$, we have*

$$\sum_{i \in I} \lambda_i + \sum_{j \in J} \mu_j \geq \sum_{k \in K} \nu_k.$$

- (3) *There exist $r \times r$ Hermitian matrices A, B, C with eigenvalues λ, μ, ν such that $A + B \geq C$.*

Theorem 6.2 states that the main inequalities controlling nonvanishing of $C_{\lambda, \mu}^\nu$ are just Horn's inequalities (17).

⁷In our notation, $C_{\lambda, \mu}^\nu$ depends on the indices k and n of the Grassmannian $\text{Gr}_k(\mathbb{C}^n)$. However, the edge labeled rule of Theorem 4.3 has the property that $C_{\lambda, \mu}^\nu$ is that for fixed k , the polynomial is independent of the choice of n provided both are sufficiently large so that $\lambda, \mu, \nu \subseteq k \times (n - k)$. Indeed, the coefficients $C_{\lambda, \mu}^\nu$ for such values are the structure constants for the Schubert basis in the graded inverse limit of equivariant cohomology rings under the standard embedding $\iota : \text{Gr}_k(\mathbb{C}^n) \hookrightarrow \text{Gr}_k(\mathbb{C}^{n+1})$; see [3, Section 1.2].

Remark 6.3. The second condition in (18) is unnecessary in Theorem 5.2 since it is already implied by (16) combined with (17). The condition is not required for the equivalence (2) \iff (3). However it is needed for the equivalence with (1). For example, the 1×1 matrices $A = [1], B = [1], C = [0]$ satisfy $A + B \geq C$, but $C_{(1),(1)}^{(0)} = 0$. This is not in contradiction with Theorem 6.2 since $\max\{1, 1\} \leq 0$ is violated. \square

Just as in Remark 5.3, the equivalence (1) \iff (2) and the homogeneity of the Friedland–Fulton inequalities implies Theorem 6.1. On the other hand the equivalence (1) \iff (2) relies on the classical Horn theorem (Theorem 5.2) and the edge labeled ballot tableau rule Theorem 4.6. To give the reader a sense of the proof we now provide:

Sketch of proof that (1) \implies (2): Using Theorem 4.6, it is an exercise to show that

Claim 6.4. *If $C_{\lambda, \mu}^\nu \neq 0$ and $|\nu| < |\lambda| + |\mu|$, then for any s such that $|\nu| - |\lambda| \leq s < |\mu|$, there is a $\mu^\downarrow \subset \mu$ with $|\mu^\downarrow| = s$ and $C_{\lambda, \mu^\downarrow}^\nu \neq 0$.*

Since $C_{\lambda, \mu}^\nu \neq 0$, by the claim (and induction) there exists $\lambda^\downarrow \subseteq \lambda$ such that $|\lambda^\downarrow| + |\mu| = |\nu|$ and $C_{\lambda^\downarrow, \mu}^\nu \neq 0$. This latter number is a classical Littlewood–Richardson coefficient, we can apply Theorem 5.2 to conclude that for any triple (I, J, K) with $c_{\tau(I), \tau(J)}^{\tau(K)} \neq 0$ one has

$$\sum_{i \in I} \lambda_i^\downarrow + \sum_{i \in J} \mu_i \geq \sum_{k \in K} \nu_k.$$

Now we are done since $\sum_{i \in I} \lambda_i \geq \sum_{i \in I} \lambda_i^\downarrow$. \square

The converse (2) \implies (1) uses another exercise that can be proved using Theorem 4.6:

Claim 6.5. *If $C_{\lambda, \mu}^\nu \neq 0$ then $C_{\lambda, \mu^\uparrow}^\nu \neq 0$ for any $\mu \subset \mu^\uparrow \subseteq \nu$.*

See [3, Section 2] for proofs of Claim 6.4 and Claim 6.5.

Remark 6.6. Just as with Theorem 5.2, the list of inequalities in Theorem 6.2(2) contain redundancies. Building from the results discussed in Remark 5.5, W. Fulton [19] shows that one can also replace the “ $c_{\tau(\lambda), \tau(\mu)}^{\tau(\nu)} \neq 0$ ” with “ $c_{\tau(\lambda), \tau(\mu)}^{\tau(\nu)} = 1$ ”. W. Fulton’s work shows that if the second condition in (18) is ignored, the inequalities are minimal for the equivalence (2) \iff (3). \square

Let $\text{EqLR}_r = \{(\lambda, \mu, \nu) : \ell(\lambda), \ell(\mu), \ell(\nu) \leq r, C_{\lambda, \mu}^\nu \neq 0\}$.

Corollary 6.7. *EqLR_r is a semigroup.*

Proof. Just as Corollary 3.11 clearly follows from the first two equivalences of Theorem 5.2 (Remark 5.4), the present claim holds by the first two equivalences of Theorem 6.2. However, we can also prove this directly from Claims 6.4 and 6.5: By applying Claim 6.4 there exists $\lambda^\circ \subseteq \lambda$ and $\alpha^\circ \subseteq \alpha$ such that $C_{\lambda^\circ, \mu}^\nu = c_{\lambda^\circ, \mu}^\nu \neq 0$ and $C_{\alpha^\circ, \beta}^\gamma = c_{\alpha^\circ, \beta}^\gamma \neq 0$. By Corollary 3.11, $(\lambda^\circ + \alpha^\circ, \mu + \beta, \nu + \gamma) \in \text{LR}_r$. Now, since $\lambda^\circ + \alpha^\circ \subseteq \lambda + \alpha \subseteq \nu + \gamma$, we can apply Claim 6.5 to conclude $(\lambda + \alpha, \mu + \beta, \nu + \gamma) \in \text{EqLR}_r$, as desired. \square

Proposition 6.8. *EqLR_r is finitely generated.*

The argument we give is based on discussion with S. Fomin and A. Knutson. It applies *mutatis mutandis* to prove that LR_r is finitely generated:

Proof. Since the inequalities from Theorem 6.2 are finite in number, and each inequality has its bounding hyperplane containing the origin, the set $\mathcal{C} \subseteq \mathbb{R}^{3r}$ they define is a polyhedral cone. Since the inequalities have rational coefficients, by definition, \mathcal{C} is rational. Moreover, \mathcal{C} is also clearly pointed, i.e., $\mathcal{C} \cap -\mathcal{C} = \{0\}$. Now apply [57, Theorem 16.4]. \square

Naturally, one would like a solution for the generalization of Problem 3.12 to EqLR_r .

Using Theorem 6.1 and Theorem 4.6, A. Adve together with the first and third authors [3] prove a generalization of Theorem 5.7:

Theorem 6.9 ([3]). *The decision problem of determining $C_{\lambda,\mu}^\nu \neq 0$ is in the class P of polynomial problems.*

Sketch of proof: Using Theorem 5.7 one can construct a polytope $\mathcal{Q}_{\lambda,\mu}^\nu$ analogous to $\mathcal{P}_{\lambda,\mu}^\nu$. The main property is that $C_{\lambda,\mu}^\nu \neq 0$ if and only if $\mathcal{Q}_{\lambda,\mu}^\nu$ contains a lattice point (in particular, in contrast to Theorem 3.9, the *number* of lattice points of $\mathcal{Q}_{\lambda,\mu}^\nu$ is not well-understood). The remainder of the proof proceeds exactly as in the proof of Theorem 5.7, except that we use Theorem 6.1 in place of Theorem 5.1. \square

For the remainder of this section, assume $C_{\lambda,\mu}^\nu$ is expressed (uniquely) as a polynomial in the variables $\beta_i := t_i - t_{i+1}$. We now state a few open problems/conjectures introduced in [55], for the special case of Grassmannians.

A refinement of the nonvanishing question is:

Question 6.10. *What is the computational complexity of determining if $[\beta_1^{i_1} \cdots \beta_{n-1}^{i_{n-1}}] C_{\lambda,\mu}^\nu \neq 0$?*

This question concerns the Newton polytope of $C_{\lambda,\mu}^\nu$. Recall, the *Newton polytope* of

$$f = \sum_{(n_1, \dots, n_r) \in \mathbb{Z}_{\geq 0}^r} c_{n_1, \dots, n_r} \prod_{j=1}^r \alpha_j^{n_j} \in \mathbb{R}[\alpha_1, \dots, \alpha_r]$$

is $\text{Newton}(f) := \text{conv}\{(n_1, \dots, n_r) : c_{n_1, \dots, n_r} \neq 0\} \subseteq \mathbb{R}^r$. f has *saturated Newton polytope* (SNP) [45] if $c_{n_1, \dots, n_r} \neq 0 \iff (n_1, \dots, n_r) \in \text{Newton}(f)$.

Conjecture 6.11. $C_{\lambda,\mu}^\nu$ has SNP.

This raises the question:

Problem 6.12. *Give a half space description of $\text{Newton}(C_{\lambda,\mu}^\nu)$.*

A proof of Conjecture 6.11 together with any reasonable solution to Problem 6.12 would imply that the decision problem in Question 6.10 is in the computational complexity class $\text{NP} \cap \text{coNP}$. This would strongly suggest that the decision problem is not NP-complete, and in fact suggest the problem is in P. We refer the reader to [1, Section 1] for elaboration on these points.

7. MAXIMAL ORTHOGONAL AND LAGRANGIAN GRASSMANNIANS

7.1. Goals in the sequel. Beyond Grassmannians $\text{Gr}_k(\mathbb{C}^n)$, the *maximal orthogonal Grassmannians* and *Lagrangian Grassmannians* have been of significant interest. Their classical (non-equivariant) Schubert calculus shares many analogies with the Grassmannian case. They concern the Q -Schur polynomials of I. Schur [59], and the tableau combinatorics of

D. Worley [72], B. Sagan [56] and J. Stembridge [63]. Although these combinatorial results were originally developed to study projective representations of the symmetric group, the connection to Schubert calculus of these spaces was established by P. Pragacz [50].

It is therefore natural to seek extensions of the results from Sections 1-6. Discussion of efforts toward this goal occupy the remainder of this work.

7.2. Definition of the spaces. Consider the two classical Lie groups of non-simply laced type: $G = \mathrm{SO}_{2n+1}(\mathbb{C})$ and $G = \mathrm{Sp}_{2n}(\mathbb{C})$. These are the automorphism groups of a non-degenerate bilinear form $\langle \cdot, \cdot \rangle$. In the former case, $\langle \cdot, \cdot \rangle$ is symmetric, and on $W = \mathbb{C}^{2n+1}$. In the latter case, $\langle \cdot, \cdot \rangle$ is skew-symmetric, and on $W = \mathbb{C}^{2n}$. A subspace $V \subseteq W$ is called *isotropic* if $\langle v_1, v_2 \rangle = 0$ for all $v_1, v_2 \in V$. The maximum dimension of an isotropic subspace of W is n .

Let $Y = \mathrm{OG}(n, 2n+1)$ be the *maximal orthogonal Grassmannian* of n -dimensional isotropic subspaces of \mathbb{C}^{2n+1} ; this space has an action of $G = \mathrm{SO}_{2n+1}(\mathbb{C})$. Similarly, let $Z = \mathrm{LG}(n, 2n)$ be the *Lagrangian Grassmannian* of n -dimensional isotropic subspaces of \mathbb{C}^{2n} . In either case, the (opposite) Borel subgroup $B_- \leq G$ consists of the lower triangular matrices in G . The maximal torus T are the diagonal matrices in G . Just as in the case of the Grassmannian, the corresponding B_- acts on Y (resp. Z) with finitely many orbits Y_λ° (resp. Z_λ°); these are the *Schubert cells*.

In both cases, the Schubert cells and thus Schubert varieties $Y_\lambda = \overline{Y_\lambda^\circ}$ (resp. $Z_\lambda = \overline{Z_\lambda^\circ}$) are indexed by strict partitions fitting inside the shifted staircase

$$\rho_n = (n, n-1, n-2, \dots, 3, 2, 1).$$

A *strict partition* is an integer partition $\lambda = (\lambda_1 > \lambda_2 > \dots > \lambda_\ell > 0)$. Identify λ with its shifted shape, which is the usual Young diagram (in English notation) but where the i -th row from the top is indented $i-1$ many spaces. We refer to, e.g., [28, Section 6] and the references therein for additional details.

Let $\sigma_\lambda(Y) \in H^{2|\lambda|}(Y)$ be the Poincaré dual to Y_λ . These *Schubert classes* form a \mathbb{Z} -linear basis of $H^*(Y)$, and we define the structure constants by

$$\sigma_\lambda(Y) \smile \sigma_\mu(Y) = \sum_{\nu \subseteq \rho_n} o_{\lambda, \mu}^\nu \sigma_\nu(Y).$$

Similarly, the Schubert classes $\sigma_\lambda(Z)$ form a \mathbb{Z} -linear basis of $H^*(Z)$, and

$$\sigma_\lambda(Z) \smile \sigma_\mu(Z) = \sum_{\nu \subseteq \rho_n} l_{\lambda, \mu}^\nu \sigma_\nu(Z).$$

7.3. Schur P - and Q - functions; P. Pragacz's theorem. Let

$$q_r(x_1, \dots, x_n) = 2 \sum_{i=1}^n x_i^r \prod_{i \neq j} \frac{x_i + x_j}{x_i - x_j}.$$

This is clearly symmetric and in fact a polynomial. Next, set

$$Q_{(r,s)} = q_r q_s + 2 \sum_{i=1}^s (-1)^i q_{r+i} q_{s-i}.$$

Recall that the *Pfaffian* of a $2t \times 2t$ skew-symmetric matrix $M = (m_{ij})$ is

$$\text{Pf}(M) = \sum_{\sigma \in \mathfrak{S}_{2t}} \text{sgn}(\sigma) \prod_{i=1}^t m_{\sigma(2i-1), \sigma(2i)},$$

where σ satisfies $\sigma(2i-1) < \sigma(2i)$ for $1 \leq i \leq m$ and $\sigma(1) < \sigma(3) < \dots < \sigma(2i-3) < \sigma(2i-1)$. Then for $\lambda = (\lambda_1 > \lambda_2 > \dots > \lambda_\ell > 0)$, the *Schur Q-function* [59] is

$$(19) \quad Q_\lambda = \text{Pf}(Q_{(\lambda_i, \lambda_j)}).$$

If $\ell(\lambda)$ is odd, we add a 0 at the end of λ . The Schur Q -functions linearly span the subalgebra $\Gamma \subset \text{Sym}$ generated by the q_i 's. In fact, $Q_{(i)} = q_i$. The *Schur P-function* is

$$(20) \quad P_\lambda := 2^{-\ell(\lambda)} Q_\lambda.$$

P. Pragacz [50] proved that the Schur P - and Schur Q -polynomials represent the Schubert classes of Y and Z respectively. That is,

$$H^*(Y) \cong \Gamma/J,$$

where J is the ideal $\langle P_\lambda : \lambda \not\subseteq \rho_n \rangle$, and $\sigma_\lambda(Y)$ maps to $P_\lambda + J$ under this isomorphism. Moreover,

$$P_\lambda P_\mu = \sum_{\nu} o_{\lambda, \mu}^{\nu} P_\nu.$$

Similarly,

$$Q_\lambda Q_\mu = \sum_{\nu} l_{\lambda, \mu}^{\nu} Q_\nu.$$

J. R. Stembridge [63] proved

$$P_\lambda = \sum_T x^T,$$

where the sum is over semistandard fillings of the shifted shape λ . That is, fill each box of λ with a label from the ordered set $1' < 1 < 2' < 2 < 3' < 3 < \dots$ such that the rows and columns are weakly increasing, two i' -s cannot appear in the same row and two i -s cannot appear in the same column. Moreover, there are no primed entries on the main diagonal. For example, if $\lambda = (2, 1)$, the shifted semistandard tableaux are

$$\begin{array}{|c|c|} \hline 1 & 1 \\ \hline & 2 \\ \hline \end{array} \quad \begin{array}{|c|c|} \hline 1 & 2' \\ \hline & 2 \\ \hline \end{array} \quad \begin{array}{|c|c|} \hline 1 & 1 \\ \hline & 3 \\ \hline \end{array} \quad \begin{array}{|c|c|} \hline 1 & 2' \\ \hline & 3 \\ \hline \end{array} \quad \begin{array}{|c|c|} \hline 1 & 2 \\ \hline & 3 \\ \hline \end{array} \quad \begin{array}{|c|c|} \hline 1 & 3' \\ \hline & 3 \\ \hline \end{array} \quad \begin{array}{|c|c|} \hline 2 & 2 \\ \hline & 3 \\ \hline \end{array} \quad \begin{array}{|c|c|} \hline 2 & 3' \\ \hline & 3 \\ \hline \end{array} \quad \dots$$

Hence $P_{(2,1)}(x_1, x_2, x_3) = x_1^2 x_2 + x_1 x_2^2 + x_1^2 x_3 + x_1 x_2 x_3 + x_1 x_2 x_3 + x_1 x_3^2 + x_2^2 x_3 + x_2 x_3^2 + \dots$. If we allow primed entries on the diagonal, we get the Schur Q -function. It is easy to see that this definition satisfies (20).

7.4. Shifted Littlewood–Richardson rules. D. Worley [72] introduced a *jeu de taquin* theory for shifted shapes. A standard tableau T of shifted skew shape ν/λ is a filling of λ with the labels $1, 2, 3, \dots, |\nu/\lambda|$ that is increasing along rows and columns. Let $\text{shSYT}(\nu/\lambda)$ denote the set of these tableaux. The notions of slides and rectification are just as in the unshifted case, using (J1) and (J2). With this, the exact analogues of Theorems 3.1 and 3.3 hold and one can define ShEjdt and shRect etc, in the obvious manner. Indeed, one has the following combinatorial rule for $o_{\lambda, \mu}^{\nu}$:

Theorem 7.1 (Shifted jeu de taquin Littlewood–Richardson rule). *Fix $U \in \text{shSYT}(\mu)$. Then*

$$o_{\lambda, \mu}^{\nu} = \#\{T \in \text{shSYT}(\nu/\lambda) : \text{shRect}(T) = U\}.$$

Example 7.2. Let $\lambda = (3, 1)$, $\mu = (3, 1)$, $\nu = (4, 3, 1)$. The following are the 2 shifted tableaux of shape ν/λ that rectify to $U = \begin{array}{|c|c|c|} \hline 1 & 2 & 3 \\ \hline & & \\ \hline 4 & & \\ \hline \end{array}$. Thus, $o_{\lambda, \mu}^{\nu} = 2$.

			1
		2	3
		4	

			3
		1	4
		2	

Define the *reading word* of a (possibly skew) shifted tableau T to be the word obtained by reading the rows of T from left to right starting with the bottom row. For a word $w = w_1 w_2 \dots w_n$, define $\text{JS}_i(j)$ for $1 \leq j \leq 2n$, $i \geq 1$, depending on w :

$$\text{JS}_i(j) := \text{number of occurrences of } i \text{ among } w_{n-j+1}, \dots, w_n, \quad 0 \leq j \leq n,$$

and

$$\text{JS}_i(n+j) := \text{JS}_i(n) + \text{number of occurrences of } i' \text{ among } w_1, \dots, w_j, \quad 0 < j \leq n.$$

The word w is *proto-ballot* if, when $\text{JS}_i(j) = \text{JS}_{i-1}(j)$, both of these statements hold:

$$\begin{aligned} w_{n-j} &\neq i, i', \text{ if } 0 \leq j < n; \\ w_{j-n+1} &\neq i-1, i', \text{ if } n \leq j < 2n. \end{aligned}$$

Let $|w|$ be the word obtained from w by removing all primes. Now, w is *ballot* if it is *proto-ballot* and the leftmost i of $|w|$ is unprimed in w for all i . In his work on projective representation theory of symmetric groups, J. Stembridge [63, Theorem 8.3] gave the following semistandard analogue of Theorem 7.1.

Theorem 7.3 (Shifted ballot Littlewood–Richardson rule). *$o_{\lambda, \mu}^{\nu}$ counts the number of shifted semistandard tableaux of shape ν/λ of content μ that are ballot.*

Example 7.4. Let $\lambda = (3, 1)$, $\mu = (3, 1)$, $\nu = (4, 3, 1)$, then the following are the only 2 shifted semistandard tableaux of shape ν/λ of content μ that are ballot.

			1'
		1	1
		2	

			1
		1'	2
		1	

It follows from (20) that

$$(21) \quad l_{\lambda, \mu}^{\nu} = 2^{\ell(\nu) - \ell(\lambda) - \ell(\mu)} o_{\lambda, \mu}^{\nu}.$$

Thus the above rules give a rule to compute $l_{\lambda, \mu}^{\nu}$ in a manifestly positive manner, as well.

7.5. Nonvanishing. K. Purbhoo-F. Sottile [53, 54] gave an extension of the Horn recursion to describe when $o_{\lambda, \mu}^{\nu}$ (or equivalently $l_{\lambda, \mu}^{\nu}$) is nonzero. Fix n and r . Suppose α is a (ordinary) partition whose (unshifted) Young diagram is contained in $r \times (n-r)$. Let


$$I_n(\alpha) := \{n-r+1-\alpha_1, n-r+2-\alpha_2, \dots, n-\alpha_r\}.$$

Index the corners of ρ_n top to bottom from 1 to n . For $\lambda \subseteq \rho_n$, $0 < r < n$ and $\alpha \subseteq r \times (n-r)$ let $[\lambda]_{\alpha}$ be the number of boxes of λ that survive after crossing out the rows to the right and columns above the corners indexed by $I_n(\alpha)$. Define λ^c to be the complement of λ in ρ_n (after reflecting), whereas $\alpha^c = \alpha^{\vee}$ is the rotation of the complement of α in $r \times (n-r)$.

Example 7.5. Suppose that $n = 6, r = 3$ and $\alpha = (3, 2, 1)$. Then

$$I_n(\alpha) = \{6 - 3 + 1 - 3, 6 - 3 + 2 - 2, 6 - 3 + 3 - 1\} = \{1, 3, 5\}.$$

Suppose $\lambda = (6, 4, 3, 1) \subseteq \rho_n$. In the figure below, yellow boxes are the ones that are crossed out. Thus, $[\lambda]_\alpha = 3$.



Theorem 7.6 (K. Purbhoo-F. Sottile's theorem). For $\lambda, \mu, \nu \subseteq \rho_n$, $o_{\lambda, \mu}^{\nu^c} \neq 0$ if and only if

- $|\lambda| + |\mu| + |\nu| = \dim Y = \binom{n+1}{2}$, and
- for all $0 < r < n$ and all $\alpha, \beta, \gamma \subset r \times (n-r)$ such that $c_{\alpha, \beta}^{\gamma^c} \neq 0$, one has $[\lambda]_\alpha + [\mu]_\beta + [\nu]_\gamma \leq \binom{n+1-r}{2}$.

Remark 7.7. The obvious analogue of saturation does *not* hold. For example, take $\lambda = (2, 1), \mu = (2), \nu = (3, 2)$. Then $o_{\lambda, \mu}^{\nu} \neq 0$, but $o_{2\lambda, 2\mu}^{2\nu} = o_{(4, 2), (4)}^{(6, 4)} = 0$. \square

Inspired by the complexity results concerning $c_{\lambda, \mu}^{\nu}$, we take this opportunity to pose:

Problem 7.8. Is the decision problem of determining if $o_{\lambda, \mu}^{\nu} \neq 0$ in the class P of polynomial time problems?

Remark 7.7 implies that the argument used in the proofs of Theorems 5.7 and 6.9 cannot work.

Problem 7.9. Is counting $o_{\lambda, \mu}^{\nu}$ in the class of #P-complete problems?

8. EQUIVARIANT SCHUBERT CALCULUS OF Y AND Z

One is interested in the equivariant cohomology of Y and Z . As with Grassmannians (Section 2), one has structure constants with respect to the Schubert basis,

$$\xi_\lambda(Y) \cdot \xi_\mu(Y) = \sum_{\nu \subseteq \rho_n} O_{\lambda, \mu}^{\nu} \xi_\nu(Y) \quad \text{and} \quad \xi_\lambda(Z) \cdot \xi_\mu(Z) = \sum_{\nu \subseteq \rho_n} L_{\lambda, \mu}^{\nu} \xi_\nu(Z).$$

If $|\lambda| + |\mu| = |\nu|$ then $O_{\lambda, \mu}^{\nu} = o_{\lambda, \mu}^{\nu}$ and $L_{\lambda, \mu}^{\nu} = l_{\lambda, \mu}^{\nu}$. For sake of brevity, we refer to [28] and the references therein.

Suppose that $H_T^*(pt) = \mathbb{Z}[t_1, \dots, t_n]$. The general form of Theorem 2.1 (see the attached footnote to that result) states that if $\gamma_1 = t_1$ and for $i > 1$, $\gamma_i = t_i - t_{i-1}$, then

$$O_{\lambda, \mu}^{\nu} \in \mathbb{Z}_{\geq 0}[\gamma_1, \gamma_2, \dots, \gamma_n].$$

Similarly, if $\alpha_1 = 2t_1, \alpha_2 = t_2 - t_1, \dots, \alpha_n = t_n - t_{n-1}$ then

$$L_{\lambda, \mu}^{\nu} \in \mathbb{Z}_{\geq 0}[\alpha_1, \alpha_2, \dots, \alpha_n].$$

Problem 8.1. Give a combinatorial rule for $O_{\lambda, \mu}^{\nu}$ and/or $L_{\lambda, \mu}^{\nu}$.

λ	μ	ν	$O_{\lambda,\mu}^\nu$	$L_{\lambda,\mu}^\nu$
[1]	[1]	[1]	γ_1	α_1
[1]	[1]	[1]	γ_1	α_1
[1]	[1]	[2]	1	2
[1]	[2]	[2]	$\gamma_1 + \gamma_2$	$\alpha_1 + 2\alpha_2$
[2]	[1]	[2]	$\gamma_1 + \gamma_2$	$\alpha_1 + 2\alpha_2$
[2]	[2]	[2]	$\gamma_1\gamma_2 + \alpha_2^2$	$\alpha_1\alpha_2 + 2\alpha_2^2$
[1]	[2]	[2, 1]	1	1
[1]	[2, 1]	[2, 1]	$2\gamma_1 + \gamma_2$	$2\alpha_1 + 2\alpha_2$
[2]	[1]	[2, 1]	1	1
[2]	[2]	[2, 1]	$2\gamma_1 + 2\gamma_2$	$\alpha_1 + 2\alpha_2$
[2]	[2, 1]	[2, 1]	$2\gamma_1^2 + 3\gamma_1\gamma_2 + \gamma_2^2$	$\alpha_1^2 + 3\alpha_1\alpha_2 + 2\alpha_2^2$
[2, 1]	[1]	[2, 1]	$2\gamma_1 + \gamma_2$	$2\alpha_1 + 2\alpha_2$
[2, 1]	[2]	[2, 1]	$2\gamma_1^2 + 3\gamma_1\gamma_2 + \gamma_2^2$	$\alpha_1^2 + 3\alpha_1\alpha_2 + 2\alpha_2^2$
[2, 1]	[2, 1]	[2, 1]	$2\gamma_1^3 + 3\gamma_1^2\gamma_2 + \gamma_1\gamma_2^2$	$\alpha_1^3 + 3\alpha_1^2\alpha_2 + 2\alpha_1\alpha_2^2$

TABLE 1. Table of products for $n = 2$

Naturally, we desire a rule in terms of shifted edge labeled tableaux. Such a rule (or any combinatorial rule) has eluded us. The reader wishing to give Problem 8.1 a try might find Table 1 useful.

Let $\tilde{L}_{\lambda,\mu}^\nu$ be the polynomial obtained from $L_{\lambda,\mu}^\nu$ after substitutions $\alpha_1 \mapsto 2\gamma_1$ and $\alpha_i \mapsto \gamma_i$ for $i > 1$. The following is a refinement of (21):

Theorem 8.2 (cf. Theorem 1.1 of [55]). $O_{\lambda,\mu}^\nu = 2^{\ell(\nu)-\ell(\lambda)-\ell(\mu)} \tilde{L}_{\lambda,\mu}^\nu$.

Thus, the $O_{\lambda,\mu}^\nu$ and $L_{\lambda,\mu}^\nu$ versions of Problem 8.1 are essentially equivalent.

Theorem 8.2 was stated in a weaker form as a conjecture in C. Monical's doctoral thesis [44, Conjecture 5.1]. A proof of a generalization was given in [55, Theorem 1.1]. Below, we offer another proof that uses a variation of the associativity recurrence alluded to at the end of Section 2.3. This recurrence should be useful to prove any guessed rule for $O_{\lambda,\mu}^\nu$ or $L_{\lambda,\mu}^\nu$, so we wish to explicate it here.

Proof. This will serve as the base case of the associativity recurrence below:

Lemma 8.3. $O_{\lambda,\mu}^\lambda = 2^{\ell(\nu)-\ell(\lambda)-\ell(\mu)} \tilde{L}_{\lambda,\mu}^\lambda$.

Proof. By the same reasoning as the derivation of (8), we have $L_{\lambda,\mu}^\lambda = \xi_\mu(Z)|_\lambda$ and $O_{\lambda,\mu}^\lambda = \xi_\mu(Y)|_\lambda$. The lemma thus holds since, by [28, Theorem 3], $\xi_\mu(Y)|_\lambda = 2^{-\ell(\mu)} \xi_\mu(Z)|_\lambda$. \square

Assign weights to each box of the staircase ρ_n as follows. For Y , the boxes on the main diagonal are assigned weight γ_1 . The boxes on the next diagonal are assigned γ_2 , etc. For Z , the boxes on the main diagonal are assigned α_1 whereas the boxes on the second diagonal are assigned $2\alpha_2$, and the third diagonal $2\alpha_3$, etc. Let $\beta_Y := \rho_n \rightarrow \{\gamma_i\}$ and $\beta_Z : \rho_n \rightarrow \{\alpha_1, 2\alpha_2, \dots, 2\alpha_n\}$ be these two assignments.

Thus, when $n = 3$ the assignment is

$$\begin{array}{|c|c|c|} \hline \gamma_1 & \gamma_2 & \gamma_3 \\ \hline & \gamma_1 & \gamma_2 \\ \hline & & \gamma_1 \\ \hline \end{array} \quad (\text{for } Y) \quad \text{and} \quad \begin{array}{|c|c|c|} \hline \alpha_1 & 2\alpha_2 & 2\alpha_3 \\ \hline & \alpha_1 & 2\alpha_2 \\ \hline & & \alpha_1 \\ \hline \end{array} \quad (\text{for } Z).$$

For a straight shape $\lambda \subseteq \rho_n$, define

$$\mathbf{wt}_Y(\lambda) = \sum_{x \in \lambda} \beta_Y(x).$$

For a skew shape $\nu/\lambda \subseteq \rho_n$,

$$\mathbf{wt}_Y(\nu/\lambda) := \mathbf{wt}_Y(\nu) - \mathbf{wt}_Y(\lambda).$$

Similarly, one defines $\mathbf{wt}_Z(\nu/\lambda)$.

Let λ^+ be λ with a box added. Also let ν^- be ν with a box removed. We claim that

$$(22) \quad \sum_{\lambda^+} O_{\lambda^+, \mu}^\nu = O_{\lambda, \mu}^\nu \mathbf{wt}_Y(\nu/\lambda) + \sum_{\nu^-} O_{\lambda, \mu}^{\nu^-}.$$

This is proved by the considering the associativity relation

$$(\xi_\lambda(Y) \cdot \xi_{(1)}(Y)) \cdot \xi_\mu(Y) = \xi_\lambda(Y) \cdot (\xi_{(1)}(Y) \cdot \xi_\mu(Y)),$$

and using the Pieri rule for Y :

$$(23) \quad \xi_{(1)}(Y) \cdot \xi_\lambda(Y) = \mathbf{wt}_Y(\lambda) \xi_\lambda(Y) + \sum_{\lambda^+} \xi_{\lambda^+}(Y).$$

The proof of (23) can be obtained starting with the same reasoning as the derivation of (9). Alternatively, it can be deduced by specializing more general formulas such as C. Lenart-A. Postnikov's [40, Corollary 1.2].

Similarly, the Pieri rule for Z reads

$$\xi_{(1)}(Z) \cdot \xi_\lambda(Z) = \mathbf{wt}_Z(\lambda) \xi_\lambda(Z) + \sum_{\lambda^+} 2^{\ell(\lambda)+1-\ell(\lambda^+)} \xi_{\lambda^+}(Z).$$

Consequently, by the same reasoning we obtain

$$(24) \quad \sum_{\lambda^+} L_{\lambda^+, \mu}^\nu 2^{\ell(\lambda)+1-\ell(\lambda^+)} = L_{\lambda, \mu}^\nu \mathbf{wt}_Z(\nu/\lambda) + \sum_{\nu^-} L_{\lambda, \mu}^{\nu^-} 2^{\ell(\nu^-)+1-\ell(\nu)}.$$

Now, to complete the proof by induction we start from (24). This is an identity of polynomials and remains so after the substitution $\alpha_1 \mapsto 2\gamma_1$ and $\alpha_i \mapsto \gamma_i$ for $i > 1$. That is,

$$(25) \quad \sum_{\lambda^+} \tilde{L}_{\lambda^+, \mu}^\nu 2^{\ell(\lambda)+1-\ell(\lambda^+)} = \tilde{L}_{\lambda, \mu}^\nu \widetilde{\mathbf{wt}}_Z(\nu/\lambda) + \sum_{\nu^-} \tilde{L}_{\lambda, \mu}^{\nu^-} 2^{\ell(\nu^-)+1-\ell(\nu)},$$

where $\widetilde{\mathbf{wt}}_Z(\nu/\lambda)$ is $\mathbf{wt}_Z(\nu/\lambda)$ with the same substitution. Note that

$$(26) \quad \frac{1}{2} \widetilde{\mathbf{wt}}_Z(\nu/\lambda) = \mathbf{wt}_Y(\nu/\lambda).$$

Now multiply both sides of (25) by $\frac{1}{2} \times 2^{\ell(\nu)-\ell(\lambda)-\ell(\mu)}$. This gives

$$(27) \quad \sum_{\lambda^+} \tilde{L}_{\lambda^+, \mu}^\nu 2^{\ell(\nu)-\ell(\lambda^+)-\ell(\mu)} = 2^{\ell(\nu)-\ell(\lambda)-\ell(\mu)} \tilde{L}_{\lambda, \mu}^\nu \mathbf{wt}_Y(\nu/\lambda) + \sum_{\nu^-} \tilde{L}_{\lambda, \mu}^{\nu^-} 2^{\ell(\nu^-)-\ell(\lambda)-\ell(\mu)}.$$

By induction,

$$(28) \quad \sum_{\lambda^+} O_{\lambda^+, \mu}^\nu = 2^{\ell(\nu) - \ell(\lambda) - \ell(\mu)} \tilde{L}_{\lambda, \mu}^\nu \text{wt}_Y(\nu/\lambda) + \sum_{\nu^-} O_{\lambda, \mu}^{\nu^-}.$$

Comparing (28) and (22) we deduce that $O_{\lambda, \mu}^\nu = 2^{\ell(\nu) - \ell(\lambda) - \ell(\mu)} \tilde{L}_{\lambda, \mu}^\nu$, as needed. \square

Turning to nonvanishing, clearly:

Corollary 8.4. $[\gamma_1^{i_1} \cdots \gamma_n^{i_n}] O_{\lambda, \mu}^\nu \neq 0 \iff [\alpha_1^{i_1} \cdots \alpha_n^{i_n}] L_{\lambda, \mu}^\nu \neq 0$; in particular $O_{\lambda, \mu}^\nu \neq 0 \iff L_{\lambda, \mu}^\nu \neq 0$.

Moreover, C. Monical [44] gave a conjectural equivariant extension of Theorem 7.6.

Conjecture 8.5 (C. Monical's Horn-type conjecture). *For $\lambda, \mu, \nu \subseteq \rho_n$ (and not a smaller staircase), $O_{\lambda, \mu}^{\nu^c} \neq 0$ if and only if for $k = |\lambda| + |\mu| + |\nu| - \binom{n+1}{2}$,*

- $k \geq 0$, and
- for all $0 < r < n$ and all $\alpha, \beta, \gamma \subseteq r \times (n-r)$ with $|\alpha| + |\beta| + |\gamma| = r(n-r)$ and $c_{\alpha, \beta}^{\gamma^c} \neq 0$ we have $[\lambda]_\alpha + [\mu]_\beta + [\nu]_\gamma - k \leq \binom{n+1-r}{2}$.

In *loc. cit.*, C. Monical reports checking this conjecture for all $\lambda, \mu, \nu \subseteq \rho_5$. Now, from Corollary 8.4 we obtain:


Corollary 8.6 (cf. Conjecture 5.3 of [44]). *C. Monical's inequalities characterize $O_{\lambda, \mu}^\nu \neq 0$ if and only if they characterize $L_{\lambda, \mu}^\nu \neq 0$.*

9. SHIFTED EDGE LABELED TABLEAUX

In this section, we define *shifted edge labeled tableaux*. At present, we do not know a good theory when edge labels are permitted on arbitrary horizontal edges. However, our central new idea is to *restrict edge labels to diagonal boxes*. This restriction gives rise to a combinatorial rule which defines a commutative and (conjecturally) associative ring.

9.1. Main definitions. If $\mu \subseteq \lambda$, then λ/μ is the skew-shape consisting of boxes of λ not in μ . The boxes in matrix position (i, i) are the *diagonal boxes*. A *diagonal edge* of λ/μ refers to the southern edge of a diagonal box of λ . If $\mu = \emptyset$, we call $\lambda = \lambda/\mu$ a *straight shape*.

For example if $\lambda = (6, 3, 1)$ and $\mu = (3, 1)$, the shape λ/μ consists of the six unmarked boxes shown below



This has one diagonal box but three diagonal edges.

A *shifted edge labeled tableau* of shape λ/μ is a filling of the boxes of λ/μ and southern edges of the diagonal boxes with the labels $[N] = \{1, 2, 3, \dots, N\}$ such that:

- (S1) Every box of λ/μ is filled.
- (S2) Each *diagonal edge* contains a (possibly empty) subset of $[N]$.
- (S3) $1, 2, \dots, N$ appears exactly once.
- (S4) The labels strictly increase left to right along rows and top to bottom along columns. In particular, each label of a diagonal edge is strictly larger than the box labels in the same column.

These conditions imply that $N \geq |\lambda/\mu|$. Let $\text{eqShSYT}(\lambda/\mu, N)$ be the set of all such tableaux. If we restrict to tableaux satisfying only (S1), (S3) and (S4), then $N = |\lambda/\mu|$ and we obtain the notion of shifted standard Young tableaux from Section 7.

An *inner corner* c of λ/μ is a maximally southeast box of μ . For $T \in \text{eqShSYT}(\lambda/\mu, N)$, we define a (*shifted, edge labeled*) *jeu de taquin slide* $\text{shEjdt}_c(T)$, obtained as follows. Initially place \bullet in c , and apply one of the following *slides*, depending on what T looks like locally around c :

$$(J1) \begin{array}{|c|c|} \hline \bullet & a \\ \hline b & \\ \hline \end{array} \mapsto \begin{array}{|c|c|} \hline b & a \\ \hline \bullet & \\ \hline \end{array} \text{ (if } b < a, \text{ or } a \text{ does not exist)}$$

$$(J2) \begin{array}{|c|c|} \hline \bullet & a \\ \hline b & \\ \hline \end{array} \mapsto \begin{array}{|c|c|} \hline a & \bullet \\ \hline b & \\ \hline \end{array} \text{ (if } a < b, \text{ or } b \text{ does not exist)}$$

$$(J3') \begin{array}{|c|c|} \hline \bullet & a \\ \hline S & \\ \hline \end{array} \mapsto \begin{array}{|c|c|} \hline a & \bullet \\ \hline S & \\ \hline \end{array} \text{ (if } c \text{ is a diagonal box and } a < \min(S))$$

$$(J4') \begin{array}{|c|c|} \hline \bullet & a \\ \hline S & \\ \hline \end{array} \mapsto \begin{array}{|c|c|} \hline s & a \\ \hline S' & \\ \hline \end{array} \text{ (if } c \text{ is a diagonal box, } s := \min(S) < a \text{ and } S' := S \setminus \{s\})$$

Repeat the above sliding procedure on the new box c' containing the new position \bullet until \bullet arrives at a box or diagonal edge d of λ that has no labels immediately south or east of it. Then $\text{shEjdt}_c(T)$ is obtained by erasing \bullet .

A *rectification* of $T \in \text{eqShSYT}(\lambda/\mu, N)$ is defined as usual: Choose an inner corner c_0 of λ/μ and compute $T_1 := \text{shEjdt}_{c_0}(T)$, which has shape $\lambda^{(1)}/\mu^{(1)}$. Now let c_1 be an inner corner of $\lambda^{(1)}/\mu^{(1)}$ and compute $T_2 := \text{shEjdt}_{c_1}(T_1)$. Repeat $|\mu|$ times, arriving at a standard tableau of straight shape. Let $\text{shEqRect}_{\{c_i\}}(T)$ be this tableau.

In general, shEqRect is not independent of rectification order, when $N > |\lambda/\mu|$:

Example 9.1. The reader can check that if one uses column rectification order (picking the rightmost inner corner at each step) then

$$\begin{array}{|c|c|c|} \hline & & \\ \hline & & 1 \\ \hline & & 2 \\ \hline & & 3 \\ \hline \end{array}$$

rectifies to $\begin{array}{|c|c|c|} \hline 1 & 2 & 3 \\ \hline \end{array}$ while row rectification (choosing the southmost inner corner at each step) gives $\begin{array}{|c|c|} \hline 1 & 2 \\ \hline & 3 \\ \hline \end{array}$. □

We will define $\text{shEqRect}(T)$ to be the rectification under row rectification order.

9.2. A (putative) commutative ring structure. Let S_μ be the superstandard tableau of shifted shape μ , which is obtained by filling the boxes of μ in English reading order with $1, 2, 3, \dots$. For example,

$$S_{(5,3,1)} = \begin{array}{|c|c|c|c|c|} \hline 1 & 2 & 3 & 4 & 5 \\ \hline & 6 & 7 & 8 & \\ \hline & & 9 & & \\ \hline \end{array}$$

Define

$$d_{\lambda, \mu}^{\nu} := \#\{T \in \text{eqShSYT}(\nu/\lambda, |\mu|) : \text{shEqRect}(T) = S_\mu\}.$$

λ	μ	$\lambda \star \mu$
[1]	[1]	$z[1] + 2[2]$
[1]	[2]	$z[2] + [2, 1] + 2[3]$
[1]	[2, 1]	$2z[2, 1] + 2[3, 1]$
[1]	[3]	$z[3] + [3, 1]$
[1]	[3, 1]	$2z[3, 1] + 2[3, 2]$
[1]	[3, 2]	$2z[3, 2] + [3, 2, 1]$
[1]	[3, 2, 1]	$3z[3, 2, 1]$
[2]	[2]	$z[2, 1] + z[3] + 2[3, 1]$
[2]	[2, 1]	$z^2[2, 1] + 3z[3, 1] + 2[3, 2]$
[2]	[3]	$z[3, 1] + [3, 2]$
[2]	[3, 1]	$z^2[3, 1] + 3z[3, 2] + [3, 2, 1]$
[2]	[3, 2]	$z^2[3, 2] + 2z[3, 2, 1]$
[2]	[3, 2, 1]	$3z^2[3, 2, 1]$
[2, 1]	[2, 1]	$z^3[2, 1] + 3z^2[3, 1] + 6z[3, 2]$
[2, 1]	[3]	$z^2[3, 1] + z[3, 2] + [3, 2, 1]$
[2, 1]	[3, 1]	$z^3[3, 1] + 3z^2[3, 2] + 3z[3, 2, 1]$
[2, 1]	[3, 2]	$z^3[3, 2] + 3z^2[3, 2, 1]$
[2, 1]	[3, 2, 1]	$4z^3[3, 2, 1]$
[3]	[3]	$z[3, 2]$
[3]	[3, 1]	$z^2[3, 2] + z[3, 2, 1]$
[3]	[3, 2]	$z^2[3, 2, 1]$
[3]	[3, 2, 1]	$z^3[3, 2, 1]$
[3, 1]	[3, 1]	$z^3[3, 2] + 3z^2[3, 2, 1]$
[3, 1]	[3, 2]	$2z^3[3, 2, 1]$
[3, 1]	[3, 2, 1]	$2z^4[3, 2, 1]$
[3, 2]	[3, 2]	$z^4[3, 2, 1]$
[3, 2]	[3, 2, 1]	$z^5[3, 2, 1]$
[3, 2, 1]	[3, 2, 1]	$z^6[3, 2, 1]$

TABLE 2. Table of products for $n = 3$

Let

$$\Delta(\nu; \lambda, \mu) := |\lambda| + |\mu| - |\nu| \text{ and } L(\nu; \lambda, \mu) := \ell(\lambda) + \ell(\mu) - \ell(\nu).$$

Introduce an indeterminate z and set

$$D_{\lambda, \mu}^\nu := 2^{L(\nu; \lambda, \mu) - \Delta(\nu; \lambda, \mu)} z^{\Delta(\nu; \lambda, \mu)} d_{\lambda, \mu}^\nu.$$


Next we define formal symbols $[\lambda]$ for each $\lambda \subseteq \rho_n$. Let R_n be the free $\mathbb{Z}[z]$ -module generated by these. We declare a product structure on R_n by

$$[\lambda] \star [\mu] = \sum_{\nu} D_{\lambda, \mu}^\nu [\nu].$$

While positivity of $D_{\lambda, \mu}^\nu$ is immediate from the definition, the following is not:

Conjecture 9.2. $D_{\lambda, \mu}^\nu \in \mathbb{Z}[z]$.

Example 9.3. Suppose that $\lambda = (2, 1), \mu = (3, 1), \nu = (3, 1)$. Then $\Delta(\nu; \lambda, \mu) = 3 + 4 - 4 = 3$, $L(\nu; \lambda, \mu) = 2 + 2 - 2 = 2$. Also, $d_{\lambda, \mu}^\nu = 2$ because the following are the only 2 shifted edge labeled tableaux which rectify to S_μ .




Thus $D_{\lambda, \mu}^\nu = 2^{2-3} \times z^3 \times 2 = z^3$. □

In the previous example, $2^{L(\nu; \lambda, \mu) - \Delta(\nu; \lambda, \mu)} = 2^{-1}$. Further, in the tableaux counting $d_{\lambda, \mu}^\nu$, only the edge labels differed. In the case that $L(\nu; \lambda, \mu) - \Delta(\nu; \lambda, \mu) = -k < 0$ one might wonder if the tableaux counting $d_{\lambda, \mu}^\nu$, namely

$$F(\lambda, \mu; \nu) = \{T \in \text{shEqSYT}(\nu/\lambda, |\mu|) : \text{shEqRect}(T) = S_\mu\}$$

can be sorted into equivalence classes of size 2^k by ignoring edge labels. The following example shows this is false in general:

Example 9.4. Suppose that $\lambda = (3), \mu = (3, 2, 1), \nu = (4, 2, 1)$. Then $\Delta(\nu; \lambda, \mu) = 3 + 6 - 7 = 2$ and $L(\nu; \lambda, \mu) = 1 + 3 - 3 = 1$, so $k = 1$. Below a $T \in F_{(3), (3, 2, 1); (4, 2, 1)}$. Any $T' \in \text{eqShSYT}((4, 2, 1)/(3), 6)$ formed by moving the edge labels of T is not in $F_{(3), (3, 2, 1); (4, 2, 1)}$.



While Conjecture 9.2 is a purely combinatorial question, it would also follow from a conjectural connection to equivariant Schubert calculus, through work of D. Anderson-W. Fulton presented in Section 10.


The next result gives a further consistency check of our combinatorics. It was suggested by H. Thomas (private communication):

Theorem 9.5. R_n is commutative, i.e., $D_{\lambda, \mu}^\nu = D_{\mu, \lambda}^\nu$.


Proof. The argument is based on a variation of S. Fomin's *growth diagram* formulation of *jeu de taquin*; see, e.g., [62, Appendix 1].

Given a tableau $T \in \text{eqShSYT}(\lambda/\theta, n)$, define the corresponding *e-partition* (e for “edge”) to be $\text{epart}(T) := (\lambda_1^{i_1}, \lambda_2^{i_2}, \dots)$ where i_k = number of edge labels on the k^{th} diagonal edge. Each such T can be encoded as a sequence of e-partitions starting with the (usual) partition $\theta = (\theta_1^0, \theta_2^0, \dots)$: If 1 appears in a box in row i then the next e-partition has an extra box in this position, i.e., we replace θ_i^0 with $(\theta_i^0 + 1)^0$. Otherwise 1 appears on the edge of a diagonal box in row i , in which case, the one-larger e-partition has θ_i^0 replaced by θ_i^1 . Repeat this process by looking at the position of 2 in T etc. Evidently, such an encoding of T is unique.

Example 9.6.



Rectifying the left tableau above:



Each of these four tableaux also has an associated sequence of e-partitions. Place these atop of one another as below. The result is a *tableau rectification diagram*:

$(2^0, 1^0)$	$(2^1, 1^0)$	$(2^1, 1^1)$	$(3^1, 1^1)$	$(3^1, 1^2)$
(2^0)	(2^1)	$(2^1, 1^0)$	$(3^1, 1^0)$	$(3^1, 1^1)$
(1^0)	(1^1)	(2^1)	(3^1)	$(3^1, 1^0)$
\emptyset	(1^0)	(2^0)	(3^0)	$(3^0, 1^0)$

Given two e-partitions $\lambda = (\lambda_1^{i_1}, \lambda_2^{i_2}, \dots)$ and $\mu = (\mu_1^{j_1}, \mu_2^{j_2}, \dots)$, we will say μ *covers* λ if:

- (i) there exists unique m such that $\lambda_m + 1 = \mu_m$ and $\lambda_k = \mu_k$ for $k \neq m$, and $i_k = j_k$ for all k ; or
- (ii) $\lambda_k = \mu_k$ for all k and there exists a unique m such that $i_m + 1 = j_m$ and $i_k = j_k$ for $k \neq m$.

In the case that μ covers λ we define μ/λ to be the extra box added in row m (if in cases (i) above), or the m^{th} diagonal edge (in case (ii)). If x is a diagonal edge, define $\text{shEjdt}_x(T) = T$. For two e-partitions, $\lambda = (\lambda_1^{i_1}, \lambda_2^{i_2}, \dots)$ and $\mu = (\mu_1^{j_1}, \mu_2^{j_2}, \dots)$, let

$$\lambda \vee \mu = (\max\{\lambda_1, \mu_1\}^{i_1+j_1}, \max\{\lambda_2, \mu_2\}^{i_2+j_2}, \dots).$$

Consider the following local conditions on any 2×2 subsquare $\begin{smallmatrix} \alpha & \beta \\ \gamma & \delta \end{smallmatrix}$ on a grid of e-partitions:

- (G1) Each e-partition covers the e-partition immediately to its left or below.
- (G2) $\delta = \gamma \vee \text{epart}(\text{shEjdt}_{\alpha/\gamma}(T))$, where T is the filling of β/α by 1. Similarly $\alpha = \gamma \vee \text{epart}(\text{shEjdt}_{\delta/\gamma}(T))$ where T is the filling of β/δ by 1.

Call any rectangular table of e-partitions satisfying (G1) and (G2) a *growth diagram*. By the symmetry in the definition of (G1) and (G2), if \mathcal{G} is a growth diagram, then so is \mathcal{G} reflected about its antidiagonal. The following is straightforward from the definitions:

Claim 9.7. If $\begin{smallmatrix} \alpha & \beta \\ \gamma & \delta \end{smallmatrix}$ is a 2×2 square in the tableau rectification diagram, then (G1) and (G2) hold.

Proof. Fix any two rows of the tableau rectification diagram; call this $2 \times (n+1)$ subdiagram \mathcal{R} . The higher of the two rows corresponds to some shifted edge labeled tableau U and the other row corresponds to $\text{shEjdt}_c(U)$ where c is a box (determined by the shapes in the leftmost column). Now U is filled by $1, 2, \dots, n$. Notice that if we consider the submatrix \mathcal{R}' of \mathcal{R} consisting of the leftmost $k+1$ columns, then \mathcal{R}' corresponds to the computation of $\text{shEjdt}_c(U')$ where U' is U with labels $k+1, k+2, \dots, n$ removed. The upshot is that it suffices to prove the claim for the rightmost 2×2 square in \mathcal{R} , which we will label with shapes $\begin{array}{|c|c|} \hline \alpha & \beta \\ \hline \gamma & \delta \\ \hline \end{array}$. Let U_β be the tableau associated to the chain of e-partitions ending at β . Similarly define U_α . As well we have

$$(29) \quad U_\gamma = \text{shEjdt}_c(U_\alpha)$$

and

$$(30) \quad U_\delta = \text{shEjdt}_c(U_\beta).$$

Insofar as (G1) is concerned, it is obvious that β covers α and δ covers γ . That β covers δ follows from (30). Similarly, α covers γ because of (29).

Now we turn to the proof of the first sentence of (G2). Suppose that $S \in \text{eqShSYT}(\lambda/\theta, n)$. Define \bar{S} to be the tableau obtained by forgetting the entries $1, 2, \dots, n-1$ in S and replacing the n by 1. Also define \tilde{S} to be the tableau obtained by forgetting the entry n in S . Then, it is clear that

$$(31) \quad \text{epart}(S) = \text{epart}(\tilde{S}) \vee \text{epart}(\bar{S})$$

By definition, $\tilde{U}_\delta = U_\gamma$. This combined with (31) applied to $S = U_\delta$ shows that to prove the claim it suffices to show that $\text{shEjdt}_{\alpha/\gamma}(T) = \bar{U}_\delta$.

Case 1: (α covers γ by (i)) In the computation of $\text{shEjdt}_c(U_\alpha)$, the \bullet (that starts at c) arrives at the outer corner box α/γ . By definition, U_β contains n at β/α . Therefore, if α/γ is not adjacent to β/α , clearly U_δ is U_γ with n adjoined at β/α . Thus, $\text{shEjdt}_{\alpha/\gamma}(T) = \bar{U}_\delta$ as desired.

Otherwise α/γ is adjacent to β/α . Then by the definition of shEjdt , the position of 1 in $\text{shEjdt}_{\alpha/\gamma}(T)$ is the same as the position of n in U_δ . Thus, $\text{shEjdt}_{\alpha/\gamma}(T) = \bar{U}_\delta$.

Case 2: (α covers γ by (ii)) Then in the computation of $\text{shEjdt}_c(U_\alpha)$, the \bullet must have arrived at a diagonal box, and $k(< n)$ is the smallest edge label of this same box, resulting in a (J4') slide. Now regardless of where n is placed in U_β , it is clear that U_δ is U_γ with n adjoined in the same place as n 's place in U_β , i.e., β/α . In other words, $\text{shEjdt}_{\alpha/\gamma}(T) = \bar{U}_\delta$.

Proof of the second sentence of (G2): For a pair of e-partitions λ, μ with μ covering λ , define $U_{\mu/\lambda}$ to be the tableau with 1 placed in the location μ/λ in μ . Clearly, $\alpha = \text{epart}(U_\alpha) = \text{epart}(U_\gamma) \vee \text{epart}(U_{\alpha/\gamma})$. Thus, it suffices to show

$$(32) \quad \text{shEjdt}_{\delta/\gamma}(T) = U_{\alpha/\gamma}.$$

Case 1: (δ covers γ by (i)) Then in U_δ , n occupies box δ/γ . If δ/γ is not adjacent to β/δ , it is clear that U_β and U_δ have n in the same place, i.e., δ/γ . Thus β/δ and α/γ are the same box or edge position. Thus Equation (32) follows.

Otherwise δ/γ is adjacent to β/δ . Then,

$$(33) \quad \text{shEjdt}_{\delta/\gamma}(T) = U_{\delta/\gamma}.$$

By the definition of shEjdt , it follows that in the computation of $\text{shEjdt}_c(U_\alpha)$, the \bullet arrived at an outer corner α/γ . Using this, combined with the fact that δ/γ is adjacent to β/δ , we conclude that α/γ is a box in the same position as the box δ/γ . Now (33) is precisely (32).

Case 2: (δ covers γ by (ii)) Then δ/γ is an edge, so n occupies an edge in U_δ . Thus in $\text{shEjdt}_c(U_\beta)$, n is never moved. So, $\beta/\delta = \alpha/\gamma$. Since δ/γ is a diagonal edge, $\text{shEjdt}_{\delta/\gamma}(T) = T := U_{\beta/\delta} = U_{\alpha/\gamma}$. \square

Let $\text{Growth}(\lambda, \mu; \nu)$ be the set of growth diagrams such that:

- the leftmost column encodes the superstandard tableau of shape λ ;
- the bottom-most row encodes the superstandard tableau of shape μ ;
- the shape of the e-partition in the top right corner is ν .

Claim 9.8. $\#\text{Growth}(\lambda, \mu; \nu) = \#F(\lambda, \mu; \nu)$.

Proof. Given $T \in F(\lambda, \mu; \nu)$, form the tableau rectification diagram $\mathcal{G}(T)$. Notice that since we are using row rectification order, the left side of the diagram will be the sequence for S_λ . Since T is assumed to rectify to S_μ , the bottom row of the diagram will be the sequence for S_μ . Hence by Claim 9.7, $\mathcal{G}(T) \in \text{Growth}(\lambda, \mu; \nu)$, and thus $T \mapsto \mathcal{G}(T)$ is an injection implying $\#F(\lambda, \mu; \nu) \leq \#\text{Growth}(\lambda, \mu; \nu)$.

For the reverse inequality, given any $\mathcal{G} \in \text{Growth}(\lambda, \mu; \nu)$, by (G1), the top row defines $T(\mathcal{G}) \in \text{eqShSYT}(\nu/\lambda, |\mu|)$. Then $T(\mathcal{G})$ has a tableau rectification diagram \mathcal{G}' . By Claim 9.7, \mathcal{G}' is uniquely determined by its left and top borders together with (G2). Thus, since \mathcal{G} and \mathcal{G}' share the same left and top borders and both satisfy (G2), $\mathcal{G} = \mathcal{G}'$. In particular, $T(\mathcal{G}) \in F(\lambda, \mu; \nu)$. Thus, $\mathcal{G} \mapsto T(\mathcal{G})$ is an injection proving $\#\text{Growth}(\lambda, \mu; \nu) \leq \#F(\lambda, \mu; \nu)$. \square

To conclude, we must show that $d_{\lambda, \mu}^\nu = d_{\mu, \lambda}^\nu$. Since

$$d_{\lambda, \mu}^\nu = \#F(\lambda, \mu; \nu) = \#\text{Growth}(\lambda, \mu; \nu),$$

it suffices to show that $\#\text{Growth}(\lambda, \mu; \nu) = \#\text{Growth}(\mu, \lambda; \nu)$. Reflecting along the antidiagonal defines a bijection between $\text{Growth}(\lambda, \mu; \nu)$ and $\text{Growth}(\mu, \lambda; \nu)$. \square

Example 9.9. Under column rectification, Theorem 9.5 is false. Suppose $\lambda = (4, 3)$, $\mu = (3, 2, 1)$ and $\nu = (4, 3, 2, 1)$. The number of tableaux of shape ν/λ that column rectify to S_μ is 20 while the number of those with shape ν/μ column rectifying to S_λ is 16. \square

Conjecture 9.10. (R_n, \star) is an associative ring.

Additional support for Conjecture 9.10 comes from a conjectural connection to a commutative, associative ring studied by D. Anderson-W. Fulton, as described in the next section.

10. CONJECTURAL CONNECTION TO WORK OF W. FULTON-D. ANDERSON AND EQUIVARIANT SCHUBERT CALCULUS

10.1. Results of W. Fulton-D. Anderson. For a strict shape $\lambda \subseteq \rho_n$, let $\sigma_\lambda = \text{Pf}(c_{\lambda_i, \lambda_j})$ where

$$c_{p,q} = \sum_{0 \leq a \leq b \leq q} (-1)^b \left(\binom{b}{a} + \binom{b-1}{a} \right) z^a c_{p+b-a} c_{q-b}.$$

If $\ell = \ell(\lambda)$ is odd, define $\lambda_{\ell+1} = 0$.

Recently, D. Anderson-W. Fulton have studied a $\mathbb{Z}[z]$ -algebra

$$\mathfrak{P} = \mathbb{Z}[z, c_1, c_2, \dots] / (c_{p,p} = 0, \forall p > 0)$$

and shown it has a basis over $\mathbb{Z}[z]$ of σ_λ . Define structure constants by

$$\sigma_\lambda \cdot \sigma_\mu = \sum_{\nu \subseteq \rho_n} \mathfrak{D}_{\lambda,\mu}^\nu \sigma_\nu.$$

Also let

$$\mathfrak{d}_{\lambda,\mu}^\nu := \frac{\mathfrak{D}_{\lambda,\mu}^\lambda}{2^{L(\lambda;\lambda,\mu) - \Delta(\lambda;\lambda,\mu)} z^{\Delta(\lambda;\lambda,\mu)}}.$$

Conjecture 10.1. *There is a ring isomorphism $\phi : R_n \rightarrow \mathfrak{P}$ that sends $[\lambda] \mapsto \sigma_\lambda$. Therefore, $\mathfrak{D}_{\lambda,\mu}^\nu = D_{\lambda,\mu}^\nu$ (equivalently $\mathfrak{d}_{\lambda,\mu}^\nu = d_{\lambda,\mu}^\nu$).*

We have exhaustively checked Conjecture 10.1 for all $n \leq 4$ and many $n = 5$ cases.

D. Anderson-W. Fulton (private communication) connected the above ring to equivariant Schubert calculus of Z . That is,

$$(34) \quad L_{\lambda,\mu}^\nu(\alpha_1 \mapsto z, \alpha_2 \mapsto 0, \dots, \alpha_n \mapsto 0) = \mathfrak{D}_{\lambda,\mu}^\nu.$$

Proposition 10.2. *Conjecture 10.1 \implies Conjecture 9.2.*

Proof. By Theorem 2.1, $L_{\lambda,\mu}^\nu$ is a nonnegative integer polynomial in $\alpha_1, \dots, \alpha_n$, the simples of the type C root system. Now apply (34). \square

In turn, Conjecture 10.1 should follow from a proof that $D_{\lambda,(p)}^\nu = \mathfrak{D}_{\lambda,(p)}^\nu$, together with Conjecture 9.10, by a variation of the “associativity argument” of Section 3.

10.2. Two numerologically nice cases of Conjecture 10.1.

Theorem 10.3. $d_{\lambda,(p)}^\lambda = \binom{\ell(\lambda)}{p} 2^{p-1} = \mathfrak{d}_{\lambda,(p)}^\lambda$.

Proof. We will use some results of T. Ikeda-H. Naruse [28] that we now recall. For a strict partition $\lambda = (\lambda_1 > \dots > \lambda_r > 0)$, let D_λ denote the associated shifted shape. Explicitly,


$$D_\lambda = \{(i, j) \in \mathbb{Z}^2 \mid 1 \leq i \leq r, i \leq j < \lambda_i + i\}.$$

For instance, $D_{(3,1)} = \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & & \\ \hline \end{array}$. Given an arbitrary subset $C \subset D_\lambda$, if a box $x \in C$ satisfies either of the following conditions:

- (I) $x = (i, i)$ and $(i, i+1), (i+1, i+1) \in D_\lambda \setminus C$
- (II) $x = (i, j), j \neq i$ and $(i+1, i), (i, i+1), (i+1, i+1) \in D_\lambda \setminus C$

then set $C' = C \cup \{x + (1, 1)\} \setminus \{x\}$. The procedure $C \rightarrow C'$ is called an *elementary excitation occurring at $x \in C$* . Any subset $C' \subset D_\lambda$ obtained from C by an application of successive elementary excitations is called an *excited Young diagram (EYD) of C* . Denote by $\mathcal{E}_\lambda(\mu)$ the set of all EYDs of D_μ contained in D_λ .

Example 10.4. Suppose $\lambda = (4, 2, 1), \mu = (2)$, then $\mathcal{E}_\lambda(\mu)$ consists of the following EYDs



Lemma 10.5. $\mathfrak{D}_{\lambda,\mu}^\lambda = \#\mathcal{E}_{\rho_{\ell(\lambda)}}(\mu) \times z^{|\mu|}$.

Proof. This follows from [28, Theorem 3] which gives a formula for $\xi_\lambda(Z)|_\mu = L_{\lambda,\mu}^\mu$ in terms of EYDs, combined with (34). We omit the details, which amount mostly to translating from the appropriate Weyl group elements to the associated partitions. \square

By Lemma 10.5,

$$(35) \quad \mathfrak{d}_{\lambda,\mu}^\lambda = \frac{\mathfrak{D}_{\lambda,\mu}^\lambda}{2^{L(\lambda;\lambda,\mu) - \Delta(\lambda;\lambda,\mu)} z^{\Delta(\lambda;\lambda,\mu)}} = \frac{\#\mathcal{E}_{\rho_{\ell(\lambda)}}(\mu) \times z^{|\mu|}}{2^{\ell(\mu) - |\mu|} z^{|\mu|}} = \#\mathcal{E}_{\rho_{\ell(\lambda)}}(\mu) \times 2^{|\mu| - \ell(\mu)}.$$

As all boxes in $D_{(p)}$ are in distinct columns, they stay in distinct columns even after the application of the excitation moves (I) and (II). Therefore, $\mathcal{E}_{\rho_{\ell(\lambda)}}((p))$ contains at most $\binom{\ell(\lambda)}{p}$ elements, since, there are $\ell(\lambda)$ columns in $D_{\rho_{\ell(\lambda)}}$. It is not hard to see from (I) and (II) this upper bound is an equality. This proves the second equality of the theorem.

Let $N = \ell(\lambda)$; we now prove the first equality of the theorem statement by induction on $N + p$. When $N + p \leq 1$ the claim is obvious. When $N + p = 2$, there is one case, namely, $\lambda = (1)$, $p = 1$ and $d_{\lambda,(p)}^\lambda = 1 = \binom{1}{1} 2^{1-1}$, as desired. Now suppose $N + p = k > 2$ and the claim holds for smaller $N + p$.

Let $F(\lambda, (p); \lambda)$ be the tableaux enumerated by $d_{\lambda,(p)}^\lambda$. If $T \in F(\lambda, (p); \lambda)$ we say that a label q appears in row r if q is an edge label on the southern edge of the diagonal box in row r . Let \bar{T} be T with the first row removed, this is of shape $\bar{\lambda}$. There are three disjoint cases that T can fall into:

- (1) (1 does not appear in row 1 of T and $\bar{T} \in F(\bar{\lambda}, (p); \bar{\lambda})$): Then there are $d_{\bar{\lambda},(p)}^{\bar{\lambda}}$ many such choices; this equals $\binom{N-1}{p} 2^{p-1}$, by induction.
- (2) (1 does not appear in row 1 of T and $\bar{T} \notin F(\bar{\lambda}, (p); \bar{\lambda})$): Then it is straightforward to check (from the assumption that $T \in F(\lambda, (p); \lambda)$) that \bar{T} row rectifies to \bar{S} of shape $(p-1)$ where the first row consists of box labels $1, 3, 4, \dots, p-1$ and has a 2 in the south edge of the first box. Notice that the choices for \bar{T} are in bijection with $F(\bar{\lambda}, (p-1); \bar{\lambda})$ where the map is to remove the edge label 2 and shift the labels $3, 4, 5, \dots, p$ down by one. This, combined with induction asserts that there are $d_{\bar{\lambda},(p-1)}^{\bar{\lambda}} = \binom{N-1}{p-1} 2^{p-2}$ many choices.
- (3) (1 appears in row 1): No other label appears in row 1 of T . Let U be \bar{T} with every entry decremented by one. It is straightforward that $U \in F(\bar{\lambda}, (p-1); \bar{\lambda})$, and that the map $T \mapsto U$ is bijective. Thus, there are $d_{\bar{\lambda},(p-1)}^{\bar{\lambda}} = \binom{N-1}{p-1} 2^{p-2}$ many tableaux in this case, by induction.

By Pascal's identity,

$$\binom{N}{p} 2^{p-1} = \binom{N-1}{p} 2^{p-1} + \binom{N-1}{p-1} 2^{p-2} + \binom{N-1}{p-1} 2^{p-2}.$$

This, combined with cases (1)-(3), completes the induction. \square

Theorem 10.6. $d_{\rho_n, \rho_n}^{\rho_n} = 2^{\binom{n}{2}} = \mathfrak{d}_{\rho_n, \rho_n}^{\rho_n}$.

Proof sketch: In the special case when $\ell(\lambda) = \ell(\mu)$, it is easy to observe that $|\mathcal{E}_\mu^\lambda| = 1$. Thus in this case,

$$(36) \quad \mathfrak{d}_{\lambda, \mu}^\lambda = 2^{|\mu| - \ell(\mu)}$$

Further when $\mu = \lambda = \rho_n$,

$$\mathfrak{d}_{\rho_n, \rho_n}^{\rho_n} = 2^{|\mu| - \ell(\mu)} = 2^{\binom{n}{2}}.$$

This proves the rightmost equality.

For the remaining equality, consider $T \in \text{shEqSYT}(\rho_n/\rho_n, N)$ where $N = |\rho_n| = \binom{n+1}{2}$. For $1 \leq i \leq n$, let

$$E_i(T) = \{k \mid k \text{ lies on the } i\text{th diagonal edge of } T\}.$$

For $T \in \text{shSYT}(\nu/\lambda)$, let $T(i, j)$ be the entry in box (i, j) (in matrix coordinates). Define $U_n \in \text{shEqSYT}(\rho_n/\rho_n, N)$ by the requirement that $E_i(U_n) = \bigcup_{r=1}^i S_{\rho_n}(r, i)$. That is, the labels on the i th diagonal edge of U_n are precisely the labels appearing in column i of S_{ρ_n} .

For $T \in \text{shEqSYT}(\rho_n/\rho_n, N)$ and $I \subseteq E_i(T)$ for some $i \in [n-1]$, define the I -slide of T , $\text{Sl}_I(T) \in \text{shEqSYT}(\rho_n/\rho_n, N)$, by

$$E_k(\text{Sl}_I(T)) := \begin{cases} E_k(T) & \text{if } k \in [n] \setminus \{i, i+1\}, \\ E_k(T) \setminus I & \text{if } k = i, \\ E_k(T) \cup I & \text{if } k = i+1. \end{cases}$$

Example 10.7. Let $n = 4$. Taking $I = \{6\} \subseteq E_3(U_4) = \{3, 6, 8\}$, below we illustrate $\text{Sl}_{\{6\}}(U_4)$.

$$S_{\rho_4} = \begin{array}{|c|c|c|c|} \hline 1 & 2 & 3 & 4 \\ \hline & 5 & 6 & 7 \\ \hline & & 8 & 9 \\ \hline & & & 10 \\ \hline \end{array} \quad U_4 = \begin{array}{|c|c|c|c|} \hline 1 & & & \\ \hline & 25 & & \\ \hline & & 368 & \\ \hline & & & 47910 \\ \hline \end{array} \quad \text{Sl}_{\{6\}}(U_4) = \begin{array}{|c|c|c|c|} \hline 1 & & & \\ \hline & 25 & & \\ \hline & & 38 & \\ \hline & & & 467910 \\ \hline \end{array}$$

For $T \in \text{shSYT}(\nu/\lambda)$, let

$$\text{row}_k(T) = \{\text{entries in row } k \text{ of } T\}.$$

Say $I \subseteq E_i(T)$ is n -slidable if $1 \leq i < n$,


$$(37) \quad I \subseteq \bigcup_{k=1}^i \{\min(E_i(T) \cap \text{row}_k(S_{\rho_n}))\},$$

and for $i < k \leq n$, $E_k(T) = E_k(U_n)$.

Example 10.8. Consider T below and $i = 3$. Then to the right we have S_{ρ_4} with

$$\bigcup_{k=1}^3 \{\min(E_3(T) \cap \text{row}_k(S_{\rho_4}))\} = \bigcup_{k=1}^3 \{\min(\{1, 3, 5, 6, 8\} \cap \text{row}_k(S_{\rho_4}))\} = \{1, 5, 8\}$$

shaded yellow and $\{3, 6\}$, the remainder of entries of $E_3(T)$, shaded gray. Thus any $I \subseteq \{1, 5, 8\}$ is 4-slidable, so $\{1, 8\}$ is 4-slidable but $\{1, 3, 8\}$ is not.



The proof of this claim is lengthy and will appear elsewhere:

Claim 10.9. Fix $T \in \text{shEqSYT}(\rho_n/\rho_n, N)$. Then $\text{shEqRect}(T) = S_{\rho_n}$ if and only if $T = \text{Sl}_{I_{n-1}} \circ \text{Sl}_{I_{n-2}} \circ \dots \circ \text{Sl}_{I_1}(U_n)$ where each $I_i \subseteq E_i(\text{Sl}_{I_{i-1}} \circ \dots \circ \text{Sl}_{I_1}(U_n))$ is n -slidable.

By Claim 10.9, $d_{\rho_n, \rho_n}^{\rho_n}$ equals the number of sequences $\{I_i\}_{i=1}^{n-1}$ where

$$I_i \subseteq E_i(\text{Sl}_{I_{i-1}} \circ \dots \circ \text{Sl}_{I_1}(U_n))$$


is n -slidable. We assert that

$$i = \# \bigcup_{k=1}^i \{\min(E_i(\text{Sl}_{I_{i-1}} \circ \dots \circ \text{Sl}_{I_1}(U_n)) \cap \text{row}_k(S_{\rho_n}))\}.$$


Indeed, to see this, note that $i = \# \bigcup_{k=1}^i \{\min(E_i(U_n) \cap \text{row}_k(S_{\rho_n}))\}$ and, by definition of I -slidable, $E_i(\text{Sl}_{I_{i-1}} \circ \dots \circ \text{Sl}_{I_1}(U_n)) \supseteq E_i(U_n)$. Hence, by (37), there are 2^i choices for each n -slidable I_i , so $d_{\rho_n, \rho_n}^{\rho_n} = 2^{\binom{n}{2}}$, as desired. \square

We illustrate Claim 10.9 with the following example:

Example 10.10. Below is $T = \text{Sl}_{I_3} \circ \text{Sl}_{I_2} \circ \text{Sl}_{I_1}(U_4)$ with the choices of I_i given above each arrow. Beneath each arrow, entries in $\bigcup_{k=1}^i \{\min(E_i(\text{Sl}_{I_{i-1}} \circ \dots \circ \text{Sl}_{I_1}(U_4)) \cap \text{row}_k(S_{\rho_n}))\}$ are shaded yellow in S_{ρ_4} and the remaining entries of $E_i(T)$ are shaded gray. Thus I_i is 4-slidable if and only if all entries of I_i are yellow. Thus in the example below, I_1, I_2 , and I_3 are all 4-slidable. Therefore by Lemma 10.9, $\text{shEqRect}(T) = S_{\rho_4}$.



However, in the example below, I'_1, I'_2 are 4-slidable, but I'_3 is not. Thus by Claim 10.9, $\text{shEqRect}(T') \neq S_{\rho_4}$.



While $2^{\binom{n}{2}}$ is the number of labeled graphs on n vertices, consulting the Online Encyclopedia of Integer Sequences [60], one also finds that it counts the number of

- perfect matchings of order n Aztec diamond [61],
- Gelfand-Zeitlin patterns with bottom row $[1, 2, 3, \dots, n]$ [73], and
- certain domino tilings [60, A006125]

among other things.

We end with a problem of enumerative combinatorics:

Problem 10.11. Give bijections between the shifted edge labeled tableaux counted by $d_{\rho_n, \rho_n}^{\rho_n}$ and the equinumerous objects above.

ACKNOWLEDGEMENTS

AY's thanks to Bill Fulton goes back to 1999. In May 2018, the authors attended a conference on Schubert calculus held at Ohio State University where Bill kindly shared the fine points of his ongoing work with David Anderson. This was the stimulus for this chapter. We also thank David Anderson, Soojin Cho, Sergey Fomin, Allen Knutson, William Linz, Alun Morris, Gidon Orelowitz, John Stembridge, Hugh Thomas, Brian Shin, and the anonymous referees for helpful remarks. We thank Anshul Adve, David Anderson, Cara Monical, Oliver Pechenik, Ed Richmond, and Hugh Thomas for their contributions reported here. AY was partially supported by an NSF grant, a UIUC Campus research board grant, and a Simons Collaboration Grant. This material is based upon work of CR supported by the National Science Foundation Graduate Research Fellowship Program under Grant No. DGE – 1746047.

REFERENCES

- [1] A. Adve, C. Robichaux and A. Yong, *Vanishing of Littlewood–Richardson polynomials is in P*, Comput. Complexity 28 (2019), no. 2, 241–257.
- [2] D. Anderson, *Introduction to Equivariant Cohomology in Algebraic Geometry*, Contributions to Algebraic Geometry: Impanga Lecture Notes, 2012.
- [3] D. Anderson, E. Richmond and A. Yong, *Eigenvalues of Hermitian matrices and equivariant cohomology of Grassmannians*, Compos. Math. 149 (2013), no. 9, 1569–1582.
- [4] H. Andersen, J. Jantzen and W. Soergel., *Representations of quantum groups at a p th root of unity and of semisimple groups in characteristic p : independence of p* , Asterisque No. 220 (1994), 321 pp.
- [5] A. Arabia, *Cohomologie T -équivariante de la variété de drapeaux d'un groupe de Kac–Moody*, Bull. Math. Soc. France 117 (1989), 129–165.

- [6] P. Belkale, *Local systems on $\mathbb{P}^1 \setminus S$ for a finite set*, Ph. D. thesis, University of Chicago, 1999.
- [7] ———, *Local systems on $\mathbb{P}^1 \setminus S$ for S a finite set*, *Compositio Math.* 129 (2001), no. 1, 67–86.
- [8] ———, *Geometric proofs of Horn and saturation conjectures*, *J. Algebraic Geom.* 15 (2006), no. 1, 133–173.
- [9] ———, *Extremal rays in the Hermitian eigenvalue problem*, *Math. Ann.* 373 (2019), no. 3-4, 1103–1133.
- [10] R. Bhatia, *Linear algebra to quantum cohomology: the story of Alfred Horn’s inequalities*, *Amer. Math. Monthly* 108 (2001), no. 4, 289–318.
- [11] A. S. Buch, *The saturation conjecture (after A. Knutson and T. Tao)*. With an appendix by William Fulton. *Enseign. Math.* (2) 46 (2000), no. 1-2, 43–60.
- [12] A. S. Buch, A. Kresch and H. Tamvakis, *Littlewood–Richardson rules for Grassmannians*, *Adv. Math.* 185 (2004), no. 1, 80–90.
- [13] J. A. De Loera and T. B. McAllister, *On the computation of Clebsch–Gordan coefficients and the dilation effect*, *Experiment. Math.*, 15(1):7–19, 2006.
- [14] H. Derksen and J. Weyman, *Semi-invariants of quivers and saturation for Littlewood–Richardson coefficients*, *J. Amer. Math. Soc.* 13 (2000), no. 3, 467–479.
- [15] A. G. Elashvili, *Invariant algebras*, *Advances in Soviet Math.*, 8 (1992), 57–64.
- [16] S. Fomin and C. Greene, *Noncommutative Schur functions and their applications*, *Discrete Mathematics* Volume 193, Issues 1-3, 28 November 1998, 179–200.
- [17] S. Friedland, *Finite and infinite dimensional generalizations of Klyachko’s theorem*, *Linear Algebra Appl.*, 319 (2000), 3–22.
- [18] W. Fulton, *Young tableaux*. With applications to representation theory and geometry. London Mathematical Society Student Texts, 35. Cambridge University Press, Cambridge, 1997.
- [19] ———, *Eigenvalues of majorized Hermitian matrices and Littlewood–Richardson coefficients*, *Linear Algebra Appl.* 319 (2000), no. 1–3, 23–36.
- [20] ———, *Eigenvalues, invariant factors, highest weights, and Schubert calculus*, *Bull. Amer. Math. Soc. (N.S.)* 37 (2000), no. 3, 209–249 (electronic).
- [21] M. Goresky, R. Kottwitz and R. MacPherson, *Equivariant cohomology, Koszul duality, and the localization theorem*, *Invent. Math.* 131 (1998), no. 1, 25–83.
- [22] W. Graham, *Positivity in equivariant Schubert calculus*, *Duke Math. J.* 109 (2001), no. 3, 599–614.
- [23] M. Grotschel, L. Lovasz and A. Schrijver, *Geometric algorithms and combinatorial optimization*, Springer Verlag, 1993.
- [24] M. D. Haiman, *Dual equivalence with applications, including a conjecture of Proctor*, *Discrete Math.* 99 (1992), no. 1-3, 79–113.
- [25] A. Horn, *Eigenvalues of sums of Hermitian matrices*, *Pacific J. Math.*, 12 (1962), 225–241.
- [26] T. Ikeda, *Schubert classes in the equivariant cohomology of the Lagrangian Grassmannian*, *Adv. Math.* 215 (2007), no. 1, 1–23.
- [27] T. Ikeda, L. Mihalcea, and H. Naruse, *Double Schubert polynomials for the classical groups*, *Adv. Math.* 226 (2011), no. 1, 840–886.
- [28] T. Ikeda and H. Naruse, *Excited Young diagrams and equivariant Schubert calculus*, *Trans. Amer. Math. Soc.* 361 (2009), no. 10, 5193–5221.
- [29] S. L. Kleiman, *The transversality of a general translate*, *Compositio Math.* 28 (1974), 287–297.
- [30] A. A. Klyachko, *Stable vector bundles and Hermitian operators*, *Selecta Math. (N.S.)* 4 (1998), 419–445.
- [31] A. Knutson, E. Miller and A. Yong, *Gröbner geometry of vertex decompositions and of flagged tableaux*, *J. Reine Angew. Math.* 630 (2009), 1–31.
- [32] A. Knutson and T. Tao, *The honeycomb model of $GL_n(\mathbb{C})$ tensor products I: proof of the saturation conjecture*, *J. Amer. Math. Soc.* 12 (1999), 1055–1090.
- [33] ———, *Puzzles and (equivariant) cohomology of Grassmannians*, *Duke Math. J.* 119 (2003), no. 2, 221–260.
- [34] A. Knutson, T. Tao and C. Woodward, *The honeycomb model of $GL_n(\mathbb{C})$ tensor products. II. Puzzles determine facets of the Littlewood–Richardson cone*, *J. Amer. Math. Soc.* 17 (2004), no. 1, 19–48.
- [35] ———, *A positive proof of the Littlewood–Richardson rule using the octahedron recurrence*, *Electron. J. Combin.* 11 (2004), no. 1, Research Paper 61, 18 pp.
- [36] A. Knutson and P. Zinn-Justin, *Schubert puzzles and integrability I: invariant trilinear forms*, preprint, 2017. arXiv:1706.10019
- [37] S. Kumar, *Kac–Moody groups, their flag varieties and representation theory*, *Progress in Mathematics*, 204. Birkhäuser Boston, Inc., Boston, MA, 2002. xvi+606 pp.

- [38] B. Kostant and S. Kumar, *T-equivariant K-theory of generalized flag varieties*, J. Differential Geom. 32 (1990), no. 2, 549–603.
- [39] V. Kreiman, *Equivariant Littlewood–Richardson skew tableaux*, Trans. Amer. Math. Soc. 362 (2010), no. 5, 2589–2617.
- [40] C. Lenart and A. Postnikov, *Affine Weyl groups in K-theory and representation theory*, Int. Math. Res. Not. IMRN 2007, no. 12, Art. ID rnm038, 65 pp.
- [41] E. Miller and B. Sturmfels, *Combinatorial commutative algebra*. Graduate Texts in Mathematics, 227. Springer-Verlag, New York, 2005. xiv+417 pp.
- [42] A. I. Molev, *Littlewood–Richardson polynomials*, J. Algebra 321 (2009), no. 11, 3450–3468.
- [43] A. Molev and B. Sagan, *A Littlewood–Richardson rule for factorial Schur functions*, Trans. Amer. Math. Soc. 351 (1999), no. 11, 4429–4443.
- [44] C. Monical, *Polynomials in algebraic combinatorics*, Ph.D thesis, University of Illinois at Urbana-Champaign, 2018.
- [45] C. Monical, N. Tokcan and A. Yong, *Newton polytopes in algebraic combinatorics*, preprint, 2017. arXiv:1703.02583
- [46] K. D. Mulmuley, H. Narayanan and M. Sohoni, *Geometric complexity theory III: on deciding nonvanishing of a Littlewood–Richardson coefficient*, J. Algebraic Combin. 36 (2012), no. 1, 103–110.
- [47] H. Narayanan, *On the complexity of computing Kostka numbers and Littlewood–Richardson coefficients*, J. Algebraic Combin. 24 (2006), no. 3, 347–354.
- [48] O. Pechenik and A. Yong, *Equivariant K-theory of Grassmannians II: the Knutson–Vakil conjecture*, Compos. Math. 153 (2017), no. 4, 667–677.
- [49] ———, *Equivariant K-theory of Grassmannians*, Forum Math. Pi, 5 (2017), e3, 128 pp.
- [50] P. Pragacz, *Algebro-geometric applications of Schur S- and Q-polynomials*. Topics in invariant theory (Paris, 1989/1990), 130–191, Lecture Notes in Math., 1478, Springer, Berlin, 1991.
- [51] K. Purbhoo, *Root games on Grassmannians*, J. Algebraic Combin. 25 (2007), no. 3, 239–258.
- [52] ———, *Vanishing and nonvanishing criteria in Schubert calculus*, Int. Math. Res. Not. 2006, Art. ID 24590, 38 pp.
- [53] K. Purbhoo and F. Sottile, *The Horn recursion for Schur P- and Q- functions*, FPSAC 2006.
- [54] ———, *The recursive nature of cominuscule Schubert calculus*, Adv. Math., 217(2008), 1962–2004.
- [55] C. Robichaux, H. Yadav and A. Yong, *The A·B·C·Ds of Schubert calculus*, preprint, 2019. arXiv:1906.03646
- [56] B. E. Sagan, *Shifted tableaux, Schur Q-functions, and a conjecture of R. Stanley*, J. Combin. Theory Ser. A 45 (1987), no. 1, 62–103.
- [57] A. Schrijver, *Theory of linear and integer programming*, Wiley-Interscience Series in Discrete Mathematics. A Wiley-Interscience Publication. John Wiley & Sons, Ltd., Chichester, 1986. xii+471 pp.
- [58] M.-P. Schützenberger, *La correspondance de Robinson*. (French) Combinatoire et représentation du groupe symétrique (Actes Table Ronde CNRS, Univ. Louis-Pasteur Strasbourg, Strasbourg, 1976), pp. 59–113. Lecture Notes in Math., Vol. 579, Springer, Berlin, 1977.
- [59] J. Schur, *Über die Darstellung der symmetrischen und der alternierenden Gruppe durch gebrochene lineare Substitutionen*, (German) J. Reine Angew. Math. 139 (1911), 155–250.
- [60] N. J. A. Sloane, editor, *The On-Line Encyclopedia of Integer Sequences*, published electronically at <https://oeis.org>
- [61] D. Speyer, *Perfect matchings and the octahedron recurrence*, J. Algebraic Combin. 25 (2007), no. 3, 309–348.
- [62] R. P. Stanley, *Enumerative combinatorics. Vol. 2*. With a foreword by Gian-Carlo Rota and appendix 1 by Sergey Fomin. Cambridge Studies in Advanced Mathematics, 62. Cambridge University Press, Cambridge, 1999. xii+581.
- [63] J. R. Stembridge, *Shifted Tableaux and the Projective Representations of Symmetric Groups*, Adv. Math. 74, 87–134 (1989)
- [64] É. Tardos, *A Strongly Polynomial Algorithm to Solve Combinatorial Linear Programs*, Operations Research, Vol. 34, No. 2 (Mar. - Apr., 1986), 250–256.
- [65] H. Thomas and A. Yong, *A combinatorial rule for (co)minuscule Schubert calculus*, Adv. Math. 222 (2009), no. 2, 596–620.
- [66] ———, *Equivariant Schubert calculus and jeu de taquin*, Ann. Inst. Fourier (Grenoble) 68 (2018), no. 1, 275–318.
- [67] ———, *A combinatorial rule for (co)minuscule Schubert calculus*, Adv. Math. 222 (2009), no. 2, 596–620.

- [68] B. Totaro, *Tensor products of semistables are semistable*, in Geometry and Analysis on Complex Manifolds, Festschrift for Professor S. Kobayashi's 60th Birthday, ed. T. Noguchi, J. Noguchi, and T. Ochiai, World Scientific Publ. Co., Singapore, 1994, 242–250.
- [69] L. W. Tu, *What is ... equivariant cohomology?* Notices Amer. Math. Soc. 58 (2011), no. 3, 423–426.
- [70] J. S. Tymoczko, *An introduction to equivariant cohomology and homology, following Goresky, Kottwitz, and Macpherson* preprint, 2005. arXiv:0503369
- [71] L. G. Valiant, *The complexity of computing the permanent*, Theoret. Comput. Sci. 8 (1979), no. 2, 189–201.
- [72] D. R. Worley, *A theory of shifted Young tableaux*, Thesis (Ph.D.), Massachusetts Institute of Technology. ProQuest LLC, Ann Arbor, MI, 1984.
- [73] D. Zeilberger, *Dave Robbins's Art of Guessing*, Adv. in Appl. Math. 34 (2005), 939–954.
- [74] A. Zelevinsky, *Littlewood–Richardson semigroups*, New perspectives in algebraic combinatorics (Berkeley, CA, 1996–97), 337–345, Math. Sci. Res. Inst. Publ., 38, Cambridge Univ. Press, Cambridge, 1999.
- [75] P. Zinn-Justin, *Littlewood–Richardson coefficients and integrable tilings*, Electron. J. Combin. 16 (2009), no. 1, Research Paper 12, 33 pp.

DEPT. OF MATHEMATICS, UNIVERSITY OF ILLINOIS AT URBANA-CHAMPAIGN, URBANA, IL 61801

Email address: cer2@illinois.edu, yadav7@illinois.edu, ayong@illinois.edu