

# A Correction Note: Attractive Nearest Neighbor Spin Systems on the Integers

Jeffrey Lin (UCLA)

October 4, 2014

## Abstract

In this note, we discuss a proof in T. Liggett's work on attractive translation invariant nearest neighbor spin systems on the integers. A correction to a wrong estimate is provided.

## 1 Introduction

For the purposes of this note, an attractive translation invariant nearest neighbor spin system is a certain kind of Feller process defined on the compact state space  $X = \{0, 1\}^{\mathbb{Z}}$  with rates satisfying the attractiveness inequalities, and depending only on the nearest neighbors. (See [2], p.144-145 for more details.)

T. Liggett proves in [1] and [2], (p. 152 Theorem 3.13) the following theorem:

**Theorem 1.** *All attractive translation invariant nearest neighbor spin systems on the integers have only the minimal and maximal invariant measures (ordered stochastically) as the extremal invariant measures.*

We discuss the estimate in [2] of lemma 3.7 part (e). It is wrong, but the similar estimate  $\epsilon \int g_{m,n}^{l+1} d\nu \leq (4Kl + 2\epsilon) \int g_{m,n}^l d\nu$  is valid. This change does not affect the proof moving forward, because the only time the estimate is used is in the proof of lemma 3.10 in [2], where it is used to derive that  $\sup_{m \leq n} \int g_{m,n}^l d\nu < \infty$ . This fact still holds with the new estimate. The same line of reasoning appears in [1] within the proof of lemma 2.2 there.

We will refer to [2] from now on.

## 2 A Correction

The problem with the estimate as written is at the top of p.150 of [2] in the discussion of bounding below the positive contribution to  $\tilde{\Omega} g_{m,n}^l$ . The argument there fails to consider the possibility that change in the  $\gamma$  coordinate at any of the  $x_i$  among the left and right endpoints in the  $l + 1$  length intervals may not only create a length  $l$  interval, but also destroy an adjacent length  $l$  interval. But this can only happen at at most  $2g_{m,n}^l$  such sites, i.e. the left and right

neighbors of intervals of length  $l$ . The bound below on the rate of the type of flip in question is still correctly stated as  $\epsilon$  so as long as we replace  $\epsilon(g_{m,n}^{l+1})$  by  $\epsilon(g_{m,n}^{l+1} - 2g_{m,n}^l)$  the estimate is correct. This results in the display reading

$$\tilde{\Omega}g_{m,n}^l \geq \epsilon(g_{m,n}^{l+1} - 2g_{m,n}^l) - 4Klg_{m,n}^l.$$

The author would like to thank T. Liggett for his input on these matters and NSF, for part of the funding of this project.

## References

- [1] Liggett, T. (1978). Attractive Nearest Neighbor Spin Systems on the Integers. *Ann. Probability* Vol. 6 No. 4, p.629-636.
- [2] Liggett, T. (1985). *Interacting Particle Systems*. Springer Berlin Heidelberg 2005. p.143-152 and Ch. 1.