

Endomorphisms of Fano 3-folds and log Bott vanishing

Burt Totaro

A basic problem of algebraic dynamics is to determine which projective varieties admit an endomorphism of degree greater than 1. To avoid degenerate cases, we focus on *int-amplified* endomorphisms $f: X \rightarrow X$, meaning that there is an ample divisor H such that $f^*H - H$ is ample [Men20, MZ20]. Fakhruddin, Meng, Zhang, and Zhong conjectured that a smooth complex projective rationally connected variety that admits an int-amplified endomorphism must be a toric variety [Fak03, Question 4.4], [MZZ22, Question 1.1]. Kawakami and I introduced a new approach to this problem, showing that a variety with an int-amplified endomorphism must satisfy *Bott vanishing* [KT23, Theorem C].

In this paper, we extend that result to a logarithmic version of Bott vanishing for an endomorphism with a totally invariant divisor (Theorem 3.1). We also connect log Bott vanishing with some related problems: which varieties are images of toric varieties (Theorem 5.1)? Which varieties admit morphisms of unbounded degree from some other variety (Theorem 4.1)?

We apply these results to Fano 3-folds. Meng, Zhang, and Zhong showed that the only smooth complex Fano 3-folds that admit an int-amplified endomorphism are the toric ones [MZZ22, Theorem 1.4]. Also, Achinger, Witaszek, and Zdanowicz showed that the only smooth complex Fano 3-folds that are images of toric varieties are the toric ones [AWZ21, proof of Theorem 4.4.1], [AWZ23, Theorems 6.9 and 7.7]. Using log Bott vanishing, we reprove both results and extend them to characteristic p , for morphisms of degree prime to p (Theorems 6.1 and 7.2). This resolves [KT23, Question 1.6]. For Fano 3-folds with Picard number 1, these extensions already appeared in [KT23, Theorem A and Proposition 3.10].

1 Notation

A Weil divisor (with integer coefficients) on a normal projective variety is called *ample* if some positive multiple is an ample Cartier divisor. A *contraction* of a normal variety X over a field k is a proper morphism $\pi: X \rightarrow Y$ with $\pi_*\mathcal{O}_X = \mathcal{O}_Y$. For a projective variety X over k , $N_1(X)$ is the vector space of 1-cycles with real coefficients modulo numerical equivalence, which has finite dimension. The *Néron-Severi* space $N^1(X)$ is the space of **R**-Cartier divisors modulo numerical equivalence, and so $N^1(X) = N_1(X)^*$.

A morphism of varieties $f: Y \rightarrow X$ over a field k is *separable* if it is dominant and the function field $k(Y)$ is a separable field extension of $k(X)$. For k algebraically closed, f is separable if and only if the derivative of f is surjective at some smooth point of Y . For an endomorphism f of a variety X , a closed subset S of X is *totally invariant* under f if $f^{-1}(S) = S$.

For a normal variety X over a field k and $i \geq 0$, we write Ω_X^i for $\Omega_{X/k}^i$. The sheaf of *reflexive differentials* $\Omega_X^{[i]}$ is the double dual $(\Omega_X^i)^{**}$. For a reduced divisor D on X , $\Omega_X^{[i]}(\log D)$ (reflexive differentials with *log poles* along D) denotes the sheaf of i -forms α on the smooth locus of $X - D$ such that both α and $d\alpha$ have at most a simple pole along each component of D . For a reflexive sheaf M and a Weil divisor E on X , we write $M(E)$ for the reflexive sheaf $(M \otimes_{\mathcal{O}_X} \mathcal{O}_X(E))^{**}$. If X is smooth over k , then $\mathcal{O}_X(E)$ is a line bundle, and $M(E)$ is just the tensor product $M \otimes_{\mathcal{O}_X} \mathcal{O}_X(E)$.

2 The trace of a differential form with log poles

Here is the key lemma for this paper, extending the proof of [KT23, Theorem C] to allow log poles.

Lemma 2.1. *Let $f: Y \rightarrow X$ be a finite surjective morphism of normal varieties over a perfect field k . Let A be a Weil divisor on X . Let E_X and D_X be reduced divisors on X with $0 \leq E_X \leq D_X$. Let D_Y be the sum of the components of $f^{-1}(D_X)$ along which the ramification degree e of f is invertible in k , and let E_Y be the corresponding divisor inside $f^{-1}(E_X)$. Then, for every $i \geq 0$, the pullback and pushforward of differential forms give maps of reflexive sheaves*

$$\Omega_X^{[i]}(\log D_X)(A - E_X) \rightarrow f_* (\Omega_Y^{[i]}(\log D_Y)(f^* A - E_Y)) \rightarrow \Omega_X^{[i]}(\log D_X)(A - E_X),$$

with composition equal to multiplication by $\deg(f)$.

In characteristic p , even if the morphism f in Lemma 2.1 has degree prime to p , it may be wildly ramified along some divisors. That requires the careful choice of D_Y and E_Y in the statement, in order to get something true.

Proof. Since k is perfect, the normal varieties X and Y over k are geometrically normal [SPA24, Tag 038O], hence smooth over k in codimension 1.

As in the statement, let D_Y be the sum of the components of $f^{-1}(D_X)$ along which the ramification degree e of f is invertible in k . (The *ramification degree* e along a component D_2 of $f^{-1}(D_X)$ is the coefficient of D_2 in the pullback Weil divisor $f^* D_X$, which is defined since f is finite and D_X is generically Cartier. The condition that e is invertible in k is not the same as f being tamely ramified along D_2 ; tame ramification would mean that e is invertible in k and that $f: D_2 \rightarrow f(D_2)$ is separable.) Likewise, let E_Y be the sum of the components of $f^{-1}(E_X)$ along which the ramification degree of f is invertible in k .

We want to construct pullback and pushforward maps of reflexive sheaves:

$$\Omega_X^{[i]}(\log D_X)(-E_X) \rightarrow f_* \Omega_Y^{[i]}(\log D_Y)(-E_Y) \rightarrow \Omega_X^{[i]}(\log D_X)(-E_X).$$

First consider the pullback map. Since we are mapping to a reflexive sheaf, it suffices to define this map outside the subset $X^{\text{sing}} \cup f(Y^{\text{sing}})$, which has codimension at least 2 in X . The map is the usual pullback of differential forms outside D_X . So it suffices to show that the pullback sends $\Omega_X^i(\log D_X)(-E_X)$ into $\Omega_Y^i(\log D_Y)(-E_Y)$ near the generic point of each component D_2 of $f^{-1}(D_X)$. Let $D_1 = f(D_2)$.

There are four cases.

Case 1 D_1 not in E_X , $e \in k^*$. Then we need to show that $\Omega_X^i(\log D_1)$ pulls back into $\Omega_Y^i(\log D_2)$ near the generic point of D_2 . Let g be a local defining function of D_1 in X and h a local defining function of D_2 in Y . We have $f^*(g) = h^e u$ for some unit $u \in \mathcal{O}_{Y, D_2}^*$. Then $f^*(\frac{dg}{g}) = e\frac{dh}{h} + \frac{du}{u}$, and so it is clear that forms in $\Omega_X^i(\log D_1) = \Omega_X^i + \frac{dg}{g}\Omega_X^{i-1}$ pull back to $\Omega_Y^i(\log D_2) = \Omega_Y^i + \frac{dh}{h}\Omega_Y^{i-1}$ near the generic point of D_2 .

Case 2 D_1 not in E_X , $e = 0 \in k$. Then we need to show that forms in $\Omega_X^i(\log D_1)$ pull back to Ω_Y^i near the generic point of D_2 . This is clear by the formulas in Case 1, using that $e = 0 \in k$.

Case 3 D_1 in E_X , $e \in k^*$. Then we need to show that $\Omega_X^i(\log D_1)(-D_1)$ pulls back into $\Omega_Y^i(\log D_2)(-D_2)$ near the generic point of D_2 . This follows from Case 1, using that g pulls back to a unit times a positive power of h .

Case 4 D_1 in E_X , $e = 0 \in k$. Then we need to show that $\Omega_X^i(\log D_1)(-D_1)$ pulls back into Ω_Y^i near the generic point of D_2 . This is clear, using that $\Omega_X^i(\log D_1)(-D_1)$ is contained in Ω_X^i .

Next, we want to define the *pushforward* (or *trace*) map

$$f_*\Omega_Y^{[i]}(\log D_Y)(-E_Y) \rightarrow \Omega_X^{[i]}(\log D_X)(-E_X).$$

Outside $D_X \cup X^{\text{sing}} \cup f(Y^{\text{sing}})$, this trace map was defined by Garel and Kunz [Gar84], [Kun86, section 16], [SPA24, Tag 0FLC]. (They assume that the finite morphism f is flat with lci fibers; that holds on the open set mentioned, since both X and Y are smooth there [SPA24, Tags 00R4 and 09Q7].) Since we are mapping into a reflexive sheaf, it remains to check that the trace map above is regular near the generic point of each component D_1 of D_X . It suffices to check this after replacing X by an elementary étale neighborhood [SPA24, Tag 02LE] of the generic point of D_1 , in such a way that Y becomes a disjoint union of varieties containing the different components D_2 of $f^{-1}(D_1)$. (Then the trace from Y to X is the sum of the traces for these different varieties.) We can work on one of these varieties; that is, we can assume that $f^{-1}(D_1)$ is an irreducible divisor D_2 .

There are four cases.

Case 1 D_1 not in E_X , $e \in k^*$. Then we need to show that the pushforward of a form in $\Omega_Y^i(\log D_2)$ lies in $\Omega_X^i(\log D_1)$ near the generic point of D_1 . We have $\Omega_Y^i(\log D_2) = \Omega_Y^i + \frac{dh}{h}\Omega_Y^{i-1}$. Since $f^*(g) = h^e u$ in the notation above, we have $e\frac{dh}{h} = f^*(\frac{dg}{g}) - \frac{du}{u}$. Since $e \in k^*$, it follows that $\Omega_Y^i(\log D_2) = \Omega_Y^i + f^*(\frac{dg}{g})\Omega_Y^{i-1}$. By the projection formula [SPA24, Tag 0FLC], the trace of these forms lies in $\Omega_X^i + \frac{dg}{g}\Omega_X^{i-1} = \Omega_X^i(\log D_1)$.

Case 2 D_1 not in E_X , $e = 0 \in k$. Then we need to show that forms in Ω_Y^i push forward to $\Omega_X^i(\log D_1)$ near the generic point of D_1 . This is easy, since the pushforward is contained in $\Omega_X^i \subset \Omega_X^i(\log D_1)$.

For cases 3 and 4, we use the following property of the trace map on differential forms [Hüb89, Remark 5.7]:

Proposition 2.2. *Let $f: Y \rightarrow X$ be a finite flat morphism with lci fibers over a field k . Suppose that D_1 is an irreducible divisor in X such that $D_2 := f^{-1}(D_1)$ is also irreducible. Let e be the ramification degree of f along D_2 . Then the following diagram commutes near the generic point of D_1 :*

$$\begin{array}{ccc} \Omega_Y^* & \longrightarrow & \Omega_{D_2}^* \\ \downarrow \text{tr}_f & & \downarrow e \text{tr}_f \\ \Omega_X^* & \longrightarrow & \Omega_{D_1}^*. \end{array}$$

In our situation (above), the flatness and lci assumptions of Proposition 2.2 hold near the generic points of D_1 and D_2 , because X and Y are smooth there.

Case 3 D_1 in E_X , $e \in k^*$. Then we need to show that all the forms in $\Omega_Y^i(\log D_2)(-D_2)$ push forward into $\Omega_X^i(\log D_1)(-D_1)$ near the generic point of D_1 . That is, we want to show that the trace of $h\Omega_Y^i + h\frac{dh}{h}\Omega_Y^{i-1}$ is contained in $g\Omega_X^i + g\frac{dg}{g}\Omega_X^{i-1}$. Proposition 2.2 gives that the trace of $(h, dh)\Omega_Y^*$ is contained in $(g, dg)\Omega_X^*$, which proves the claim.

Case 4 D_1 in E_X , $e = 0 \in k$. Then we need to show that forms in Ω_Y^i push forward to $\Omega_X^i(\log D_1)(-D_1)$ near the generic point of D_1 . That is, we want $\text{tr}_f(\Omega_Y^i)$ to be contained in $(g, dg)\Omega_X^*$. This follows from Proposition 2.2, using that $e = 0 \in k$. Thus we have constructed the pushforward map in all cases.

We now return to the original morphism $f: Y \rightarrow X$ (before we restricted to an étale neighborhood of X). The composition of pullback and pushforward is multiplication by $\deg(f)$, as one can check on an open subset where X and Y are smooth, by the projection formula: for a form α on Y , $\text{tr}_f(f^*\alpha) = \text{tr}_f(1)\alpha = \deg(f)\alpha$ [SPA24, Tag 0FLC].

Tensoring the maps above with $\mathcal{O}_X(A)$ and taking double duals, we have pullback and pushforward maps

$$\Omega_X^{[i]}(\log D_X)(A - E_X) \rightarrow f_*(\Omega_Y^{[i]}(\log D_Y)(f^*A - E_Y)) \rightarrow \Omega_X^{[i]}(\log D_X)(A - E_X),$$

with composition equal to multiplication by $\deg(f)$, as we want. (Note that f^*A is a Weil divisor (with integer coefficients), because f is a finite surjective morphism between normal varieties.) \square

3 Endomorphisms and log Bott vanishing

Kawakami and I showed that a projective variety with a suitable endomorphism must satisfy Bott vanishing [KT23, Theorem C]. (The endomorphism is assumed to be int-amplified and of degree invertible in the base field.) We now prove a logarithmic generalization, when the endomorphism has a totally invariant divisor. This generalizes Fujino's work on the case of toric varieties [Fuj08, Theorem 1.3]. Indeed, every toric variety has an action of the multiplicative monoid of positive integers, and every toric divisor is totally invariant under those endomorphisms.

Theorem 3.1. *Let X be a normal projective variety over a perfect field k . Suppose that X admits an int-amplified endomorphism f whose degree is invertible in k . Let*

D be a reduced divisor on X such that $f^{-1}(D) \subset D$. Then (X, D) satisfies log Bott vanishing. That is,

$$H^j(X, \Omega_X^{[i]}(\log D)(A - E)) = 0$$

for every reduced divisor E with $0 \leq E \leq D$, $i \geq 0$, $j > 0$, and A an ample Weil divisor.

It may seem artificially strong to assume that an endomorphism has a totally invariant divisor. But in fact, this property comes up naturally in classifying varieties with endomorphisms. See [MZZ22] or the proof of Theorem 6.1.

Theorem 3.1 is the most general vanishing property I know how to prove, for a variety with an endomorphism. The idea of allowing any divisor $0 \leq E \leq D$ was suggested by Fujino's result for toric varieties [Fuj08, Theorem 1.3].

Remark 3.2. Bott vanishing can fail if we only assume that f is int-amplified and separable (rather than of degree invertible in k), by [KT23, Proposition 5.1]. A fortiori, log Bott vanishing can fail in that situation. (Bott vanishing clearly fails for inseparable endomorphisms, since *every* projective variety over a finite field has the Frobenius endomorphism, which is int-amplified.)

Proof. (Theorem 3.1) Since f is int-amplified, there is an ample Cartier divisor H on X such that $f^*H - H$ is ample. In particular, f^*H is ample, and so f does not contract any curves. So $f: X \rightarrow X$ is finite. Since $f^{-1}(D) \subset D$ and f is surjective, we have $f^{-1}(D) = D$.

Since $f^{-1}(D) = D$, f permutes the (finitely many) irreducible components of D . After replacing f by a positive iterate, we can assume that f maps each component of D to itself. So $f^{-1}(D_1) = D_1$, for each component D_1 of D . In particular, since E is a reduced divisor with $0 \leq E \leq D$, we now have that $f^{-1}(E) = E$. (In what follows, we use only that $f^{-1}(E) = E$, not that f maps each component of D to itself. This makes the proof clearer, in terms of notation.)

We are given that the degree of f is invertible in k . The inverse image of each irreducible component D_1 of D is a single irreducible component D_2 of D . Therefore, the degree of $f: X \rightarrow X$ is the product of the degree of $f: D_2 \rightarrow D_1$ and the ramification degree of f along D_2 . It follows that this ramification degree is invertible in k and that $k(D_2)$ is a finite separable extension of $k(D_1)$ via f . That is, f is tamely ramified over each component of D .

Let A be an ample Weil divisor on X . By Lemma 2.1, we have pullback and pushforward maps

$$\Omega_X^{[i]}(\log D)(A - E) \rightarrow f_*(\Omega_X^{[i]}(\log D)(f^*A - E)) \rightarrow \Omega_X^{[i]}(\log D)(A - E),$$

with composition equal to multiplication by $\deg(f)$. (Note that f^*A is a Weil divisor (with integer coefficients), because f is a finite surjective morphism between normal varieties.)

Given this, the proof of [KT23, Theorem C] works. Namely, since $\deg(f)$ is invertible in k , it follows that the pullback map is split injective as a map of O_X -modules. Let $j > 0$. Taking cohomology (and using that f is finite), it follows that $H^j(X, \Omega_X^{[i]}(\log D)(A - E)) \rightarrow H^j(X, \Omega_X^{[i]}(\log D)(f^*A - E))$ is split injective. The same argument works for any iterate f^e with $e \geq 1$.

Using that f is int-amplified, we showed that the iterates $(f^e)^*(A)$ become arbitrarily large in the ample cone of X , as e goes to infinity [KT23, proof of Theorem C]. By Fujita vanishing for Weil divisors [KT23, Lemma 2.6], it follows that there is an $e \geq 1$ such that $H^j(X, \Omega_X^{[i]}(\log D)((f^e)^*A - E)) = 0$. By the previous paragraph, we have $H^j(X, \Omega_X^{[i]}(\log D)(A - E)) = 0$, as we want. \square

4 Morphisms

Let X and Y be projective varieties of the same dimension with Picard number 1. If X does not satisfy Bott vanishing, then there is an upper bound on the degrees of all morphisms $Y \rightarrow X$ with degree invertible in k [KT23, Proposition 2.7]. We now prove a log version of that result.

Theorem 4.1. *Let X and Y be normal projective varieties of the same dimension over a perfect field k . Assume that X and Y have Picard number 1. Let $D_X \subset X$ and $B_Y \subset Y$ be reduced divisors. Suppose that there are morphisms $f: Y \rightarrow X$ of arbitrarily high degree such that the degree is invertible in k and $f^{-1}(D_X) \subset B_Y$. Then (X, D_X) satisfies log Bott vanishing. That is, we have*

$$H^j(X, \Omega_X^{[i]}(\log D_X)(A - E)) = 0$$

for every reduced divisor $0 \leq E \leq D_X$, $i \geq 0$, $j > 0$, and A an ample Weil divisor on X .

The proof of Theorem 4.1 shows that one can drop the assumption that X and Y have Picard number 1 if one replaces “ f of arbitrarily high degree” by “ f^*H arbitrarily large in the ample cone of Y ”, for a fixed ample Cartier divisor H on X .

Proof. Since each morphism $f: Y \rightarrow X$ has degree invertible in k , the degree is positive. Since Y has Picard number 1, the pullback of an ample Cartier divisor on X is ample on Y , and so f is finite.

For each morphism $f: Y \rightarrow X$, let D_Y be the sum of the components of $f^{-1}(D_X)$ along which the ramification degree of f is invertible in k , and likewise define $E_Y \subset f^{-1}(E_X)$. Since D_Y and E_Y are contained inside the fixed divisor B_Y , we can assume (after passing to a subsequence of the morphisms f) that D_Y and E_Y are the same for all the morphisms $f: Y \rightarrow X$ that we consider.

Let A be an ample Weil divisor on X . By Lemma 2.1, we have pullback and pushforward maps

$$\Omega_X^{[i]}(\log D_X)(A - E) \rightarrow f_*(\Omega_Y^{[i]}(\log D_Y)(f^*A - E_Y)) \rightarrow \Omega_X^{[i]}(\log D_X)(A - E).$$

The composition is multiplication by $\deg(f)$, and so the pullback map is split injective. It follows that $H^j(X, \Omega_X^{[i]}(\log D_X)(A - E))$ injects into $H^j(Y, \Omega_Y^{[i]}(\log D_Y)(f^*A - E_Y))$. Since we have morphisms f of arbitrarily large degree, f^*A becomes arbitrarily large in the ample cone of Y (here just one ray). Therefore, for f of sufficiently large degree, Fujita vanishing for Weil divisors gives that $H^j(Y, \Omega_Y^{[i]}(\log D_Y)(f^*A - E_Y))$ is equal to zero [KT23, Lemma 2.6]. It follows that $H^j(X, \Omega_X^{[i]}(\log D_X)(A - E)) = 0$, as we want. \square

5 Images of toric varieties: log Bott vanishing

A projective variety that is an image of a toric variety (by a morphism of degree invertible in k) must satisfy Bott vanishing [KT23, Proposition 3.10]. We now prove a log version of that result.

Theorem 5.1. *Let X be a normal projective variety over a perfect field k . Suppose that there is a morphism f from a proper toric variety Y onto X . If $Y \rightarrow Y_1 \rightarrow X$ is the Stein factorization of f , assume that the degree of $Y_1 \rightarrow X$ is invertible in k . Let $D_X \subset X$ be a reduced divisor such that $f^{-1}(D_X)$ is a union of toric divisors. Then (X, D_X) satisfies log Bott vanishing. That is, we have*

$$H^j(X, \Omega_X^{[i]}(\log D_X)(A - E)) = 0$$

for every reduced divisor $0 \leq E \leq D_X$, $i \geq 0$, $j > 0$, and A an ample Weil divisor on X .

It may seem unrealistically strong to assume that $f^{-1}(D_X)$ is a union of toric divisors. Nonetheless, the proof of Theorem 7.2 shows how Theorem 5.1 can be used for classifying images of toric varieties.

Proof. Every contraction of a toric variety is toric [Tan22, Proposition 2.7]. (This was known earlier for projective toric varieties [CLS11, Theorem 6.28 and exercise 7.2.3].) So, replacing Y by Y_1 , we can assume that the surjection $f: Y \rightarrow X$ is finite. By our assumptions, the degree of f is invertible in k . Let D_Y be the sum of the components of $f^{-1}(D_X)$ along which the ramification degree e of f is invertible in k . Likewise, let E_Y be the sum of the components of $f^{-1}(E)$ along which the ramification degree of f is invertible in k . By our assumptions, D_Y and E_Y are sums of toric divisors.

Let A be an ample Weil divisor on X . By Lemma 2.1, we have pullback and pushforward maps

$$\Omega_X^{[i]}(\log D_X)(A - E) \rightarrow f_*(\Omega_Y^{[i]}(\log D_Y)(f^*A - E_Y)) \rightarrow \Omega_X^{[i]}(\log D_X)(A - E),$$

with composition equal to multiplication by $\deg(f)$. Since $\deg(f)$ is invertible in k , it follows that the pullback map is split injective as a map of \mathcal{O}_X -modules. Let $j > 0$. Taking cohomology (and using that f is finite), it follows that $H^j(X, \Omega_X^{[i]}(\log D_X)(A - E)) \rightarrow H^j(Y, \Omega_Y^{[i]}(\log D_Y)(f^*A - E_Y))$ is split injective. Since f^*A is ample on Y , the latter cohomology group is zero by log Bott vanishing for toric varieties [Fuj08, Theorem 1.3]. (Alternatively, this follows from Theorem 3.1, using the multiplication maps on a toric variety.) It follows that $H^j(X, \Omega_X^{[i]}(\log D_X)(A - E)) = 0$, as we want. \square

6 Endomorphisms of del Pezzo surfaces and Fano 3-folds

Theorem 6.1. *Let X be a smooth Fano 3-fold over an algebraically closed field k . If X has an int-amplified endomorphism of degree invertible in k , then X is toric.*

The proof uses Tanaka's theorem that the classification of smooth Fano 3-folds has essentially the same form in every characteristic [Tan23b, Theorem 1.1], [Tan23a, Theorem 1.1]. Without that, our proof applies to Fano 3-folds in any characteristic that are given by the same construction as one of the Fano 3-folds over \mathbf{C} (classified by Iskovskikh and Mori-Mukai).

In characteristic zero, Theorem 6.1 was proved earlier by Meng, Zhang, and Zhong [MZZ22, Theorem 1.4]. Here we give a new proof which works in any characteristic. For Fano 3-folds with Picard number 1, this is [KT23, Theorem A]. We also prove an analogous result for del Pezzo surfaces (Proposition 6.4).

To prove Theorem 6.1, our basic idea is to use that X satisfies Bott vanishing, but that is not enough by itself: among the smooth complex Fano 3-folds, 18 are toric while 19 others also satisfy Bott vanishing [Tot24]. In proving Theorem 6.1, one key point is that when X has an endomorphism as above, not only X but also every contraction of X satisfies Bott vanishing (Lemma 6.2).

Proof. (Theorem 6.1) Assume that X has an int-amplified endomorphism of degree invertible in k . Then X satisfies Bott vanishing [KT23, Theorem C]. Among all smooth Fano 3-folds, exactly 19 non-toric Fano 3-folds (up to isomorphism) satisfy Bott vanishing [Tot24, Theorem 0.1]. In Mori-Mukai's numbering [MM82, MM03, IP99, Bel24], these are (2.26), (2.30), (3.15)–(3.16), (3.18)–(3.24), (4.3)–(4.8), (5.1), and (6.1). (To be precise, the answer is a subset of this in characteristic 2, where only 9 non-toric Fano 3-folds on the known list satisfy Bott vanishing [Tot24, section 2].) We need to show that none of these 19 varieties has an int-amplified endomorphism f of degree invertible in k . Table 1 describes these 19 varieties, using information from Mori-Mukai or the web site Fanography [MM82, Bel24]. In the table, V_5 is the smooth quintic del Pezzo 3-fold $\mathrm{Gr}(2, 5) \cap \mathbf{P}^6 \subset \mathbf{P}^9$, Q is the smooth quadric 3-fold, W is the flag manifold $GL(3)/B$ (or equivalently, a smooth divisor of degree (1, 1) in $\mathbf{P}^2 \times \mathbf{P}^2$), (3.17) is a smooth divisor of degree (1, 1, 1) in $\mathbf{P}^1 \times \mathbf{P}^1 \times \mathbf{P}^2$, and S_5 is the quintic del Pezzo surface.

For each of these 19 Fano 3-folds, the nef cone of X is the same as the (known) nef cone in characteristic zero. In particular, it is rational polyhedral. (The nef cone for every smooth complex Fano 3-fold is given in [CCGK16]. The nef cone was re-computed for the 19 varieties above in any characteristic, yielding the same result [Tot24]. The argument also shows that all nef divisors are semi-ample, and hence the contractions of these varieties have the same description in every characteristic.)

Lemma 6.2. *Let X be a normal projective variety over a perfect field k . Suppose that X has only finitely many contractions (for example, this holds if X is a Mori dream space, or if the nef cone is rational polyhedral). Assume that X has an int-amplified endomorphism of degree invertible in k . Then:*

- (i) *Every contraction of X admits an int-amplified endomorphism of degree invertible in k .*
- (ii) *Every contraction of X satisfies Bott vanishing for ample Weil divisors.*

If we only assume that X has a separable int-amplified endomorphism, then every contraction of X also has a separable int-amplified endomorphism.

Fano 3-fold	Description
(2.26)	the blow-up of $V_5 \subset \mathbf{P}^6$ along a general line in it
(2.30)	the blow-up of $Q \subset \mathbf{P}^4$ at a point
(3.15)	the blow-up of Q along a disjoint line and conic
(3.18)	the blow-up of Q along a conic and then along a fiber in the exceptional divisor
(3.19)	the blow-up of Q at 2 non-collinear points
(3.20)	the blow-up of Q along 2 disjoint lines
(3.23)	the blow-up of Q along a line and then along a fiber of the exceptional divisor
(4.4)	the blow-up of Q along a conic and then along 2 fibers of the exceptional divisor
(5.1)	the blow-up of Q along a conic and then along 3 fibers of the exceptional divisor
(3.16)	the blow-up of $W \subset \mathbf{P}^2 \times \mathbf{P}^2$ along a curve of degree $(2, 1)$
(3.24)	the blow-up of W along a curve of degree $(1, 0)$
(4.7)	the blow-up of W along disjoint $(1, 0)$ and $(0, 1)$ curves
(4.3)	the blow-up of (3.17) in $\mathbf{P}^1 \times \mathbf{P}^1 \times \mathbf{P}^2$ along a curve of degree $(1, 1, 0)$
(3.21)	the blow-up of $\mathbf{P}^1 \times \mathbf{P}^2$ along a curve of degree $(2, 1)$
(3.22)	the blow-up of $\mathbf{P}^1 \times \mathbf{P}^2$ along a curve of degree $(0, 2)$
(4.5)	the blow-up of $\mathbf{P}^1 \times \mathbf{P}^2$ along disjoint $(2, 1)$ and $(1, 0)$ curves
(4.6)	the blow-up of \mathbf{P}^3 along 3 disjoint lines
(4.8)	the blow-up of $(\mathbf{P}^1)^3$ along a curve of degree $(0, 1, 1)$
(6.1)	$\mathbf{P}^1 \times S_5$

Table 1: The 19 non-toric Fano 3-folds that satisfy Bott vanishing

One can get around the assumption that X has only finitely many contractions by considering only contractions that reduce the Picard number by 1 [MZ20, Definition 5.2 and Theorem 7.9].

Proof. Let $f: X \rightarrow X$ be a separable int-amplified endomorphism, and let $\pi: X \rightarrow Z$ be a contraction. Every curve on X is the image of a curve under f . It follows that the pullback linear map $f^*: N^1(X) \rightarrow N^1(X)$ is injective, hence an isomorphism. It also follows that $f^*(\text{Nef}(X)) = \text{Nef}(X)$. A contraction is determined by a face of the cone of curves, or equivalently by a face of the nef cone in $N^1(X)$; so f permutes the set of contractions of X . Since X has only finitely many contractions, after replacing f by a positive iterate, we can assume that f preserves the contraction π . That is, we have a commutative diagram

$$\begin{array}{ccc} X & \xrightarrow{f} & X \\ \downarrow \pi & & \downarrow \pi \\ Z & \xrightarrow{g} & Z, \end{array}$$

for an endomorphism g of Z . Here g is separable since f is.

An endomorphism f of a normal projective variety over a field k is int-amplified if and only if all eigenvalues of f^* on $N^1(X)_{\mathbf{C}}$ have absolute value greater than 1 [Men20, Theorem 1.1]. (Meng assumes that k has characteristic zero, but his proof works in any characteristic.) In our situation, the pullback $\pi^*: N^1(Z) \rightarrow N^1(X)$ is injective (because every curve on Z is the image of some curve on X). Since all eigenvalues of f^* on $N^1(X)_{\mathbf{C}}$ have absolute value greater than 1, the same holds for the eigenvalues of g^* on $N^1(Z)_{\mathbf{C}}$. That is, g is an int-amplified endomorphism of Z . That completes the proof for a separable int-amplified endomorphism f .

Suppose in addition that f has degree invertible in k . The degree of g divides the degree of f and hence is invertible in k . It follows that Z satisfies Bott vanishing by [KT23, Theorem C] (generalized as Theorem 3.1 above). \square

This immediately rules out 13 of the 19 Fano 3-folds above, since they contract to other varieties where Bott vanishing fails, by Table 1. Namely, by [Tot24, section 2] (or earlier work), Bott vanishing fails for the quintic del Pezzo 3-fold V_5 , the quadric 3-fold Q , the flag manifold $W = GL(3)/B$, and the 3-fold (3.17), a smooth divisor in $\mathbf{P}^1 \times \mathbf{P}^1 \times \mathbf{P}^2$ of degree $(1, 1, 1)$.

That leaves: (3.21), (3.22), (4.5), (4.6), (4.8), and (6.1). To analyze these, we will use:

Lemma 6.3. *Let $f: Y \rightarrow X$ be a finite surjective morphism of normal varieties over an algebraically closed field k . Let $\pi_X: X \rightarrow X_1$ and $\pi_Y: Y \rightarrow Y_1$ be contractions, and suppose that there is a morphism $g: Y_1 \rightarrow X_1$ making the diagram commute:*

$$\begin{array}{ccc} Y & \xrightarrow{f} & X \\ \downarrow \pi_Y & & \downarrow \pi_X \\ Y_1 & \xrightarrow{g} & X_1. \end{array}$$

Let Δ_X be the set of points $P \in X_1(k)$ with $\pi_X^{-1}(P)$ reducible, and likewise define $\Delta_Y \subset Y_1(k)$. Then $g^{-1}(\Delta_X) \subset \Delta_Y$.

Proof. For each k -point P in Y_1 , I claim that the fiber of Y over P maps onto the fiber of X over $g(P)$. This is fairly clear geometrically; to be precise, we can imitate the proof of [CMZ20, Lemma 7.3]. Namely, suppose that $Y_P \rightarrow X_{g(P)}$ is not surjective. Let $S = g^{-1}(g(P)) - \{P\}$, a finite set. Then $S \neq \emptyset$ and $U := Y - \pi_Y^{-1}(S)$ is an open dense subset of Y not equal to Y , using that Y is irreducible. Since X is normal, the finite surjection $f: Y \rightarrow X$ is an open map [SPA24, Tag 0F32], and so $f(U)$ is an open dense subset of X . In particular, $f(U) \cap X_{g(P)}$ is open in $X_{g(P)}$. Note that $f(U) = (X - X_{g(P)}) \cup f(Y_P)$. So $f(Y_P)$ is open in $X_{g(P)}$. It is not empty, since Y_P is not empty. Since f is proper, $f(Y_P)$ is also closed in $X_{g(P)}$. Since $\pi_X: X \rightarrow X_1$ is a contraction, $X_{g(P)}$ is connected, and so $f(Y_P) = X_{g(P)}$, as we want.

An irreducible scheme cannot map onto a reducible scheme. Therefore, $g^{-1}(\Delta_X)$ is contained in Δ_Y . \square

Let us continue the proof of Theorem 6.1, that a Fano 3-fold with an int-amplified endomorphism of degree invertible in k is toric. Of the 6 remaining cases, we start with (6.1), which is \mathbf{P}^1 times the quintic del Pezzo surface X . By Lemma 6.2, it suffices to show that X has no int-amplified endomorphism of degree invertible in k . This was shown by Nakayama [Nak10, Proposition 4.4]. We prove a bit more, namely that X has no separable int-amplified endomorphism.

Proposition 6.4. *A smooth del Pezzo surface over an algebraically closed field k that admits a separable int-amplified endomorphism must be toric.*

Proof. The classification of smooth del Pezzo surfaces has the same form in any characteristic [Kol96, section III.3]. In particular, a smooth del Pezzo surface X is toric if and only if its degree $(-K_X)^2$ is at least 6. (Then X is $\mathbf{P}^1 \times \mathbf{P}^1$ or else the blow-up of \mathbf{P}^2 at 3 or fewer points in general position.) And every smooth del Pezzo surface of degree at most 5 is a blow-up of the quintic del Pezzo surface (the blow-up of \mathbf{P}^2 at 4 points in general position). By Lemma 6.2, it suffices to show that the quintic del Pezzo surface X does not have a separable int-amplified endomorphism.

Suppose that the quintic del Pezzo surface X has a separable int-amplified endomorphism f . Let π_X be one of the contractions of X to \mathbf{P}^1 . By the proof of Lemma 6.2, after replacing f by a positive iterate, the contraction π_X is f -equivariant, giving a separable int-amplified endomorphism g of \mathbf{P}^1 .

The point is that the contraction π_X from the quintic del Pezzo surface to \mathbf{P}^1 has 3 singular fibers, and these fibers are reducible (namely, the union of two copies of \mathbf{P}^1). Let $\Delta_X \subset \mathbf{P}^1$ be the discriminant locus of π_X , consisting of 3 points. By Lemma 6.3, $g^{-1}(\Delta_X)$ is contained in Δ_X . Then pulling back differential forms gives a map

$$\Omega_{\mathbf{P}^1}^1(\log \Delta_X) \rightarrow g_* \Omega_{\mathbf{P}^1}^1(\log \Delta_X).$$

Equivalently, we can view this as a map of line bundles from $g^*(K_{\mathbf{P}^1} + \Delta_X)$ to $K_{\mathbf{P}^1} + \Delta_X$, hence as a section of $K_{\mathbf{P}^1} + \Delta_X - g^*(K_{\mathbf{P}^1} + \Delta_X)$. Since g is separable, this section is not identically zero. But $K_{\mathbf{P}^1} + \Delta_X$ has degree 1, so $K_{\mathbf{P}^1} + \Delta_X - g^*(K_{\mathbf{P}^1} + \Delta_X)$ has degree $1 - \deg(g) < 0$, a contradiction. Thus the quintic del Pezzo surface does not have a separable int-amplified endomorphism. \square

Since the 3-fold (6.1) is \mathbf{P}^1 times the quintic del Pezzo surface, Lemma 6.2 and Proposition 6.4 give that (6.1) does not admit a separable int-amplified endomorphism. A fortiori, it does not have an int-amplified endomorphism of degree invertible in k , as considered in Theorem 6.1.

The proof for (4.8), the blow-up X of $(\mathbf{P}^1)^3$ along a curve F of degree $(0, 1, 1)$, is somewhat similar. We can take F to be a point in \mathbf{P}^1 times the diagonal $\Delta_{\mathbf{P}^1}^2$ in $(\mathbf{P}^1)^2$. Suppose that X has an int-amplified endomorphism of degree invertible in k . The contraction π of X to $(\mathbf{P}^1)^2$ (corresponding to the last two \mathbf{P}^1 factors) has discriminant locus the diagonal $\Delta_{\mathbf{P}^1}$, and the fibers over $\Delta_{\mathbf{P}^1}$ are reducible. After replacing f by an iterate, π is f -equivariant, by Lemma 6.2. Write $g: (\mathbf{P}^1)^2 \rightarrow (\mathbf{P}^1)^2$ for the resulting endomorphism of $(\mathbf{P}^1)^2$, which is int-amplified and has degree invertible in k . By Lemma 6.3, $\Delta_{\mathbf{P}^1}$ is totally invariant under g .

By Theorem 3.1, it follows that $((\mathbf{P}^1)^2, \Delta_{\mathbf{P}^1})$ satisfies log Bott vanishing. In particular, taking $A = \mathcal{O}(1, 1)$ and $E = \Delta_{\mathbf{P}^1} \sim A$, we have $H^1(X, \Omega_{(\mathbf{P}^1)^2}^1(\log \Delta_{\mathbf{P}^1})) = 0$. But in fact, this cohomology group is not zero. Indeed, for any smooth divisor D in a smooth variety X over k , we have an exact sequence of coherent sheaves [SPA24, Tag 0FMW]:

$$0 \rightarrow \Omega_X^1 \rightarrow \Omega_X^1(\log D) \xrightarrow{\text{Res}} \mathcal{O}_D \rightarrow 0.$$

So we have an exact sequence of cohomology groups $H^0(D, \mathcal{O}_D) \rightarrow H^1(X, \Omega_X^1) \rightarrow H^1(X, \Omega_X^1(\log D))$. For $X = (\mathbf{P}^1)^2$ and $D = \Delta_{\mathbf{P}^1}$, we have $h^0(D, \mathcal{O}_D) = 1$ and $h^1(X, \Omega_X^1) = 2$, and so $H^1(X, \Omega_X^1(\log D)) \neq 0$ as claimed. This contradiction shows that the Fano 3-fold (4.8) does not have an int-amplified endomorphism of degree invertible in k .

Next, we exclude the 3-fold (4.6), the blow-up of \mathbf{P}^3 along three pairwise disjoint lines L_1, L_2, L_3 over k . Here X has three contractions to \mathbf{P}^1 , using that the blow-up of \mathbf{P}^3 along each line L_i is a \mathbf{P}^2 -bundle over \mathbf{P}^1 . Let $\pi: X \rightarrow (\mathbf{P}^1)^2$ be the contraction given by the morphisms to \mathbf{P}^1 associated to L_1 and L_2 . For clarity, first consider the blow-up Y of \mathbf{P}^3 along L_1 and L_2 ; then the contraction $Y \rightarrow (\mathbf{P}^1)^2$ is a \mathbf{P}^1 -bundle. The line $L_3 \subset Y$ maps to a curve of degree $(1, 1)$ in $(\mathbf{P}^1)^2$, which we can take to be the diagonal $\Delta_{\mathbf{P}^1}$. Since X is the blow-up of Y along L_3 , the discriminant locus of $\pi: X \rightarrow (\mathbf{P}^1)^2$ is the curve $\Delta_{\mathbf{P}^1}$. The fibers of π over that curve are reducible (the union of two copies of \mathbf{P}^1).

If X has an int-amplified endomorphism of degree invertible in k , then the pair $((\mathbf{P}^1)^2, \Delta_{\mathbf{P}^1})$ satisfies log Bott vanishing. But this is false, as shown above. So the Fano 3-fold (4.6) does not have an int-amplified endomorphism of degree invertible in k .

Next, we rule out (3.22), the blow-up X of $\mathbf{P}^1 \times \mathbf{P}^2$ along a conic in $p \times \mathbf{P}^2$, for a k -point p in \mathbf{P}^1 . The contraction of X to \mathbf{P}^2 has discriminant locus a conic F in \mathbf{P}^2 . As in the arguments above, if X has an int-amplified endomorphism of degree invertible in k , then so does \mathbf{P}^2 , and F is totally invariant. Therefore, the pair (\mathbf{P}^2, F) satisfies log Bott vanishing. But this is false. Namely, let $A = \mathcal{O}(1)$ and $E = F \sim \mathcal{O}(2)$; then we will show that $H^1(\mathbf{P}^2, \Omega_{\mathbf{P}^2}^1(\log F)(A - E)) = H^1(\mathbf{P}^2, \Omega_{\mathbf{P}^2}^1(\log F)(-1))$ is not zero. Use the exact sequence of coherent sheaves on \mathbf{P}^2 :

$$0 \rightarrow \Omega_{\mathbf{P}^2}^1 \rightarrow \Omega_{\mathbf{P}^2}^1(\log F) \xrightarrow{\text{Res}} \mathcal{O}_F \rightarrow 0.$$

By the exact sequence $0 \rightarrow \Omega_{\mathbf{P}^2}^1(-1) \rightarrow \mathcal{O}(-2)^{\oplus 3} \rightarrow \mathcal{O}(-1) \rightarrow 0$ on \mathbf{P}^2 , $\Omega_{\mathbf{P}^2}^1(-1)$

has zero cohomology in all degrees. So we have an isomorphism

$$H^1(\mathbf{P}^2, \Omega_{\mathbf{P}^2}^1(\log F)(-1)) \cong H^1(F, O(-1)|_F).$$

Since the conic F is isomorphic to \mathbf{P}^1 and $O(-1)$ has degree -2 on F , $h^1(F, O(-1)|_F)$ is 1, not 0. So log Bott vanishing fails for (\mathbf{P}^2, F) . It follows that the 3-fold (3.22) does not have an int-amplified endomorphism of degree invertible in k .

The last cases are (3.21) and (4.5). These are handled by:

Lemma 6.5. *The Fano 3-folds (3.21) and (4.5) contract to the quintic del Pezzo 3-fold with one node, which does not satisfy Bott vanishing.*

Proof. Here (4.5) is a blow-up of the Fano 3-fold of type (3.21). Next, the Fano 3-fold X of type (3.21) is the blow-up of $\mathbf{P}^1 \times \mathbf{P}^2$ along a curve F of degree $(2, 1)$. There is a contraction from X to the quintic del Pezzo 3-fold Y with one node [Pro13, section 5.4.2]. It remains to show that Y does not satisfy Bott vanishing. Indeed, I claim that the Euler characteristic $\chi(Y, \Omega_Y^{[2]}(1))$ is $-2 < 0$. This is harder than previous cases because Y is singular, but it is still manageable.

To prove this, it is convenient to know how the sheaf Ω_Y^2 is related to its reflexive hull $\Omega_Y^{[2]}$, for a 3-fold Y with a node. More generally, let Y be a normal hypersurface in a smooth variety W with $\dim(W) = n + 1$, so that locally Y is the zero locus of a regular function g . Define the *torsion* and *cotorsion* of a coherent sheaf M to be the kernel and cokernel of the natural map $M \rightarrow M^{**}$. Consider the complex K :

$$0 \rightarrow \Omega_W^0|_Y \rightarrow \Omega_W^1|_Y \rightarrow \cdots \rightarrow \Omega_W^{n+1}|_Y \rightarrow 0,$$

with differentials given by $\wedge dg$. Graf described the torsion and cotorsion of the sheaves Ω_Y^j in terms of this complex: we have $H^j(K) \cong \text{tor } \Omega_Y^j$ and $H^j(K) \cong \text{cotor } \Omega_Y^{j-1}$ [Gra15, Theorem 1.11]. For Y the 3-fold node, we can take $W = A^4$ and $g = xy - zw$. The complex above is the Koszul complex for the regular sequence $\partial g / \partial x = y$, $\partial g / \partial y = x$, $\partial g / \partial z = -w$, $\partial g / \partial w = -z$, tensored over O_W with O_Y . These four functions generate the ideal of the origin P in $W = A^4$; so $\text{tor } \Omega_Y^i \cong \text{Tor}_{4-i}^{O_W}(O_Y, O_P)$ and $\text{cotor } \Omega_Y^i \cong \text{Tor}_{3-i}^{O_W}(O_Y, O_P)$. These Tor groups are easy to compute, using the free resolution $0 \rightarrow O_W \xrightarrow{g} O_W \rightarrow O_Y \rightarrow 0$ of O_Y as an O_W -module. Namely, it follows that $\text{Tor}_*^{O_W}(O_Y, O_P)$ is the homology of the complex

$$0 \rightarrow O_P \xrightarrow{0} O_P \rightarrow 0,$$

which is a 1-dimensional k -vector space in degrees 1 and 0, and zero in other degrees. Therefore, Ω_Y^1 is reflexive, while Ω_Y^2 is torsion-free and its cotorsion is a 1-dimensional vector space, supported at the node P . That is, we have a short exact sequence of sheaves on a nodal 3-fold Y :

$$0 \rightarrow \Omega_Y^2 \rightarrow \Omega_Y^{[2]} \rightarrow O_P \rightarrow 0.$$

Hence, for Y the quintic del Pezzo 3-fold with one node, we have $\chi(Y, \Omega_Y^{[2]}(1)) = 1 + \chi(Y, \Omega_Y^2(1))$. It remains to show that $\chi(Y, \Omega_Y^2(1)) = -3$.

For a nodal 3-fold Y in a smooth 4-fold W over k (as for any effective Cartier divisor), we have an exact sequence of coherent sheaves on Y [SPA24, Tag 00RU]:

$$O(-Y)|_Y \rightarrow \Omega_W^1|_Y \rightarrow \Omega_Y^1 \rightarrow 0.$$

Taking exterior powers, we have an exact sequence $\Omega_Y^{j-1}(-Y) \rightarrow \Omega_W^j|_Y \rightarrow \Omega_Y^j \rightarrow 0$ for any j . For $j = 2$, the map $\Omega_Y^1(-Y) \rightarrow \Omega_W^2|_Y$ is clearly injective outside the node. Since $\Omega_Y^1(-Y)$ is torsion-free (by the analysis above), this map is actually injective as a map of sheaves on all of Y . That is, we have an exact sequence $0 \rightarrow \Omega_Y^1(-Y) \rightarrow \Omega_W^2|_Y \rightarrow \Omega_Y^2 \rightarrow 0$. Likewise, we have an exact sequence $0 \rightarrow \mathcal{O}_Y(-Y) \rightarrow \Omega_W^1|_Y \rightarrow \Omega_Y^1 \rightarrow 0$ on Y . Finally, we can tensor the exact sequence $0 \rightarrow \mathcal{O}_W(-Y) \rightarrow \mathcal{O}_W \rightarrow \mathcal{O}_Y \rightarrow 0$ with any vector bundle on W , giving $0 \rightarrow \Omega_W^1(-Y) \rightarrow \Omega_W^1 \rightarrow \Omega_W^1|_Y \rightarrow 0$ and $0 \rightarrow \Omega_W^2(-Y) \rightarrow \Omega_W^2 \rightarrow \Omega_W^2|_Y \rightarrow 0$.

Apply this to the quintic del Pezzo 3-fold $Y \subset \mathbf{P}^6$ with a node, which is a codimension-3 linear section of the Grassmannian $\mathrm{Gr}(2, 5) \subset \mathbf{P}^9$. The quintic del Pezzo 3-fold with one node is unique up to isomorphism, with explicit equations labelled $X_{5,2,4}$ in [KP23, section A.1.6]. By the equations, Y is a hyperplane section of a smooth codimension-2 linear section $W \subset \mathbf{P}^7$. By the previous paragraph, we can rewrite $\chi(Y, \Omega_Y^2(1))$ in terms of Euler characteristics on W ; but this formula would be exactly the same with Y replaced by a smooth hyperplane section $V_5 \subset \mathbf{P}^6$. Therefore, $\chi(Y, \Omega_Y^2(1)) = \chi(V_5, \Omega_{V_5}^2(1)) = -3$ [Tot24, section 2]. As discussed above, it follows that $\chi(Y, \Omega_Y^{[2]}(1)) = -2$, and so Y does not satisfy Bott vanishing. Lemma 6.5 is proved. \square

It follows that neither (3.21) nor (4.5) admits an int-amplified endomorphism of degree invertible in k . Theorem 6.1 is proved. \square

7 Images of toric varieties: del Pezzo surfaces and Fano 3-folds

Occhetta and Wiśniewski conjectured that a smooth complex projective variety X that admits a surjective morphism from a proper toric variety must be toric [OW02]. (This is known for contractions, and so it suffices to consider a finite surjective morphism.) Occhetta-Wiśniewski's conjecture was proved by Achinger, Witaszek, and Zdanowicz for X of dimension at most 2, and also for X a Fano 3-fold [AWZ21, proof of Theorem 4.4.1], [AWZ23, Theorems 6.9 and 7.7]. Using Bott vanishing, we will reprove this result for Fano 3-folds and extend it to positive characteristic (Theorem 7.2). Kawakami and I proved this extension earlier for Fano 3-folds with Picard number 1 [KT23, Proposition 3.10].

Another approach to Theorem 7.2 appeared after the first version of this paper. Namely, Kawakami and Takamatsu used Lemma 2.1 above to show that the image of an F -liftable variety in characteristic p by a morphism of degree prime to p is F -liftable [KT24, Corollary 3.13]. In particular, the image of a toric variety by a morphism of degree prime to p is F -liftable. But Achinger-Witaszek-Zdanowicz showed that a smooth Fano 3-fold that is F -liftable must be toric. So a Fano 3-fold that is an image of a toric variety by a morphism of degree prime to p must be toric.

We will prove Theorem 7.2 using log Bott vanishing rather than F -liftability. Thus our method for analyzing images of toric varieties closely parallels our approach to finding which varieties have nontrivial endomorphisms (Theorem 6.1).

We start with the analogous result in dimension 2.

Proposition 7.1. *A smooth del Pezzo surface over an algebraically closed field k that is the image of a proper toric variety must be toric.*

Surprisingly, Proposition 7.1 works in any characteristic with no assumption on the degree of the morphism f ; f may even be inseparable. The proof does not use Bott vanishing.

Proof. A smooth del Pezzo surface X is toric if and only if its degree $(-K_X)^2$ is at least 6. And every smooth del Pezzo surface of degree at most 5 is a blow-up of the quintic del Pezzo surface (the blow-up of \mathbf{P}^2 at 4 points in general position). So it suffices to show that there is no surjection f from a proper toric variety Y to the quintic del Pezzo surface X .

Let $Y \rightarrow Z \rightarrow X$ be the Stein factorization of f ; then Z is also toric, since every contraction of a toric variety is toric [Tan22, Proposition 2.7]. Replacing Y by Z , we can assume that $f: Y \rightarrow X$ is finite and surjective.

Let π_X be one of the contractions of X to \mathbf{P}^1 . Let $Y \rightarrow Y_1 \rightarrow \mathbf{P}^1$ be the Stein factorization of the composition $Y \rightarrow X \rightarrow \mathbf{P}^1$, where we write $\pi_Y: Y \rightarrow Y_1$ and $g: Y_1 \rightarrow \mathbf{P}^1$. Then Y_1 is a proper toric curve, hence isomorphic to \mathbf{P}^1 , and $g: Y_1 \rightarrow \mathbf{P}^1$ is finite and surjective.

As we used in the proof of Proposition 6.4, the contraction π_X from the quintic del Pezzo surface to \mathbf{P}^1 has 3 singular fibers, and these fibers are reducible (namely, the union of two copies of \mathbf{P}^1). Let $\Delta_X \subset \mathbf{P}^1$ be the discriminant locus of π_X , consisting of 3 points. By Lemma 6.3, $g^{-1}(\Delta_X)$ is contained in the discriminant locus Δ_Y of $\pi_Y: Y \rightarrow Y_1$. But $Y \rightarrow Y_1$ is a toric morphism, so its discriminant locus is contained in the toric divisor of Y_1 , which consists of 2 points. Since $g: Y_1 \rightarrow \mathbf{P}^1$ is surjective, this is a contradiction (2 points cannot map onto 3 points). We have shown that the quintic del Pezzo surface is not the image of a proper toric variety. \square

Theorem 7.2. *Let X be a smooth Fano 3-fold over an algebraically closed field k . If X is the image of a proper toric 3-fold by a morphism of degree invertible in k , then X is toric.*

As in Theorem 6.1, the proof uses Tanaka's theorem that the classification of smooth Fano 3-folds has essentially the same form in every characteristic [Tan23b, Theorem 1.1], [Tan23a, Theorem 1.1]. Without that, our proof applies to Fano 3-folds in any characteristic that are given by the same construction as one of the Fano 3-folds over \mathbf{C} (classified by Iskovskikh and Mori-Mukai).

Proof. Assume that X is the image of a proper toric 3-fold Y by a morphism f of degree invertible in k . After replacing f by its Stein factorization, we can also assume that $f: Y \rightarrow X$ is finite. Here Y is still toric, because every contraction of a toric variety is toric [Tan22, Proposition 2.7]. Then X satisfies Bott vanishing, by Theorem 5.1. There are exactly 19 non-toric Fano 3-folds (up to isomorphism) that satisfy Bott vanishing [Tot24, Theorem 0.1]. In Mori-Mukai's numbering, these are (2.26), (2.30), (3.15)–(3.16), (3.18)–(3.24), (4.3)–(4.8), (5.1), and (6.1). (To be precise, the answer is a subset of this in characteristic 2, where only 9 non-toric Fano 3-folds on the known list satisfy Bott vanishing [Tot24, section 2].) We need to show that none of these 19 varieties can be the image of a toric variety Y by a finite morphism f of degree invertible in k .

Of these 19 varieties, 15 have contractions that do not satisfy Bott vanishing, by the proof of Theorem 6.1, including Lemma 6.5. This excludes all the cases except four: (3.22), (4.6), (4.8), and (6.1).

Here (6.1) is \mathbf{P}^1 times the quintic del Pezzo surface. That surface is not an image of a toric variety, by Proposition 7.1. So (6.1) is not the image of a toric variety.

Next, we rule out (3.22), the blow-up X of $\mathbf{P}^1 \times \mathbf{P}^2$ along a conic in $q \times \mathbf{P}^2$, for a k -point q in \mathbf{P}^1 . The contraction of X to $X_1 := \mathbf{P}^2$ has discriminant locus a conic $F \subset \mathbf{P}^2$, and the fibers over points of F are reducible (a union of two \mathbf{P}^1 's). Suppose that there is a proper toric 3-fold Y over k with a morphism $f: Y \rightarrow X$ such that $\deg(f)$ is invertible in k .

Let $Y \rightarrow Y_1 \rightarrow X_1$ be the Stein factorization of the composition $Y \rightarrow X \rightarrow X_1$:

$$\begin{array}{ccc} Y & \xrightarrow{\quad f \quad} & X \\ \downarrow & & \downarrow \\ Y_1 & \longrightarrow & X_1. \end{array}$$

Then Y_1 is a proper toric surface over k , and $h: Y_1 \rightarrow X_1$ is finite, of degree invertible in k . The discriminant locus of $Y \rightarrow Y_1$ is contained in the union of the toric divisors. By Lemma 6.3, $h^{-1}(F)$ is contained in the discriminant locus of $Y \rightarrow Y_1$, hence in the union of the toric divisors in Y_1 .

By Theorem 5.1, it follows that the pair $(X_1, F) = (\mathbf{P}^2, F)$ satisfies log Bott vanishing. But that is false, by the proof of Theorem 6.1. Thus (3.22) is not the image of a toric 3-fold by a morphism of degree invertible in k .

A similar argument rules out (4.8), the blow-up X of $(\mathbf{P}^1)^3$ along a curve of degree $(0, 1, 1)$. The contraction of X to $X_1 := (\mathbf{P}^1)^2$ (corresponding to the last two \mathbf{P}^1 factors) has discriminant locus a curve F of degree $(1, 1)$, which we can take to be the diagonal $\Delta_{\mathbf{P}^1}$ in $(\mathbf{P}^1)^2$. The fibers over points in F are reducible (a union of two \mathbf{P}^1 's). Let $Y \rightarrow Y_1 \rightarrow X_1$ be the Stein factorization of the composition $Y \rightarrow X \rightarrow X_1$:

$$\begin{array}{ccc} Y & \xrightarrow{\quad f \quad} & X \\ \downarrow & & \downarrow \\ Y_1 & \longrightarrow & X_1. \end{array}$$

Then Y_1 is a proper toric surface over k , and $h: Y_1 \rightarrow X_1$ is finite, of degree invertible in k .

By the same argument as in the previous case, $h^{-1}(F)$ must be a toric divisor in Y_1 . By Theorem 5.1, it follows that the pair $(X_1, F) = ((\mathbf{P}^1)^2, \Delta_{\mathbf{P}^1})$ satisfies log Bott vanishing. But that is false, by the proof of Theorem 6.1. Thus (4.8) is not the image of a toric 3-fold by a morphism of degree invertible in k .

Finally, let X be the Fano 3-fold (4.6), the blow-up of \mathbf{P}^3 along three pairwise disjoint lines L_1, L_2, L_3 over k . Suppose that there is a proper toric 3-fold Y over k with a morphism $f: Y \rightarrow X$ such that $\deg(f)$ is invertible in k . By the proof of Theorem 6.1, X has a contraction to $X_1 := (\mathbf{P}^1)^2$ with discriminant locus the diagonal $F := \Delta_{\mathbf{P}^1}$ in $(\mathbf{P}^1)^2$. The fibers over F are reducible (the union of two \mathbf{P}^1 's). Then the same argument in the previous case (using that $((\mathbf{P}^1)^2, \Delta_{\mathbf{P}^1})$ does not satisfy log Bott vanishing) yields a contradiction. So the Fano 3-fold (4.6) is not the image of a toric 3-fold by a morphism of degree invertible in k . \square

Acknowledgements

This work was supported by NSF grant DMS-2054553, the National Center for Theoretical Sciences (NCTS), Simons Foundation grant SFI-MPS-SFM-00005512, and the Charles Simonyi Endowment at the Institute for Advanced Study. Thanks to Louis Esser, Tatsuro Kawakami, Supravat Sarkar, and the referee.

References

- [AWZ21] Piotr Achinger, Jakub Witaszek, and Maciej Zdanowicz. Global Frobenius liftability I. *J. Eur. Math. Soc. (JEMS)*, 23(8):2601–2648, 2021. 1, 14
- [AWZ23] Piotr Achinger, Jakub Witaszek, and Maciej Zdanowicz. Global Frobenius liftability II: surfaces and Fano threefolds. *Ann. Sc. Norm. Super. Pisa Cl. Sci.*, 24:329–366, 2023. 1, 14
- [Bel24] Pieter Belmans. Fanography: a tool to visually study the geography of Fano 3-folds. <https://www.fanography.info>, 2024. 8
- [CCGK16] Tom Coates, Alessio Corti, Sergey Galkin, and Alexander Kasprzyk. Quantum periods for 3-dimensional Fano manifolds. *Geom. Topol.*, 20(1):103–256, 2016. 8
- [CLS11] David A. Cox, John B. Little, and Henry K. Schenck. *Toric varieties*, volume 124 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2011. 7
- [CMZ20] Paolo Cascini, Sheng Meng, and De-Qi Zhang. Polarized endomorphisms of normal projective threefolds in arbitrary characteristic. *Math. Ann.*, 378(1-2):637–665, 2020. 11
- [Fak03] Najmuddin Fakhruddin. Questions on self maps of algebraic varieties. *J. Ramanujan Math. Soc.*, 18(2):109–122, 2003. 1
- [Fuj08] Osamu Fujino. Vanishing theorems for toric polyhedra. In *Higher dimensional algebraic varieties and vector bundles*, RIMS Kôkyûroku Bessatsu, B9, pages 81–95. Res. Inst. Math. Sci. (RIMS), Kyoto, 2008. 4, 5, 7
- [Gar84] Emmanuelle Garel. An extension of the trace map. *J. Pure Appl. Algebra*, 32(3):301–313, 1984. 3
- [Gra15] Patrick Graf. The generalized Lipman-Zariski problem. *Math. Ann.*, 362(1-2):241–264, 2015. 13
- [Hüb89] Reinhold Hübl. *Traces of differential forms and Hochschild homology*, volume 1368 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1989. 3

- [IP99] V. A. Iskovskikh and Yu. G. Prokhorov. Fano varieties. In *Algebraic geometry, V*, volume 47 of *Encyclopaedia Math. Sci.*, pages 1–247. Springer, Berlin, 1999. 8
- [Kol96] János Kollár. *Rational curves on algebraic varieties*, volume 32 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer-Verlag, Berlin, 1996. 11
- [KP23] A. G. Kuznetsov and Yu. G. Prokhorov. On higher-dimensional del Pezzo varieties. *Izv. Ross. Akad. Nauk Ser. Mat.*, 87(3):75–148, 2023. 14
- [KT23] Tatsuro Kawakami and Burt Totaro. Endomorphisms of varieties and Bott vanishing. [arXiv:2302.11921](#), 2023. To appear in *J. Algebraic Geom.* 1, 2, 4, 5, 6, 7, 8, 10, 14
- [KT24] Tatsuro Kawakami and Teppei Takamatsu. On Frobenius liftability of surface singularities. [arXiv:2402.08152](#), 2024. 14
- [Kun86] Ernst Kunz. *Kähler differentials*. Advanced Lectures in Mathematics. Friedr. Vieweg & Sohn, Braunschweig, 1986. 3
- [Men20] Sheng Meng. Building blocks of amplified endomorphisms of normal projective varieties. *Math. Z.*, 294(3-4):1727–1747, 2020. 1, 10
- [MM03] Shigefumi Mori and Shigeru Mukai. Erratum: “Classification of Fano 3-folds with $B_2 \geq 2$ ” [Manuscripta Math. **36** (1981/82), no. 2, 147–162; MR0641971 (83f:14032)]. *Manuscripta Math.*, 110(3):407, 2003. 8
- [MM82] Shigefumi Mori and Shigeru Mukai. Classification of Fano 3-folds with $B_2 \geq 2$. *Manuscripta Math.*, 36(2):147–162, 1981/82. 8
- [MZ20] Sheng Meng and De-Qi Zhang. Normal projective varieties admitting polarized or int-amplified endomorphisms. *Acta Math. Vietnam.*, 45(1):11–26, 2020. 1, 10
- [MZZ22] Sheng Meng, De-Qi Zhang, and Guolei Zhong. Non-isomorphic endomorphisms of Fano threefolds. *Math. Ann.*, 383(3-4):1567–1596, 2022. 1, 5, 8
- [Nak10] Noboru Nakayama. Separable endomorphisms of surfaces in positive characteristic. In *Algebraic geometry in East Asia—Seoul 2008*, volume 60 of *Adv. Stud. Pure Math.*, pages 301–330. Math. Soc. Japan, Tokyo, 2010. 11
- [OW02] Gianluca Occhetta and Jarosław A. Wiśniewski. On Euler-Jaczewski sequence and Remmert-van de Ven problem for toric varieties. *Math. Z.*, 241(1):35–44, 2002. 14
- [Pro13] Yuri Prokhorov. G -Fano threefolds, I. *Adv. Geom.*, 13(3):389–418, 2013. 13

- [SPA24] The Stacks Project Authors. The Stacks Project. <https://stacks.math.columbia.edu/>, 2024. 2, 3, 4, 11, 12, 13
- [Tan22] Hiromu Tanaka. Kawamata-Viehweg vanishing for toric varieties. [arXiv:2208.09680](https://arxiv.org/abs/2208.09680), 2022. 7, 15
- [Tan23a] Hiromu Tanaka. Fano threefolds in positive characteristic II. [arXiv:2308.08122](https://arxiv.org/abs/2308.08122), 2023. 8, 15
- [Tan23b] Hiromu Tanaka. Fano threefolds in positive characteristic IV. [arXiv:2308.08127](https://arxiv.org/abs/2308.08127), 2023. 8, 15
- [Tot24] Burt Totaro. Bott vanishing for Fano threefolds. *Math. Z.*, 307:14, 2024. 8, 10, 14, 15